UC Riverside

UC Riverside Previously Published Works

Title

Epidemic Spread of Symbiotic and Non-Symbiotic Bradyrhizobium Genotypes Across California.

Permalink

<https://escholarship.org/uc/item/05g090tf>

Journal

Microbial ecology, 71(3)

ISSN

0095-3628

Authors

Hollowell, AC
Regus, JU
Gano, KA
et al.

Publication Date

2016-04-01

DOI

10.1007/s00248-015-0685-5

Peer reviewed

Epidemic Spread of Symbiotic and Non-Symbiotic *Bradyrhizobium* Genotypes Across California

A. C. Hollowell¹ · J. U. Regus¹ · K. A. Gano¹ · R. Bantay¹ · D. Centeno¹ · J. Pham¹ · J.Y. Lyu¹ · D. Moore¹ · A. Bernardo¹ · G. Lopez¹ · A. Patil¹ · S. Patel¹ · Y. Lii¹ · J. L. Sachs^{1,2}

Received: 8 April 2015 / Accepted: 28 September 2015 / Published online: 14 October 2015
© Springer Science+Business Media New York 2015

Abstract The patterns and drivers of bacterial strain dominance remain poorly understood in natural populations. Here, we cultured 1292 *Bradyrhizobium* isolates from symbiotic root nodules and the soil root interface of the host plant *Acmispon strigosus* across a >840-km transect in California. To investigate epidemiology and the potential role of accessory loci as epidemic drivers, isolates were genotyped at two chromosomal loci and were assayed for presence or absence of accessory “symbiosis island” loci that encode capacity to form nodules on hosts. We found that *Bradyrhizobium* populations were very diverse but dominated by few haplotypes—with a single “epidemic” haplotype constituting nearly 30 % of collected isolates and spreading nearly statewide. In many *Bradyrhizobium* lineages, we inferred presence and absence of the symbiosis island suggesting recurrent evolutionary gain and or loss of symbiotic capacity. We did not find statistical phylogenetic evidence that the symbiosis island acquisition promotes strain dominance and both symbiotic and non-symbiotic strains exhibited population dominance and spatial spread. Our dataset reveals that a strikingly few *Bradyrhizobium* genotypes can rapidly spread to dominate a landscape and suggests that these epidemics are not driven by

the acquisition of accessory loci as occurs in key human pathogens.

Keywords *Rhizobia* · *Symbiosis* · *Epidemic* · Population genetics · Evolution

Introduction

A critical goal in bacteriology is to understand patterns of genotypic abundance and epidemic spread. Of particular interest are host-associated bacteria, including pathogens and symbionts. These diverse bacterial lineages colonize host surfaces, inhabit specific tissues or cells, and can often persist free in soils and or aquatic habitats between phases of host infection [1, 2]. The capacity of bacteria to thrive in host tissues is often modulated by the presence of plasmids and genomic islands, cassettes of accessory loci that can get transmitted among genomes. Horizontal transfer of these accessory loci often engenders bacteria with suites of fitness-enhancing traits including host infection capacity, pathogenicity, multi-drug resistance, and metabolic flexibility [3–5]. The acquisition of plasmids and genomic islands has been implicated in epidemic outbreaks, including *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Yersinia* spp. [6–8]. But in natural settings, we understand little about how bacterial strains vary in their capacity to dominate local sites or host populations or to spread among sites across ecological barriers. In particular, almost nothing is known about patterns of dominance and epidemic spread in symbiotic bacteria, which are important for human health, the success of crops, and other ecosystem services.

Rhizobia are proteobacteria that can exhibit capacity to infect leguminous plants and fix atmospheric nitrogen for their hosts [9]. Globally, rhizobia are responsible for the fixation of

Electronic supplementary material The online version of this article (doi:10.1007/s00248-015-0685-5) contains supplementary material, which is available to authorized users.

✉ J. L. Sachs
joels@ucr.edu

¹ Department of Biology, University of California, Riverside, CA 92521, USA

² Institute for Integrative Genome Biology, University of California, Riverside, CA 95616, USA

~150 Tg of nitrogen per year [10] and their symbiosis with legumes represents the largest input source of nitrogen into terrestrial ecosystems [11]. In agriculture, legumes account for ~27 % of global crop production [12] and are valued for their capacity to grow in nitrogen depauperate soils. Similar to bacterial pathogens, rhizobia can acquire accessory DNA elements that confer the capability to colonize and infect hosts. Rhizobial genomes are subdivided into genome regions specific for their life stages, with chromosomal loci expressed during free-living phases in the soil and symbiosis loci expressed inside of host cells [13, 14]. Symbiosis loci required for host nodulation and nitrogen fixation are clustered onto large plasmids or genomic islands [15–19], that can be transferred among lineages, presumably via conjugation [20–22]. Non-nodulating rhizobia are also common [23, 24], and these strains often lack some or all of the characterized symbiosis loci [23–28].

Bradyrhizobium is the most cosmopolitan rhizobial lineage and is found free-living in soils and in aquatic environments as well as in symbiotic association with plant and animal hosts, including humans [29–37]. Recent work suggests that there are ~19 species of *Bradyrhizobium* that form symbiotic associations with legumes [36], although non-symbiotic strains are also common in soils [37]. *Bradyrhizobium* can nodulate diverse wild legumes as well as staple crops such as soybeans (*Glycine*), peanuts (*Arachis*), and cowpea (*Vigna*) [38, 39]. In the model genome, *Bradyrhizobium diazoefficiens* strain USDA110 (previously *B. japonicum*), symbiosis-specific genes are clustered within a 410-kb symbiosis island in which the G+C content differs from the rest of the genome [40]. Some divergent *Bradyrhizobium* spp. (e.g., BTAi1 and ORS278) lack a symbiosis island and use a different mechanism to nodulate hosts [41]. Previous studies have revealed that *Bradyrhizobium* populations can exhibit patterns consistent with epidemics such as a few genotypes existing at high frequency at a single site [33] or genotypes isolated at multiple locations [34, 35, 37]. But these patterns remain poorly understood, and it is unclear what role the symbiosis island might play as a driver of increased abundance and epidemic spread.

Here, we investigated the population genetic structure of *Bradyrhizobium* spp. cultured from *Acmispon strigosus* (formerly *Lotus strigosus*), a native annual legume common across the Pacific Southwest of the United States. We cultured 850 *A. strigosus* nodules from 14 natural sites across California encompassing 185 plants collected over a >840-km transect. In parallel, we isolated 442 root surface *Bradyrhizobium* from three focal host populations within this range, which includes isolates that lack symbiosis islands and cannot infect *Acmispon* hosts. All 1292 isolates were sequenced at two hypervariable chromosomal loci, and we used a combination of greenhouse inoculation assays and PCR to test for presence of the symbiosis island in all the root surface isolates. We assigned haplotypes and symbiotic capacity information to

all isolates and examined the frequency and spatial spread of epidemic rhizobial genotypes within and among host populations. Our goals were to (i) investigate genotype frequency and spatial spread of *Bradyrhizobium* in native *A. strigosus* hosts, (ii) infer the presence or absence of the symbiosis island in *Bradyrhizobium*, (iii) test for the role of symbiosis island acquisition as a driver of *Bradyrhizobium* strain dominance, and (iv) test for community structure of rhizobial isolates due to other abiotic or biotic factors.

Methods

Collection of *Bradyrhizobium* Isolates *Bradyrhizobium* was isolated from the nodules and the soil root interface of *A. strigosus*, and clonal cultures were grown and archived for genotyping following published protocols [33]. Briefly, whole plants were excavated from native sites and transported in sealed plastic bags to the laboratory where they were washed to remove soil with tap water and sterilized tools were used to remove root nodules. Nodules were surface sterilized with bleach and rinsed with sterile water before being crushed with glass rods and the contents plated on a modified arabinose gluconate medium (MAG) [33]. For root surface isolates, the roots were dissected into ~1-cm sections, were divided into root tips and “old” roots, and were vortexed in a sterile solution of 0.01 % Tween 20 (Fisher Scientific Fair Lawn, NJ). The wash solution was then serially diluted and plated on glucose-based rhizobium-defined medium (GRDM) with cyclohexamide as an antifungal and bromothymol blue as a pH indicator [33]. Among the resultant colonies, we selected for *Bradyrhizobium* based on growth rate, color, and ability to grow on MAG and GRDM but not on Luria–Bertani (LB) medium [33]. Plant hosts for culturing were collected from 14 field collection sites across California covering an ~840-km transect. Collection sites included University of California Natural Reserves (Bodega Marine Reserve, Burns Piñon Ridge Reserve, and Motte Rimrock Reserve); an undeveloped site in the hills above University of California–Riverside, a biological field station in Claremont, CA (Robert J. Bernard Biological Field Station); natural preserves (Madrona Marsh Preserve, Pismo Dunes Natural Preserve, and Whitewater Preserve); a wildlife refuge (Guadalupe–Nipomo Dunes National Wildlife Refuge); two separate sites within a large state park (Anza Borrego Desert State Park); a municipal park (Griffith Park); a site adjacent to the San Dimas Reservoir; and an undeveloped site adjacent to human development (San Dimas Canyon) (Table S1). Nodule isolates were collected from plants at all sites, but root surface isolates were only collected from plants at the Bodega Marine Reserve, Motte Rimrock Reserve, and the undeveloped site in the hills above the University of California–Riverside.

Sequencing and Haplotype Analysis

Partial sequences from two chromosomal loci, glutamine synthetase (*glnII*) and recombinase (*recA*), totaling ~1 kb were PCR amplified and sequenced at the Institute for Integrative Genome Biology of UC Riverside using published protocols [34]. Only sequences with unambiguous bases were utilized leading to a total of 1292 sequenced isolates. Sequences from each locus were aligned separately using Clustal Omega, and isolates with identical sequences for each locus were determined using the “find redundant” command within the MacClade program [42]. Unique sequences, or haplotypes, were defined for each locus separately and for the concatenated dataset. Abundance was calculated for the concatenated dataset as the number of times each haplotype was isolated.

Symbiotic Capacity Assessment

We conducted a combination of assays on root surface isolates to test for symbiotic capacity. A subset of isolates (75) were previously assessed using greenhouse nodulation assays on *A. strigosus*, which has already been shown to be a permissive host on diverse *Bradyrhizobium* [24, 43, 44]. Here, we conducted greenhouse nodulation experiments on an additional 55 isolates, using identical procedures. Briefly, at least five seedlings per tested *Bradyrhizobium* isolate were grown in sterile conditions and were inoculated clonally with 5×10^8 cells, and parallel control hosts were inoculated with sterile water. At 8-week postinoculation, all hosts were unpotted, roots and shoots were weighed, and roots were checked for nodules. In all cases, controls lacked nodules. Hosts given the same inoculated strains either all became nodulated or were all lacking nodules. The remaining 342 root surface isolates were classified as symbiotic or non-symbiotic based on success or failure of PCR amplification of at least one symbiosis island locus (*nifD*, *nodD-A*, *nodZ*, and *nolL*) [22, 24, 45]. Earlier analyses showed that successful amplification of these loci, giving a band of the correct size, is a reliable indicator of presence of the symbiosis island [24]. Many isolates were tested at two or more loci (160/342; Table S1).

Phylogenetic Reconstruction and Species Designation

Phylogenies were reconstructed using the concatenated *glnII* and *recA* sequences from the cultured isolates and from the following reference strains: *Bradyrhizobium arachidis* (CCBAU33067), *B. betae* (PL7HG1), *B. canariense* (SEMIA928), *B. cytisi* (LMG25866), *B. diazoefficiens* (SEMIA5080), *B. elkanii* (USDA46), *B. iriomotense* (EK05), *B. diazoefficiens* (USDA110), *B. lablabi* (CCBAU61434), *B. liaoningense* (SEMIA5025), *B. retamae* (Ro19), and *B. yuanmingense* (R2m). Reference strains were chosen to represent all known species of *Bradyrhizobium* that

aligned fully with our sequenced *glnII* and *recA* regions (National Center for Biotechnology Information (NCBI) as of November 18, 2014), and *Mesorhizobium loti* (MAFF303099) was used as an outgroup. All sequences were blasted against the reference strains to confirm their identity as *Bradyrhizobium* spp. The GTR model of evolution was selected from the Akaike information criterion in jModelTest2 [46], and the phylogenetic tree was reconstructed in PhyML 3.0 [47] utilizing a BioNJ starting tree and subtree pruning and regrafting (SPR). Branch support was estimated using the fast approximate likelihood ratio test (aLRT) and the Shimodaira–Hasegawa-like (SH-like) procedure [48]. To be consistent with other studies, *Bradyrhizobium* species were defined as the monophyletic clades including no more than one type strain with branch support ≥ 0.90 [49] and attempting to adhere to past species demarcations that utilized some of the same loci [35]. We analyzed inter-species variation using the ratio of fixed to shared polymorphisms using DnaSP [50].

Sequence Statistics

Using the concatenated dataset, we calculated strain richness (number of unique haplotypes/number of isolates) and strain dominance (abundance of each haplotype/number of isolates) analogues of species richness and evenness [51]. For each of the 14 field collection sites, haplotypes were defined as dominant if they were collected at least five times and represented at least 10 % of the total isolates at that site. Spatial spread was defined as the maximum distance between any individual isolates with the same haplotype. GPS coordinates for distances used the midpoint of each of the 14 field collection sites, because distances within sites were small compared to between site differences. We also calculated Hd (haplotype diversity—probability that two haplotypes drawn uniformly at random from the population are not the same), π (nucleotide diversity—average number of nucleotide differences per site between two sequences), k (average number of nucleotide differences), linkage disequilibrium (average absolute D'), recombination (R), and the minimum number of recombination events [52–56] using DnaSP [50].

Trait Analysis

We tested for significant phylogenetic signal for the traits of symbiotic capacity, abundance, and spatial spread, which is a prerequisite for quantitative analysis of traits in a phylogenetic framework. We used Pagel's lambda, estimated with the “fit discrete” function in the “Geiger” package [57], and for symbiotic capacity, we also used Fritz Purvis' D , which was estimated using the “phylo.d” function in the “Caper” package [58]. The Mk1 model of maximum likelihood as well as parsimony were used for ancestral state reconstruction of symbiotic capacity with a modified phylogenetic tree in Mesquite

[59]. Duplicate taxa were added to the phylogenetic tree whenever a single haplotype encompassed both symbiotic and non-symbiotic isolates to avoid the ambiguity of multiple character states being assigned to a single taxon (i.e., haplotype). We tested for correlated evolution between symbiotic capacity and haplotype abundance with the `phy.anova` command in the Geiger package in R [57]. We took several steps to avoid bias caused by sampling in this analysis. Firstly, we used the subset of isolates collected from plants where both root surface and nodule collections had been made (to avoid sampling bias). Secondly, among these plants, we equally sampled across the whole root system representing samples from nodules and from the soil-root interface with equal probability per sampled root. The resultant dataset included 442 root surface isolates and 116 nodule isolates from three field locales. We also used a standard ANOVA in JMP [60] to examine variation between symbiotic and non-symbiotic isolates in terms of abundance and spatial spread. This latter analysis does not take phylogenetic relationships into account and thus assumes that trait data are independent of strain relatedness.

Community Structure

We analyzed isolation by distance with a Mantel test correlating F_{st} and physical distance matrices within PASSaGE [61]. We used Fast UniFrac [62] to test for significant differentiation among *Bradyrhizobium* communities at different collection sites. The “cluster sample” tool was used to cluster the collection sites based on the phylogenetic lineages they contained, and the “jackknife sample cluster” tool was used to assess confidence in the collection site clusters. We employed the “sample distance matrix” to numerically compare distances among collection sites. Abundance was incorporated into Fast UniFrac analyses whenever possible. The jackknife analysis was performed with the number of sequences kept equal to the smallest sample size with 100 permutations. We used the “exact test of population differentiation” in Arlequin [63] to assess differentiation among collection sites and to investigate other drivers of *Bradyrhizobium* community structure including root isolate types (nodule, root-tip surface, and old root surface), symbiotic statuses (symbiotic and non-symbiotic), and collection year.

Results

Haplotype Designation, Abundance, and Spatial Spread

The 1292 concatenated *glnII* and *recA* sequences resulted in 290 haplotypes, all of which were classified as *Bradyrhizobium*. Most of the haplotypes were unique (isolated a single time, 184/290; Table S1). Among the remaining

haplotypes, 13 were defined as dominant in at least one site and these 13 haplotypes constituted the majority of collected isolates (706/1292). We found dominant haplotypes at all but the least sampled field collection site (Anza Borrego Desert State Park–Palm Canyon; Table 1). Most haplotypes (257/290) were only found at a single collection site. Conversely, among the dominant haplotypes, most were also found to be epidemic (7/13; collected at sites spanning ≥ 10 km). Spatial spread for epidemic haplotypes varied from ~ 100 to 750 km, and we collected epidemic haplotypes at all but the two least sampled sites (Anza Borrego Desert State Park–Palm Canyon and Pismo Dunes Natural Preserve) (Table 1). One epidemic haplotype (G03_R01) encompassed 27 % of all isolates collected (355 isolates) and was found at all but four collection sites (Pismo Dunes Natural Preserve, Griffith Park, and Anza Borrego State Park–Road/Palm Canyon sites; Fig. 1).

Symbiotic Capacity Assessment

Using the combined dataset of greenhouse inoculation assays and PCR assays, we inferred 886 isolates to be symbiotic and 406 to be non-symbiotic (Table S1). For ~ 2 % of the 1292 isolates, there was at least one piece of conflicting information about symbiotic capacity, most often including nodule isolates that failed to amplify one or more of the four symbiosis island loci (17) but also including conflicting results between PCR amplification assays (9) (Table S1).

Most dominant haplotypes only encompassed isolates that were inferred to be symbiotic, but a single dominant haplotype (G64_R29) only had non-symbiotic isolates. Of the dominant haplotypes that included both symbiotic and non-symbiotic isolates, mean abundance was higher for symbiotic (36) versus non-symbiotic isolates (15.6) but the difference was not significant ($t=1.48$, $df=4$, $p=0.214$). Among the epidemic haplotypes, most encompassed symbiotic and non-symbiotic isolates (5/7), with symbiotic isolates being more frequent on average than non-symbiotic ones, but without a significant difference (90.6 vs 18.4; $t=2.05$, $df=4$, $p=0.110$).

Phylogenetic Analysis

The *Bradyrhizobium* phylogeny encompassed 20 deeply diverged, monophyletic lineages (clades) that are consistent with species-level divergence, including six that were previously identified (*B. betae*, *B. canariense*, *B. cytisi*, *B. liaoningense*, and *B. retamae*) and 14 clades that did not encompass one of the type strains (Figs. 2 and S1). Most (161) comparisons between these 20 clades uncovered more fixed than shared polymorphisms, also consistent with species designation. Two comparisons had the same number of fixed and shared polymorphisms, and 27 had more shared than fixed polymorphisms (Table S2). Almost half of these clades (8/20) were only collected at a single site. However, nearly

Table 1 Dominant and epidemic haplotypes

		Number of isolates from collection site	Total			Symbiotic		Non-symbiotic	
			Haplotype	Abundance	Spatial spread	Abundance	Spatial spread	Abundance	Spatial spread
Dominant haplotypes	Anza Borrego State Park– Palm Canyon	9	None			None		None	
	Anza Borrego State Park– Roadside	36	G58_R27	23	135.7	23	135.7	None	
			G58_R34	9	135.7	9	135.7	None	
	Bodega Marine Reserve	211	G03_R01	72	750.6	71	750.6	1	723.8
			G05_R02	24	0	22	0	2	0
	Burns Piñon Ridge Reserve	39	G03_R01	24	750.6	24	750.6	None	
			G11_R07	9	175.9	9	175.9	None	
	Gaudalupe-Nipomo Dunes National Wildlife Refuge	100	G40_R38	5	168.8	5	168.8	None	
			G03_R01	18	750.6	18	750.6	None	
	Griffith Park	69	G05_R65	35	0	35	0	None	
			G11_R07	7	175.9	7	175.9	None	
			G40_R38	50	168.8	50	168.8	None	
	Madrona Marsh Preserve	86	G03_R01	24	750.6	24	750.6	None	
			G11_R01	25	335.3	25	335.3	None	
	Motte Rimrock Reserve	120	G36_R35	10	100.2	10	100.2	None	
			G03_R01	35	750.6	23	750.6	12	723.8
			G11_R01	22	335.3	19	335.3	3	18.86
	Pismo Dunes Natural Preserve	21	G64_R29	13	0	None		13	0
			G99_R62	5	0	5	0	None	
			G03_R01	26	750.6	26	750.6	None	
	Robert J. Bernard Biological Field Station	68	G11_R01	11	335.3	11	335.3	None	
			G71_R32	11	7.04	11	7.04	None	
	San Dimas Canyon	67	G03_R01	11	750.6	11	750.6	None	
			G36_R35	9	100.2	9	100.2	None	
			G71_R39	8	0	8	0	None	
	San Dimas Reservoir	63	G03_R01	18	750.6	18	750.6	None	
			G11_R07	16	175.9	16	175.9	None	
	UC Riverside Hills	356	G03_R01	105	750.6	45	750.6	60	723.8
	Whitewater Preserve	47	G03_R01	22	750.6	22	750.6	None	
Epidemic haplotype	Epidemic haplotypes		G03_R01	355	750.6	282	750.6	73	723.8
			G11_R01	81	335.3	70	335.3	11	18.86
			G11_R07	62	175.9	58	175.9	4	0
			G36_R35	20	100.2	19	100.2	1	0
			G40_R38	55	168.8	55	168.8	None	
			G58_R27	27	135.7	24	135.7	3	0
			G58_R34	10	135.7	9	135.7	None	

All dominant haplotypes (collected ≥ 5 times per site and encompassing at least 10 % of the isolates at the site) are listed for each collection site followed by epidemic haplotypes (dominant haplotypes that have spread at least 10 km away). Abundance indicates the number of times the haplotype was collected at a given site, whereas the spatial spread is the greatest distance between collection sites the haplotype was found at. The information is then broken down by symbiotic capacity

all collection sites (13/14) were inhabited by multiple clades of *Bradyrhizobium* (Table S3). *Bradyrhizobium canariense* was particularly widespread and was collected at 11 sites. Population genetic statistics were analyzed for all *Bradyrhizobium* clades that encompassed multiple isolates

(Table 2). Linkage was high between all SNPs for all clades (>0.9). Strain richness, Hd, π , and recombination varied widely between clades, likely because of variation in sampling. When only clades with over 40 isolates were assessed, Hd, π , and recombination were comparable.

Fig. 1 Map of California indicates collection sites with black dots. Pie charts connected to black dots illustrate the relative frequencies of five focal haplotypes. The five haplotypes chosen incorporate the four haplotypes with the highest abundance and the four haplotypes with the greatest spatial spread. The distribution of epidemic haplotype G03_R01 is hypothesized based on presence or absence of data from all collection sites. Starting from the upper left and moving clockwise, the collection sites are Bodega Marine Reserve, San Dimas Canyon, San Dimas Reservoir, Bernard Field Station, UC Riverside, Motte Rimrock Reserve, Whitewater, Burns Pinon Ridge, Anza Borrego Canyon, Anza Borrego Roadsize, Madrona Marsh, Griffith Park, Nipomo Dunes, and Pismo Dunes

Ancestral state reconstructions were similar for parsimony and likelihood models (Figs. 2 and S2), and both infer gains and or losses of symbiotic capacity across multiple *Bradyrhizobium* clades (Table S4). Because many of the nodes on the phylogenetic tree are poorly resolved, it is impossible to estimate the number of gain or loss events with confidence. Considering only the 20 deeply diverged clades with likelihood support values >0.90 , 10 of them encompassed both symbiotic and non-symbiotic isolates consistent with multiple gain and loss events (Fig. S1). Using both parsimony and a conservative maximum likelihood estimate (likelihood decision threshold of 2.0), we inferred about 1.5–2 $\times$ as many loss events as gains (Table S4).

Symbiotic capacity exhibited significant phylogenetic signal, but abundance and spatial spread did not (Table S5). When we assessed symbiotic capacity and abundance for correlated evolution using chi-squared tests in JMP, we found a significant positive correlation; hence, that symbiotic clades on average exhibit higher abundance than non-symbiotic clades. When we tested for correlated evolution of these traits, we did not find a relationship (Tables S6 and S7); hence, that acquisition of a symbiosis island is not statistically associated with increase in haplotype abundance. Moreover, we did not find evidence for correlated evolution of symbiotic capacity with spatial spread using either the chi-squared tests in JMP or the phy.anova in Geiger.

Community Analyses

We found significantly different *Bradyrhizobium* communities among most collection sites (Table S8), among different isolate types (nodule, root-tip surface, and old root surface), symbiotic statuses (symbiotic and non-symbiotic), and collection years (Table S9). Non-symbiotic populations were significantly differentiated from isolates that were symbiotic within the same site at the Bodega Marine Reserve and Motte Rimrock reserve (Table S10). Two of four sites that were sampled multiple times exhibited population differentiation from year to year, San Dimas Canyon and Burns Piñon Ridge Reserve (Table S11). We did not find evidence for isolation by distance using the Mantel test. This is supported by the Fast UniFrac analyses in which clustering closely follows species makeup rather than geographical location (Figs. S3 and S4).

Discussion

Our study uncovered an epidemic distribution of *Bradyrhizobium* haplotypes across California with a striking divide between rare and dominant haplotypes. Although we recovered 290 haplotypes, the majority of isolates were encompassed within 13 dominant haplotypes (707/1292). Among the dominant haplotypes, six were endemic and seven were epidemic including haplotype G03_R01 that was dominant at most sites (10/14), exhibited a spatial spread of

Fig. 2 PhyML 3.0 phylogenetic tree reconstructed from concatenated glutamine synthetase (*glnII*) and recombinase (*recA*) loci. Non-symbiotic taxa are indicated with red dots and symbiotic taxa with black dots. Inferred ancestral states of non-symbiotic and symbiotic are indicated by red and black branches, respectively, and ambiguous nodes are colored grey (estimated likelihood proportion was <66.66 %). The relative abundance of a haplotype and the spatial spread are indicated by stacked blue and green bars. Major clades are indicated with brackets. Reference strains can be identified by the lack of symbiotic capacity,

abundance, and spatial spread data. The strains include (1) *Mesorhizobium loti* MAFF303099, (2) *Bradyrhizobium retamae* Ro19, (3) *Bradyrhizobium elkanii* USDA46, (4) *Bradyrhizobium lablabi* CCBAU61434, (5) *B. diazoefficiens* SEMIA5080, (6) *B. diazoefficiens* (*japonicum*) USDA110, (7) *Bradyrhizobium betae* PL7HG1, (8) *Bradyrhizobium iriomotense* EK05, (9) *Bradyrhizobium arachidis* CCBAU33067, (10) *Bradyrhizobium yuanmingense* R2m, (11) *Bradyrhizobium liaoningense* SEMIA5025, (12) *Bradyrhizobium cytisi* LMG25866, and (13) *Bradyrhizobium canariense* SEMIA928

750 km, and constituted nearly 30 % of the total isolates (355/1292) (Table 1). Several studies have uncovered rhizobial genotypes that achieve high frequency at localized sites [33, 51, 64], but haplotype G03_R01 dominates both locally and also across major ecological boundaries including mountain ranges and areas of inhospitable habitat. It is striking that haplotype G03_R01 was found in sites that vary markedly in their patterns of rainfall, temperature, plant community, and key soil nutrients [65, 66], especially given that most of the *Bradyrhizobium* lineages that we uncovered were relatively localized in their distribution. Recent work that focused on non-symbiotic lineages of *Bradyrhizobium* in soils has also found evidence of widespread genotypes, suggesting that this might be a common pattern for these bacteria [37].

We uncovered a broad diversity of *Bradyrhizobium* species nodulating and inhabiting the root surfaces of *A. strigosus*. The most recent efforts in defining symbiotic lineages of *Bradyrhizobium* propose 19 species [36]. Using a

conservative species definition (monophyletic clades including no more than one type strain with branch support ≥ 0.90), we recovered 20 species including 14 lineages that were not represented in the NCBI database using these common loci (Fig. S1). No single method has been agreed upon for defining species of bacteria, but most definitions use phylogenetic clusters of multiple conserved loci, as we did here [67]. Despite the diversity that we uncovered, six out of the seven epidemic haplotypes are found within the *B. canariense* and *B. novel I* species (Table 2). Important differences can exist between rhizobial lineages including the diversity of host plants infected, metabolite utilization, and antibiotic resistance [68–70], and these sources of variation may be driving differences in epidemic distributions among *Bradyrhizobium* taxa. We also found that *Bradyrhizobium* communities from different collection sites varied in species diversity (Table S3). We found the greatest diversity at the three sites where we collected root surface samples, consistent with previous work uncovering

Table 2 Species statistics for 19 *Bradyrhizobium* clades calculated in DnaSP

	Number of isolates	Number of haplotypes	Number of variable sites	Strain richness (number of haplotypes/number of isolates)	Hd (haplotype diversity)	π (nucleotide diversity)	k (average number of nucleotide differences)	Linkage disequilibrium Average $ D' $	Recombination		
									Estimate of R of R /gene	Estimate between adjacent sites	Minimum number of recombination events
<i>B. betae</i>	15	7	77	0.47	0.84	0.03	28	1.00	0.001	0.000	0
<i>B. canariense</i>	798	105	215	0.13	0.78	0.01	8	0.97	0.001	0.000	41
<i>B. cytisi</i>	40	14	92	0.35	0.85	0.03	30	0.98	0.300	0.000	9
<i>B. lablabi</i>	9	7	101	0.78	0.94	0.03	32	0.99	1.200	0.001	8
<i>B. liaoningense</i>	9	6	125	0.67	0.89	0.04	39	1.00	0.001	0.000	4
<i>B. novel I</i>	134	17	54	0.13	0.78	0.01	8	0.91	0.001	0.000	5
<i>B. novel II</i>	73	22	111	0.30	0.87	0.02	16	0.95	0.001	0.000	17
<i>B. novel III</i>	12	6	189	0.50	0.82	0.09	84	0.95	5.800	0.006	16
<i>B. novel IV</i>	3	2	4	0.67	0.67	0.00	3	1.00	7.300	0.008	0
<i>B. novel V</i>	6	2	33	0.33	0.60	0.02	20	1.00	0.700	0.001	0
<i>B. novel VI</i>	6	5	180	0.83	0.93	0.10	93	0.95	5.500	0.006	10
<i>B. novel VII</i>	3	2	67	0.67	0.67	0.05	45	1.00	1.500	0.002	0
<i>B. novel VIII</i>	15	4	48	0.27	0.67	0.01	11	1.00	0.001	0.000	3
<i>B. novel IX</i>	8	4	4	0.50	0.64	0.00	1	1.00	54.600	0.056	0
<i>B. novel X</i>	70	18	40	0.26	0.74	0.01	6	0.96	1.700	0.002	5
<i>B. novel XI</i>	60	45	80	0.75	0.99	0.01	12	0.93	19.800	0.020	12
<i>B. novel XII</i>	4	4	51	1.00	1.00	0.03	32	1.00	>10.000	Not applicable (n/a)	0
<i>B. novel XIII</i>	2	2	4	1.00	1.00	0.00	4	0.96	20.700	0.021	6
<i>B. novel XIV</i>	2	2	7	1.00	1.00	0.01	7	1.00	>10.000	n/a	0

Bradyrhizobium retamae was excluded from analyses as it included only a single isolate. Analyses included strain richness (number of unique haplotypes/number of isolates), Hd (haplotype diversity—probability that two haplotypes drawn uniformly at random from the population are not the same), π (nucleotide diversity—average number of nucleotide differences per site between two sequences), k (average number of nucleotide differences), linkage disequilibrium (average absolute D'), recombination (R), and the minimum number of recombination events

greater genotypic diversity among root surface isolates as opposed to nodule populations [33].

Genomic island acquisition can facilitate novel bacterial traits that confer fitness benefits in an environment specific manner [71]. However, we did not find support for the hypothesis that symbiosis island acquisition is an evolutionary driver of strain abundance or spatial spread in *Bradyrhizobium* (Table S7). Instead, we uncovered epidemics of only a few extremely abundant symbiotic haplotypes within *B. canariense*, and in this clade, we found evidence consistent with multiple events of symbiosis island gain and loss. Our dataset shows that both acquisition and loss of the symbiosis island appear to occur frequently across the sampled *Bradyrhizobium* populations. Both maximum likelihood and parsimony analyses suggest many more loss than gain transitions. Many types of mutations can result in the conversion of symbiotic strain to a non-symbiotic. But only the whole-scale horizontal transmission of the symbiosis island has been associated with gain of nodulation and nitrogen fixation in *Bradyrhizobium*. Given the more frequent loss events, some additional diversification must occur in the symbiotic lineages relative to the lineages without symbiosis islands.

The capacity to infect *A. strigosus* appears to have a significant affect on structuring *Bradyrhizobium* populations. We found support for differentiation in 2/3 comparisons of symbiotic and non-symbiotic communities (Table S10), and we also found that 8/9 comparisons of *Bradyrhizobium* communities collected from different plant parts exhibited differentiation (Table S9). We also found evidence for local differences among collection sites that could be responsible for *Bradyrhizobium* population differentiation; however, drift cannot be ruled out by these analyses (Table S8). Finally, we found evidence for temporal differentiation in 3/5 comparisons (Table S11). Taken together, these data are consistent with previous evidence showing that plant and soil factors affect population structure in the rhizosphere [72].

In summary, our analysis found that native *Bradyrhizobium* populations across California are dominated by a very small handful of haplotypes. We found that on average, symbiotic strains are more common than non-symbiotic ones and that symbiotic strains are also more likely to be dominant and to spread spatially. Nonetheless, we found evidence of both symbiotic and non-symbiotic strains achieving high abundance and spreading across great distances. Our ancestral state reconstruction inferred that the symbiosis island has been repeatedly gained and lost across the *A. strigosus* sampling range, and we did not find support for the hypothesis that acquisition of symbiosis islands serves as a driver of strain dominance or spread. We suggest that traits encoded on the *Bradyrhizobium* chromosome, that expresses traits important for soil survival [13, 14], are most likely responsible for variation in the capacity to spread epidemically.

Acknowledgments The following grants supported this study: to ACH a Herbert Kraft Scholarship and a UC Riverside Graduate Research Mentorship Fellowship and to JLS NSF DEB 0816663 and NSF DEB 1150278.

References

- Bright M, Bulgheresi S (2010) A complex journey: transmission of microbial symbionts. *Nat Rev Microbiol* 8:218–230. doi:10.1038/nrmicro2262
- Sachs JL, Skophammer RG, Bansal N, Stajich JE (2013) Evolutionary origins and diversification of proteobacterial mutualists. *Proc R Soc B Biol Sci* 281:20132146–20132146. doi:10.1098/rspb.2013.2146
- Groisman EA, Ochman H (1996) Pathogenicity islands: bacterial evolution in quantum leaps. *Cell* 87:791–794. doi:10.1016/S0092-8674(00)81985-6
- Jain R, Rivera MC, Moore JE, Lake JA (2003) Horizontal gene transfer accelerates genome innovation and evolution. *Mol Biol Evol* 20:1598–1602. doi:10.1093/molbev/msg154
- Gal-Mor O, Finlay BB (2006) Pathogenicity islands: a molecular toolbox for bacterial virulence. *Cell Microbiol* 8:1707–1719. doi:10.1111/j.1462-5822.2006.00794.x
- Bach S (2000) The *Yersinia* high-pathogenicity island is present in different members of the family Enterobacteriaceae. *FEMS Microbiol Lett* 183:289–294. doi:10.1016/S0378-1097(00)00005-7
- Diep BA, Gill SR, Chang RF et al (2006) Complete genome sequence of USA300, an epidemic clone of community-acquired methicillin-resistant *Staphylococcus aureus*. *Lancet* 367:731–739. doi:10.1016/S0140-6736(06)68231-7
- Qiu X, Gurkar AU, Lory S (2006) Interstrain transfer of the large pathogenicity island (PAPI-1) of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci* 103:19830–19835. doi:10.1073/pnas.0606810104
- Sprent JI (2001) Nodulation in legumes. Royal Botanic Gardens, Kew
- Sugawara M, Epstein B, Badgley BD et al (2013) Comparative genomics of the core and accessory genomes of 48 *Sinorhizobium* strains comprising five genospecies. *Genome Biol* 14:R17. doi:10.1186/gb-2013-14-2-r17
- Cleveland CC, Townsend AR, Schimel DS et al (1999) Global patterns of terrestrial biological nitrogen (N₂) fixation in natural ecosystems. *Glob Biogeochem Cycles* 13:623–645. doi:10.1029/1999GB900014
- Graham PH, Vance CP (2003) Legumes: importance and constraints to greater use. *Plant Physiol* 131:872–877. doi:10.1104/pp.017004
- Uchiumi T, Ohwada T, Itakura M et al (2004) Expression islands clustered on the symbiosis island of the *Mesorhizobium loti* genome. *J Bacteriol* 186:2439–2448. doi:10.1128/JB.186.8.2439-2448.2004
- Pessi G, Ahrens CH, Rehrauer H et al (2007) Genome-wide transcript analysis of *Bradyrhizobium japonicum* bacteroids in soybean root nodules. *Mol Plant Microbe Interact* 20:1353–1363. doi:10.1094/MPMI-20-11-1353
- Kaneko T (2000) Complete genome structure of the nitrogen-fixing symbiotic bacterium *Mesorhizobium loti*. *DNA Res* 7:331–338. doi:10.1093/dnares/7.6.331
- Galibert F (2001) The composite genome of the legume symbiont *Sinorhizobium meliloti*. *Science* 293:668–672. doi:10.1126/science.1060966
- Kaneko T, Nakamura Y, Sato S et al (2002) Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110. *DNA Res* 9:189–197. doi:10.1093/dnares/9.6.189

18. Young JPW, Crossman LC, Johnston AWB et al (2006) The genome of *Rhizobium leguminosarum* has recognizable core and accessory components. *Genome Biol* 7:R34. doi:10.1186/gb-2006-7-4-r34
19. Lee K-B, De Backer P, Aono T et al (2008) The genome of the versatile nitrogen fixer *Azorhizobium caulinodans* ORS571. *BMC Genomics* 9:271. doi:10.1186/1471-2164-9-271
20. Young JPW (1996) Diversity and phylogeny of rhizobia. *New Phytol* 133:87–94
21. Perret X, Staehelin C, Broughton WJ (2000) Molecular basis of symbiotic promiscuity. *Microbiol Mol Biol Rev* 64:180–201. doi:10.1128/MMBR.64.1.180-201.2000
22. Moulin L, Béna G, Boivin-Masson C, Stepkowski T (2004) Phylogenetic analyses of symbiotic nodulation genes support vertical and lateral gene co-transfer within the *Bradyrhizobium* genus. *Mol Phylogenet Evol* 30:720–732. doi:10.1016/S1055-7903(03)00255-0
23. Segovia L, Piñero D, Palacios R, Martínez-Romero E (1991) Genetic structure of a soil population of *Rhizobium leguminosarum*. *Appl Environ Microbiol* 57:426–433
24. Sachs JL, Ehinger MO, Simms EL (2010) Origins of cheating and loss of symbiosis in wild *Bradyrhizobium*. *J Evol Biol* 23:1075–1089. doi:10.1111/j.1420-9101.2010.01980.x
25. Sullivan JT, Eardly BD, van Berkum P, Ronson CW (1996) Four unnamed species of nonsymbiotic rhizobia isolated from the rhizosphere of *Lotus corniculatus*. *Appl Environ Microbiol* 62:2818–2825
26. Saito A, Mitsui H, Hattori R et al (1998) Slow-growing and oligotrophic soil bacteria phylogenetically close to *Bradyrhizobium japonicum*. *FEMS Microbiol Ecol* 25:277–286. doi:10.1111/j.1574-6941.1998.tb00480.x
27. Pongsilp N, Teamroong N, Nuntagij A et al (2002) Genetic structure of indigenous non-nodulating and nodulating populations of *Bradyrhizobium* in soils from Thailand. *Symbiosis* 33:39–58
28. Okubo T, Tsukui T, Maita H et al (2012) Complete genome sequence of *Bradyrhizobium* sp. S23321: insights into symbiosis evolution in soil oligotrophs. *Microbes Environ JSME* 27:306–315
29. Bhatt AS, Freeman SS, Herrera AF et al (2013) Sequence-based discovery of *Bradyrhizobium enterica* in cord colitis syndrome. *N Engl J Med* 369:517–528. doi:10.1056/NEJMoal211115
30. Chaintreuil C, Giraud E, Prin Y et al (2000) Photosynthetic bradyrhizobia are natural endophytes of the African wild rice *Oryza breviligulata*. *Appl Environ Microbiol* 66:5437–5447. doi:10.1128/AEM.66.12.5437-5447.2000
31. Costello EK, Carlisle EM, Bik EM, et al. (2013) Microbiome assembly across multiple body sites in low-birthweight infants. *mBio* 4:e00782–13–e00782–13. doi: 10.1128/mBio.00782-13
32. Hunt KM, Foster JA, Forney LJ et al (2011) Characterization of the diversity and temporal stability of bacterial communities in human milk. *PLoS ONE* 6, e21313. doi:10.1371/journal.pone.0021313
33. Sachs JL, Kembel SW, Lau AH, Simms EL (2009) In situ phylogenetic structure and diversity of wild *Bradyrhizobium* communities. *Appl Environ Microbiol* 75:4727–4735. doi:10.1128/AEM.00667-09
34. Vinuesa P, Silva C, Werner D, Martínez-Romero E (2005) Population genetics and phylogenetic inference in bacterial molecular systematics: the roles of migration and recombination in *Bradyrhizobium* species cohesion and delineation. *Mol Phylogenet Evol* 34:29–54. doi:10.1016/j.ympev.2004.08.020
35. Vinuesa P, Rojas-Jimenez K, Contreras-Moreira B et al (2008) Multilocus sequence analysis for assessment of the biogeography and evolutionary genetics of four *Bradyrhizobium* species that nodulate soybeans on the Asiatic continent. *Appl Environ Microbiol* 74:6987–6996. doi:10.1128/AEM.00875-08
36. Azevedo A, Martins-Lopes F, Silla RP, Hungria M (2015) A database for the taxonomic and phylogenetic identification of the genus *Bradyrhizobium* using multilocus sequence analysis. *BMC Genomics* 16:S10
37. VanInsberghe D, Maas KR, Cardenas E et al (2015) Non-symbiotic *Bradyrhizobium* ecotypes dominate North American forest soils. *ISME J*. doi:10.1038/ismej.2015.54
38. Rivas R, Martens M, de Lajudie P, Willems A (2009) Multilocus sequence analysis of the genus *Bradyrhizobium*. *Syst Appl Microbiol* 32:101–110. doi:10.1016/j.syapm.2008.12.005
39. Parker MA (2015) The spread of *Bradyrhizobium* lineages across host legume clades: from *Abarema* to *Zygia*. *Microb Ecol*. doi:10.1007/s00248-014-0503-5
40. Göttfert M, Röthlisberger S, Kündig C et al (2001) Potential symbiosis-specific genes uncovered by sequencing a 410-kilobase DNA region of the *Bradyrhizobium japonicum* chromosome. *J Bacteriol* 183:1405–1412. doi:10.1128/JB.183.4.1405-1412.2001
41. Giraud E, Moulin L, Vallenet D et al (2007) Legumes symbioses: absence of Nod genes in photosynthetic bradyrhizobia. *Science* 316:1307–1312. doi:10.1126/science.1139548
42. Maddison WP, Maddison DR (2005) MacClade: analysis of phylogeny and character evolution
43. Sachs JL, Russell JE, Hollowell AC (2011) Evolutionary instability of symbiotic function in *Bradyrhizobium japonicum*. *PLoS ONE* 6, e26370. doi:10.1371/journal.pone.0026370
44. Ehinger M, Mohr TJ, Starcevich JB et al (2014) Specialization-generalization trade-off in a *Bradyrhizobium* symbiosis with wild legume hosts. *BMC Ecol* 14:8. doi:10.1186/1472-6785-14-8
45. Parker M (2000) Divergent *Bradyrhizobium* symbionts on *Tachigali versicolor* from Barro Colorado Island, Panama. *Syst Appl Microbiol* 23:585–90
46. Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods* 9:772. doi:10.1038/nmeth.2109
47. Guindon S, Dufayard JF, Lefort V et al (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59:307–321. doi:10.1093/sysbio/syq010
48. Anisimova M, Gascuel O (2006) Approximate likelihood-ratio test for branches: a fast, accurate, and powerful alternative. *Syst Biol* 55:539–552. doi:10.1080/10635150600755453
49. Shimodaira H, Hasegawa M (1999) Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol Biol Evol* 16:1114
50. Librado P, Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25:1451–1452. doi:10.1093/bioinformatics/btp187
51. McInnes A (2004) Structure and diversity among rhizobial strains, populations and communities—a review. *Soil Biol Biochem* 36: 1295–1308. doi:10.1016/j.soilbio.2004.04.011
52. Lewontin RC (1964) The interaction of selection and linkage. I. General considerations; heterotic models. *Genetics* 49:49–67
53. Tajima F (1983) Evolutionary relationship of DNA sequences in finite populations. *Genetics* 105:437–460
54. Hudson RR, Kaplan NL (1985) Statistical properties of the number of recombination events in the history of a sample of DNA sequences. *Genetics* 111:147–164
55. Hudson RR (1987) Estimating the recombination parameter of a finite population model without selection. *Genet Res* 50:245–250
56. Nei M (1987) Molecular evolutionary genetics. Columbia University Press, New York
57. Harmon LJ, Weir JT, Brock CD et al (2007) GEIGER: investigating evolutionary radiations. *Bioinformatics* 24:129–131. doi:10.1093/bioinformatics/btm538
58. Orme D (2012) The caper package: comparative analysis of phylogenetics and evolution in R
59. Maddison, W. P., Maddison, D. R. (2011) Mesquite
60. SAS Institute Inc (1989) JMP. SAS Institute Inc., Cary, NC

61. Rosenberg MS, Anderson CD (2011) PASSaGE: pattern analysis, spatial statistics and geographic exegesis. Version 2: PASSaGE. *Methods Ecol Evol* 2:229–232. doi:[10.1111/j.2041-210X.2010.00081.x](https://doi.org/10.1111/j.2041-210X.2010.00081.x)
62. Hamady M, Lozupone C, Knight R (2010) Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data. *ISME J* 4:17–27. doi:[10.1038/ismej.2009.97](https://doi.org/10.1038/ismej.2009.97)
63. Excoffier L, Lischer HEL (2010) Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* 10:564–567. doi:[10.1111/j.1755-0998.2010.02847.x](https://doi.org/10.1111/j.1755-0998.2010.02847.x)
64. Silva C, Eguiarte LE, Souza V (1999) Reticulated and epidemic population genetic structure of *Rhizobium etli* biovar *phaseoli* in a traditionally managed locality in Mexico. *Mol Ecol* 8:277–287. doi:[10.1046/j.1365-294X.1999.00564.x](https://doi.org/10.1046/j.1365-294X.1999.00564.x)
65. Baldwin BG, Goldman DH (2012) The Jepson manual: vascular plants of California, 2nd edn. University of California Press, Berkeley, Calif
66. Regus JU, Gano KA, Hollowell AC, Sachs JL (2014) Efficiency of partner choice and sanctions in *Lotus* is not altered by nitrogen fertilization. *Proc R Soc B Biol Sci* 281:20132587–20132587. doi:[10.1098/rspb.2013.2587](https://doi.org/10.1098/rspb.2013.2587)
67. Thompson CC, Amaral GR, Campeao M et al (2015) Microbial taxonomy in the post-genomic era: rebuilding from scratch? *Arch Microbiol* 197:359–370. doi:[10.1007/s00203-014-1071-2](https://doi.org/10.1007/s00203-014-1071-2)
68. Dupuy N, Willems A, Pot B et al (1994) Phenotypic and genotypic characterization of bradyrhizobia nodulating the leguminous tree *Acacia albida*. *Int J Syst Bacteriol* 44:461–473. doi:[10.1099/00207713-44-3-461](https://doi.org/10.1099/00207713-44-3-461)
69. Geurts R (1996) Signal transduction in *Rhizobium*-induced nodule formation. *Plant Physiol* 112:447–453. doi:[10.1104/pp.112.2.447](https://doi.org/10.1104/pp.112.2.447)
70. Hollowell AC, Gano KA, Lopez G et al (2015) Native California soils are selective reservoirs for multidrug-resistant bacteria. *Environ Microbiol Rep*. doi:[10.1111/1758-2229.12269](https://doi.org/10.1111/1758-2229.12269)
71. Dobrindt U, Hochhut B, Hentschel U, Hacker J (2004) Genomic islands in pathogenic and environmental microorganisms. *Nat Rev Microbiol* 2:414–424. doi:[10.1038/nrmicro884](https://doi.org/10.1038/nrmicro884)
72. Berg G, Smalla K (2009) Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere: plant species, soil type and rhizosphere communities. *FEMS Microbiol Ecol* 68:1–13. doi:[10.1111/j.1574-6941.2009.00654.x](https://doi.org/10.1111/j.1574-6941.2009.00654.x)