

# UCLA

## UCLA Previously Published Works

### Title

Impact of Cognitive Impairment Across Specialties: Summary of a Report From the U13 Conference Series

### Permalink

<https://escholarship.org/uc/item/0683t0cr>

### Journal

Journal of the American Geriatrics Society, 67(10)

### ISSN

0002-8614

### Authors

Carpenter, Christopher R  
McFarland, Frances  
Avidan, Michael  
et al.

### Publication Date


2019-10-01

### DOI

10.1111/jgs.16093

Peer reviewed

# Impact of Cognitive Impairment Across Specialties: Summary of a Report From the U13 Conference Series

Christopher R. Carpenter, MD, MSc,\*  Frances McFarland, PhD, MA,<sup>†</sup>  
 Michael Avidan, MBBCh, MD,<sup>‡</sup> Miles Berger, MD,<sup>§</sup> Sharon K. Inouye, MD, MPH,<sup>¶</sup>  
 Jason Karlawish, MD,<sup>||</sup> Frank R. Lin, MD,\*\* Edward Marcantonio, MD, SM,<sup>††</sup>  
 John C. Morris, MD,<sup>‡‡</sup> David B. Reuben, MD,<sup>§§</sup> Raj C. Shah, MD,<sup>¶¶</sup>  
 Heather E. Whitson, MD, MHS,<sup>|||</sup>\*\*\* Sanjay Asthana, MD,<sup>†††</sup> and Joe Verghese, MBBS, MS<sup>‡‡‡</sup>

Although declines in cognitive capacity are assumed to be a characteristic of aging, increasing evidence shows that it is age-related disease, rather than age itself, that causes cognitive impairment. Even so, older age is a primary risk factor for cognitive decline, and with individuals living longer as a result of medical advances, cognitive impairment and dementia are increasing in prevalence. On March 26 to 27, 2018, the American Geriatrics Society convened a conference in Bethesda, MD, to explore cognitive impairment across the subspecialties. Bringing together representatives from several subspecialties, this was the third of three conferences, supported by a U13 grant from the National Institute on Aging, to aid recipients of Grants for Early Medical/Surgical

Specialists' Transition to Aging Research (GEMSSTAR) in integrating geriatrics into their subspecialties. Scientific sessions focused on the impact of cognitive impairment, sensory contributors, comorbidities, links between delirium and dementia, and issues of informed consent in cognitively impaired populations. Discussions highlighted the complexity not only of cognitive health itself, but also of the bidirectional relationship between cognitive health and the health of other organ systems. Thus, conference participants noted the importance of multidisciplinary team science in future aging research. This article summarizes the full conference report, "The Impact of Cognitive Impairment Across Specialties," and notes areas where GEMSSTAR scholars can contribute to progress as they embark on their careers in aging research. *J Am Geriatr Soc* 67:2011-2017, 2019.

From the \*Division of Emergency Medicine, Washington University School of Medicine, St Louis, Missouri; <sup>†</sup>Science Writer, Independent Consultant, Annapolis, Maryland; <sup>‡</sup>Department of Anesthesiology, Washington University School of Medicine, St Louis, Missouri; <sup>§</sup>Anesthesiology Department, Duke University Medical Center, Durham, North Carolina; <sup>¶</sup>Marcus Institute for Aging Research, Aging Brain Center, Harvard Medical School and Hebrew SeniorLife, Boston, Massachusetts; <sup>||</sup>University of Pennsylvania, Philadelphia, Pennsylvania; \*\*Department of Otolaryngology, The Johns Hopkins University, Baltimore, Maryland; <sup>††</sup>Divisions of General Medicine and Gerontology, Harvard Medical School, Beth Israel Deaconess Medical Center, Boston, Massachusetts; <sup>‡‡</sup>Department of Neurology, Washington University School of Medicine, St Louis, Missouri; <sup>§§</sup>Department of Medicine, Division of Geriatrics, David Geffen School of Medicine at UCLA, Los Angeles, California; <sup>¶¶</sup>Family Medicine and Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, Illinois; <sup>|||</sup>Department of Medicine, Division of Geriatrics, Duke University School of Medicine, Durham, North Carolina; \*\*\*Geriatrics Research Education and Clinical Center, Durham VA Health Care System, Durham, North Carolina; <sup>†††</sup>Department of Medicine, Division of Geriatrics and Gerontology, University of Wisconsin School of Medicine and Public Health; and the <sup>‡‡‡</sup>Departments of Neurology and Medicine, Albert Einstein College of Medicine, Bronx, New York.

Address correspondence to Christopher R. Carpenter, MD, MSc, Washington University School of Medicine, 660 S Euclid Ave, Campus Box 8072, St Louis, MO 63110. E-mail: carpenterc@wustl.edu.

Drs Sanjay Asthana and Joe Verghese are co-senior authors of this report.  
 DOI: 10.1111/jgs.16093

**Key words:** Aging; delirium; dementia; interdisciplinary research; research ethics

Traditionally, adults are assumed to lose cognitive capacity as they age. However, increasing evidence shows that it is age-related disease, rather than age itself, that causes cognitive impairment. Even so, older age is a primary risk factor for cognitive decline. For example, approximately 10% of individuals aged 65 years and older have Alzheimer disease (AD)-associated dementia. With medical advances leading to individuals living longer, cognitive impairment and dementia are increasing in prevalence and threaten to become a public health crisis.

Increasing evidence suggests that the pathology underlying AD and other dementias begins long before symptoms appear. For example, 50% of neurons in certain brain regions, such as the hippocampus, are already dead<sup>1</sup>; and

other copathologies are present by the time the first symptoms of AD appear.<sup>2</sup> Thus, AD can be viewed as a continuous process of synaptic and neuronal deterioration that can be divided roughly into an asymptomatic, preclinical stage, and a symptomatic stage, with the preclinical stage constituting the bulk of the illness. Efforts toward prevention of dementia are under way, and by 2030, there will likely be more preventive options, including risk factor identification and monitoring. Although the transition from the preclinical stage and relative cognitive normality to the initial onset of symptoms is difficult to detect at present, researchers are now characterizing the presymptomatic stage. Biomarkers have been identified, including cerebrospinal fluid SNAP-25, VILIP-1, and YKL-40.<sup>3</sup> In addition, race-dependent mechanisms for AD are being explored, with lower cerebrospinal fluid tau noted in African Americans than in white participants.<sup>4</sup> Several trials are exploring interventions to prevent or delay the onset of symptoms. Among potential markers of presymptomatic illness are physical function and performance. In a longitudinal study of 444 cognitively normal older adults, 134 of whom were later diagnosed with dementia, individuals who later developed dementia did not perform as well as those who remained stable, even when they appeared to be normal.<sup>5</sup>

Despite the promise of these efforts, the only currently available options for management of dementia include managing the disease while caring for the patient, family, and caregivers. Disease management typically relies on the use of cholinesterase inhibitors and memantine, which have only modest benefits.<sup>6</sup> Caring for the patient includes the management of symptoms, which is best addressed by behavioral approaches, and the management of both dementia-related and non-dementia-related comorbidities. New models of comprehensive care focusing on both patients and caregivers, including community-based models, such as Benjamin Rose Institute (BRI)Care Consultation™ Care Consultation<sup>7</sup> and Maximizing Independence at Home(MIND at Home™) at Home,<sup>8</sup> and health system-based models, such as the multidisciplinary Healthy Aging Brain Center at Indiana University<sup>9</sup> and the University of California, Los Angeles, Alzheimer's and Dementia Care Program,<sup>10</sup> have been shown to reduce caregiver strain and nursing home placements.<sup>7,9-12</sup>

The Grants for Early Medical/Surgical Specialists' Transition to Aging Research (GEMSSTAR) award supports early-career physician-scientists and dentist-scientists who have recently completed their medical, surgical, or dental training in any specialty and are launching careers in clinical aging research in that specialty. The program builds on the success of two programs: the T. Franklin Williams Scholar program, previously funded by the Atlantic Philanthropies; and the Dennis W. Jahnigen Scholar program, funded by the John A. Hartford Foundation. On March 26 to 27, 2018, the American Geriatrics Society convened a conference in Bethesda, MD, to explore cognitive impairment across the subspecialties. This was the third of three conferences, supported by a U13 grant from the National Institute on Aging (NIA), to aid GEMSSTAR awardees in integrating geriatrics into their subspecialties. Drs Molly Wagster, of the NIA, Sanjay Asthana, of the University of Wisconsin-Madison School of Medicine and Public Health, and Joe Verghese, of the Albert Einstein College of

**Table 1. Represented Specialties**

Anesthesiology	Internal Medicine
Bioethics	Neurology
Cardiology	Oncology
Critical Care	Ophthalmology
Emergency Medicine	Otolaryngology
Family Medicine	Psychology
Geriatric Medicine	

Medicine, served as co-chairs for the meeting. Scientific sessions focused on the impact of cognitive impairment, sensory contributors, comorbidities, links between delirium and dementia, and issues of informed consent in cognitively impaired populations.

As with the previous two conferences in this series, this conference brought together representatives of several subspecialties (Table 1). It aimed to introduce GEMSSTAR awardees to key concepts and approaches to consider and to help them build collaborative networks as they embark on their careers in aging research. This article provides a high-level overview of the full conference report, "The Impact of Cognitive Impairment Across Specialties," ([https://www.americangeriatrics.org/sites/default/files/inline-files/AGS\\_GEMSSTAR\\_Conference\\_Report\\_Cognitive\\_Impairment\\_2.pdf](https://www.americangeriatrics.org/sites/default/files/inline-files/AGS_GEMSSTAR_Conference_Report_Cognitive_Impairment_2.pdf)) and notes opportunities for GEMSSTAR awardees to contribute to research progress in this area.

## IMPACT OF COGNITIVE IMPAIRMENT ACROSS SPECIALTIES

The conference included a specific focus on the important interrelationship and clinical coexistence between delirium and dementia. Despite exclusive diagnostic criteria that can be distinguished by onset, duration, attention, consciousness, speech, and psychomotor subtypes,<sup>13</sup> one systematic review has found that the presence of dementia at baseline is a strong risk factor for the development of incident delirium and that delirium is a risk factor for subsequent dementia.<sup>14</sup> Another systematic review has shown an association between delirium and increased institutionalization and mortality,<sup>15</sup> and clinical studies have observed a link between delirium and long-term cognitive decline.<sup>16-18</sup> Thus, the prevention of delirium may offer the unprecedented opportunity to prevent or ameliorate future cognitive decline.

The *Diagnostic and Statistical Manual of Mental Disorders (DSM)* definition of delirium is difficult to apply at the bedside, and how to prevent and treat complicated delirium is not clear. Who is at risk, the causes and mechanisms, and the relationships between vulnerability and precipitating factors for complicated delirium, which leads to long-term cognitive decline, are areas of ongoing research. Mechanistic studies in animal models suggest links between anesthesia and surgery and molecular processes associated with AD,<sup>19,20</sup> and changes in AD-associated biomarkers have been associated with increased risk for delirium and postoperative cognitive dysfunction.<sup>21-23</sup> However, whether anesthesia and surgery increases the risk for AD is controversial. A 20-item diagnostic assessment that is based on the Confusion Assessment Method and can be completed in

3 minutes has been found to identify patients with delirium, with a sensitivity of 93% and a specificity of 96% among normal patients and those with mild cognitive impairment and a sensitivity of 96% and a specificity of 86% among patients with dementia<sup>24</sup>; some items from this assessment can be used as brief screeners.<sup>25</sup>

Other presentations at the conference discussed several anatomical, neuroimaging, and cerebrospinal fluid biomarkers of AD and noted that changes in some of these biomarkers precede the onset of AD symptoms by at least 20 years.<sup>26</sup> As suggested by a recently published research framework,<sup>27</sup> the clinical diagnosis of AD will likely be confirmed through measures of amyloid and tau deposition and neurodegeneration. However, this paradigm-shifting framework will have to be confirmed through large, prospective, clinical studies before it is adopted for AD diagnosis in clinical practice.

Yet, as one speaker noted, the brain is not an island. Because this conference focused on the impact of cognitive impairment across the medical specialties, an overarching theme was the complex, bidirectional relationship between cognition and other organ systems. A large proportion of intensive care unit survivors experience some degree of cognitive impairment on hospital discharge, and some experience measurable impairment for years afterward.<sup>28</sup> These effects are particularly pronounced among patients aged 65 years and older.<sup>29</sup> Likewise, strong evidence from longitudinal studies indicates associations between sensory impairments, such as olfactory impairment and hearing loss, and cognitive decline and dementia. In fact, a 2017 report from the Lancet Commission concluded that hearing loss was the single modifiable risk factor for dementia, accounting for the greatest proportion of attributable risk compared with all other known modifiable risk factors.<sup>30</sup> Whereas the link between olfactory impairment and cognitive impairment or dementia is likely mediated by a common cause, hearing loss is likely an etiologic contributor to cognitive impairment and dementia. Although some evidence from epidemiological studies suggests an association between visual impairment and cognitive decline and dementia,<sup>31-33</sup> the overall evidence of associations between visual impairments and cognitive decline and dementia is limited and sometimes conflicting.<sup>34-37</sup>

Cognitive impairment can serve as a biomarker of or influence the course of various diseases and conditions. For example, a single-center study in patients with heart failure found that cognitive impairment, as measured by performance on the Mini-Cog, was the strongest independent predictor of a composite outcome of readmission and mortality.<sup>38,39</sup> At the same time, medical comorbidities can affect cognitive health and brain aging. Evidence also suggests that cognitive impairment and a medical condition can have additive or synergistic effects on function. For example, one study has found that the risk for disability is higher with worsening vision loss or cognitive impairment, but that the combination of vision loss and cognitive impairment is associated with even higher risk for disability.<sup>40</sup> Thus, cognitive impairment can, therefore, be considered as a risk factor or outcome measure for disease. Cognitive impairment may also be a mediator of outcomes, as many aspects of managing medical morbidities, such as taking medications, driving to the physician's office, and

managing diet, depend on cognition. Although it is becoming evident that damage to any organ system affects the brain, the mechanisms underlying this link are not clear. As suggested by one pathoetiologic model of delirium,<sup>41</sup> it is likely that mechanisms differ by organ systems.

## CHALLENGES AND STRATEGIES IN RESEARCH ON COGNITIVE IMPAIRMENT

As noted by conference participants, a major barrier to such research is the exclusion of cognitively impaired patients from studies because of their inability to consent, adhere to protocols, and provide longitudinal outcomes. However, several studies among older adults indicate a high willingness to participate in research in the event that they become impaired and unable to give consent, depending on the risk-benefit profile.<sup>42-46</sup> Safeguards that can protect against unwanted participation include enrollment of impaired individuals only when the research cannot be done as well with individuals who can give consent, enrollment of impaired individuals in research that poses minimal risk or offers a benefit that justifies the risk, the designation of a surrogate decision maker who can decide based on substituted judgment or the best interests of a participant, and obtaining the impaired patient's assent. Beyond the designation of a legally authorized representative, however, there are no specific guidelines in the federal regulations with respect to these safeguards.

One institutional review board (IRB) suggested that, rather than exclude individuals with dementia, investigators should screen for decisional capacity, with more rigorous and detailed capacity assessments for studies with greater risk, and request an IRB-appointed proxy for those unable to consent.<sup>47</sup> Decision-making capacity includes the ability to communicate a choice, comparative and consequential reasoning, understanding, and appreciation.<sup>48</sup> Although existing assessments of decision-making abilities often ask individuals to make a choice and why they made that choice (eg, their reasoning), they seldom ask about the individuals' understanding or appreciation of a decision. Moreover, tests such as the Mini-Mental State Examination (MMSE), which is copyright protected, do not offer an adequate assessment of decision-making capacity. How best to assess decisional capacity among patients in the "gray zone," where it is not clear whether an individual is able to give consent, is not clear.

Conference speakers and participants therefore suggested that investigators tailor safeguards to the types of studies they propose to do. Intervention protocols and materials should be adapted for the cognitively impaired (eg, by incorporating alternative outcomes that have been adapted for the this population).<sup>47</sup> Corrective feedback and teach back can be used to ensure participants' understanding. Staff who will obtain consent can be trained to conduct assessments of potential study participants' capacity to give consent. Memory aids can be used to emphasize the most important information participants should know about the research being conducted. In one study among patients with AD and an MMSE score of 20 or higher, those who received a one-page summary of key elements at a sixth-grade reading level performed better on understanding and appreciation and were more likely to be judged capable of

providing informed consent, compared with those who received a standard consent alone.<sup>49</sup>

### THE FUTURE OF AGING RESEARCH: HOW GEMSSTAR SCHOLARS CAN CONTRIBUTE TO PROGRESS


The aging of the US population and the growing burden of dementia make this an area of critical research focus at the National Institutes of Health (NIH). Cognitive impairment and dementia are themselves complex and multifactorial, and as highlighted during this conference, the relationship between cognition and other organ systems is complex and bidirectional. Thus, research exploring mechanisms, methods of identification, prevention, and treatment of cognitive impairment, particularly delirium and dementia, is highly relevant across the medical subspecialties. Likewise, continued research is needed to develop and optimize comprehensive care models focusing on patients with dementia and their caregivers.

Conference participants noted that understanding and addressing cognitive health and its relationship with the health of other organ systems will require multidisciplinary team science. Such teams can approach this complexity by exploring a framework, such as the multihit model, which describes the theoretical effects of comorbidities on brain structure and function (Figure 1). Or these teams can explore a clinical and scientific problem as a continuum based on an underlying biologic mechanism. Such an approach could draw research questions from problems seen in the clinic and focus on patient-, disease-, care setting-, institution-, and environment-level factors simultaneously. Multidisciplinary team science is the best approach to complex, multifaceted conditions, such as cognitive impairment and dementia. Bringing together the right set of collaborators, including individuals on the front lines of the clinical research problem, is important in determining which questions are the most important to study. At

Washington University (St Louis, MO), teams are formed when someone has an idea and the requisite expertise is assembled around that idea. Generating questions from problems seen in clinical care can also add value. In addition, paradigms are shifting, and team science is increasingly recognized as a factor in promotions. Team science also can drive individual members' research into previously unanticipated directions. Thus, it possible to work in multidisciplinary teams and still maintain independence.

Specific research questions discussed at the conference are listed in Table 2. With shifting research, promotion, and tenure paradigms, increasing recognition of the value of team science, and the ability of team science to drive members' individual research projects in unanticipated directions, investigators can participate in multidisciplinary teams and still maintain independence. Thus, GEMSSTAR awardees were encouraged to seek out prospective team members to discuss ideas.

Because the population of older adults and those at risk for cognitive impairment is becoming more diverse, researchers should think proactively of how to develop research questions and study designs that will generate knowledge applicable to a wide range of individuals older than 65 years and the experiences they have. Although racial/ethnic gaps in life expectancy have improved from a 13-year gap between white and black individuals in 1900 to a 3-year gap in the present, full equity has not been reached. In addition, the US population is becoming more diverse. Frameworks, such as the Centers for Disease Control and Prevention's Health Impact Pyramid and the NIA Health Disparities Research Framework, present fundamental factors and levels of analyses to consider when including diverse populations in research. Other mechanisms have also been proposed to explain how sociocultural factors influence behavior and biology.<sup>50,51</sup> The NGAGE model<sup>52</sup> provides a systematic approach to engaging diverse populations in research: Networking (eg, by attending community boards and one-on-one leader meetings); Giving



**Figure 1.** Multiple comorbidities affect the cognitive function of the aging brain. Cognitive decline is caused by neurodegenerative diseases, such as Alzheimer disease, but other factors that may affect cognitive function in the patient with comorbidities include depression, anxiety, impairments in vision or hearing, medications, cerebrovascular disease, and pain. Adapted from Lin FR, Albert M. Hearing loss and dementia – who's listening? *Aging Ment Health*. 2014;18(6):671-673.

**Table 2. Topics that Nongeriatric Subspecialty Researchers Can Explore in Aging Research**

Topic	Future contributions from specialty clinician educators
The bidirectional relationship between cognitive health and the health of other organ systems	<ul style="list-style-type: none"> <li>• Studies of the role of sensory function as biomarkers of vs contributors to cognitive impairment</li> <li>• Studies of possible synergistic effects between sensory impairments and established risk factors for brain aging, cognitive impairment, and dementia</li> <li>• Development and standardization of neurocognitive testing protocols that account for sensory impairments</li> <li>• Knowledge of how post-ICU trajectories can be modified</li> <li>• Knowledge of how post-ICU and postoperative cognitive dysfunction or delirium-associated cognitive decline can be distinguished from trajectories associated with healthy aging</li> <li>• Identification of objective risk factors for post-ICU impairment</li> </ul>
Delirium and dementia	<ul style="list-style-type: none"> <li>• The role of inflammation and neuroinflammation in delirium and dementia</li> <li>• The magnitude by which blood-brain barrier dysfunction in older adults is accelerated after surgery</li> <li>• How disturbances in circadian rhythms mediate delirium and sundowning</li> </ul>

Abbreviations: ICU, intensive care unit.

**Table 3. Examples of Data Sets for Dementia Researchers**

Public data sets	Data sets used in psychometrics conferences
Health and Retirement Study ( <a href="https://hrs.isr.umich.edu/data-products">https://hrs.isr.umich.edu/data-products</a> ) National AD Coordinating Center ( <a href="https://www.alz.washington.edu">https://www.alz.washington.edu</a> ) AD Neuroimaging Initiative ( <a href="http://adni.loni.usc.edu/data-samples/access-data">http://adni.loni.usc.edu/data-samples/access-data</a> )	Rush Religious Orders Study ( <a href="https://www.maelstrom-research.org/mica/individual-study/ros#/">https://www.maelstrom-research.org/mica/individual-study/ros#/</a> ) and Memory and Aging Project ( <a href="https://knightadrc.wustl.edu/Volunteer/MAP.htm">https://knightadrc.wustl.edu/Volunteer/MAP.htm</a> ) Washington Heights Inwood Columbia Aging Project ( <a href="https://www.maelstrom-research.org/mica/individual-study/whicap#/">https://www.maelstrom-research.org/mica/individual-study/whicap#/</a> ) UC-Davis Diversity Cohort ( <a href="https://health.ucdavis.edu/alzheimers/">https://health.ucdavis.edu/alzheimers/</a> ) Reasons for Geographical and Race Disparities in Stroke ( <a href="https://www.uab.edu/soph/regardsstudy/">https://www.uab.edu/soph/regardsstudy/</a> ) Advanced Cognitive Training for Independent and Vital Elderly ( <a href="https://clinicaltrials.gov/ct2/show/NCT00298558">https://clinicaltrials.gov/ct2/show/NCT00298558</a> ) Framingham Heart Study ( <a href="https://www.framinghamheartstudy.org/">https://www.framinghamheartstudy.org/</a> ) Adult Changes in Thought ( <a href="https://www.maelstrom-research.org/mica/individual-study/act#/">https://www.maelstrom-research.org/mica/individual-study/act#/</a> ) Integrative Analysis of Longitudinal Studies on Aging– ( <a href="https://www.maelstrom-research.org/mica/network/ialsa#/">https://www.maelstrom-research.org/mica/network/ialsa#/</a> )

Abbreviations: AD, Alzheimer disease; UC, University of California.

first, or building trust by listening to what the community needs; Advocating, or describing the proposed research as trust is built; Giving back, or providing study findings and learning for the community to use once the research is done; and Evaluating how well the study team has done in engaging diverse populations over time.

Several resources are available for junior investigators interested in cognitive research. The NIH TOOLBOX for Assessment of Behavioral and Neurological Function is a multidimensional set of brief, well-validated, psychometrically sound measures to assess cognitive, sensory, motor, and emotional function across diverse study designs and settings. By using NIH TOOLBOX measures as an adjunct to their own studies, investigators can facilitate data sharing and comparison of findings across studies and clinical settings. Investigators also can take advantage of existing data sets (Table 3). However, they should be aware of these databases' limitations, particularly with respect to data on cognition, and tailor their research questions to the amount,

type, and quality of available data. Ways to overcome challenges that hamper data sharing, such as difficulties in managing data-use agreements and IRB-approved consent forms, must also be identified.

**ACKNOWLEDGMENTS**

The authors would like to thank Frances McFarland, PhD, for her assistance in the preparation of this manuscript. The authors would also like to thank Sue Ziemann, MD, PhD, Basil Eldadah, MD, PhD, Robin Barr, PhD, Molly Wagster, PhD, and Dave Wendler, PhD, for their support of the conference and manuscript. This report has been reviewed by all workshop speakers.

**Conflicts of Interests:**

1. C.C. was a speaker for Emergency Medical Abstracts and Best Evidence in Emergency Medicine. C.C. also serves as a Board Member for Schwartz-Reisman Emergency Medicine Institute International Advisory Board

Chair, is Deputy Editor-in-Chief of *Academic Emergency Medicine*, and is Associate Editor of *Journal of the American Geriatrics Society*.

2. M.B. receives grant funding from National Institute on Aging (NIA) K76 AG057022 and NIA R03 AG050918, additional funding support from NIA AG028716, and funding from the Alzheimer's Drug Discovery Foundation. M.B. serves as a board member for Early Stage Anesthesiology Scholars and as a liaison from the American Society of Anesthesiology to the American Geriatrics Society.
3. S.I.'s mentoring time was covered in part by grants K07AG041835, R24AG054259, and 2P01AG031720 from the NIA.
4. J.K. has received funding from the National Institutes of Health (NIH) and foundations and industry for sponsored projects. J.K. was a consultant for Squintmetrics, under \$5000 in the past 12 months and coming 12 months. J.K. is a board member of the Greenwall Foundation.
5. J.M. is currently participating in clinical trials of antimentia drugs from Eli Lilly and Company and Biogen. J.M. receives research support from Eli Lilly/Avid Radiopharmaceuticals and is funded by NIH grants P50AG005681, P01AG003991, P01AG026276, and UF01AG032438.
6. R.S. reports grants for clinical research from NIH, the Centers for Medicare and Medicaid Services, the Department of Defense, and the Illinois Department of Public Health; being a noncompensated board member of the Alzheimer's Association, Illinois Chapter; and being the site principal investigator or subinvestigator for clinical trials for which his institution (Rush University Medical Center) is compensated (Amylyx Pharmaceuticals, Inc, Eli Lilly & Co, Inc, Genentech, Inc, Merck & Co, Inc, Navidea Biopharmaceuticals, Novartis Pharmaceuticals, Inc, Roche Holdings AG, and Takeda Development Center Americas, Inc).
7. S.A. received grants from NIA/NIH to conduct Alzheimer disease (AD) research. S.A. received grants from Merck Pharmaceutical, Eisai, Toyama Chemical, and Lundbeck to serve as a site principal investigator for clinical trials involving patients with mild cognitive impairment and AD. Please note these grants were awarded to University of Wisconsin–Madison. S.A. received royalties from McGraw Hill Education as an Associate Editor for the textbook entitled *Hazzard's Geriatric Medicine and Gerontology*.
8. J.V. received grant funding from NIH. J.V. participated in a consultancy for Saint Care Corporation, Japan.
9. Authors F.M., M.A., F.L., E.M., D.R., and H.W. had no conflicts of interest to disclose.

**Author Contributions:** Authors C.C., F.M., M.A., M.B., S.I., J.K., F.L., E.M., J.M., H.W., and S.A. worked on concept and design of the manuscript. Authors C.C., F.M., M.B., S.I., J.K., J.M., and H.W. worked on analysis and interpretation of data of the manuscript. Authors C.C., F.M., M.B., S.I., J.K., F.L., E.M., D.R., R.S., H.W., S.A., and J.V. worked on the preparation of the manuscript.

**Sponsor's Role:** Funding for this conference was made possible, in part, by 5 U13 AG 048721-04 from the National Institute on Aging.

The views expressed in written conference materials or publications and by speakers and moderators do not necessarily reflect the official policies of the Department of Health and Human Services; nor does mention of trade names, commercial practices, or organizations imply endorsement by the US Government.

**Dedication:** We humbly dedicate this article to Arti Hurria, Grants for Early Medical/Surgical Specialists' Transition to Aging Research U13 principal investigator, friend, and hero, who lost her life on November 7, 2018. Dr Hurria was a tireless champion advancing geriatrics concepts across medical specialties. Her exceptional leadership was the driving force behind the successful U13 conferences.

## REFERENCES

1. Price JL, Ko AI, Wade MJ, Tsou SK, McKeel DW, Morris JC. Neuron number in the entorhinal cortex and CA1 in preclinical Alzheimer disease. *Arch Neurol.* 2001;58(9):1395-1402.
2. Boyle PA, Yu L, Wilson RS, Leurgans SE, Schneider JA, Bennett DA. Person-specific contribution of neuropathologies to cognitive loss in old age. *Ann Neurol.* 2018;83(1):74-83.
3. Schindler SE, Yan L, Todd KW, et al. Emerging cerebrospinal fluid biomarkers in autosomal dominant Alzheimer's disease. *Alzheimers Dement.* 2019;15(5):655-665.
4. Morris JC, Schindler SE, McCue LM, et al. Assessment of racial disparities in biomarkers for Alzheimer disease. *JAMA Neurol.* 2019;76(3):264-273.
5. Storandt M, Morris JC. Ascertainment bias in the clinical diagnosis of Alzheimer disease. *Arch Neurol.* 2010;67(11):1364-1369.
6. Buckley JS, Salpeter SR. A risk-benefit assessment of dementia medications: systematic review of the evidence. *Drugs Aging.* 2015;32(6):453-467.
7. Bass DM, Judge KS, Snow AL, et al. Caregiver outcomes of partners in dementia care: effect of a care coordination program for veterans with dementia and their family members and friends. *J Am Geriatr Soc.* 2013;61(8):1377-1386.
8. Samus QM, Johnston D, Black BS, et al. A multidimensional home-based care coordination intervention for elders with memory disorders: the maximizing independence at home (MIND) pilot randomized trial. *Am J Geriatr Psychiatry.* 2014;22(4):398-414.
9. Boustani MA, Sachs GA, Alder CA, et al. Implementing innovative models of dementia care: the Healthy Aging Brain Center. *Aging Ment Health.* 2011;15(1):13-22.
10. Reuben DB, Everson LC, Wenger NS, et al. The University of California at Los Angeles Alzheimer's and Dementia Care Program for comprehensive, coordinated, patient-centered care: preliminary data. *J Am Geriatr Soc.* 2013;61(12):2214-2218.
11. Bass DM, Judge KS, Snow AL, et al. A controlled trial of partners in dementia care: veteran outcomes after six and twelve months. *Alzheimers Res Ther.* 2014;6(1):9.
12. Tanner JA, Black BS, Johnston D, et al. A randomized controlled trial of a community-based dementia care coordination intervention: effects of MIND at home on caregiver outcomes. *Am J Geriatr Psychiatry.* 2015;23(4):391-402.
13. Oh ES, Fong TG, Hshieh TT, Inouye SK. Delirium in older persons: advances in diagnosis and treatment. *JAMA.* 2017;318(12):1161-1174.
14. Fong TG, Davis D, Growdon ME, Albuquerque A, Inouye SK. The interface between delirium and dementia in elderly adults. *Lancet Neurol.* 2015;14(8):823-832.
15. Witlox J, Eurelings LS, de Jonghe JF, Kalisvaart KJ, Eikelenboom P, van Gool WA. Delirium in elderly patients and the risk of postdischarge mortality, institutionalization, and dementia: a meta-analysis. *JAMA.* 2010;304(4):443-451.
16. Saczynski JS, Marcantonio ER, Quach L, et al. Cognitive trajectories after postoperative delirium. *N Engl J Med.* 2012;367(1):30-39.
17. Inouye SK, Marcantonio ER, Kosar CM, et al. The short-term and long-term relationship between delirium and cognitive trajectory in older surgical patients. *Alzheimers Dement.* 2016;12(7):766-775.
18. Vasunilashorn SM, Fong TG, Albuquerque A, et al. Delirium severity post-surgery and its relationship with long-term cognitive decline in a cohort of patients without dementia. *J Alzheimers Dis.* 2018;61(1):347-358.
19. Berger M, Nadler JW, Friedman A, et al. The effect of propofol versus isoflurane anesthesia on human cerebrospinal fluid markers of Alzheimer's disease: results of a randomized trial. *J Alzheimers Dis.* 2016;52(4):1299-1310.

20. Berger M, Burke J, Eckenhoff R, Mathew J. Alzheimer's disease, anesthesia, and surgery: a clinically focused review. *J Cardiothorac Vasc Anesth*. 2014;28(6):1609-1623.
21. Evered L, Silbert B, Scott DA, Ames D, Maruff P, Blennow K. Cerebrospinal fluid biomarker for Alzheimer disease predicts postoperative cognitive dysfunction. *Anesthesiology*. 2016;124(2):353-361.
22. Xie Z, McAuliffe S, Swain CA, et al. Cerebrospinal fluid  $A\beta$  to tau ratio and postoperative cognitive change. *Ann Surg*. 2013;258(2):364-369.
23. Xie Z, Swain CA, Ward SA, et al. Preoperative cerebrospinal fluid beta-amyloid/tau ratio and postoperative delirium. *Ann Clin Transl Neurol*. 2014;1(5):319-328.
24. Marcantonio ER, Ngo LH, O'Connor M, et al. 3D-CAM: derivation and validation of a 3-minute diagnostic interview for CAM-defined delirium: a cross-sectional diagnostic test study. *Ann Intern Med*. 2014;161(8):554-561.
25. Fick DM, Inouye SK, Guess J, et al. Preliminary development of an ultrabrief two-item bedside test for delirium. *J Hosp Med*. 2015;10(10):645-650.
26. Bateman RJ, Xiong C, Benzinger TL, et al. Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N Engl J Med*. 2012;367(9):795-804.
27. Jack CR Jr, Bennett DA, Blennow K, et al. NIA-AA research framework: toward a biological definition of Alzheimer's disease. *Alzheimers Dement*. 2018;14(4):535-562.
28. Wilcox ME, Lim AS, McAndrews MP, et al. A study protocol for an observational cohort investigating COGNitive outcomes and WELLness in survivors of critical illness: the COGWELL study. *BMJ Open*. 2017;7(7):e015600.
29. Pandharipande PP, Girard TD, Jackson JC, et al. Long-term cognitive impairment after critical illness. *N Engl J Med*. 2013;369(14):1306-1316.
30. Livingston G, Sommerlad A, Orgeta V, et al. Dementia prevention, intervention, and care. *Lancet*. 2017;390(10113):2673-2734.
31. Chen SP, Bhattacharya J, Pershing S. Association of vision loss with cognition in older adults. *JAMA Ophthalmol*. 2017;135(9):963-970.
32. Tay T, Wang JJ, Kifley A, Lindley R, Newall P, Mitchell P. Sensory and cognitive association in older persons: findings from an older Australian population. *Gerontology*. 2006;52(6):386-394.
33. Lin MY, Gutierrez PR, Stone KL, et al. Vision impairment and combined vision and hearing impairment predict cognitive and functional decline in older women. *J Am Geriatr Soc*. 2004;52(12):1996-2002.
34. Keenan TD, Goldacre R, Goldacre MJ. Associations between age-related macular degeneration, Alzheimer disease, and dementia: record linkage study of hospital admissions. *JAMA Ophthalmol*. 2014;132(1):63-68.
35. Klaver CC, Ott A, Hofman A, Assink JJ, Breteler MM, de Jong PT. Is age-related maculopathy associated with Alzheimer's disease? the Rotterdam study. *Am J Epidemiol*. 1999;150(9):963-968.
36. Reyes-Ortiz CA, Kuo YF, DiNuzzo AR, Ray LA, Raji MA, Markides KS. Near vision impairment predicts cognitive decline: data from the Hispanic Established Populations for Epidemiologic Studies of the Elderly. *J Am Geriatr Soc*. 2005;53(4):681-686.
37. Baker ML, Wang JJ, Rogers S, et al. Early age-related macular degeneration, cognitive function, and dementia: the Cardiovascular Health Study. *Arch Ophthalmol*. 2009;127(5):667-673.
38. Patel A, Parikh R, Howell EH, Hsich E, Landers SH, Gorodeski EZ. Mini-cog performance: novel marker of post discharge risk among patients hospitalized for heart failure. *Circ Heart Fail*. 2015;8(1):8-16.
39. Walsh MN. Assessment of cognitive impairment: the holy grail of risk prediction? *Circ Heart Fail*. 2015;8(1):2-4.
40. Whitson HE, Cousins SW, Burchett BM, Hybels CF, Pieper CF, Cohen HJ. The combined effect of visual impairment and cognitive impairment on disability in older people. *J Am Geriatr Soc*. 2007;55(6):885-891.
41. Maldonado JR. Pathoetiological model of delirium: a comprehensive understanding of the neurobiology of delirium and an evidence-based approach to prevention and treatment. *Crit Care Clin*. 2008;24(4):789-856. ix.
42. Wendler D, Martinez RA, Fairclough D, Sunderland T, Emanuel E. Views of potential subjects toward proposed regulations for clinical research with adults unable to consent. *Am J Psychiatry*. 2002;159(4):585-591.
43. Muthappan P, Forster H, Wendler D. Research advance directives: protection or obstacle? *Am J Psychiatry*. 2005;162(12):2389-2391.
44. Kim SY, Kim HM, Langa KM, Karlawish JH, Knopman DS, Appelbaum PS. Surrogate consent for dementia research: a national survey of older Americans. *Neurology*. 2009;72(2):149-155.
45. De Vries R, Ryan KA, Stanczyk A, et al. Public's approach to surrogate consent for dementia research: cautious pragmatism. *Am J Geriatr Psychiatry*. 2013;21(4):364-372.
46. Abdoler E, Wendler D. Using data to improve surrogate consent for clinical research with incapacitated adults. *J Empir Res Hum Res Ethics*. 2012;7(2):37-50.
47. Prusaczyk B, Cherney SM, Carpenter CR, DuBois JM. Informed consent to research with cognitively impaired adults: transdisciplinary challenges and opportunities. *Clin Gerontol*. 2017;40(1):63-73.
48. Appelbaum PS. Clinical practice: assessment of patients' competence to consent to treatment. *N Engl J Med*. 2007;357(18):1834-1840.
49. Rubright J, Sankar P, Casarett DJ, Gur R, Xie SX, Karlawish J. A memory and organizational aid improves Alzheimer disease research consent capacity: results of a randomized, controlled trial. *Am J Geriatr Psychiatry*. 2010;18(12):1124-1132.
50. Adler NE, Stewart J. Health disparities across the lifespan: meaning, methods, and mechanisms. *Ann N Y Acad Sci*. 2010;1186:5-23.
51. Adler NE, Stewart J. Preface to the biology of disadvantage: socioeconomic status and health. *Ann N Y Acad Sci*. 2010;1186:1-4.
52. Barnes LL, Shah RC, Aggarwal NT, Bennett DA, Schneider JA. The Minority Aging Research Study: ongoing efforts to obtain brain donation in African Americans without dementia. *Curr Alzheimer Res*. 2012;9(6):734-745.