UC Davis

UC Davis Previously Published Works

Title

A Promising Treatment for Osteoarthritis?

Permalink

<https://escholarship.org/uc/item/0855p9ch>

Journal

Annals of internal medicine, 173(7)

ISSN

0003-4819

Authors

Lane, Nancy
Felson, David

Publication Date

2020-10-01

DOI

10.7326/m20-4938

Peer reviewed

A Promising Treatment for Osteoarthritis?

Osteoarthritis, the most common type of arthritis, increases in prevalence with age and is a leading cause of joint pain and disability worldwide (1). Currently, no treatments exist that can either prevent or slow progression of the disease. Osteoarthritis is characterized by loss of articular cartilage, bone remodeling, and synovial inflammation. Studies suggest that both inflammation and abnormal joint loading trigger local release of inflammatory cytokines, including interleukin-1 β (IL-1 β), IL-6, and tumor necrosis factor. These inflammatory cytokines alter the chromatin landscape and activate the genes that transcribe proteins catabolically to cartilage and bone, including matrix metalloproteinases, ADAMTS5, and MAP kinases. Also, inflammatory cytokines can induce other disease-related genes within the inflamed synovium, including nerve growth factor and prostaglandin E2, key pain sensitizers in osteoarthritis (2).

The role of systemic inflammation in osteoarthritis has been a topic of great interest. The relation of these inflammatory mediators to risk for osteoarthritis may identify patients who would benefit from specific therapies. One such biomarker, C-reactive protein (CRP), reflects systemic and localized inflammation and is induced by inflammatory cytokines in the liver, in inflammatory cells within the synovium, and at other sites of inflammation. High-sensitivity CRP (hs-CRP) is associated with an increased risk for painful osteoarthritis (3, 4). However, the precise role of CRP in osteoarthritis development is unclear. Because CRP production is activated by inflammatory cytokines, including IL-6 and IL-1 β , the serum level of CRP may also reflect inflammation within the joint.

Trials testing IL-1 inhibitors for knee osteoarthritis (Table) have ranged from 4 to 50 weeks in duration and have tested monoclonal antibodies that block IL-1 activity by interfering with either IL-1 or its receptor. All the trials focused on pain outcomes, and although none showed a significant reduction in pain in patients randomly assigned to IL-1 inhibition versus the placebo groups, results in the non-canakinumab studies trended toward modest efficacy, with 5% to 10% lower pain scores on average in the active-treatment versus the placebo groups. No studies tested the effect of an IL-1 inhibitor on radiographic disease progression or cartilage loss. Findings of trials testing inhibitors of tumor necrosis factor- α ,

another inflammatory cytokine, have also been null—although, like the IL-1 studies, most have shown small, non-significant reductions in pain for patients receiving active treatment versus those receiving placebo.

CANTOS (Canakinumab Anti-Inflammatory Thrombosis Outcomes Study) was a trial that tested IL-1 inhibition as a treatment for secondary prevention of cardiovascular disease. It included 10 061 patients with a history of myocardial infarction and an hs-CRP level of 2 mg/L or greater. The trial compared 3 doses (50, 150, and 300 mg) of canakinumab—a monoclonal antibody targeting IL-1 β —or placebo, administered subcutaneously every 3 months for up to 5 years. Patients who received canakinumab had a significantly lower rate of recurrent cardiovascular events and related mortality than those in the placebo group (9). However, patients in the canakinumab groups also had a greater number of serious infections and higher infection-related mortality, which canceled out any overall reduction in mortality; therefore, canakinumab was not recommended for secondary prevention of cardiovascular disease.

As part of CANTOS, data were collected on all adverse events (AEs). Because total hip or total knee replacement (THR/TKR) requires hospitalization and all hospitalized events are reported as serious AEs in a trial, a comprehensive count of THR/TKRs exists. More than 90% of replacements were performed for osteoarthritis. Joint pains and other potential osteoarthritis symptoms were queried as AEs, but information on these outcomes was nonspecific and likely inconsistently collected.

In their article, Schieker and colleagues (10) present a secondary analysis of osteoarthritis data from the CANTOS trial. The primary end point was time to first THR/TKR, with a median follow-up of 3.7 years. The combined incidence rates for THR/TKR were 40% to 47% lower with canakinumab treatment; the hazard ratios were 0.54 (95% CI, 0.36 to 0.81) among men and 0.66 (CI, 0.38 to 1.12) among women. All canakinumab doses reduced the number of joint replacements similarly. The reduction in joint replacements among patients who received canakinumab versus the placebo group became apparent after only 1 year of treatment. The effects of canakinumab remained statistically significant when CANTOS participants with a history of crystalline or inflammatory arthritis were excluded. How-

Table. Randomized Placebo-Controlled Trials of IL-1 Inhibitors for Knee Osteoarthritis

Study, Year (Reference)	IL-1 Treatment	Patients, <i>n</i>	Outcome	Duration, <i>wk</i>	Results
Chevalier et al, 2009 (5)	Anakinra (IL-1RA)*	170	Pain	4	Null
Fleischmann et al, 2019 (6)	Antibody to IL-1 α/β	350	Pain/synovitis	50	Null for both outcomes
Cohen et al, 2011 (7)	mAb to IL-1R1	160	Pain	12	Null
NCT01160822 (8)	Canakinumab (mAb to IL-1 β)*	136	Pain	4	Null

IL-1 = interleukin-1; IL-1R1 = IL-1 receptor, type 1; IL-1RA = IL-1 receptor antagonist; mAb = monoclonal antibody.

* Intra-articular administration of treatment and placebo.

ever, the number of women in this trial was small, which is a concern given that knee osteoarthritis predominantly affects older women.

The results of this exploratory trial are both unexpected and exciting, and novel aspects of the trial design deserve comment. The investigators used elevated hs-CRP level as an entry criterion. This requirement, not routine in osteoarthritis trials, may have identified a subgroup of persons with osteoarthritis in whom inflammatory cytokines activate pathways that accelerate joint degeneration.

Most osteoarthritis trials, even those focusing on radiographic osteoarthritis progression, are of much shorter duration than this study. The primary study end point was time to joint replacement. Total hip or total knee replacement is performed in patients who have unacceptably high levels of joint pain or functional impairment despite medical and rehabilitative treatment, and they usually have advanced radiographic deterioration in the affected joint. Joint replacement is a robust end point and one that might be adapted by the U.S. Food and Drug Administration for evaluating disease-modifying osteoarthritis drugs.

On the basis of our understanding of osteoarthritis pathophysiology, IL-1 is a central player in both cartilage degradation and the enhancement of joint pain. This secondary analysis of the CANTOS trial presents intriguing results that may offer promise for IL-1 inhibition as a treatment. Further studies should evaluate the importance of elevated CRP levels as a factor affecting response to treatment, include more women to better reflect the osteoarthritis population, explore how to minimize infections, and try to better define the duration of therapy needed to detect treatment effects. This unexpected finding deserves additional investigation in developing potential disease-modifying osteoarthritis drugs.

Nancy Lane, MD
UC Davis Health
Sacramento, California

David Felson, MD, MPH
Boston University School of Medicine
Boston, Massachusetts

Grant Support: By National Institutes of Health grants AR072571 and P50AR063043.

Disclosures: Disclosures can be viewed at www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum=M20-4938.

Corresponding Author: Nancy Lane, MD, Department of Internal Medicine and Division of Rheumatology, UC Davis Health, 4625 Second Avenue, Sacramento, CA 95817; e-mail, nelane@ucdavis.edu.

Current author addresses are available at Annals.org.

Ann Intern Med. 2020;173:580-581. doi:10.7326/M20-4938

References

1. Neogi T. The epidemiology and impact of pain in osteoarthritis. *Osteoarthritis Cartilage*. 2013;21:1145-53. [PMID: 23973124] doi:10.1016/j.joca.2013.03.018
2. Vincent TL. IL-1 in osteoarthritis: time for a critical review of the literature. *F1000Res*. 2019;8. [PMID: 31249675] doi:10.12688/f1000research.18831.1
3. Jin X, Beguerie JR, Zhang W, et al. Circulating C reactive protein in osteoarthritis: a systematic review and meta-analysis. *Ann Rheum Dis*. 2015;74:703-10. [PMID: 24363360] doi:10.1136/annrheumdis-2013-204494
4. Pearle AD, Scanzello CR, George S, et al. Elevated high-sensitivity C-reactive protein levels are associated with local inflammatory findings in patients with osteoarthritis. *Osteoarthritis Cartilage*. 2007;15:516-23. [PMID: 17157039]
5. Chevalier X, Goupille P, Beaulieu AD, et al. Intraarticular injection of anakinra in osteoarthritis of the knee: a multicenter, randomized, double-blind, placebo-controlled study. *Arthritis Rheum*. 2009;61:344-52. [PMID: 19248129] doi:10.1002/art.24096
6. Fleischmann RM, Bliddal H, Blanco FJ, et al. A phase II trial of lutikizumab, an anti-interleukin-1 α/β dual variable domain immunoglobulin, in knee osteoarthritis patients with synovitis. *Arthritis Rheumatol*. 2019;71:1056-1069. [PMID: 30653843] doi:10.1002/art.40840
7. Cohen SB, Proudman S, Kivitz AJ, et al. A randomized, double-blind study of AMG 108 (a fully human monoclonal antibody to IL-1R1) in patients with osteoarthritis of the knee. *Arthritis Res Ther*. 2011;13:R125. [PMID: 21801403] doi:10.1186/ar3430
8. U.S. National Library of Medicine. To Determine the Safety, Tolerability, Pharmacokinetics and Effect on Pain of a Single Intra-articular Administration of Canakinumab in Patients With Osteoarthritis in the Knee. Accessed at <https://clinicaltrials.gov/ct2/show/NCT01160822> on 29 June 2020.
9. Ridker PM, Everett BM, Thuren T, et al; CANTOS Trial Group. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med*. 2017;377:1119-1131. [PMID: 28845751] doi:10.1056/NEJMoa1707914
10. Schieker M, Conaghan PG, Mindeholm L, et al. Effects of interleukin-1 β inhibition on incident hip and knee replacement. Exploratory analyses from a randomized, double-blind, placebo-controlled trial. *Ann Intern Med*. 2020;173:509-15. [PMID: 32744862] doi:10.7326/M20-0527

Current Author Addresses: Dr. Lane: Department of Internal Medicine and Division of Rheumatology, UC Davis Health, 4625 Second Avenue, Sacramento, CA 95817.
Dr. Felson: Evans Biomedical Research Center, 650 Albany Street, Boston, MA 02118.