

# Lawrence Berkeley National Laboratory

## Environ Genomics & Systems Bio

### Title

Peatland microbial community responses to plant functional group and drought are depth-dependent

### Permalink

<https://escholarship.org/uc/item/08d147rh>

### Journal

Molecular Ecology, 30(20)

### ISSN

0962-1083

### Authors

Lamit, Louis J  
Romanowicz, Karl J  
Potvin, Lynette R  
[et al.](#)

### Publication Date

2021-10-01

### DOI

10.1111/mec.16125

Peer reviewed

1 **Peatland microbial community responses to plant functional group and drought are depth-**  
2 **dependent**

3

4 **Running title:** Peat microbial communities and global change

5

6 Louis J. Lamit<sup>1,\*</sup>, Karl J. Romanowicz<sup>1,\*</sup>, Lynette R. Potvin<sup>2,\*</sup>, Jay T. Lennon<sup>3</sup>, Susannah G.

7 Tringe<sup>4</sup>, Rodney A. Chimner<sup>1</sup>, Randall K. Kolka<sup>5</sup>, Evan S. Kane<sup>1,2</sup>, Erik A. Lilleskov<sup>2</sup>

8

9 <sup>1</sup>College of Forest Resources and Environmental Science, Michigan Technological University,  
10 Houghton, MI 49931, USA.

11 <sup>2</sup>USDA Forest Service Northern Research Station, Houghton, MI 49931, USA.

12 <sup>3</sup>Department of Biology, Indiana University, Bloomington, IN 47405, USA.

13 <sup>4</sup>DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA 94720,  
14 USA.

15 <sup>5</sup>USDA Forest Service Northern Research Station, Grand Rapids, MN 55744, USA

16 \*Current Addresses. Louis J. Lamit: Department of Biology, Syracuse University, Syracuse, NY  
17 13244, USA, and Department of Environmental and Forest Biology, State University of New  
18 York College of Environmental Science and Forestry, Syracuse, New York 13210, USA.

19 Karl J. Romanowicz: Department of Ecology and Evolutionary Biology, University of  
20 Michigan, Ann Arbor, MI 48109, USA. Lynette R. Potvin: Isle Royale National Park,  
21 Houghton, MI 49931, USA.

22 Corresponding author: LJ Lamit. 315-446-4368, ljlamit@syr.edu, ORCID Identifier 0000-0002-  
23 0385-6010

24 **Abstract** - Peatlands store one-third of Earth's soil carbon, the stability of which is uncertain due  
25 to climate change-driven shifts in hydrology and vegetation, and consequent impacts on  
26 microbial communities that mediate decomposition. Peatland carbon cycling varies over steep  
27 physicochemical gradients characterizing vertical peat profiles. However, it is unclear how  
28 drought-mediated changes in plant functional groups (PFGs) and water table (WT) levels affect  
29 microbial communities at different depths. We combined a multi-year mesocosm experiment  
30 with community sequencing across a 70 cm depth gradient, to test the hypotheses that vascular  
31 PFGs (Ericaceae vs. sedges) and WT (high vs. low) structure peatland microbial communities in  
32 depth-dependent ways. Several key results emerged. 1) Both fungal and prokaryote (bacteria and  
33 archaea) community structure shifted with WT and PFG manipulation, but fungi were much  
34 more sensitive to PFG whereas prokaryotes were much more sensitive to WT. 2) PFG effects  
35 were largely driven by Ericaceae, although sedge effects were evident in specific cases (e.g.,  
36 methanotrophs). 3) Treatment effects varied with depth: the influence of PFG was strongest in  
37 shallow peat (0-10, 10-20 cm), whereas WT effects were strongest at the surface and middle  
38 depths (0-10, 30-40 cm), and all treatment effects waned in the deepest peat (60-70 cm). Our  
39 results underscore the depth-dependent and taxon-specific ways that plant communities and  
40 hydrologic variability shape peatland microbial communities, pointing to the importance of  
41 understanding how these factors integrate across soil profiles when examining peatland  
42 responses to climate change.

43

44 **Keywords** - Ericaceae, microbial community, peatlands, plant functional group, soil depth, water  
45 table

46

47 **Introduction**

48 Hydrology is the main driver of wetland ecosystem structure and function (Mitsch &  
49 Gosselink, 2015), and climate change-driven alterations to hydrology are having extensive  
50 impacts on earth's wetlands (Junk et al., 2013; Moomaw et al., 2018). Because wetlands are  
51 globally important carbon storage reservoirs and methane sources, their responses to climate  
52 change will likely feed back to further modulate climate (Bardgett, Freeman, & Ostle, 2008;  
53 Davidson & Janssens, 2005; Zhang et al., 2017). This is a particularly important issue for  
54 carbon-accumulating wetlands (peatlands) which contain approximately one-third of Earth's soil  
55 carbon, more than twice the carbon stored aboveground in Earth's tropical rain forests (Joosten  
56 & Couwenberg, 2008). Drier conditions in peatlands can alter carbon cycles and expose carbon  
57 formerly sequestered below the water table (WT) to aerobic microbial oxidation (Bragazza,  
58 Parisod, Buttler, & Bardgett, 2013; Bridgham, Pastor, Dewey, Weltzin, & Updegraff, 2008;  
59 Davidson & Janssens, 2005; Freeman, Ostle, & Kang, 2001; Kane et al., 2019).

60 In addition to direct effects on hydrology, climate change can also alter peatland plant  
61 communities, which has extended consequences for carbon cycling (Bragazza et al., 2013;  
62 Dieleman, Branfireun, McLaughlin, & Lindo, 2016; Jassey et al., 2018; Potvin, Kane, Chimner,  
63 Kolka, & Lileskov, 2015). Bogs and poor fens of the northern hemisphere are dominated by  
64 *Sphagnum* mosses whose highly recalcitrant tissues form the bulk of peat-building organic  
65 matter (van Breeman, 1995; Rydin & Jeglum, 2013). Growing in the *Sphagnum* matrix are dwarf  
66 shrubs in the Ericaceae and graminoids in the Cyperaceae (hereafter Sedges), two vascular plant  
67 functional groups (PFGs) with distinctive chemical, morphological and carbon allocation traits  
68 that influence ecosystem processes (Crow & Wieder, 2005; Dorrepaal, Cornelissen, Aerts,  
69 Wallén, & van Logtestijn, 2005; Rydin & Jeglum, 2013; Ward et al. 2015). Ericaceae lack

70 aerenchyma but form adventitious roots as their stems are buried in accumulating peat (Rydin &  
71 Jeglum, 2013). This allows Ericaceae to proliferate in the acrotelm, the frequently oxic, upper  
72 part of peat soil profiles (Moore, Bubier, Froking, Lafleur, & Roulet, 2002; Wallén, 1987). In  
73 contrast, aerenchyma allows sedges to dominate wetter sites, and to extend active roots into the  
74 catotelm, the deeper anoxic peat below the WT (Moore et al., 2002; Rydin & Jeglum, 2013).  
75 Strong evidence supports a shift towards dominance by Ericaceae when peatlands become drier  
76 (Bragazza et al., 2013; Breeuwer et al., 2009; Chimner, Pypker, Hribljan, Moore, & Waddington,  
77 2017; Malhotra et al., 2020; Potvin et al., 2015; Weltzin, Bridgham, Pastor, Chen, & Harth,  
78 2003), a pattern driven in part by the divergent root traits of Ericaceae and sedges.

79         The roots of PFGs interact with peatland microbial communities to further modulate  
80 ecosystem processes. Ericaceae roots form mutualistic symbioses with ericoid mycorrhizal fungi  
81 (ErMF) which have the enzymatic capacity to degrade some forms of complex organic matter to  
82 access immobilized nutrients (Cairney & Burke, 1998; Martino et al., 2018; Read, Leake, &  
83 Perez-Moreno, 2004). Ready access to host photosynthate should make ErMF taxa strong  
84 competitors with free-living saprotrophs (Verbruggen, Pena, Fernandez, & Soong, 2017). The  
85 differential abilities of ErMF vs saprotrophic fungi to degrade organic matter, coupled with the  
86 high phenolic content of ericaceous litter (Bragazza et al., 2013; Dorrepaal et al., 2005), have  
87 consequences for decomposition rates and ecosystem carbon storage (Bragazza et al., 2013;  
88 Orwin, Kirschbaum, St John, & Dickie, 2011; Ward et al., 2015; Verbruggen et al., 2017) that  
89 potentially carryover to affect microbial taxa that utilize byproducts of decomposition. Although  
90 both PFGs are colonized by endophytic fungi with poorly known functions, peatland sedges do  
91 not typically host mycorrhizal fungi (Thormann, Currah, & Bayley, 1999; Weishampel &  
92 Bedford, 2006). Therefore, some of the most important effects of sedges on microbial

93 communities are likely mediated through their influence on free-living taxa. Sedge roots can be  
94 sources of labile carbon that fuel processes such as fermentation and methanogenesis, while  
95 aerenchyma-enabled rhizosphere oxygenation may allow aerobic microorganisms to be active  
96 around roots in water saturated conditions and promote unique biogeochemical processes at the  
97 sharp oxygen concentration gradients associated with rhizospheres (Chanton et al., 2008; Lamers  
98 et al., 2012; Rupp, Kane, Dieleman, Keller, & Turetsky, 2019). A number of studies provide  
99 important experimental evidence that PFGs uniquely influence aspects of peatland microbial  
100 communities or associated carbon cycling processes (e.g., Robroek et al., 2015; Rupp et al.,  
101 2019; Ward et al., 2015). However, there remains a large gap in our understanding of who in the  
102 microbial communities is affected, because tools that provide the resolution necessary to fully  
103 characterize the composition of diverse fungal, bacterial and archaeal communities (e.g., high  
104 throughput amplicon sequencing) have not been extensively applied.

105         Several key points highlight the importance of accounting for WT and sampling depth  
106 when understanding PFG effects on peatland microbial communities. First, there is clear  
107 evidence that microbial communities shift with changes in WT (Emsens et al., 2020; Jassey et  
108 al., 2018; Urbanová & Barta, 2016), however concomitant shifts in vegetation can sometimes  
109 make it difficult to fully decouple WT from PFG effects. Second, microbial communities can  
110 change dramatically with increasing depth in peat profiles, in part a direct result of WTs  
111 excluding obligate aerobes from living in anoxic conditions deep in peat profiles (Artz et al.,  
112 2007; Asemaninejad, Thorn, & Lindo, 2017; Asemaninejad et al., 2019; Andersen, Chapman, &  
113 Artz, 2013; Emsens et al., 2020; Kotiaho et al., 2013; Lamit et al., 2017; Lin et al., 2014). This is  
114 one reason why the abundance of fungi, most of which prefer oxic conditions (Kavanagh, 2011),  
115 drops dramatically with depth in peatlands (Golovchenko, Dobrovol'skaya, & Inisheva, 2002;

116 Lamit et al. 2017; Lin et al. 2014). Third, microbial community depth stratification should also  
117 be a product of the unique effects of plant roots along peat profiles, with different PFGs having  
118 distinct depth effects based on their differences in rooting depth. Taken together, these points  
119 emphasize the need for experimental decoupling of WT and PFG effects on peatland microbial  
120 communities, within the context of depth.

121 We conducted a multi-year mesocosm experiment to examine how microbial  
122 communities are shaped by seasonal drought and contrasting PFGs. We hypothesized, **H1)**  
123 manipulation of PFGs will shift microbial community structure, with these effects being unique  
124 to each PFG and strongly depth-dependent. We specifically predicted that Ericaceae removal  
125 will have its strongest effect in the upper peat profile, while sedge removal should influence  
126 communities along a greater length of the peat profile. Next, we hypothesized, **H2)** WT  
127 manipulation will shift microbial community structure, with these effects also being strongly  
128 depth-dependent. Although WT impacts are likely broad, the most distinct effects of WT should  
129 occur at depths where the WT is the most dynamic and distinct between the treatments, and is  
130 known to manifest a strong influence on peat and porewater chemistry (Kane et al., 2019; Lin et  
131 al., 2014). Given that the depth-dependent effects of each PFG are in part a consequence of the  
132 different ways that sedges and Ericaceae interact with oxic vs anoxic conditions, we further  
133 hypothesized, **H3)** the responses of microbial communities to manipulation of one factor (PFG or  
134 WT) will be dependent on the level of the other factor and these interactive effects will in turn be  
135 dependent on depth in the peat profile. Understanding how climate change mediated changes in  
136 WT and PFGs impact microorganisms along peat depth gradients is important because these are  
137 some of the earth's most taxonomically and functionally diverse groups of organisms and  
138 because their activities influence the most carbon-rich ecosystems on earth, peatlands.

139

## 140 **Materials and Methods**

### 141 *Experimental study system*

142 PEATcosm was a mesocosm experiment designed to test the influence of seasonal  
143 drought and PFG on peatland ecosystems. Detailed descriptions of the experimental design, peat  
144 characteristics, porewater chemistry and vegetation can be found in Kane et al. (2019), Lamit et  
145 al. (2017), and Potvin et al. (2015). The experiment contained twenty-four ~1 m<sup>3</sup> peat monoliths  
146 excavated from an oligotrophic acidic (pH = ~ 4) *Sphagnum* peatland in Minnesota, USA  
147 (N47.07278°, W92.73167°), in May 2010. The monoliths were installed in the Houghton  
148 Mesocosm Facility, USDA Forest Service, Northern Research Station, Forestry Sciences  
149 Laboratory in Houghton, Michigan (N47.11469°, W88.54787°). Monoliths were naturally  
150 vegetated by a continuous layer of *Sphagnum* mosses (primarily *Sphagnum rubellum*, but some  
151 *S. magellanicum* and *S. fuscum*), with *Polytrichum strictum* also present, and a vascular  
152 community of Ericaceae (primarily *Chamaedaphne calyculata*, *Kalmia polifolia*, *Vaccinium*  
153 *oxycoccus*) and sedges (*Carex oligosperma*, *Eriophorum vaginatum*). PFG manipulation was  
154 initiated in June 2011 and included Ericaceae removal, sedge removal and unmanipulated  
155 vegetation treatments (n = 8 per PFG treatment). PFG treatments were carried out by a  
156 combination of gentle removal of target species' stems plus roots when avoidance of moss  
157 damage was possible, followed by clipping all remaining aboveground tissues. PFG treatments  
158 were subsequently maintained by clipping growth of excluded PFGs on a weekly basis, as  
159 needed. WT manipulations were imposed by maintaining 12 mesocosms at average (high WT)  
160 and 12 at summer drought (low WT) conditions (n = 4 replicates per WT x PFG treatment). WT  
161 manipulation was carried out with rain-out shelters, artificial rainwater addition and controlled



162 drainage in the spring and after heavy rains at the acrotelm-catotelm boundary (~25 cm depth).  
163 The depth separation of WT treatments was small in year one (2011), intermediate in year two  
164 (2012), and the greatest in years three and four (2013, 2014) to simulate strong summer drought  
165 (Fig S1).

166

### 167 *Peat sampling and molecular methods*

168 Peat for the focal dataset presented here was collected from four depth increments below  
169 the peat surface (0-10, 10-20, 30-40, 60-70 cm) in late August/early September in year three of  
170 the experiment, using a 5.08 cm diameter circular corer fitted to an electric drill. Additional  
171 samples for complementary datasets were collected in late August/early September of year one  
172 and in late July of year four using a 2.54 cm (year one) or 7.62 cm (year four) diameter corer  
173 from two depth increments (10-20, 30-40 cm). Different diameter corers and sampling dates  
174 were necessary to accommodate a variety of intended uses for the peat, which varied depending  
175 on the year, and coring multiple times a season with a smaller corer would have created  
176 excessive disturbance in the mesocosms. However, in all years each 10 cm depth increment was  
177 not homogenized but instead a vertical split representative of approximately 25 ml of peat was  
178 subsampled from each for DNA work, ensuring the volume represented by the sampled material  
179 remained comparable among years. Upon collection, samples were flash frozen in liquid nitrogen  
180 and stored at -80°C.

181 Samples were pulverized using a mortar and pestle under liquid nitrogen, followed by a  
182 coffee grinder. DNA was extracted from 0.5 g of ground peat from each sample using a  
183 PowerSoil DNA Isolation kit, cleaned with a PowerClean DNA Clean-Up kit (MoBio  
184 Laboratories; now Qiagen, Germantown, MD, USA), and quantified with a Qubit Fluorometer

185 (Invitrogen, Life Technologies, Carlsbad, CA, USA). DNA amplicon sequencing was conducted  
186 at the U.S. Department of Energy Joint Genome Institute (JGI, Walnut Creek, California, USA;  
187 now Berkeley, California) following Caporaso et al. (2012), with small modifications (see  
188 Coleman-Derr et al., 2016; Tremblay et al., 2015). PCR amplification utilized the primers 515F  
189 and 806R (Caporaso et al., 2012) targeting the bacterial and archaeal 16S V4 region, and fITS9  
190 (Ihrmark et al., 2012) and ITS4 (White, Bruns, Lee, & Taylor, 1990) targeting the fungal ITS2  
191 region. Primers were fitted with Illumina adaptors and the reverse primer contained an 11 bp  
192 barcode. Samples were pooled into equimolar portions and sequenced on an Illumina MiSeq  
193 platform (Illumina, Inc., San Diego, CA) using 2 x 250 bp (year one) or 2 x 300 bp (years three,  
194 four) chemistry.

195

## 196 *Bioinformatics*

197 Processing of DNA sequence reads proceeded as follows. Illumina adapters and PhiX  
198 174 were removed with BBDuk ([sourceforge.net/projects/bbmap/](http://sourceforge.net/projects/bbmap/)), and 3' and 5' PCR primers  
199 were trimmed with Cutadapt 1.18 (Martin, 2011). Paired reads were merged with BBmerge  
200 (Bushnell, Rood, & Singer, 2017), and those with expected error rate >1 and/or ambiguous bases  
201 were removed with VSEARCH 2.5.1 (Rognes, Flouri, Nichols, Quince, & Mahé, 2016). The  
202 5.8S (94 bases) and 28S (35 bases) flanks were trimmed from ITS2 reads with Cutadapt, and  
203 resulting amplicons < 95 bases long were filtered. 16S V4 reads were not trimmed, but those  
204 exceeding  $\pm 8$  bases from the median length (253 bases) were excluded. Chimera detection and  
205 removal was implemented with the VSEARCH plug-in for QIIME 2 (Bolyen et al., 2019) using  
206 the UNITE UCHIME ITS2 reference dataset (v. 7.2; Nilsson et al., 2015), and the SILVA 16S  
207 dataset (128 QIIME release; Quast et al., 2013). De novo operational taxonomic units (OTUs)

208 were created with the QIIME 2 VSEARCH plug-in by first clustering at 98.5% similarity, then  
209 clustering the resulting OTU reference sequences at 97% similarity. 97% OTUs were curated  
210 with LULU (min match = 90, min relative cooccurrence = 0.95; Frøslev et al., 2017), followed  
211 by the removal of OTUs with < 10 reads in the dataset (Lamit et al., 2017).

212 The OTU matrices were further filtered and annotated. The QIIME 2 naive Bayes  
213 feature-classifier plug-in (Bokulich et al., 2018; Pedregosa et al., 2011) was used to assign  
214 taxonomy with the UNITE all eukaryote dynamic species hypothesis dataset for fungi (v. 8.0,  
215 released 02.02.2019; Kõljalg et al., 2013) and the SILVA 16S +18S dataset for prokaryotes (128  
216 QIIME release; Quast et al., 2013). OTUs not assigned the taxonomy of target lineages were  
217 excluded. Next, ITS2 OTU representative sequences were aligned to the UNITE species  
218 hypothesis dynamic fungal dataset (v. 8.0, released 02.02.2019) using BLAST in QIIME 2, and  
219 reads that did not match at least  $\geq 70\%$  of their length to fungi with a similarity of  $\geq 75\%$  were  
220 filtered (Tedersoo et al., 2015). To further remove potential non-target sequences, the above 16S  
221 V4 pipeline was run with 28 marine samples originally sequenced on the same plate as the year  
222 one peat samples. Although rare in the dataset, OTUs that occurred in year one peat samples and  
223 the marine samples, and had BLAST matches to salt-tolerant taxa reported from marine/saline  
224 systems in NCBI DNA sequence database (<http://blast.ncbi.nlm.nih.gov>) were removed as  
225 potential contaminants. The datasets were then rarified to 5000 reads per sample prior to all  
226 statistical analyses. Tentative functional groups were assigned with FAPROTAX (Louca,  
227 Parfrey, & Doebeli, 2016) and FUNGuild (Nguyen et al., 2016), with further refinement based  
228 on literature searches.

229

230 *Statistical analyses*

231 Our first set of analyses examined overall patterns of OTU composition over the course  
232 of the experiment, with the aim of identifying when treatment effects began to manifest most  
233 distinctly and if the focal dataset (year three) was representative of the overall treatment effects  
234 of the experiment. PerMANOVA (Anderson, 2001) was used to separately examine the two peat  
235 depths sampled in all years (10-20 cm, 30-40 cm), with each analysis including the fixed effects  
236 of WT (high, low), presence/absence of Ericaceae, presence/absence of sedges, year, and all  
237 possible interactions among these effects. Each model also included block as a fixed effect, and  
238 individual mesocosm as a random effect to account for non-independence of samples from the  
239 same mesocosm. Modeling treatments in this way allowed for assessment of the presence of each  
240 vascular PFG (as opposed to the removal of the other PFG) because each PFG was present in  
241 mesocosms with and without the other PFG, but a direct Ericaceae by sedge interaction could not  
242 be tested. For all PerMANOVAs, we also report the square root of the estimated component of  
243 variation for each factor; these are in Bray-Curtis units (scaled between 0 and 100), and can be  
244 used for comparing the relative importance of terms in a model for explaining the overall  
245 variation in community composition (Andersen, Gorley & Clarke, 2008). Removal or pooling of  
246 model terms with negative estimates for components of variation (Andersen, Gorley & Clarke,  
247 2008) had little effect on the significance tests or estimates of components of variation for the  
248 remaining terms in any of the PerMANOVA we ran, therefore all terms were always retained for  
249 simplicity. These analyses were complemented with canonical analysis of principal coordinates  
250 (CAP; Andersen & Willis 2003) to visualize communities. PerMANOVA and CAP were run  
251 using Bray-Curtis dissimilarity. Prior to analyses, OTU matrices were relativized as proportions  
252 of sample read totals followed by 4<sup>th</sup> root transformation to down-weight dominant OTUs.  
253 PerMANOVAs were conducted with Type III sums of squares, using permutation of residuals

254 from partial models. PerMANOVA was run in Primer 6.1.15 with PERMANOVA+ 1.0.5  
255 (PRIMER-E, Plymouth, UK), and other analyses utilized *Vegan* (Oksanen et al., 2019) in R 3.6.3  
256 (R core team, 2020).

257 To gain deeper insight into the depth-specific microbial community responses to PFG and  
258 WT manipulations, we focused our more detailed analyses on microbial communities along the  
259 70 cm depth gradient sampled in year three. First, PerMANOVAs were run using the equivalent  
260 model structure as those for the multi-year analysis described above, substituting a depth effect  
261 for the year effect. Second, PerMANOVAs were also run individually for each depth using the  
262 fixed effects of WT, presence/absence of Ericaceae, presence/absence of sedges, block and two-  
263 way interactions between WT and the presence/absence of each PFG. Third, CAP ordinations  
264 were used to visualize OTU composition. Fourth, indicator species analysis was used to identify  
265 the 25 strongest indicator OTUs associated with each depth and treatment using the R package  
266 *indicspecies* (De Cáceres & Legendre, 2009). Fifth, linear mixed models were used to examine  
267 the responses of OTU richness and total archaea, and the relative abundances of a limited set of  
268 select functional groups with known relevance to carbon cycling (ErMF, lignocellulose-  
269 degrading fungi, methanotrophic bacteria, methanogenic archaea). These models included the  
270 fixed effects of depth, WT, presence/absence of sedges and Ericaceae, and all possible  
271 interactions among these fixed effects, plus the fixed effect of block and the random effect of  
272 mesocosm. Linear mixed models were fit in R with *lmerTest* (Kuznetsova, Brockhoff, &  
273 Christensen, 2017) using the Kenward-Roger approximation for *F*-tests. The *emmeans* package  
274 (Russell, 2020) was used to generate marginal means from linear mixed models, and the  
275 *effectsize* package (Ben-Shachar, Makowski & Lüdtke, 2020) was used to obtain partial- $\omega^2$   
276 values to use as effects sizes for comparison between model terms.

277           The final set of analyses utilized structural equation modeling with the depth gradient  
278 data from year three (Fig. S2). Our specific goals were to: 1) test if the effect of PFG  
279 manipulation was primarily due to the alteration of plant species composition as opposed to  
280 unmeasured variables affected by PFG manipulation, 2) examine the potential for WT to  
281 modulate PFG effects through modification of the plant community, and 3) measure how the  
282 strength of PFG and WT effects changed with depth. Separate models were created for each  
283 microbial community, at each depth. Variables used in the models included a vector for WT  
284 treatment (0 = high WT, 1 = low WT), a matrix coding for PFG treatment, matrices representing  
285 Ericaceae and sedge species composition, and the fungal or prokaryote matrices utilized in all  
286 previous analyses. The PFG treatment matrix had three columns, each representing a different  
287 treatment group, that were coded as 1 or 0 denoting the treatment applied to the mesocosm a  
288 sample originated from. The sedge and Ericaceae matrices were point-intercept data from Potvin  
289 et al., (2015) measured in year three, and expressed as the % of intercepts represented by each  
290 species to emphasize shifts in absolute abundances. A small constant was added as a dummy  
291 species to each plant community matrix to account for some mesocosms lacking members of  
292 sedges or Ericaceae. Model sub-components were tested with multiple regression on distance  
293 matrices (Lichstein, 2007) with ranked data using *ecodist* (Goslee & Urban, 2007) in R, and  
294 model fit was assessed with directional separation tests (Shiple, 2000) calculated manually.

295

## 296 **Results**

297           Diverse communities were recovered through amplicon sequencing. The rarefied fungal  
298 ITS2 dataset contained 189 samples (sample size: year one = 48, year three =96, year four = 45;  
299 945,000 sequences), and in all years was dominated by Helotiales (Ascomycota), followed by

300 Agaricales, Sebaciniales and Polyporales (Basidiomycota) (Fig. S3A). In total, there were 1,193  
301 fungal OTUs, with an average of 64.1 OTUs per sample (stdev = 17.8, range = 25-117 OTUs).  
302 The rarefied prokaryote dataset contained 191 samples (sample size: year one = 48, year three =  
303 96, year four = 47; 955,000 sequences). In all years, bacteria were dominated by Acidobacteria  
304 and Proteobacteria, while Archaea were dominated by Euryarchaeota, followed by  
305 Thaumarchaeota and Bathyarchaeota (Fig. S3B). There was a total of 7,353 prokaryote OTUs,  
306 with an average of 606.0 OTUs per sample (stdev = 158.1, range = 235-897 OTUs).

307

### 308 ***Response of community composition across four years of WT and PFG manipulation***

309 Changes in OTU composition over time supported the hypotheses that PFG and WT  
310 manipulation alter microbial communities (H1, H2) but provided no evidence for their  
311 interaction (H3). At the 10-20 cm depth, fungal and prokaryote composition were influenced by  
312 the presence of Ericaceae and WT manipulation (Table 1; Fig. 1A, 1B). The components of  
313 variation from the PerMANOVA models indicate that the main effect of Ericaceae on fungal  
314 composition was ~50% greater than that of WT at 10-20 cm, while the main effect of Ericaceae  
315 on prokaryotes was 25% less than for WT (Table 1). At the 30-40 cm depth, WT was a slightly  
316 stronger influence on fungi than was the presence of Ericaceae, while the influence of WT on  
317 prokaryote composition was more than twice the strength of the marginally significant influence  
318 of Ericaceae (Table 1; Fig. 1C, 1D). At both depths, many of the treatment effects manifested  
319 most strongly in years three and four (significant treatment by year interactions; Table 1), which  
320 is visually apparent in CAP ordinations (Fig. 1). When integrating over the course of the  
321 experiment there was no evidence for a sedge effect on OTU composition at either depth (Table  
322 1). Importantly, patterns in OTU composition across years confirmed that our focal depth

323 gradient dataset (year three) was representative of the broader PFG and WT effects over the  
324 course of the experiment (Fig. 1).

325

326 ***Community responses over the year three 70 cm peat depth gradient***

327 *Community depth stratification* — Depth had the largest influence on OTU composition  
328 of any factor included in the full depth gradient PerMANOVA models (Table 1), and all depths  
329 showed uniqueness in fungal and prokaryote communities relative to other depths (Fig 2, S3).  
330 All depths had indicator OTUs that were members of the Helotiales, while 0-10 and 60-70 cm  
331 depths also had many indicator OTUs representing a broader set of additional fungal lineages  
332 (Table S1). Fungal indicators of the 0-10 and 10-20 cm depths included ErMF, plant pathogens,  
333 general saprotrophs and lignocellulose degraders, while indicators of the 30-40 and 60-70 cm  
334 depths included non-mycorrhizal root associates, general saprotrophs and lignocellulose  
335 degraders (Table S1). Many prokaryote indicators, especially in the 0-10, 10-20 and 30-40 cm  
336 depths, were from acid tolerant groups (e.g., Acidobacteriaceae, Acetobacteraceae). In the  
337 deepest depth (60-70 cm) there was an increase in Deltaproteobacteria and archaeal indicator  
338 OTUs, many of which are adapted to reduced conditions (e.g., methanogens, sulfate reducers;  
339 Table S1). Prokaryote and fungal OTU richness decreased with depth, while total archaea  
340 relative abundance increased (Table 2; Fig. S4).

341 *Community composition responses to PFG and WT across the 70 cm depth gradient* — In  
342 support of H1, fungal and prokaryote OTU composition exhibited depth-specific responses to  
343 PFG manipulation, although responses were only driven by Ericaceae (Table 1, 3). The influence  
344 of Ericaceae on fungi and prokaryotes was distinctive in the upper depths (0-10, 10-20 cm) and  
345 disappeared in the deeper depths (30-40, 60-70 cm; Table 1, 3; Fig. 2). Microbial communities in



346 mesocosms with and without Ericaceae exhibited depth-specific differences in their top indicator  
347 OTUs, but there were also general patterns. ErMF OTUs were some of the top fungal indicators  
348 of mesocosms with Ericaceae, while root endophytes and lignocellulose degraders were top  
349 indicators of mesocosms lacking Ericaceae (Table S1). Of particular note is the lignocellulose-  
350 degrading genus *Galerina* whose members are top indicators of the absence of Ericaceae at 0-10  
351 and 10-20 cm (Table S1), and have high relative abundances in these depths (Fig. S5).  
352 Mesocosms lacking Ericaceae included indicators from 10 different bacteria phyla and indicators  
353 of mesocosms containing Ericaceae were primarily Proteobacteria, but both treatments included  
354 some Acidobacteria indicators (Table S1).

355 Fungal and prokaryote OTU composition also exhibited depth-dependent responses to  
356 WT manipulation, in support of H2, but there was no evidence for WT x PFG interactions (H3).  
357 The WT effect in the upper depths was stronger in the surface (0-10 cm) than subsurface (10-20  
358 cm) peat for both communities. Despite this, at the upper two depths the strength of the WT  
359 effect on fungal OTU composition was subordinate to the Ericaceae effect, whereas this pattern  
360 was reversed for prokaryotes (Table 3, Fig. 2). Prokaryote composition was the most divergent  
361 between WT groups at the 30-40 cm peat depth, while the influence of WT on fungi at this depth  
362 was clearly far weaker than for prokaryotes (Table 3; Fig. 2). Neither fungal nor prokaryote  
363 composition responded to WT at 60-70 cm (Table 3). The responses of composition were driven  
364 by large depth-specific shifts in indicator OTUs for both fungi and prokaryotes (Table S1). For  
365 example, fungal indicators of the high WT treatment in the surface peat (0-10 cm) were primarily  
366 non-ErMF taxa likely associated with living or recently dead plant tissues, and indicators of the  
367 low WT treatment included a higher proportion of ErMF (Table S1). In contrast, indicators of the  
368 high WT treatment at 30-40 cm represented a very broad range of functions whereas the 30-40

369 cm low WT indicators were primarily non-mycorrhizal root-associates (Table S1). Interestingly,  
370 an OTU assigned to the Methanomicrobia (hydrogenotrophic methanogens) was an indicator of  
371 the high WT treatment in the three upper depths (Table S1), and three known methanotrophs  
372 were top indicators of the 10-20 cm depth high WT treatment.

373 *Microbial functional group and OTU richness responses to PFG and WT across the 70*  
374 *cm depth gradient* — Microbial functional groups and OTU richness exhibited complex  
375 responses to PFG, WT and/or PFG x WT interactions, lending support to all hypotheses (Table  
376 2; Fig. 3). Total ErMF relative abundance was strongly suppressed by Ericaceae removal and the  
377 high WT treatment in the upper two depths (0-10, 10-20 cm; Table 2, Fig. 3A). Total  
378 lignocellulose degrading fungi increased in relative abundance in the absence of Ericaceae,  
379 especially in the upper depths (Table 2; Fig. 3B). The response of lignocellulose degraders  
380 remained significant after including total ErMF relative abundance as a covariate in the mixed  
381 model, indicating that the response was not solely driven by the removal of ErMF from the DNA  
382 pool (Ericaceae effect:  $F_{(1,21.3)} = 4.87$ ,  $P = 0.038$ ; marginal means  $\pm 1$  SE : Ericaceae present =  
383  $0.08 \pm 0.01$ , Ericaceae absent =  $0.17 \pm 0.05$ ). Methanogens and total archaea had elevated relative  
384 abundances in high WT, except at 60-70 cm, with methanogen relative abundance peaking in the  
385 30-40 cm high WT treatment (Table 2; Fig. 3C, S4C). Ericaceae removal tended to increase  
386 methanotroph relative abundance with high WT and decrease it in low WT at most depths  
387 (especially prominent at 10-20 and 30-40 cm) whereas sedge removal generally decreased  
388 methanotroph relative abundance (Table 2; Fig. 3D). The major fungal and prokaryote taxa  
389 within these functional groups often but not always followed the overall patterns of the group as  
390 a whole (Fig. S5).

391 OTU richness results partially supported H1, H2 and H3. High WTs clearly depressed

392 fungal richness in the surface peat (0-10 cm), although WT was not a significant overall effect in  
393 the model (Table 2; Fig S4A). Prokaryote richness was influenced by interactions among  
394 Ericaceae, WT, depth and sedges, with results only marginally significant in some cases (Table  
395 2; Fig. S4B). For example, in the 0-10 cm depth the highest richness occurred in mesocosms  
396 lacking Ericaceae, and at the 30-40 cm depth prokaryote OTU richness was elevated in low WT  
397 treatments but this pattern reversed at 60-70 cm (Fig. S4B). Sedge removal tended to elevate  
398 prokaryote OTU richness in the high WT treatment at the 0-10 cm depth, whereas a similar  
399 effect was evident in the low WT treatment at the 10-20 cm depth (Fig. S4B).

400       *Structural equation modeling* — SEM supported the hypotheses concerning depth-  
401 dependent effects of PFG and WT (H1, H2), with several key results. First, the variation in  
402 fungal composition explained by the models was greatest in the 0-10 and 10-20 depths, the  
403 greatest variation in prokaryote composition was explained by models for the 0-10 and 30-40 cm  
404 depths, and the models for the 60-70 cm depth explained almost no variation in either  
405 community (Fig. 4A, 4B). Second, PFG treatment effects were stronger on fungi than  
406 prokaryotes and were most pronounced at 0-10 and 10-20 cm, while WT effects were stronger on  
407 prokaryotes than fungi and were more pronounced in the 0-10 and 30-40 cm depths (Fig. 4).  
408 Third, at the 0-10 cm depth the effect of PFG manipulation primarily occurred through changes  
409 in Ericaceae composition. In contrast, initial model fit tests at 10-20 cm ( $P < 0.1$ ) indicated the  
410 need for a direct path from PFG treatment to microbial communities. This direct path to OTU  
411 composition (Fig. 4A, 4B) was equivalent or greater than the compound effect of PFGs acting  
412 through Ericaceae composition at 10-20 cm (compound effect: fungi partial  $\rho = 0.24$ ,  
413 prokaryotes partial  $\rho = 0.04$ ; Table S2.). Fourth, WT influenced Ericaceae composition, but the  
414 effect of WT on microbial communities through Ericaceae was small (Table S2).

415

416 **Discussion**

417         Our results reveal the strength and depth-dependence of WT and PFG effects on  
418 microbial communities. The strikingly greater impact of PFG on fungi near the surface  
419 contrasted with the stronger impact of WT on prokaryotes across a broader range of depths.  
420 These patterns can be explained by abiotic and biotic factors: the intolerance of most fungi to  
421 anoxic conditions (Kavanagh, 2011) constraining most taxa to shallow peat, and the colocation  
422 of the dominant ErMF with their shallowly-rooted Ericaceae hosts (Moore et al., 2002; Wallén,  
423 1987). In contrast, the broad range of moisture niches, metabolic pathways, and redox tolerance  
424 among soil prokaryotes (e.g., Bodelier & Dedysh, 2013; Lennon, Aanderud, Lehmkuhl, &  
425 Schoolmaster, 2012) and the strong sensitivity of prokaryote communities to changes in soil  
426 moisture (e.g., Bapiri, Bååth, & Rousk, 2010; Barnard, Osborne, & Firestone, 2013), explain  
427 their shift with WT treatments in both drier surface peat as well as at acrotelm/catotelm boundary  
428 where redox conditions are most dynamic (Kane et al., 2019; Tfaily et al., 2018). These depth-  
429 dependent effects indicate that WT and PFG are among the key shapers of the vertical  
430 physicochemical gradients that structure peatland microbial communities (Andersen, Chapman  
431 & Artz 2013; Artz et al., 2007; Lin et al., 2014), the activities of which then feed back to  
432 modulate carbon cycling along the peat profile (Chanton et al., 2009; Kane et al., 2019; Lin et  
433 al., 2014; Tfaily et al., 2018). Although discussions on wetland carbon cycling usually emphasize  
434 the role of anoxic reducing conditions (e.g., Schlesinger & Bernhart, 2013), most carbon inputs  
435 from primary production in bogs and poor fens derive from senesced *Sphagnum* in the largely  
436 oxic acrotelm (van Breeman, 1995; Rydin & Jeglum, 2013), making aerobic organisms  
437 instrumental as the initial transformers of peatland organic matter. As this partially degraded

438 organic matter transitions into the catotelm, anaerobic metabolism becomes paramount, which is  
439 reflected in the OTU composition in the deeper depths of PEATcosm and other studies (e.g., Lin  
440 et al., 2014; Wang et al., 2019). Hence, the microbial community present at a peat depth sets  
441 bounds on how the community can change with drought and changes in dominant PFGs, and the  
442 integrated responses of fungi and prokaryotes along the profile may influence the magnitude of  
443 CO<sub>2</sub> and CH<sub>4</sub> released from peatlands under different climate change scenarios.

444

#### 445 ***Responses to PFG manipulation***

446 As predicted, the influence of Ericaceae was greatest in the upper peat, which suggests a  
447 restructuring of the community involved in aerobic carbon cycling with changes in dominant  
448 PFGs. The presence of Ericaceae can have a strong impact on fungal communities in surface peat  
449 (Ward et al., 2015; Kennedy, Mielke, & Nguyen, 2018), and observational studies indicate that  
450 links between microbial communities and vegetation composition decline with depth in the peat  
451 profile (e.g., Artz et al., 2007; Lin et al., 2014). ErMF showed a marked decrease in abundance  
452 when Ericaceae were removed, indicating a preference for host photosynthate despite the free-  
453 living saprotrophic capabilities of some ErMF (Martino et al., 2018). Of particular interest is that  
454 relative abundance of the dominant ErMF Ascomycota, *Hyaloscypha ericae* (deprecated  
455 synonyms are *Pezoloma ericae* and *Rhizoscyphus ericae*), was depressed less by Ericaceae  
456 removal than the dominant ErMF Basidiomycota, *Serindipita* spp., possibly indicating a greater  
457 degree of host dependency in the latter or more dormant propagules in the former. Ericaceae also  
458 influenced a variety of non-ErMF fungi and prokaryotes, which may be driven by several  
459 mechanisms. ErMF may competitively suppress saprotrophs (Verbruggen et al., 2017). In  
460 particular, one of the most abundant saprotroph genera in surface depths, *Galerina*, responded

461 very positively to removal of Ericaceae. These *Galerina* species are *Sphagnum* peatland  
462 specialists (Gulden, Stensrud, Shalchian-Tabrizi, & Kauserud. 2005, Castellano 2003), and the  
463 lignocellulose degrading capability of fungi in the genus (Nagendran, Hallen-Adams, Paper,  
464 Aslam, & Walton. 2009; Riley et al., 2014) suggest that these species are adapted to *Sphagnum*  
465 as a substrate. If their activity is suppressed by ErMF it could also lead to greater accumulation  
466 of partially-degraded *Sphagnum* litter, because ErMF do not possess the complete suite of lignin-  
467 degrading enzymes, most notably class II peroxidases. Although *Sphagnum* does not technically  
468 produce lignin, it does have analogous chemical components that resist hydrolytic decomposition  
469 (Bengtsson, Rydin, & Hájek. 2018). Some taxa may also utilize the byproducts from ErMF  
470 decomposition of organic matter. Ericaceae tissues also represent a direct input of carbon into  
471 surface peat through exudates, senescence, and leaching phenolics that may act as unique  
472 microbial substrates and/or inhibitors (Weigang, Artz, & Johnson, 2008).

473         Two unexpected findings about the PFG effects are worth noting. First, SEM suggested  
474 the effect of Ericaceae in the 10-20 cm peat was partially due to factors influenced by PFG  
475 manipulation aside from direct changes in Ericaceae composition. These may include  
476 subsidence, peat accumulation rates and other physicochemical parameters influenced by PFG  
477 (Kane et al., 2019; Potvin et al., 2015). Additionally, aboveground plant community data may  
478 not fully reflect the density of some or all Ericaceae species roots (even non-relativized, as we  
479 used in the SEM), especially in the 10-20 cm depth where root density is high; this may reduce  
480 the explanatory power of the Ericaceae community matrix thus elevating the strength of the  
481 direct path from PFG manipulation to the microbial community. Second, inconsistent with our  
482 hypothesis, sedges had a limited effect on microbial communities. Oxygenation and substrates  
483 from sedge roots are important shapers of microbial-driven processes in sedge dominated

484 minerotrophic fens (Chanton et al., 2008; Rupp et al., 2019), and of diverse microbial  
485 communities associate with sedges rhizospheres (Hough et al., 2020). The modest sedge effect in  
486 PEATcosm might be due to their relatively low biomass in the ombrotrophic habitat we focused  
487 on, which became more variable through the course of the experiment (Potvin et al., 2015). Our  
488 findings contrast with a study, also in an ombrotrophic peatland, where microbial phospholipid  
489 fatty acid (PLFA) composition responded more to sedge removal than Ericaceae removal  
490 (Robroek et al., 2015). However, it is difficult to directly compare PLFA and amplicon  
491 sequencing results, and results from Ward et al. (2015) suggest no sedge influence on fungal  
492 composition. It is also possible that the *Eriophorum*-dominated sedge communities in Robroek et  
493 al. (2015) have a greater effect on microbial communities than our *Carex*-dominated sedge  
494 communities.

495

#### 496 ***Responses to WT manipulation***

497 As predicted, the microbial response to WT manipulation was depth-dependent. In both  
498 WT treatments, the upper 20 cm of peat was above the WT for a considerable time period prior  
499 to sampling, with the 0-10 cm depth being above the WT surface for most of the growing season.  
500 Distance above the WT interacts with peat density, porosity, and capillarity to drive differences  
501 in moisture availability that may have directly affected the response of microbial communities in  
502 the unsaturated upper peat to WT manipulation. Importantly, changes in taxa associated with  
503 methane cycling (methanogens, methanotrophs) and the processing of complex organic matter  
504 (lignocellulose degraders, ErMF) highlight the potential extended effects of WT on carbon  
505 cycling even in non-saturated peat. Our results contrast a recent mesocosm study by  
506 Asemaninejad et al. (2018) who did not detect an effect of WT manipulation on fungal

507 communities at any depth in the profile. However, the discrepancy in results may be due to the  
508 more limited WT depth differential between treatment levels in Asemaninejad et al. (2018),  
509 which were maintained at a stable level for the course of the experiment. Moss species  
510 composition and productivity were also influenced by WT treatment in PEATcosm (Potvin et al.  
511 2015), and we suspect that the stronger WT effect on fungi and prokaryotes in the 0-10 cm depth  
512 than the 10-20 cm depth is in part due to the role of *Sphagnum* mosses in structuring their  
513 microbiomes (Kostka et al., 2016). The potential foundational influence of moss in acid  
514 *Sphagnum* peatlands is one factor making these systems distinct from minerotrophic fens, where  
515 moss cover has a less clear influence on microbial communities in surface peat (Emsens et al.,  
516 2020).

517         Microbial community responses in the deeper peat were likely driven by WT inundation  
518 and oscillation. WT manipulation elicited some of its strongest responses (especially on  
519 prokaryotes) at the 30-40 cm depth, where peat was perennially underwater in the high WT  
520 treatment but seasonally above the WT in the low WT treatment. This should promote  
521 communities capable of aerobic decomposition during the driest part of the season and anaerobic  
522 decomposition on the shoulders of the growing season in the low WT treatment, and  
523 communities associated with slower carbon transformations typical of anoxic reduced conditions  
524 in the high WT treatment, a contention supported by our indicator species analyses. Interestingly,  
525 methanogens reached their greatest relative abundance in the high WT treatment at the 30-40 cm  
526 depth. This might reflect a preference for anoxic conditions combined with inputs of fresh, labile  
527 substrates for fermenters and syntrophs to generate H<sub>2</sub> and CO<sub>2</sub> used in hydrogenotrophic  
528 methanogenesis (Conrad, 1999). Oscillation between oxic and anoxic conditions at the interface  
529 between the acrotelm and catotelm (i.e., the mesotelm) is associated with rapid organic matter



530 transformations (Kane et al., 2019; Lin et al., 2014; Tfaily et al., 2018), and shifts in microbial  
531 composition in the low WT treatment at 30-40 cm indicate a downward extension of this  
532 biogeochemical “hotspot” during drought. In contrast, the community at the 60-70 cm depth was  
533 continuously submerged far below the WT surface, buffered from changes in the WT level and  
534 less affected by roots, suggesting that deep peat microbial communities and their influences on  
535 carbon cycling may not be strongly affected by seasonal drought, at least in the short term.

536

### 537 *Interactions between PFG and WT*

538         The response of some components of the microbial community highlighted the potential  
539 for WT and PFG interactions. The negative effect that Ericaceae tended to have on aerobic  
540 methanotrophs in the high WT treatment might be explained by dense Ericaceae roots depleting  
541 rhizosphere O<sub>2</sub> through respiration, although other mechanisms of direct interference are possible  
542 (e.g., via antibiosis; Adeoyo, Pletschke, & Dames, 2019). Additionally, at the 10-20 cm depth,  
543 the more complete decline in ErMF after Ericaceae removal in high WT mesocosms suggests  
544 that the negative effects of host removal are compounded by anoxia when this depth is flooded  
545 during the shoulders of the growing season. Given the importance of ErMF and methanotrophs  
546 to peatland carbon cycling, these results indicate that the responses of subcomponents of the  
547 community to PFG x WT interactions may cause shifts in the functioning of microbial  
548 communities even when overall OTU composition responds more slowly.

549         Climate change-driven shifts in peatland soil moisture can influence microbial  
550 communities indirectly by shifting the composition of plant communities (e.g., Bragazza et al.,  
551 2013; Jassey et al., 2018). Evidence for this in PEATcosm was not strong; although the low WT  
552 treatment promoted Ericaceae cover and productivity, with some species responding more than

553 others (Potvin et al., 2015), the compound path effects from WT through Ericaceae composition  
554 to the microbial communities were very small. Our SEMs may have underestimated the path  
555 coefficient between WT and Ericaceae composition because mesocosms lacking Ericaceae did  
556 not have a community to respond to WT treatment. However, we suspect that the strong effect of  
557 simply having or not having Ericaceae overshadowed the impact of WT on microbial  
558 communities acting through modification of the plant community. Over a longer time scale the  
559 influence of WT on microbial communities acting through vegetation change should become  
560 stronger, which would represent an indirect pathway in natural systems for climate change-  
561 related droughts to influence microbial communities.

562

### 563 *Conclusions*

564 Our results demonstrate the importance of WT and PFG in structuring peatland microbial  
565 communities along peat depth profiles. Peatlands in many regions are experiencing increased  
566 temperatures, changes in long-term precipitation patterns, and other anthropogenic disturbances  
567 that influence WT dynamics and cause shifts in plant communities; the influence of these factors  
568 on carbon cycling will be contingent on how they influence microbial communities. Importantly,  
569 microbial lineages and functional groups do not all respond equivalently to WT and PFG  
570 manipulations, and their responses are depth-dependent. The strong mutualistic interactions of  
571 fungi and Ericaceae appear to be the driver of the greater PFG effect on fungal than prokaryote  
572 communities. The dominance of ErMF in shallow oxic peat indicates a large potential for  
573 Ericaceae to shape microbial community structure and function near the surface, where most new  
574 organic matter enters the peat profile. The fact that when Ericaceae are removed lignocellulose  
575 degraders respond very positively in relative abundance begs the question of whether this

576 represents a functional release of those taxa in the absence of mycorrhizal competition.  
577 Similarly, the significant prokaryote response to PFG manipulations indicates changing  
578 resources and conditions driven by plant traits can also structure these communities, especially  
579 methanotrophs; but the weaker prokaryote (vs. fungal) response to PFG vs. WT manipulation  
580 suggests redox conditions predominate in structuring these communities. The vertical complexity  
581 in responses, likely driven by declining PFG influence and increasing influence of declines in  
582 both redox potential and organic matter quality with depth, highlights the necessity of accounting  
583 for depth stratification when understanding the responses of peatland microbial communities to  
584 global change.

585

## 586 **Acknowledgments**

587 We thank K. Griffith, E. Matthys, A. Bales, R. Schwartz, L. Theobald, J. Hribljan, T. Ontl and  
588 M. Wiedermann. This work was supported by the USDA Forest Service Northern Research  
589 Station Climate Change Program, the US National Science Foundation (DEB-1 146 149), and  
590 the U.S. Department of Energy Joint Genome Institute Community Science Program (Proposal  
591 ID 1445). The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE  
592 Office of Science User Facility, is supported by the Office of Science of the U.S. Department of  
593 Energy under Contract No. DE-AC02-05CH11231.

594

## 595 **Data Accessibility**

596 - These sequence data have been submitted to the National Center for Biotechnology Information  
597 (NCBI) Sequence Read Archive under accession number PRJNA650129.  
598 -Vegetation data is available through Pangea: [doi.pangaea.de/10.1594/PANGAEA.902313](https://doi.org/10.1594/PANGAEA.902313).

599

600 **Author Contributions**

601 E.A. Lilleskov, E.S. Kane, R.K. Kolka, J.T. Lennon, R.A. Chimner and S.G. Tringe conceived  
602 the study and obtained funding. L.R. Potvin, E.A. Lilleskov, E.S. Kane, K.J. Romanowicz, S.G.  
603 Tringe and L.J. Lamit performed the research. L.J. Lamit analyzed the data. L.J. Lamit and E.A.  
604 Lilleskov wrote the paper, with all coauthors contributing to revisions.

605

606 **References**

- 607 Adeoyo, O. R., Pletschke, B. I., & Dames, J.F. (2019). Molecular identification and antibacterial  
608 properties of an ericoid associated mycorrhizal fungus. *BMC microbiology*, *19*, 178.  
609 doi.org/10.1186/s12866-019-1555-y
- 610 Asemaninejad, A., Thorn, R.G. & Lindo, Z., 2017. Vertical distribution of fungi in hollows and  
611 hummocks of boreal peatlands. *Fungal Ecology*, *27*, 59-68.
- 612 Asemaninejad, A., Thorn, R.G., Branfireuna, B.A., & Lindo, Z. (2018). Climate change favours  
613 specific fungal communities in boreal peatlands. *Soil Biology and Biochemistry*, *120*, 28-36.
- 614 Asemaninejad, A., Thorn, R.G., Branfireun, B.A. & Lindo, Z. (2019). Vertical stratification of  
615 peatland microbial communities follows a gradient of functional types across hummock–  
616 hollow microtopographies. *Ecoscience*, *26*, 249-258.
- 617 Andersen, R., Chapman, S. J., & Artz, R. R. E. (2013). Microbial communities in natural and  
618 disturbed peatlands: A review. *Soil Biology and Biochemistry*, *57*, 979-94.
- 619 Anderson, M.J., Gorley, R.N., & Clarke KR. (2008). PERMANOVA for PRIMER: Guide to  
620 Software and Statistica Methods. Plymouth, UK: PRIMER-E.
- 621 Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of  
622 variance. *Austral Ecology*, *26*, 32–46.
- 623 Anderson, M. J., & Willis, T. J. (2003). Canonical analysis of principal coordinates: a useful  
624 method of constrained ordination for ecology. *Ecology*, *84*, 511-525.
- 625 Artz, R. R. E., Anderson, I. C., Chapman, S. J. Hagn, A., Schloter, M., Potts, J.M. & Colin D.  
626 Campbell. (2007). Changes in fungal community composition in response to vegetational  
627 succession during the natural regeneration of cutover peatlands. *Microbial Ecology*, *54*, 508–  
628 22.
- 629 Bapiri, A., Bååth, E., & Rousk J. (2010). Drying-rewetting cycles affect fungal and bacterial  
630 growth differently in an arable soil. *Microbial Ecology*, *60*, 419–428. doi: 10.1007/s00248-  
631 010-9723-5
- 632 Barnard, R. L., Osborne, C. A., & Firestone, M. K. (2013). Responses of soil bacterial and fungal  
633 communities to extreme desiccation and rewetting. *ISME Journal*, *7*, 2229–2241.  
634 doi:10.1038/ismej.2013.104
- 635 Bardgett, R. D, Freeman, C., & Ostle N. J. (2008). Microbial contributions to climate change  
636 through carbon cycle feedbacks. *ISME Journal*, *2*, 805–814.
- 637 Bengtsson, F., Rydin, H., Hájek, T. (2018). Biochemical determinants of litter quality in 15

638 species of *Sphagnum*. *Plant and Soil*, 425, 161-76.

639 Bodelier, P. L. E., & Dedysch, S.N. (2013). Microbiology of wetlands. *Frontiers in Microbiology*,  
640 4, doi: 10.3389/fmicb.2013.00079

641 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., ...  
642 Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data  
643 science using QIIME 2. *Nature Biotechnology*, 37, 852–857. doi: 10.1038/s41587-019-0209-  
644 9

645 Bokulich, N. A., Kaehler, B. D., Rideout, J. R., Dillon, M., Bolyen, E., Knight, R., ... Caporaso,  
646 J. G. (2018). Optimizing taxonomic classification of marker-gene amplicon sequences with  
647 qiime 2's q2-feature-classifier plugin. *Microbiome*, 6, 90. doi: 10.1186/s40168-018-0470-z

648 Bragazza, L., Parisod, J., Buttler, A., & Bardgett, R. D. (2013). Biogeochemical plant–soil  
649 microbe feedback in response to climate warming in peatlands. *Nature Climate Change*, 3,  
650 273–277.

651 Breeuwer, A., Robreck, B. J. M., Limpens, J., Heijmans, M. P. D., Schouten, M. G. C., &  
652 Berendse, F. (2009). Decreased summer water table depth affects peatland vegetation. *Basic  
653 and Applied Ecology*, 10, 330–339.

654 Bridgham, S. D., Pastor, J., Dewey, B., Weltzin, J. F., & Updegraff, K. (2008). Rapid carbon  
655 response of peatlands to climate change. *Ecology*, 89, 3041-3048.

656 Bushnell, B., Rood, J., & Singer, E. (2017). BBMerge – Accurate paired shotgun read merging  
657 via overlap. *PLOS*. doi: 10.1371/journal.pone.0185056

658 Cairney, J. W. G., & Burke, R. M. (1998). Extracellular enzyme activities of the ericoid  
659 mycorrhizal endophyte *Hymenoscyphus ericae* (Read) Korf & Kernan: their likely roles in  
660 decomposition of dead plant tissue in soil. *Plant and Soil*, 205, 181–192.

661 Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., ...  
662 Knight, R. (2012). Ultra-high-throughput microbial community analysis on the Illumina  
663 HiSeq and MiSeq platforms. *ISME Journal*. doi: 10.1038/nmeth.f.303

664 Carini, P., Marsden, P. J., Leff, J. W., Morgan, E. E., Strickland, M. S., & Fierer, N. (2016).  
665 Relic DNA is abundant in soil and obscures estimates of soil microbial diversity. *Nature  
666 Microbiology*, 2, 16242.

667 Castellano, M. A. (2003). Handbook to additional fungal species of special concern in the  
668 Northwest Forest Plan. US Department of Agriculture, Forest Service, Pacific Northwest  
669 Research Station.

670 Chanton, J. P., Glaser, P. H., Chasar, L. S., Burdige, D. J., Hines, M. E., Siegel, D. I., ... Cooper,  
671 W. T. (2008). Radiocarbon evidence for the importance of surface vegetation on  
672 fermentation and methanogenesis in contrasting types of boreal peatlands. *Global  
673 Biogeochemical Cycles*, 22. doi: 10.1029/2008GB003274

674 Chimner, R. A., Pypker, T. G., Hribljan, J. A., Moore, P. A., & Waddington, J. M. (2017). Multi-  
675 decadal changes in water table levels alter peatland carbon cycling. *Ecosystems*, 20, 1042–  
676 1057. doi: 10.1007/s10021-016-0092-x

677 Coleman-Derr, D., Desgarenes, D., Fonseca-Garcia, C., Gross, S., Clingenpeel, S., Woyke, T.,  
678 ... Tringe, S. G. (2016). Plant compartment and biogeography affect microbiome  
679 composition in cultivated and native *Agave* species. *New Phytologist*, 209, 798–811. doi:  
680 10.1111/nph.13697

681 Conrad, R. (1999). Contribution of hydrogen to methane production and control of hydrogen  
682 concentrations in methanogenic soils and sediments. *FEMS Microbiology Ecology*, 28, 193-  
683 202.

684 Crow, S. E. & Wieder, R. K. (2005). Sources of CO<sub>2</sub> emission from a northern peatland: root  
685 respiration, exudation and deposition. *Ecology*, 86, 1825-1834.

686 Davidson, E. A., & Janssens, I. A. (2006). Temperature sensitivity of soil carbon decomposition  
687 and feedbacks to climate change. *Nature*, 440, 165-173.

688 De Cáceres, M., & Legendre, P. (2009). Associations between species and groups of sites:  
689 indices and statistical inference. *Ecology*, 90, 3566-3574.

690 Dieleman, C. M., Branfireun, B. A., McLaughlin, J. W., & Lindo, Z. (2016). Enhanced carbon  
691 release under future climate conditions in a peatland mesocosm experiment: the role of  
692 phenolic compounds. *Plant and Soil*, 400, 81-91.

693 Dorrepaal, E. E. Cornelissen, J. H. C, Aerts, R., Wallén, B., & van Logtestijn, R. S. P. (2005).  
694 Are growth forms consistent predictors of leaf litter quality and decomposability across  
695 peatlands along a latitudinal gradient? *Journal of Ecology*, 93, 817-828.

696 Emsens, W.-J., Diggelen, R., Aggenbach, C., Cajthaml, T., Frouz, J., Klimkowska, A., ...  
697 Verbruggen, E. (2020). Recovery of fen peatland microbiomes and predicted functional  
698 profiles after rewetting. *The ISME Journal*. 10.1038/s41396-020-0639-x.

699 Freeman, C., Ostle, N., & Kang, H. (2001). An enzymic 'latch' on a global carbon store. *Nature*,  
700 409, 149. doi.org/10.1038/35051650

701 Frøslev, T. G., Kjølner, R., Bruun, H. H., Ejrnæs, R., Brunbjerg, A. K., Pietroni, C., & Hansen,  
702 A. J. (2017). Algorithm for post-clustering curation of DNA amplicon data yields reliable  
703 biodiversity estimates. *Nature communications*, 8, 1188. doi: 10.1038/s41467-017-01312-x.

704 Golovchenko, A. V., Dobrovol'skaya, N. G., Inisheva, L. I. (2002). Structure and stocks of  
705 microbial biomass in oligotrophic peat bogs of the southern Taiga in western Siberia.  
706 *Eurasian Soil Science*, 35, 1296-1301.

707 Goslee, S. C., & Urban, D. L. (2007). The ecodist package for dissimilarity-based analysis of  
708 ecological data. *Journal of Statistical Software*, 22, 1-19.

709 Gulden, G., Stensrud, Ø., Shalchian-Tabrizi, K., Kauserud, H. (2005). *Galerina Earle*: a  
710 polyphyletic genus in the consortium of dark-spored agarics. *Mycologia*, 97, 823-37.

711 Hough, M., McClure, A., Bolduc, B., Dorrepaal, E., Saleska, S., Klepac-Ceraj, V., & Rich, V.  
712 (2020). Biotic and environmental drivers of plant microbiomes across a permafrost thaw  
713 gradient. *Frontiers in Microbiology*, 11, 796. doi: 10.3389/fmicb.2020.00796

714 Ihrmark, K., Bödeker, I. T. M., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., &  
715 Lindahl, B. D. (2012). New primers to amplify the fungal ITS2 region – evaluation by 454-  
716 sequencing of artificial and natural communities. *FEMS Microbiology Ecology*, 82, 666-677.

717 Jasse, V. E. J., Reczuga, M. K., Zielińska, M., Słowińska, S., Robroek, B. J. M., Mariotte, P., ...  
718 Buttler, A. (2018). Tipping point in plant-fungal interactions under severe drought causes  
719 abrupt rise in peatland ecosystem respiration. *Global Change Biology*, 24, 972-986. doi:  
720 10.1111/gcb.13928

721 Joosten, H., & Couwenberg, J. (2008). Peatlands and carbon. In F. Parish, A. Sirin, D. Charman,  
722 H. Joosten, T. Minayeva, M. Silvius, & L. Stringer (Eds.), *Assessment on Peatlands,*  
723 *Biodiversity and Climate Change: Main Report* (pp. 99-117). Kuala Lumpur, Malaysia:  
724 Wetlands International.

725 Junk, W. J., An, S., Finlayson, C.M., Gopal, B., Květ, J., Mitchell, S.A., ... Robarts, R.D.  
726 (2013). Current state of knowledge regarding the world's wetlands and their future under  
727 global climate change: a synthesis. *Aquatic Sciences*, 75, 151-167. doi: 10.1007/s00027-012-  
728 0278-z

729 Kane, E. S., Veverica, T. J., Tfaily, M. M., Lilleskov, E. A., Meingast, K. M., Kolka, R. K., ...

730 Chimner, R.A. (2019). Reduction-oxidation potential and dissolved organic matter  
731 composition in northern peat soil: interactive controls of water table position and plant  
732 functional groups. *Geophysical Research: Biogeosciences*, *124*, 3600–3617. doi: 10.1029/  
733 2019JG005339

734 Kavanagh, K. (2011). *Fungi: Biology and Applications*. Chichester, UK: John Wiley & Sons.

735 Kennedy, P.G., Mielke, L.A., & Nguyen, N.H. (2018). Ecological responses to forest age,  
736 habitat, and host vary by mycorrhizal type in boreal peatlands. *Mycorrhiza*, *28*, 315–328.

737 Kõljalg, U., Nilsson, R. H., & Abarenkov, K. (2013). Towards a unified paradigm for sequence-  
738 based identification of fungi. *Molecular Ecology*, *22*, 5271–5277.

739 Kostka, J. E., Weston, D. J., Glass, J. B., Lilleskov, E. A., Shaw, A. J., & Turetsky, M. R.  
740 (2016). The *Sphagnum* microbiome: new insights from an ancient plant lineage. *New*  
741 *Phytologist*, *211*, 57–64

742 Kotiaho, M., Fritze, H., Merilä, P., Tuomivirta, T., Väiliranta, M., Korhola, A., ... Tuittila, E-S.  
743 (2013). Actinobacteria community structure in the peat profile of boreal bogs follows a  
744 variation in the microtopographical gradient similar to vegetation. *Plant and Soil*, *369*, 103-  
745 114.

746 Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in  
747 Linear Mixed Effects Models. *Journal of Statistical Software*. *82*: 1–26.  
748 doi:10.18637/jss.v082.i13.

749 Lamers, L. P. M., van Diggelen, J. M. H., Op den Camp, H. J. M., Visser, E. J. W., Lucassen, E.  
750 C. H. E. T., Vile, M. A., ... Roelofs, J. G. M. (2012). Microbial transformations of nitrogen,  
751 sulfur, and iron dictate vegetation composition in wetlands: a review. *Frontiers in*  
752 *Microbiology*, *3*, 156. doi: 10.3389/fmicb.2012.00156.

753 Lamit, L. J., Romanowicz, K. J., Potvin, L. R., Rivers A. R., Singh, K., Lennon, J. T., ...  
754 Lilleskov, E. A. (2017). Patterns and drivers of fungal community depth stratification in  
755 *Sphagnum* peat. *FEMS Microbiology Ecology*, *93*: doi: 10.1093/femsec/fix082

756 Legendre, P., & Anderson, M. J. (1999). Distance-based redundancy analysis: testing  
757 multispecies responses in multifactorial ecological experiments. *Ecological Monographs*, *69*,  
758 1–24.

759 Lennon J. T., Aanderud Z. T., Lehmkuhl, B. K., & Schoolmaster, D. R. (2012). Mapping the  
760 niche space of soil microorganisms using taxonomy and traits. *Ecology*, *93*, 1867–1879. doi:  
761 10.1890/11-1745.1

762 Lichstein, J. W. (2007). Multiple regression on distance matrices: A multivariate spatial analysis  
763 tool. *Plant Ecology*, *188*, 117–131.

764 Lin, X., Tfaily, M. M., Steinweg, J. M., Chanton. P., Esson. K., Yang, Z.K., ... Kostka, J. E.  
765 (2014). Microbial community stratification linked to utilization of carbohydrates and  
766 phosphorus limitation in a boreal peatland at Marcell Experimental Forest, Minnesota, USA.  
767 *Applied and Environmental Microbiology*, *80*, 3518–3530.

768 Louca, S., Parfrey, L. W., & Doebeli, M. (2016). Decoupling function and taxonomy in the  
769 global ocean microbiome. *Science*, *353*, 1272–1277. doi.org/10.1126/science.aaf4507

770 Malhotra, A., Brice, D.J., Childs, J., Graham, J.D., Hobbie, E.A., Vander Stel, H., Feron, S.C.,  
771 Hanson, P.J. & Iversen, C.M. (2020). Peatland warming strongly increases fine-root growth.  
772 *Proceedings of the National Academy of Sciences*, *117*, 17627-17634.

773 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.  
774 *EMBnet.journal*. *17*, 10-12.

775 Martino, E., Morin, E., Grelet, G-A., Kuo, A., Kohler, A., Daghino, A., ... Perotto, S. (2018).

776 Comparative genomics and transcriptomics depict ericoid mycorrhizal fungi as versatile  
777 saprotrophs and plant mutualists. *New Phytologist*, 217, 1213–1229.

778 Mitsch, W. J., & Gosselink, J. G. (2015). *Wetlands* (5<sup>th</sup> ed). Hoboken, NJ, USA: Wiley.

779 Moomaw, W. R., Chmura, G. L., Davies, G. T. Finlayson, C. M., Middleton, B. A., Natali, S.  
780 M., ... Sutton-Grier, A. E. (2018). Wetlands in a changing climate: science, policy and  
781 management. *Wetlands*, 38, 183-205.

782 Moore, T. R., Bubier, J. L., Frolking, S. E., Lafleur, P. M., & Roulet, N. T. (2002). Plant biomass  
783 and production and CO<sub>2</sub> exchange in an ombrotrophic bog. *Journal of Ecology*, 90, 25–36.

784 Nagendran, S., Hallen-Adams, H. E., Paper, J. M., Aslam, N., Walton, J. D. (2009). Reduced  
785 genomic potential for secreted plant cell-wall-degrading enzymes in the ectomycorrhizal  
786 fungus *Amanita bisporigera*, based on the secretome of *Trichoderma reesei*. *Fungal Genetics  
787 and Biology*, 46, 427-35.

788 Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., ... Kennedy, P.G.  
789 (2016). FUNGuild: an open annotation tool for parsing fungal community datasets by  
790 ecological guild. *Fungal Ecology*, 20, 241–248. doi: 10.1016/j.funeco.2015.06.006

791 Nilsson, R. H., Tedersoo, L., Ryberg, M., Kristiansson, E., Hartmann, M., Unterseher, M., ...  
792 Abarenkov, K. (2015). A comprehensive, automatically updated fungal ITS sequence dataset  
793 for reference-based chimera control in environmental sequencing efforts. *Microbes and  
794 Environments*, 30, 145-150.

795 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., ... Wagner, H.  
796 (2019). vegan: Community Ecology Package. R package version 2.5-6. [https://CRAN.R-  
797 project.org/package=vegan](https://CRAN.R-project.org/package=vegan)

798 Orwin, K. H., Kirschbaum, M. U. F., St John, M. G., & Dickie, I.A. (2011). Organic nutrient  
799 uptake by mycorrhizal fungi enhances ecosystem carbon storage: a model- based assessment.  
800 *Ecology Letters*, 14, 493–502.

801 Potvin, L., Kane, E. S., Chimner, R. A., Kolka, R. K., & Lileskov, E. A. (2015). Effects of water  
802 table position and plant functional group on plant community, aboveground production, and  
803 peat properties in a peat- land mesocosm experiment (PEATcosm). *Plant and Soil*, 387, 277–  
804 294.

805 Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., ... Duchesnay,  
806 E. (2011). Scikit-learn: machine learning in python. *Journal of Machine Learning Research*,  
807 12, 2825–2830.

808 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., ... Glöckner, F. O. (2013).  
809 The SILVA ribosomal RNA gene database project: improved data processing and web-based  
810 tools. *Nucleic Acids Research*, 41, D590-D596.

811 R Core Team. (2018). R: A language and environment for statistical computing. R Foundation  
812 for Statistical Computing, Vienna, Austria. URL. <https://www.R-project.org/>.

813 Read, D. J., Leake, J. R., & Perez-Moreno, J. (2004). Mycorrhizal fungi as drivers of ecosystem  
814 processes in heathland and boreal forest biomes. *Canadian Journal of Botany*, 82, 1243–63.

815 Riley, R., Salamov, A. A., Brown, D. W., Nagy, L. G., Floudas, D., Held, B. W., ... Lindquist  
816 EA. (2013). Extensive sampling of basidiomycete genomes demonstrates inadequacy of the  
817 white-rot/brown-rot paradigm for wood decay fungi. *Proceedings of the National Academy of  
818 Sciences*, 111, 9923-9928.

819 Robroek, B.J.M., Jassey V.E., Kox M.A.R., Berendsen R.L., Mills R.T.E., Cécillon L., ...  
820 Bodelier P.L.E. (2015). Peatland vascular plant functional types affect methane dynamics by  
821 altering microbial community structure. *Journal of Ecology*, 103, 925-934.



822 Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open  
823 source tool for metagenomics. *PeerJ*, 4, e2584. doi: 10.7717/peerj.2584

824 Rupp, D., Kane, E. S., Dieleman, C., Keller, J. K., & Turetsky, M. (2019). Plant functional group  
825 effects on peat carbon cycling in a boreal rich fen. *Biogeochemistry*, 144, 305-327.

826 Russell, L. (2020). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package  
827 version 1.4.5. <https://CRAN.R-project.org/package=emmeans>

828 Rydin, H. & Jeglum, J. K. (2013). *The Biology of Peatlands*, 2<sup>nd</sup> edition.

829 Ben-Shachar, M.S., Makowski, D., & Lüdecke, D. (2020). Compute and interpret indices of  
830 effect size. CRAN. Available from <https://github.com/easystats/effectsize>.

831 Shipley, B. (2000). *Cause and Correlation in Biology: A User's Guide to Path Analysis,*  
832 *Structural Equations and Causal Inference* (1st Edition). Cambridge, UK: Cambridge  
833 University Press.

834 Schlesinger, W. H., & Bernhart, E. S. (2013). *Biogeochemistry: An Analysis of Global Change.*  
835 Cambridge, USA: Academic Press.

836 Tfaily, M. M., Wilson, R. M., Cooper, W. T., Kostka, J. E., Hanson, P., & Chanton, J. P. (2018).  
837 Vertical stratification of peat pore water dissolved organic matter composition in a peat bog  
838 in northern Minnesota. *Journal of Geophysical Research: Biogeosciences*, 123, 479–494.  
839 <https://doi.org/10.1002/2017JG004007>

840 Tedersoo, L., Anslan, S., Bahram, M., Pölme, S., Riit, T., Liiv, I., ... Abarenkov, K. (2015).  
841 Shotgun metagenomes and multiple primer pair-barcode combinations of amplicons reveal  
842 biases in metabarcoding analyses of fungi. *MycKeys*. 10, 1-43.

843 Thormann, M. N., Currah, R. S., & Bayley, S. E. (1999). The mycorrhizal status of the dominant  
844 vegetation along a peatland gradient in southern boreal Alberta, Canada. *Wetlands*, 19, 438–  
845 450.

846 Tremblay, J., Singh, K., Fern, A., Kirton, E. S., He, S. M., Woyke, T., ... Tringe, S. G. (2015).  
847 Primer and platform effects on 16S rRNA tag sequencing. *Frontiers in Microbiology*, 6, 771.  
848 doi: 10.3389 /fmicb.2015.00771

849 Urbanová, Z., & Barta, J. (2016). Effects of long-term drainage on microbial community  
850 composition vary between peatland types. *Soil Biology and Biochemistry*, 92, 16-26.  
851 10.1016/j.soilbio.2015.09.017.

852 van Breemen, N. (1995). How *Sphagnum* bogs down other plants. *Trends in ecology &*  
853 *evolution*, 10, 270-275.

854 Verbruggen, E., Pena, R., Fernandez, C.W., & Soong, J.L. (2017). Mycorrhizal interactions with  
855 saprotrophs and impact on soil carbon storage. In N.C. Johnson, C. Gehring & J. Jansa  
856 (Eds.), *Mycorrhizal mediation of soil: Fertility, Structure, and Carbon Storage* (pp. 441-  
857 460). Elsevier Press. doi: 10.1016/B978-0-12-804312-7.00024-3

858 Wallén, B. (1987). Living roots in hummocks go down to water table, living roots in lawns go  
859 down ~15 below table, to the lowest H<sub>2</sub>O table point. *Holarctic Ecology*, 1987, 10:73–79.

860 Waddington, J. M., Morris, P. J., Kettridge, N., Granath, G., Thompson, D. K., Moore, P. A.  
861 (2015). Hydrological feedbacks in northern peatlands. *Ecohydrology*, 8, 113-127.

862 Wang, M., Tian, J., Bua, Z., Lamit, L.J., Chenc, H., Zhud, Q., & Peng, C. (2019). Structural and  
863 functional differentiation of the microbial community in the surface and subsurface peat of  
864 two minerotrophic fens in China. *Plant and Soil*, 437, 21-40.

865 Ward, S. E., Orwin, K. H., Ostle, N. J., Briones, M. J. I., Thomson, B. C., Griffiths, R. I., ...  
866 Bardgett, R. D. (2015). Vegetation exerts a greater control on litter decomposition than  
867 climate warming in peatlands. *Ecology*, 96, 113–123.

- 868 Weigang, Y., Artz, R. R. E., & Johnson, D. (2008). Species-specific effects of plants colonising  
869 cutover peatlands on patterns of carbon source utilisation by soil microorganism. *Soil*  
870 *Biology and Biochemistry*, 40, 544-549.
- 871 Weishampel, P. S., & Bedford, B. L. (2006). Wetland dicots and monocots differ in colonization  
872 by arbuscular mycorrhizal fungi and dark septate endophytes. *Mycorrhiza*, 16, 495–502. doi:  
873 10.1007/s00572-006-0064-7
- 874 Weltzin, J. F., Bridgman, S. D., Pastor, J., Chen, J., & Harth, C. (2003). Potential effects of  
875 warm- ing and drying on peatland plant community composition. *Global Change Biolog*, 9,  
876 141–51.
- 877 White, T. J., Bruns, T., Lee, S., & Taylor, J. W. (1990). Amplification and direct sequencing of  
878 fungal ribosomal RNA genes for phylogenetics. In M. Innis, D. Gelfand, J. Sninsky & T.  
879 White (Eds.). *PCR protocols: a guide to methods and applications* (pp. 315–322). Orlando,  
880 Florida: Academic Press.
- 881 Zhang, Z., Zimmermann, N. E., Stenke, A., Lin, X., Hodson, E. L., Zhu, G., ... Poulter, B.  
882 (2017). Wetland methane emissions in future climate change. *Proceedings of the National*  
883 *Academy of Sciences*, 114, 9647-9652. doi: 10.1073/pnas.1618765114

884 **Table 1.** PerMANOVA full model results of OTU composition responses treatment factors across years or depths.<sup>†‡§</sup>

Multi-year models	Ericaceae	Sedge	WT	Year	Ericaceae x WT	Sedge x WT	Ericaceae x Year	Sedge x Year	WT x Year	Ericaceae x WT x Year	Sedge x WT x Year
	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$
Fungi	4.05 (1, 15)	1.06 (1, 15)	2.20 (1, 15)	4.62 (2, 54)	0.96 (1, 15)	1.08 (1, 15)	1.39 (2, 36)	0.99 (2, 36)	1.65 (2, 36)	0.98 (2, 36)	0.93 (2, 36)
10-20 cm	<b>&lt;0.001</b> , 15.0	0.382, 2.09	<b>&lt;0.001</b> , 9.4	<b>&lt;0.001</b> , 17.1	0.545, 0.0	0.355, 3.4	<b>0.045</b> , 8.0	0.485, 0.0	<b>0.007</b> , 10.2	0.498, 0.0	0.616, 0.0
Fungi	1.8 (1, 15)	0.76 (1, 16.1)	2.47 (1, 15.4)	4.49 (2, 33)	0.78 (1, 15)	0.77 (1, 16.1)	1.06 (2, 33)	0.96 (2, 33)	1.19 (2, 33)	0.93 (2, 33)	0.89 (2, 33)
30-40 cm	<b>0.016</b> , 8.3	0.805, 0.0	<b>0.002</b> , 11.3	<b>&lt;0.001</b> , 17.0	0.780, 0.0	0.791, 0.0	0.344, 3.2	0.526, 0.0	0.200, 5.5	0.605, 0.0	0.666, 0.0
Prokaryotes	2.07 (1, 15.6)	0.96 (1, 15)	3.28 (1, 15.6)	4.2 (2, 35)	0.99 (1, 15.6)	1.05 (1, 15)	1.13 (2, 35)	0.89 (2, 35)	1.39 (2, 35)	0.93 (2, 35)	0.89 (2, 35)
10-20 cm	<b>0.002</b> , 7.4	0.575, 0.0	<b>&lt;0.001</b> , 10.8	<b>&lt;0.001</b> , 15.1	0.693, 0.0	0.379, 2.2	0.188, 4.3	0.729, 0.0	<b>0.019</b> , 7.5	0.646, 0.0	0.736, 0.0
Prokaryotes	1.74 (2, 15)	0.75 (1, 15)	4.79 (1, 15)	4.49 (2, 36)	0.81 (1, 15)	1.23 (1, 15)	0.87 (6, 36)	0.83 (2, 36)	2.06 (2, 36)	0.99 (2, 36)	0.93 (2, 36)
30-40 cm	0.059, 6.5	0.781, 0.0	<b>&lt;0.001</b> , 14.7	<b>&lt;0.001</b> , 14.8	0.674, 0.0	0.200, 5.2	0.763, 0.0	0.851, 0.0	<b>&lt;0.001</b> , 11.5	0.471, 0.0	0.6245, 0.0

70cm depth gradient models	Ericaceae	Sedge	WT	Depth	Ericaceae x WT	Sedge x WT	Ericaceae x Depth	Sedge x Depth	WT x Depth	Ericaceae x WT x Depth	Sedge x WT x Depth
	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$	<i>F</i> ( <i>df</i> ) <i>P</i> , $\sqrt{\text{Var}}$
Fungi	3.02 (1, 15)	1.22 (1, 15)	1.15 (1, 15)	12.89 (3, 54)	0.74 (1, 15)	0.76 (1, 15)	1.44 (3, 54)	0.916 (3, 54)	2.12 (3, 54)	0.80 (3, 54)	0.88 (3, 54)
	<b>&lt;0.001</b> , 13.2	0.173, 4.4	0.080, 5.5	<b>&lt;0.001</b> , 30.2	0.876, 0.0	0.840, 0.0	<b>0.009</b> , 8.2	0.684, 0.0	<b>&lt;0.001</b> , 13.1	0.910, 0.0	0.764, 0.0
Prokaryotes	1.89 (1, 15)	0.96 (1, 15)	3.37 (1, 15)	18.39 (3, 54)	0.93 (1, 15)	0.80 (1, 15)	1.30 (3, 54)	0.89 (3, 54)	2.37 (3, 54)	0.86 (3, 54)	0.89 (3, 54)
	<b>0.002</b> , 5.8	0.570, 0.0	<b>&lt;0.001</b> , 9.5	<b>&lt;0.001</b> , 33.4	0.636, 0.0	0.856, 0.0	<b>0.015</b> , 6.2	0.831, 0.0	<b>&lt;0.001</b> , 13.2	0.913, 0.0	0.852, 0.0

885  
886 <sup>†</sup> *Ericaceae* = presence/absence Ericaceae, *Sedge* = presence/absence sedges, *WT* = water table manipulation, *Year* = year sampled,  
887 *Depth* = peat sampling depth.  
888 <sup>‡</sup> Models also included individual *mesocosm* as a random effect and *Block* as a fixed effect. No hypothesis test was applied to these  
889 factors.  
890 <sup>§</sup>  $\sqrt{\text{Var}}$  = the square root of the estimated component of variation for each factor. Negative estimates are reported as zero for  
891 simplicity.  
892

893 **Table 2.** Linear mixed model results for the relative abundances of functional groups, total archaea and OTU richness in the 70 cm  
 894 depth gradient sampled in year three.<sup>†‡§</sup>  
 895

<b>Ericaceae</b>	<b>Sedge</b>	<b>WT</b>	<b>Depth</b>	<b>Ericaceae x WT</b>	<b>Sedge x WT</b>	<b>Ericaceae x Depth</b>	<b>Sedge x Depth</b>	<b>WT x Depth</b>	<b>Ericaceae x WT x Depth</b>	<b>Sedge x WT x Depth</b>
$F_{(1,15)} P$	$F_{(1,15)} P$	$F_{(1,15)} P$	$F_{(3,54)} P$	$F_{(1,15)} P$	$F_{(1,15)} P$	$F_{(3,54)} P$	$F_{(3,54)} P$	$F_{(3,54)} P$	$F_{(3,54)} P$	$F_{(3,54)} P$
$\omega_p^2$	$\omega_p^2$	$\omega_p^2$	$\omega_p^2$	$\omega_p^2$	$\omega_p^2$	$\omega_p^2$	$\omega_p^2$	$\omega_p^2$	$\omega_p^2$	$\omega_p^2$
<b>Fungi</b>										
OTU richness										
0.02, 0.890	0.22, 0.647	2.71, 0.120	7.42, <0.001	0.01, 0.910	0.00, 0.975	0.19, 0.901	0.39, 0.763	1.94, 0.134	0.22, 0.879	0.30, 0.822
0.00	0.05	0.09	0.25	0.06	0.00	0.00	0.00	0.05	0.00	0.00
Ericoid mycorrhizal fungi										
18.60, <0.001	2.24, 0.155	4.72, <b>0.046</b>	47.90, <0.001	3.95, 0.065	0.22, 0.646	2.00, 0.124	0.04, 0.989	3.80, <b>0.015</b>	0.64, 0.592	0.37, 0.778
0.51	0.07	0.18	0.71	0.15	0.00	0.05	0.00	0.13	0.00	0.00
Lignin degraders										
11.68, <b>0.004</b>	0.15, 0.708	0.06, 0.813	4.11, <b>0.011</b>	0.07, 0.798	0.11, 0.743	1.13, 0.347	1.25, 0.301	1.70, 0.177	0.34, 0.794	0.23, 0.873
0.39	0.00	0.00	0.14	0.00	0.00	0.01	0.01	0.04	0.00	0.00
<b>Prokaryotes</b>										
OTU richness										
4.13, 0.060	0.04, 0.851	0.16, 0.693	98.88, <0.001	0.85, 0.372	0.04, 0.853	4.2, <b>0.009</b>	0.49, 0.694	2.73, <b>0.053</b>	0.56, 0.647	2.50, 0.069
0.16	0.00	0.00	0.84	0.00	0.00	0.14	0.00	0.08	0.00	0.07
Methanotrophs										
0.01, 0.942	4.98, <b>0.041</b>	0.10, 0.756	7.74, <0.001	9.79, <b>0.007</b>	0.89, 0.361	0.80, 0.502	1.30, 0.284	4.86, <b>0.005</b>	2.63, 0.059	1.12, 0.348
0.00	0.19	0.00	0.26	0.34	0.00	0.00	0.02	0.17	0.08	0.01
Methanogens										
0.41, 0.534	1.88, 0.191	25.26, <0.001	39.20, <0.001	1.58, 0.228	0.94, 0.347	1.28, 0.290	0.62, 0.604	8.17, <0.001	0.60, 0.615	1.17, 0.329
0.00	0.05	0.59	0.66	0.03	0.00	0.01	0.00	0.27	0.00	0.01
Total Archaea										
2.35, 0.146	1.86, 0.193	1.95, 0.183	73.32, <0.001	0.19, 0.669	0.83, 0.378	1.75, 0.168	0.22, 0.881	5.04, <b>0.004</b>	0.76, 0.521	2.08, 0.113
0.07	0.05	0.05	0.79	0.00	0.00	0.04	0.00	0.17	0.00	0.05

896  
 897 <sup>†</sup>All response variables were log10 transformed prior to analysis, except fungal and prokaryote OTU richness.

898 <sup>‡</sup> Models also included individual *mesocosm* bin as a random effect and *Block* as a fixed effect. No hypothesis test was applied to these  
 899 factors.

900 <sup>§</sup>  $\omega_p^2$  = partial- $\omega^2$  values. Negative estimates are reported as zero for simplicity.

901 **Table 3.** PerMANOVA results for community composition responses to treatments within  
 902 individual depths of the 70 cm peat depth gradient sampled in year three.<sup>†‡</sup>  
 903


<b>Taxa and depth of model</b>	<b>Ericaceae</b> <i>F (df)</i> <i>P, √Var</i>	<b>Sedge</b> <i>F (df)</i> <i>P, √Var</i>	<b>WT</b> <i>F (df)</i> <i>P, √Var</i>	<b>Ericaceae x WT</b> <i>F (df)</i> <i>P, √Var</i>	<b>Sedge x WT</b> <i>F (df)</i> <i>P, √Var</i>
<b>Fungi</b>					
00-10cm	3.25 (1, 15) <b>&lt;0.001</b> , 20.03	1.27 (1, 15) 0.163, 6.89	2.31 (1, 15) <b>0.001</b> , 15.27	1.08 (1, 15) 0.367, 5.45	1.04 (1, 15) 0.433, 3.74
10-20cm	3.20 (1, 15) <b>0.002</b> , 20.67	1.32 (1, 15) 0.141, 7.86	1.55 (1, 15) <b>0.051</b> , 10.36	0.73 (1, 15) 0.822, 0.00	0.87 (1, 15) 0.647, 0.00
30-40cm	1.45 (1, 15) 0.105, 9.81	0.84 (1, 15) 0.675, 0.00	1.67 (1, 15) <b>0.051</b> , 12.13	0.78 (1, 15) 0.740, 0.00	0.687 (1, 15) 0.848, 0.00
60-70cm	0.92 (1, 15) 0.5.2, 0.00	0.85 (1, 15) 0.580, 0.00	1.69 (1, 15) 0.094, 12.30	0.56 (1, 15) 0.898, 0.00	0.78 (1, 15) 0.672, 0.00
<b>Prokaryotes</b>					
00-10cm	1.79 (1, 15) <b>0.005</b> , 10.19	1.00 (1, 15) 0.513, 0.00	2.73 (1, 15) <b>&lt;0.001</b> , 15.09	0.96 (1, 15) 0.579, 0.00	0.99 (1, 15) 0.536, 0.00
10-20cm	2.01 (1, 15) <b>0.002</b> , 11.72	1.01 (1, 15) 0.474, 1.05	2.11 (1, 15) <b>0.001</b> , 12.27	0.96 (1, 15) 0.554, 0.00	1.02 (1, 15) 0.462, 2.25
30-40cm	1.11 (1, 15) 0.292, 4.00	0.88 (1, 15) 0.598, 0.00	4.16 (1, 15) <b>&lt;0.001</b> , 21.28	0.92 (1, 15) 0.529, 0.00	0.86 (1, 15) 0.631, 0.00
60-70cm	0.96 (1, 15) 0.429, 0.00	0.78 (1, 15) 0.687, 0.00	1.47 (1, 15) 0.135, 7.52	0.67 (1, 15) 0.832, 0.00	0.58 (1, 15) 0.913, 0.00

904  
 905 <sup>†</sup>*Ericaceae* = presence/absence Ericaceae, *Sedge* = presence/absence sedges, *WT* = water table  
 906 manipulation, *Depth* = peat sampling depth.

907 <sup>‡</sup>  $\sqrt{\text{Var}}$  = the square root of the estimated component of variation for each factor. Negative  
 908 estimates are reported as zero for simplicity.


909  
 910  
 911

912 **Figure 1.** Canonical analysis of principal coordinates (CAP) ordinations with fungal (A, C) and  
 913 prokaryote (B, D) operational taxonomic unit (OTU) composition at the 10-20 cm (A ,B), and  
 914 30-40 cm (C, D) depths sampled in years one, three and four. Ordinations were constrained by  
 915 year, plant functional group treatment (E = Ericaceae only, S = Sedge only, U = Unmanipulated),  
 916 and water table treatment (H = high, L = low). Ellipses represent 95% confidence intervals of the  
 917 ordination points. Dashed lines are provided to show distinction between the strongest effects  
 918 observed in the latter two years (years three + four).  
 919




920

921 **Figure 2.** Canonical analysis of principal coordinates (CAP) ordinations with fungal (A, C) and  
 922 prokaryote (B, D) operational taxonomic unit composition (OTU) across the 70 cm depth  
 923 gradient sampled in year three. Ordinations were first (A, C) conducted by constraining by  
 924 sampling depth, plant functional group treatment (E = Ericaceae only, S = Sedge only, U =  
 925 Unmanipulated) and water table treatment (H = high, L = low), and only by water table and plant  
 926 functional group treatments for datasets within each depth (B, D). Ellipses represent 95%  
 927 confidence intervals of the ordination points.



928


929 **Figure 3.** Marginal means ( $\pm 1$ SE) for relative abundances (Rel. Abund.) of ericoid mycorrhizal  
 930 fungi (A), lignocellulose degrading fungi (B), methanogenic archaea (C) and methanotrophic  
 931 bacteria (D) for the presence/absence (solid lines = presence, dashed lines = absence) of  
 932 Ericaceae (left panel) or Sedges (right panel) by water table treatment (green = low water table,  
 933 blue = high water table) along the 70 cm peat depth gradient. Marginal means are estimated from  
 934 linear mixed models (see Table 2).  
 935  
 936



937  
 938  
 939  
 940



941 **Figure 4.** Structural equation model results for fungi (A) and prokaryotes (B), and associated  
 942 model fit statistics and total effects (C), linking microbial responses to water table (WT  
 943 treatment), plant functional group manipulation (PFG treatment) and the vegetation community  
 944 (Ericaceae composition, Sedge composition). Path widths are scaled proportional to their path  
 945 coefficients (*Rho* or partial-*Rho* values) and are dashed when not significant at an alpha-level of  
 946 0.05. Each variable is represented by a dissimilarity/distance matrix. Separate models were run  
 947 for each depth and each taxonomic group, using samples from year three of the experiment, and  
 948 A and B both represent the combined results of four different models. The core of the models  
 949 (black arrows) were equivalent for all models because the same plant community data was used  
 950 in each; paths and estimates specific to each depth's model are color coded. Models for the 10-20  
 951 cm depth required the addition of a direct path from PFG treatment to fungal or prokaryote  
 952 composition to obtain good model fit, but models at other depth did not require this additional  
 953 path.  
 954



C

Sampling depth (cm)	Fungi			Prokaryotes		
	Model fit (C, df, P)	PFG net effect	WT net effect	Model fit (C, df, P)	PFG net effect	WT net effect
0-10	2.44, 2, 0.294	0.50	0.27	0.08, 2, 0.986	0.17	0.64
10-20	-	0.48	0.12	-	0.17	0.34
30-40	1.07, 2, 0.587	-0.02	0.25	1.64, 2, 0.440	0.03	0.69
60-70	0.96, 2, 0.619	0.00	0.12	0.75, 2, 0.688	-0.04	0.10

955