UCLA

UCLA Previously Published Works

Title

A cross-sectional survey of coronary plaque composition in individuals on non-statin lipid lowering drug therapies and undergoing coronary computed tomography angiography.

Permalink

<https://escholarship.org/uc/item/08d4m86h>

Journal

Journal of cardiovascular computed tomography, 13(2)

ISSN

1934-5925

Authors

Al'Aref, Subhi J
Su, Amanda
Gransar, Heidi
et al.

Publication Date

2019-03-01

DOI

10.1016/j.jcct.2019.01.015

Peer reviewed

Get Access

Share

Export

Journal of Cardiovascular Computed Tomography

Available online 29 January 2019

In Press, Corrected Proof

A cross-sectional survey of coronary plaque composition in individuals on non-statin lipid lowering drug therapies and undergoing coronary computed tomography angiography

Subhi J. Al'Aref^a, Amanda Su^a, Heidi Gransar^c, Alexander R. van Rosendael^a, Asim Rizvi^{a, b}, Daniel S. Berman^c, Tracy Q. Callister^d, Augustin DeLago^e, Martin Hadamitzky^f, Joerg Hausleiter^g, Mouaz H. Al-Mallah^h, Matthew J. Budoffⁱ, Philipp A. Kaufmann^j, Gilbert L. Raff^k, Kavitha Chinnaiyan^k, Filippo Cademartiri^l, Erica Maffei^m, Todd C. Villinesⁿ ...
Fay Y. Lin^a

Show more<https://doi.org/10.1016/j.jcct.2019.01.015>[Get rights and content](#)

Abstract

Introduction

Non-statin therapy (NST) is used as second-line treatment when statin **monotherapy** is inadequate or poorly tolerated.

Objective

To determine the association of NST with **plaque** composition, alone or in combination with **statins**, in patients undergoing coronary computed tomography angiography (coronary CTA).

Methods

From the multicenter CONFIRM registry, we analyzed individuals who underwent coronary CTA with known lipid-lowering therapy status and without prior **coronary artery** disease at baseline. We created a propensity score for being on NST, followed by stepwise multivariate **linear regression**, adjusting for the propensity score as well as risk factors, to determine the association

between NST and the number of coronary artery segments with each plaque type (non-calcified (NCP), partially calcified (PCP) or calcified (CP)) and segment [stenosis](#) score (SSS).

Results

Of the 27,125 subjects in CONFIRM, 4,945 met the inclusion criteria; 371 (7.5%) took NST. At baseline, patients on NST had more prevalent risk factors and were more likely to be on concomitant cardiac medications. After multivariate and propensity score adjustment, NST was not associated with plaque composition: NCP (0.07 increase, 95% CI: -0.05, 0.20; $p=0.26$), PCP (0.10 increase, 95% CI: -0.10, 0.31; $p=0.33$), CP (0.18 increase, 95% CI: -0.10, 0.46; $p=0.21$) or SSS (0.45 increase, 95% CI: -0.02, 0.93; $p=0.06$). The absence of an effect of NST on plaque type was not modified by statin use (p for interaction >0.05 for all).

Conclusion

In this [cross-sectional study](#), non-statin therapy was not associated with differences in plaque composition as assessed by coronary CTA.

Keywords

Coronary computed tomography angiography; Coronary plaque composition; Non-statin therapy; Ezetimibe; Fibrate; Niacin

[Recommended articles](#)

Citing articles (0)

© 2019 Published by Elsevier Inc. on behalf of Society of Cardiovascular Computed Tomography

ELSEVIER [About ScienceDirect](#) [Remote access](#) [Shopping cart](#) [Contact and support](#) [Terms and conditions](#)
[Privacy policy](#)

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the [use of cookies](#).

Copyright © 2019 Elsevier B.V. or its licensors or contributors. ScienceDirect® is a registered trademark of Elsevier B.V.

RELX Group™