

UC San Diego

UC San Diego Previously Published Works

Title

Mental Health Services for Autistic Individuals Across the Lifespan: Recent Advances and Current Gaps

Permalink

<https://escholarship.org/uc/item/08q267qm>

Journal

Current Psychiatry Reports, 23(10)

ISSN

1523-3812

Authors

Maddox, Brenna B
Dickson, Kelsey S
Stadnick, Nicole A
[et al.](#)

Publication Date

2021-10-01

DOI

10.1007/s11920-021-01278-0

Peer reviewed



HHS Public Access

Author manuscript

Curr Psychiatry Rep. Author manuscript; available in PMC 2022 March 29.

Published in final edited form as:

Curr Psychiatry Rep. ; 23(10): 66. doi:10.1007/s11920-021-01278-0.

Mental Health Services for Autistic Individuals Across the Lifespan: Recent Advances and Current Gaps

Brenna B. Maddox¹, Kelsey S. Dickson^{2,3}, Nicole A. Stadnick^{2,4,5}, David S. Mandell⁶, Lauren Brookman-Frazee^{2,4,5,7}

¹Department of Psychiatry, TEACCH Autism Program, University of North Carolina-Chapel Hill, Chapel Hill, NC, USA

²Child and Adolescent Services Research Center, San Diego, CA, USA

³Department of Child and Family Development, San Diego State University, San Diego, CA, USA

⁴Department of Psychiatry, University of California San Diego, San Diego, CA, USA

⁵Dissemination and Implementation Science Center, University of California San Diego Altman Clinical and Translational Research Institute, San Diego, CA, USA

⁶Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

⁷Rady Children's Hospital, San Diego, CA, USA

Abstract

Purpose of Review—This synthesis of recent mental health services research with autistic individuals presents significant advances, current gaps, and recommendations for improving mental healthcare for this population.

Recent Findings—Recent advances include improved understanding of co-occurring mental health conditions among autistic individuals, a growing evidence base for interventions to address them, the development and implementation of new service models to support mental health for this population, and a substantial increase in mental health services and implementation research focused on autism. Ongoing challenges include a lack of mental health interventions designed for community implementation with autistic individuals, limited workforce capacity, complex and disconnected service systems, and racial, ethnic, and socioeconomic disparities in accessibility and quality of mental health services.

Summary—Despite the advances in our understanding of mental health needs and mental health services for autistic individuals, several critical gaps remain. We encourage future efforts to develop and test interventions that can be used in community settings, train and incentivize the

Under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

✉ Brenna B. Maddox, brenna_maddox@med.unc.edu.

Human and Animal Rights and Informed Consent All reported studies/experiments with human or animal subjects performed by the authors have been previously published and complied with all applicable ethical standards (including the Helsinki declaration and its amendments, institutional/national research committee standards, and international/national/institutional guidelines).

workforce to provide them, realign policies and funding with best practice, and embrace an equity-focused approach to autism research and care.

Keywords

Autism; Co-occurring psychiatric conditions; Mental health services; Lifespan; Disparities

Introduction

Autism spectrum disorder (ASD) comprises a constellation of social communication difficulties and restricted, repetitive patterns of behaviors or interests that have strong genetic underpinnings and appear early in life [1]. ASD prevalence continues to rise; the current youth estimate in the USA is 1 in 54 [2]. At the start of this review, we acknowledge the rich and important discussions in the autism research, practice, and self-advocacy communities about preferences for identify-first versus person-first language [3, 4]. To this end, we elected to use identity-first language (e.g., “autistic person”) throughout this review to honor the preferences, autonomy, and rights of the autistic community [5–8].

Autistic individuals across the lifespan commonly experience co-occurring mental health problems that require transdisciplinary care [9, 10]. Attention-deficit hyperactivity disorder, anxiety disorders, sleep-wake disorders, disruptive behavior disorders, and depressive disorders are the most common disorders among autistic individuals [11]. Prevalence is higher in clinically referred samples than in population-based or registry studies; both sets of estimates are significantly greater than what is found in the general population [11]. Co-occurring mental health conditions can be more impairing, distressing, and negatively impactful on quality of life than core autistic characteristics [12]. Although most research in this area to date has focused on autistic youth, mental health problems continue and often increase in adulthood [13–17].

Examining and addressing co-occurring mental health conditions, mental health service needs, evidence-based mental health treatments, and capacity building of mental health services for autistic individuals is an important priority of stakeholders, including autistic individuals, family members, providers, organization and system leaders, and researchers [18, 19]. In this review, we summarize recent advances and continued gaps in quality mental healthcare for autistic individuals across the lifespan. Because publicly funded systems, including Medicaid-reimbursed healthcare and public education services, are the primary payers for autism care in the USA, we focus on findings from research in these systems.

Overview of Mental Health Services

Mental health services comprise a wide range of interventions (both pharmacological and non-pharmacological approaches) and supports (which address challenges to mental health, but may not be directly therapeutic). Examples of mental health services include psychiatric diagnostic evaluation, individual/group/family psychotherapy, case management, psychological testing evaluation, crisis intervention, and consultation. Mental health service providers include psychologists, masters-level therapists, social workers, psychiatrists, psychiatric nurses, primary care physicians, and other allied health professionals. Based on

funding, acuity of mental health needs, client age, and other factors, mental health services can be provided to autistic individuals in a variety of settings (see Table 1).

Recent Research and Practice Advances

Improved Understanding of Co-occurring Mental Health Conditions Among Autistic Individuals Across the Lifespan

While many studies have demonstrated high rates of co-occurring mental health conditions, there has been increased recognition of the *range* of these conditions commonly present in autistic individuals, including the sequelae of trauma and gender dysphoria [11]. There is also growing attention to the increased rates of suicidal thoughts and behaviors in autistic individuals compared with the general population [20], with co-occurring psychiatric conditions documented as a risk factor for suicide attempts and deaths in autistic individuals [21].

As we learn more about the broad range of co-occurring mental health conditions in autistic individuals, there has been an increased focus on the mental health needs among autistic adults and autistic females, two historically understudied and underserved groups. Autistic adults are at disproportionate risk for mental health problems compared with the general population, with more than half meeting criteria for a co-occurring psychiatric condition [14]. A lifespan approach is important because primary co-occurring mental health concerns change across development. For example, challenging behaviors are more common in younger children, while depression and suicidality are more common in adolescents and adults [15, 22]. Emerging research on sex differences suggests that autistic females have higher rates of internalizing disorders than autistic males [23] and experience particularly elevated rates of suicidal thoughts and behaviors [21, 24]. Despite these high rates and associated impairment of co-occurring mental health conditions, autistic adults and autistic females face significant barriers in accessing quality mental health services [16, 25, 26, 27].

Growing Evidence Base for Mental Health Interventions for Autistic Individuals

There have been considerable efforts to develop and test interventions targeting co-occurring mental health conditions in ASD [28, 29, 30]. For youth, this work has taken two general approaches: (1) developing interventions specifically for use with autistic individuals and (2) adapting existing evidence-based interventions (EBIs) targeting mental health conditions in other groups for use with autistic youth [31]. The most common EBI adaptations apply to the intervention techniques, specifically tailoring or adding EBI elements to align with the unique characteristics and needs of the autistic individual (e.g., increasing parent involvement, adding visual supports, incorporating focused interests, using more concrete language) [31]. These modifications are important to improve intervention fit, increase engagement, and promote generalization of skills at home. In addition, the inclusion of EBI adaptations can enhance the efficacy of these interventions, with recent work demonstrating improved efficacy of an EBI (cognitive behavioral therapy) adapted for autistic youth with co-occurring anxiety compared to the standard EBI and treatment as usual [32].

Developing and Implementing New Service Models to Address the Documented Disparities in Access to Services for Autistic Individuals

Navigating mental health services for autistic individuals is extremely complicated, and many families need assistance accessing and engaging in services. Family navigation is an evidence-based case management practice that involves motivational interviewing, service navigation, and collaborative problem solving to increase access to diagnostic and intervention services within a time-limited period. Family navigation has been adapted for ASD [33, 34] and is currently being adapted for ASD and co-occurring mental health needs. Other service delivery models with goals of identifying co-occurring mental health needs and managing or linking to mental healthcare have been adopted in primary care settings. Primary care is well positioned to reach those who may be at most risk of facing health disparities. In addition, primary care may be a less stigmatizing and more holistic place to provide mental healthcare to autistic individuals, similar to collaborative care models for individuals with psychiatric conditions [35]. Recent examples include the Extension for Community Healthcare Outcomes (ECHO) Autism program [36, 37] and the Access to Tailored Autism Integrated Care model [38••, 39], which are both accumulating evidence for feasibility, acceptability, and adoption.

Substantial Increase in Mental Health Services and Implementation Research

Another recent advance is the growing application of implementation science methods, models, and measures to reduce the gap between research and practice in community mental health services. Implementation science has emerged and evolved as a transdisciplinary field to accelerate uptake of evidence-based policies, practices, and programs into routine healthcare systems. It has been applied to autism to address similar quality and access gaps [40, 41]. This has resulted in the field starting to move beyond a “train and hope” model of translating EBIs into community practice. Attention to both the characteristics of EBIs and the use of systematic strategies to implement EBIs is essential for adoption and sustainment in service contexts. A central component of this work is the inclusion of community-partnered and stakeholder-engaged approaches. The close collaboration with relevant community stakeholders helps to ensure that EBIs are not only feasible and acceptable for community settings, but that they address community-identified needs or implementation gaps, for organizations, providers, and autistic individuals. There has also been notable progress in developing and testing implementation strategies at both the provider and organizational level for EBIs with autistic youth in community-based children’s mental health settings [42, 43••, 44]. An additional important methodological feature of these efforts has been the examination of the impact of these EBIs and implementation strategies using hybrid effectiveness-implementation designs that include a simultaneous focus on examining the effects of the clinical EBI as well as key implementation factors, processes, or strategies [45]. This combination of effectiveness and implementation trial designs serves to both speed and improve the translation of research into routine practice and improve the quality of mental health services for autistic individuals. Overall, the application of implementation science methods and frameworks represents a critical advance in the science of translating research to community practice and assuring the significant efforts developing and adapting EBIs to date yield a positive public health impact.

Current Gaps

Limited Representation in EBI Efficacy Studies

Despite the advances in our understanding of mental health needs and mental health services for autistic individuals, several critical gaps remain. Chief among these is the fact that few autistic individuals receive EBIs as part of their mental healthcare [46]. Several factors contribute to the limited implementation of these practices in community settings. A recent review indicates that most mental health interventions for autistic youth were primarily tested with white individuals, those with higher cognitive functioning, and/or those with a single co-occurring mental health condition, with little emphasis on older and/or transition age youth [47]. This narrow target limits the ability to meet the mental health needs of autistic individuals, who typically present with multiple co-occurring conditions [11]. The review also found limited representation of providers typical of the clinical workforce providing community mental health services. Another recent review, specific to cognitive behavior therapy research with autistic youth, found significant overrepresentation of white youth, significant underrepresentation of Black and Latino youth, and significant underrepresentation of families from low socioeconomic status backgrounds in the intervention efficacy trials [48]. Overall, these findings illuminate the critical gap in our understanding of the treatment effects and relevance of these interventions for the clinically and socio-demographically diverse range of individuals typically served in community mental health settings.

In addition, most mental health EBIs were developed and tested in tightly controlled efficacy trials conducted in academic or medical research settings with extensive expertise and resources [31••], versus developed for community implementation by providers with limited autism training and supports. This limits the feasibility and resulting uptake of these interventions in community settings. For example, an EBI may have been designed without consideration of whether the structure (e.g., length of treatment sessions) or the delivery format (e.g., group-based) fits most community service provision models, making them difficult to implement in these settings.

Limited Community Workforce Capacity

Another major barrier to quality mental health services for autistic individuals is the shortage of mental health clinicians trained to work with this population [25•, 49••, 50, 51, 52••, 53]. Mental health clinicians who regularly treat psychiatric conditions such as anxiety and depression are not confident treating these same conditions in autistic clients, given their limited training and experience in this area [25•, 50, 53]. A small subset of mental health clinicians also specializes in autism, but these specialists are often private pay only and/or based in university or urban settings, making them less accessible [52••, 54]. A lack of autism-specific training is most pronounced in adult-focused providers [53], which is partially explained by the historical view of ASD as a childhood condition.

Complex and Disconnected Service Systems

A major challenge in improving mental healthcare for autistic individuals is that no one system has responsibility for providing this care (Table 1). Because mental health

services are delivered in multiple systems, there are complexities with understanding eligibility, accessing care, and aligning this with funding appropriateness [55], particularly as autistic individuals transition from youth-serving systems to adult-serving systems. These complexities are further amplified for autistic individuals who may be additionally eligible for autism-specific services that are typically funded differently than both general healthcare and mental healthcare [56]. For example, autism-specific services such as applied behavior analysis are often coordinated and funded by the Developmental Disability service system in states. From our collective experiences in autism research and practice, we have observed confusion by providers, caregivers, and autistic individuals about the purpose, appropriateness, and logistics of accessing different services, particularly differentiating between applied behavior analysis and mental health services. This confusion may be partially explained by service reimbursement policies that have differentiated eligibility for developmental disability and mental health services [56, 57]. A related challenge for autistic youth with co-occurring mental health needs and their families is the ongoing tension between the education system (where developmental disabilities are often supported) and the healthcare system (where mental health conditions are often managed), making it challenging to connect care [58] and raising questions about how mental health services should be financed and delivered in schools [59].

Racial, Ethnic, and Socioeconomic Disparities

Finally, the gaps and barriers described in this section are exacerbated in racial and ethnic minority and low-income communities [52••, 60–63]. A recent systematic review found racial, ethnic, and socioeconomic disparities in accessibility and quality of services for autistic children [64••]. Specifically, families of color and lower income reported reduced service access and quality and greater unmet service needs than white, higher-resourced families. In a study of Medicaid-enrolled autistic children, Black, Asian, and Native American/Pacific Islander children received fewer outpatient services relative to white children [65••]. In addition, a large statewide survey of autistic adults and their caregivers on a waiting list for home- and community-based Medicaid services found that Black autistic adults had significantly greater unmet needs for mental health services, relative to white autistic adults [66••]. These documented disparities for autistic individuals are not exclusive to mental health services (e.g., allied health services [67], healthcare transition services [68]). More research is needed to identify and test structural changes to increase equity in autistic individuals' access to evidence-based mental healthcare.

Moving Forward

Designing EBIs for Community Delivery From the Outset

To address the problem that EBIs were not designed for community implementation, we need a different approach for mental health intervention development and testing (e.g., “designing for dissemination” [69, 70]). We recommend leveraging implementation science methods, starting early in the pipeline of research-to-practice translation, in order to promote the uptake and sustainment of EBIs in community care [56, 71]. Community-academic partnerships and stakeholder engagement with end users are key aspects of this process [72]. Recent work has further highlighted the importance of listening to the voices of

Author Manuscript

autistic individuals [73] and their caregivers [74] to improve accessibility and quality of services. In addition, investigators conducting intervention research need to explicitly recruit clinically and socio-demographically diverse samples [64••], which will require overcoming barriers to minority families participating in research (e.g., distrust of the research process [75]). More work on cultural adaptations to EBIs for autistic individuals is also needed [76, 77]. These collective efforts are critical to ensure that the mental health intervention evidence base is applicable for autistic individuals and families from diverse and minoritized backgrounds.

Strengthening Mental Health Workforce Capacity

Author Manuscript

Second, to address the limited workforce capacity, we need more mental health clinicians and other professionals willing and able to work effectively with autistic individuals. We recommend adding autism-focused curricula and training opportunities for professionals across primary, specialty, and mental healthcare, at both the preservice level (e.g., in graduate programs through coursework and practicum placements) and in continuing education [25•]. Training content should cover both the assessment and treatment of co-occurring psychiatric conditions in autistic individuals, as well as the assessment and management of suicide risk [78]. Strategies at the system/policy and organizational levels are also needed to address providers' willingness and capacity to accept autistic individuals as clients or patients (e.g., broadening or clarifying service eligibility to include autism in both youth and adult service systems, increasing dissemination of mental health EBIs for autistic individuals [79]). We note that the challenge of limited workforce capacity extends far beyond autism, and improvements will likely require significant changes to preservice training, continuing education, supervision, and reimbursement models.

Restructuring Policies and Coordination Across Service Systems

Author Manuscript

Third, to address the disconnected service systems, we need to integrate autism mental healthcare into existing systems and improve communication between systems. This process will involve developing and testing policy and fiscal innovations to address the currently disjointed organization of public services [56]. Effective communication between systems is critical because autism care necessitates a multiple service system approach — there is no single autism service system [80]. The National Association for Dual Diagnosis [81] offers helpful tools and resources to improve interagency collaboration, particularly between the mental health and developmental disabilities systems.

Structural Change for Promoting Equity

Author Manuscript

Fourth, to increase equitable access to quality mental health services for autistic individuals, we must continue to recognize and address the impact of systemic racism on autism research and clinical practice. Several recent articles offer practical recommendations to enact meaningful changes in policy, research, clinical practice, and implementation science [82–86]. We need to think both about *place* (e.g., school mental health funding tied to property taxes disenfranchises low-income communities) and *race* (e.g., underrepresentation of Black researchers and clinicians in the autism field) to promote equity in autism research, service access, and care quality. By applying a social justice lens to autism research and

clinical practice, we can advocate for systemic change and reduce disparities in mental health services.

Conclusions

The recent advances in mental health services and implementation research for autistic individuals are encouraging. However, significant gaps remain, and much work is still needed to improve the accessibility and quality of mental health services for autistic individuals, particularly those from racial/ethnic minority and lower-income families. This change requires explicit attention to the organization and financing of mental health services. This review summarizes recent research and provides recommendations for addressing key problems, with a primary focus on publicly funded mental health services in the USA. Future work could focus on other service types and international efforts, particularly those in low- and middle-income countries. While there has been considerable progress in recognizing and treating co-occurring psychiatric conditions in autistic individuals, we still need work in developing interventions that can be used in community settings, training and incentivizing the workforce to provide them, and modifying policies and funding so that they align with best practice. Closing these gaps requires a systems- and equity-focused approach to autism research.

Funding

This work was supported by grants from the NIMH (K23 MH115100, PI: Dickson; K23 MH110602, R34 MH120190, PI: Stadnick).

References

Papers of particular interest, published within the past 3 years, have been highlighted as:

• Of importance

•• Of major importance

1. Lord C, Elsabbagh M, Baird G, Veenstra-Vanderweele J. Autism spectrum disorder. *Lancet*. 2018;392:508–20. 10.1016/S0140-6736(18)31129-2. [PubMed: 30078460]
2. Maenner MJ, Shaw KA, Baio J, Washington A, Patrick M, DiRienzo M, et al. Prevalence of autism spectrum disorder among children aged 8 years — Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2016. *MMWR Surveill Summ*. 2020;69:1–12. 10.15585/mmwr.ss6904a1.
3. Vivanti G Ask the editor: what is the most appropriate way to talk about individuals with a diagnosis of autism? *J Autism Dev Disord*. 2020;50:691–3. 10.1007/s10803-019-04280-x. [PubMed: 31676917]
4. Botha M, Hanlon J, Williams GL. Does language matter? identity-first versus person-first language use in autism research: a response to Vivanti. *J Autism Dev Disord*. 2021;1–9. 10.1007/s10803-020-04858-w.
5. Bury SM, Jellett R, Spoor JR, Hedley D. “It defines who I am” or “it’s something I have”: what language do [autistic] Australian adults [on the autism spectrum] prefer? *J Autism Dev Disord*. 2020;1–11. 10.1007/s10803-020-04425-3. [PubMed: 31729599]
6. Kapp SK, Gillespie-Lynch K, Sherman LE, Hutman T. Deficit, difference, or both? Autism and neurodiversity *Dev Psychol*. 2013;49:59–71. 10.1037/a0028353. [PubMed: 22545843]

7. Kenny L, Hattersley C, Molins B, Buckley C, Povey C, Pellicano E. Which terms should be used to describe autism? Perspectives from the UK autism community. *Autism Int J Res Pract*. 2016;20:442–62. 10.1177/1362361315588200.
8. Bottema-Beutel K, Kapp SK, Lester JN, Sasson NJ, Hand BN. Avoiding ableist language: suggestions for autism researchers. *Autism Adulthood*. 2021;3:18–29. 10.1089/aut.2020.0014.
9. Gotham K, Cassidy S, Weiss J. Mental health across the lifespan. *Autism*. 2020;24:805–8. 10.1177/1362361320912132. [PubMed: 32429820]
10. Hossain MM, Khan N, Sultana A, Ma P, McKyer ELJ, Ahmed HU, et al. Prevalence of comorbid psychiatric disorders among people with autism spectrum disorder: an umbrella review of systematic reviews and meta-analyses. *Psychiatry Res*. 2020;287: 112922. 10.1016/j.psychres.2020.112922. [PubMed: 32203749] This umbrella review summarizes 14 systematic reviews and 12 meta-analyses to demonstrate high prevalence rates of co-occurring psychiatric conditions in autistic individuals.
11. Lai M-C, Kasseh C, Besney R, Bonato S, Hull L, Mandy W, et al. Prevalence of co-occurring mental health diagnoses in the autism population: a systematic review and meta-analysis. *Lancet Psychiatry*. 2019;6:819–29. 10.1016/S2215-0366(19)30289-5. [PubMed: 31447415]
12. Lawson LP, Richdale AL, Haschek A, Flower RL, Vartuli J, Arnold SR, et al. Cross-sectional and longitudinal predictors of quality of life in autistic individuals from adolescence to adulthood: the role of mental health and sleep quality. *Autism*. 2020;24:954–67. 10.1177/1362361320908107. [PubMed: 32169010]
13. Hollocks MJ, Lerh JW, Magiati I, Meiser-Stedman R, Brugha TS. Anxiety and depression in adults with autism spectrum disorder: a systematic review and meta-analysis. *Psychol Med*. 2019;49:559–72. 10.1017/S0033291718002283. [PubMed: 30178724]
14. Lugo-Marín J, Magán-Maganto M, Rivero-Santana A, Cuellar-Pompa L, Alviani M, Jenaro-Rio C, et al. Prevalence of psychiatric disorders in adults with autism spectrum disorder: a systematic review and meta-analysis. *Res Autism Spectr Disord*. 2019;59:22–33. 10.1016/j.rasd.2018.12.004.
15. Uljarević M, Hedley D, Rose-Foley K, Magiati I, Cai RY, Dissanayake C, et al. Anxiety and depression from adolescence to old age in autism spectrum disorder. *J Autism Dev Disord*. 2020;50:3155–65. 10.1007/s10803-019-04084-z. [PubMed: 31190198]
16. Vogan V, Lake JK, Tint A, Weiss JA, Lunskey Y. Tracking health care service use and the experiences of adults with autism spectrum disorder without intellectual disability: a longitudinal study of service rates, barriers and satisfaction. *Disabil Health J*. 2017;10:264–70. 10.1016/j.dhjo.2016.11.002. [PubMed: 27899267]
17. Weiss JA, Isaacs B, Diepstra H, Wilton AS, Brown HK, McGarry C, et al. Health concerns and health service utilization in a population cohort of young adults with autism spectrum disorder. *J Autism Dev Disord*. 2018;48:36–44. 10.1007/s10803-017-3292-0. [PubMed: 28900771]
18. Frazier TW, Dawson G, Murray D, Shih A, Sachs JS, Geiger A. Brief report: a survey of autism research priorities across a diverse community of stakeholders. *J Autism Dev Disord*. 2018;48:3965–71. 10.1007/s10803-018-3642-6. [PubMed: 29948533]
19. Pellicano E, Dinsmore A, Charman T. What should autism research focus upon? Community views and priorities from the United Kingdom. *Autism Int J Res Pract*. 2014;18:756–70. 10.1177/1362361314529627.
20. Cassidy S. Suicidality and self-harm in autism spectrum conditions. In: White SW, Maddox BB, Mazefsky CA, editors. *The Oxford handbook of autism and co-occurring psychiatric conditions*. New York, NY: Oxford University Press; 2020. p. 349–68. 10.1093/oxfordhb/9780190910761.013.18.
21. Kølves K, Fitzgerald C, Nordentoft M, Wood SJ, Erlangsen A. Assessment of suicidal behaviors among individuals with autism spectrum disorder in Denmark. *JAMA Netw Open*. 2021;4: e2033565. 10.1001/jamanetworkopen.2020.33565. [PubMed: 33433599]
22. Brookman-Frazee L, Stadnick N, Chlebowski C, Baker-Ericzén M, Ganger W. Characterizing psychiatric comorbidity in children with autism spectrum disorder receiving publicly funded mental health services. *Autism Int J Res Pract*. 2018;22:938–52. 10.1177/1362361317712650.

23. Green RM, Travers AM, Howe Y, McDougle CJ. Women and autism spectrum disorder: diagnosis and implications for treatment of adolescents and adults. *Curr Psychiatry Rep.* 2019;21:22. 10.1007/s11920-019-1006-3. [PubMed: 30852705]
24. Kirby AV, Bakian AV, Zhang Y, Bilder DA, Keeshin BR, Coon H. A 20-year study of suicide death in a statewide autism population. *Autism Res.* 2019;12:658–66. 10.1002/aur.2076. [PubMed: 30663277]
- 25•. Maddox BB, Crabbe S, Beidas RS, Brookman-Frazee L, Cannuscio CC, Miller JS, et al. “I wouldn’t know where to start”: perspectives from clinicians, agency leaders, and autistic adults on improving community mental health services for autistic adults. *Autism.* 2020;24:919–30. 10.1177/1362361319882227. [PubMed: 31674198] This paper provides recommendations for improving community mental health services for autistic adults.
26. Tint A, Weiss JA, Lunsky Y. Identifying the clinical needs and patterns of health service use of adolescent girls and women with autism spectrum disorder. *Autism Res Off J Int Soc Autism Res.* 2017;10:1558–66. 10.1002/aur.1806.
27. Tint A, Weiss JA. A qualitative study of the service experiences of women with autism spectrum disorder. *Autism Int J Res Pract.* 2018;22:928–37. 10.1177/1362361317702561.
28. Keefer A, White SW, Vasa RA, Reaven J. Psychosocial interventions for internalizing disorders in youth and adults with ASD. *Int Rev Psychiatry.* 2018;30:62–77. 10.1080/09540261.2018.1432575. [PubMed: 29537895]
29. Weston L, Hodgekins J, Langdon PE. Effectiveness of cognitive behavioural therapy with people who have autistic spectrum disorders: a systematic review and meta-analysis. *Clin Psychol Rev.* 2016;49:41–54. 10.1016/j.cpr.2016.08.001. [PubMed: 27592496]
- 30•. White SW, Simmons GL, Gotham KO, Conner CM, Smith IC, Beck KB, et al. Psychosocial treatments targeting anxiety and depression in adolescents and adults on the autism spectrum: review of the latest research and recommended future directions. *Curr Psychiatry Rep.* 2018;20:1–10. 10.1007/s11920-018-0949-0. [PubMed: 29368239] This paper synthesizes treatment research related to anxiety and depression in autistic adolescents and adults.
- 31••. Dickson KS, Lind T, Jobin A, Kinnear M, Lok H, Brookman-Frazee L. A systematic review of mental health interventions for ASD: characterizing interventions, intervention adaptations, and implementation outcomes. *Adm Policy Ment Health.* 2021;1–27. 10.1007/s10488-021-01133-7. This review characterizes the types of interventions tested for co-occurring mental health conditions in autistic youth, describes common treatment adaptations, and informs the translation of evidence-based mental health interventions to community service settings.
32. Wood JJ, Kendall PC, Wood KS, Kerns CM, Seltzer M, Small BJ, et al. Cognitive behavioral treatments for anxiety in children with autism spectrum disorder: a randomized clinical trial. *JAMA Psychiatry.* 2020;77:474–83. 10.1001/jamapsychiatry.2019.4160. [PubMed: 31755906]
33. Broder-Fingert S, Stadnick NA, Hickey E, Goupil J, Diaz Lindhart Y, Feinberg E. Defining the core components of family navigation for autism spectrum disorder. *Autism Int J Res Pract.* 2020;24:526–30. 10.1177/1362361319864079.
34. Feinberg E, Augustyn M, Broder-Fingert S, Bennett A, Weitzman C, Kuhn J, et al. Effect of family navigation on diagnostic ascertainment among children at risk for autism: a randomized clinical trial from DBPNet. *JAMA Pediatr.* 2021;175:243–50. 10.1001/jamapediatrics.2020.5218. [PubMed: 33427861]
35. Richardson LP, McCarty CA, Radovic A, Suleiman AB. Research in the integration of behavioral health for adolescents and young adults in primary care settings: a systematic review. *J Adolesc Health.* 2017;60:261–9. 10.1016/j.jadohealth.2016.11.013. [PubMed: 28087267]
36. Mazurek MO, Brown R, Curran A, Sohl K. ECHO autism: a new model for training primary care providers in best-practice care for children with autism. *Clin Pediatr.* 2017;56:247–56. 10.1177/0009922816648288.
37. Mazurek MO, Stobbe G, Loftin R, Malow BA, Agrawal MM, Tapia M, et al. ECHO autism transition: enhancing healthcare for adolescents and young adults with autism spectrum disorder. *Autism Int J Res Pract.* 2020;24:633–44. 10.1177/1362361319879616.
- 38••. Stadnick NA, Brookman-Frazee L, Mandell DS, Kuelbs CL, Coleman KJ, Sahms T, et al. A mixed methods study to adapt and implement integrated mental healthcare for children with autism spectrum disorder. *Pilot Feasibility Stud.* 2019;5:51. 10.1186/s40814-019-0434-5.

[PubMed: 30976456] This paper highlights the need for integrated mental healthcare for youth on the autism spectrum.

39. Stadnick NA, Penalosa MG, Martinez K, Brookman-Frazee L, Gizzo DP, Sahms T, et al. Pre-implementation organizational environment associated with pediatric integrated care readiness in primary care. *Evid-Based Pract Child Adolesc Ment Health*. 2021;0:1–7. 10.1080/23794925.2021.1875344.
40. Brookman-Frazee L, Baker-Ericzen M, Chan J, Dickson K, Rieth S, Haine-Schlagel R, et al. Applying dissemination and implementation science to facilitate community implementation of evidence-based interventions. In: Matson JL, Sturmey P, editors. *Handbook of autism and pervasive developmental disorders*. 2021; in press.
41. Drahota A, Meza RD, Bustos TE, Sridhar A, Martinez JI, Brikho B, et al. Implementation-as-usual in community-based organizations providing specialized services to individuals with autism spectrum disorder: a mixed methods study. *Adm Policy Ment Health Ment Health Serv Res*. 2021;48:482–98. 10.1007/s10488-020-01084-5.
42. Brookman-Frazee L, Stahmer AC. Effectiveness of a multi-level implementation strategy for ASD interventions: study protocol for two linked cluster randomized trials. *Implement Sci*. 2018;13:66. 10.1186/s13012-018-0757-2. [PubMed: 29743090]
- 43••. Brookman-Frazee L, Chlebowski C, Suhrheinrich J, Finn N, Dickson KS, Aarons GA, et al. Characterizing shared and unique implementation influences in two community services systems for autism: applying the epis framework to two large-scale autism intervention community effectiveness trials. *Adm Policy Ment Health*. 2020;47:176–87. 10.1007/s10488-019-00931-4. [PubMed: 30905009] This study demonstrates that provider attitudes and implementation leadership are important for successful implementation in community-based service settings.
44. Stahmer A, Brookman-Frazee L. Utilizing community-based implementation trials to advance understanding of service disparities in autism spectrum disorder. *Glob Pediatr Health*. 2019;6. 10.1177/2333794X19854939.
45. Curran GM, Bauer M, Mittman B, Pyne JM, Stetler C. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care*. 2012;50:217–26. 10.1097/MLR.0b013e3182408812. [PubMed: 22310560]
46. Wood JJ, McLeod BD, Klebanoff S, Brookman-Frazee L. Toward the implementation of evidence-based interventions for youth with autism spectrum disorders in schools and community agencies. *Behav Ther*. 2015;46:83–95. 10.1016/j.beth.2014.07.003. [PubMed: 25526837]
47. Dickson K, Galligan ML, Lock H. Short report: a quantitative methodological review of the literature testing mental health interventions for youth with ASD. 2021; under review.
48. Pickard K, Reyes N, Reaven J. Examining the inclusion of diverse participants in cognitive behavior therapy research for youth with autism spectrum disorder and anxiety. *Autism*. 2019;23:1057–64. 10.1177/1362361318795678. [PubMed: 30160515]
- 49••. Adams D, Young K. A systematic review of the perceived barriers and facilitators to accessing psychological treatment for mental health problems in individuals on the autism spectrum. *Rev J Autism Dev Disord*. 2020;1–18. 10.1007/s40489-020-00226-7. This review highlights common barriers that autistic individuals face when attempting to access mental health services.
50. Brookman-Frazee L, Drahota A, Stadnick N, Palinkas LA. Therapist perspectives on community mental health services for children with autism spectrum disorders. *Adm Policy Ment Health*. 2012;39:365–73. 10.1007/s10488-011-0355-y. [PubMed: 21533846]
51. Camm-Crosbie L, Bradley L, Shaw R, Baron-Cohen S, Cassidy S. “People like me don’t get support”: autistic adults’ experiences of support and treatment for mental health difficulties, self-injury and suicidality. *Autism Int J Res Pract*. 2019;23:1431–41. 10.1177/1362361318816053.
- 52••. Cantor J, McBain RK, Kofner A, Stein BD, Yu H. Fewer than half of US mental health treatment facilities provide services for children with autism spectrum disorder. *Health Aff (Millwood)*. 2020;39:968–74. 10.1377/hlthaff.2019.01557. [PubMed: 32479238] This study demonstrates that a small percentage of US mental health treatment facilities have clinicians with specialized training in autism.
53. Maddox BB, Crabbe SR, Fishman JM, Beidas RS, Brookman-Frazee L, Miller JS, et al. Factors influencing the use of cognitive-behavioral therapy with autistic adults: a survey of community

- mental health clinicians. *J Autism Dev Disord.* 2019;49:4421–8. 10.1007/s10803-019-04156-0. [PubMed: 31385175]
54. Ning M, Daniels J, Schwartz J, Dunlap K, Washington P, Kalantarian H, et al. Identification and quantification of gaps in access to autism resources in the United States: an infodemiological study. *J Med Internet Res.* 2019;21: e13094. 10.2196/13094. [PubMed: 31293243]
55. Garland AF, Hough RL, Landsverk JA, Brown SA. Multi-sector complexity of systems of care for youth with mental health needs. *Child Serv.* 2001;4:123–40. 10.1207/S15326918CS0403_2.
56. Brookman-Frazee L, Drahota A, Chlebowski C, Koenig Y, Williams KN, Hill B, et al. Designing autism spectrum disorder interventions for community implementation: addressing children’s challenging behaviors in publicly funded mental health services. In: White SW, Maddox BB, Mazefsky CA, editors. *The Oxford handbook of autism and co-occurring psychiatric conditions.* New York, NY: Oxford University Press; 2020. p. 409–22. 10.1093/oxfordhb/9780190910761.001.0001.
57. Maddox BB, Gaus VL. Community mental health services for adults: good news and bad news. *Autism Adulthood.* 2019;1:13–7. 10.1089/aut.2018.0006.
58. Malik-Soni N, Shaker A, Luck H, Mullin AE, Wiley RE, Suzanne Lewis ME, et al. Tackling healthcare access barriers for individuals with autism from diagnosis to adulthood. *Pediatr Res.* 2021. 10.1038/s41390-021-01465-y.
59. Kang-Yi CD, Locke J, Marcus SC, Hadley TR, Mandell DS. School-based behavioral health service use and expenditures for children with autism and children with other disorders. *Psychiatr Serv.* 2016;67:101–6. 10.1176/appi.ps.201400505. [PubMed: 26278232]
60. Angell AM, Empey A, Zuckerman KE. Chapter four - a review of diagnosis and service disparities among children with autism from racial and ethnic minority groups in the United States. Hodapp RM, Fidler DJ, editors. *Int Rev Res Dev Disabil.* 2018;55:145–80. 10.1016/bs.irrdd.2018.08.003.
61. McBain RK, Kareddy V, Cantor JH, Stein BD, Yu H. Systematic review: United States workforce for autism-related child healthcare services. *J Am Acad Child Adolesc Psychiatry.* 2020;59:113–39. 10.1016/j.jaac.2019.04.027. [PubMed: 31150751]
62. Singh JS, Bunyak G. Autism disparities: a systematic review and meta-ethnography of qualitative research. *Qual Health Res.* 2019;29:796–808. 10.1177/1049732318808245. [PubMed: 30474491]
63. Drahota A, Sadler R, Hippensteel C, Ingersoll B, Bishop L. Service deserts and service oases: utilizing geographic information systems to evaluate service availability for individuals with autism spectrum disorder. *Autism Int J Res Pract.* 2020;24:2008–20. 10.1177/1362361320931265.
- 64••. Smith KA, Gehricke J-G, Iadarola S, Wolfe A, Kuhlthau KA. Disparities in service use among children with autism: a systematic review. *Pediatrics.* 2020;145:S35–46. 10.1542/peds.2019-1895G. [PubMed: 32238530] This systematic review highlights service-and treatment-related disparities in access and quality of care among children on the autism spectrum.
- 65••. Bilaver LA, Sobotka SA, Mandell DS. Understanding racial and ethnic disparities in autism-related service use among Medicaid-enrolled children. *J Autism Dev Disord.* 2020;1–15. 10.1007/s10803-020-04797-6. [PubMed: 31729599] This study found racial and ethnic disparities in the use of autism-related services, such as outpatient services and case management/care coordination services.
- 66••. Schott W, Nonnemacher S, Shea L. Service use and unmet needs among adults with autism awaiting home- and community-based Medicaid services. *J Autism Dev Disord.* 2021;51:1188–200. 10.1007/s10803-020-04593-2. [PubMed: 32671666] Findings from this state-wide survey of adults and their caregivers on a waiting list for autism waivers highlight significant unmet needs for mental and behavioral health services.
67. Dallman AR, Artis J, Watson L, Wright S. Systematic review of disparities and differences in the access and use of allied health services amongst children with autism spectrum disorders. *J Autism Dev Disord.* 2021;51:1316–30. 10.1007/s10803-020-04608-y. [PubMed: 32683545]
68. Eilenberg JS, Paff M, Harrison AJ, Long KA. Disparities based on race, ethnicity, and socioeconomic status over the transition to adulthood among adolescents and young adults on the autism spectrum: a systematic review. *Curr Psychiatry Rep.* 2019;21:32. 10.1007/s11920-019-1016-1. [PubMed: 30903399]

69. Brownson RC, Jacobs JA, Tabak RG, Hoehner CM, Stamatakis KA. Designing for dissemination among public health researchers: findings from a national survey in the United States. *Am J Public Health*. 2013;103(9):1693–9. 10.2105/AJPH.2012.301165. [PubMed: 23865659]
70. Chambers DA. Sharpening our focus on designing for dissemination: lessons from the SPRINT program and potential next steps for the field. *Transl Behav Med*. 2020;10:1416–8. 10.1093/tbm/ibz102. [PubMed: 31313812]
71. Stadnick NA, Aarons GA, Blake L, Brookman-Frazee LI, Dourgnon P, Engell T, et al. Leveraging implementation science to reduce inequities in children’s mental health care: highlights from a multidisciplinary international colloquium. *BMC Proc*. 2020;14:2. 10.1186/s12919-020-00184-2. [PubMed: 32280371]
72. Pellecchia M, Mandell DS, Nuske HJ, Azad G, Wolk CB, Maddox BB, et al. Community–academic partnerships in implementation research. *J Community Psychol*. 2018;46:941–52. 10.1002/jcop.21981. [PubMed: 30565736]
73. Benevides TW, Shore SM, Palmer K, Duncan P, Plank A, Andresen M-L, et al. Listening to the autistic voice: mental health priorities to guide research and practice in autism from a stakeholder-driven project. *Autism*. 2020;24:822–33. 10.1177/1362361320908410. [PubMed: 32429818]
74. Stahmer AC, Vejnaska S, Iadarola S, Straiton D, Segovia FR, Luelmo P, et al. Caregiver voices: cross-cultural input on improving access to autism services. *J Racial Ethn Health Disparities*. 2019;6:752–73. 10.1007/s40615-019-00575-y. [PubMed: 30859514]
75. Shaia WE, Nichols HM, Dababnah S, Champion K, Garbarino N. Brief report: participation of Black and African-American families in autism research. *J Autism Dev Disord*. 2020;50:1841–6. 10.1007/s10803-019-03926-0. [PubMed: 30805765]
76. Chlebowski C, Magaña S, Wright B, Brookman-Frazee L. Implementing an intervention to address challenging behaviors for autism spectrum disorder in publicly-funded mental health services: therapist and parent perceptions of delivery with Latinx families. *Cultur Divers Ethnic Minor Psychol*. 2018;24:552–63. 10.1037/cdp0000215. [PubMed: 30024185]
77. Chlebowski C, Hurwich-Reiss E, Wright B, Brookman-Frazee L. Using stakeholder perspectives to guide systematic adaptation of an autism mental health intervention for Latinx families: a qualitative study. *J Community Psychol*. 2020;48:1194–214. 10.1002/jcop.22296. [PubMed: 31816103]
78. Jager-Hyman S, Maddox BB, Crabbe SR, Mandell DS. Mental health clinicians’ screening and intervention practices to reduce suicide risk in autistic adolescents and adults. *J Autism Dev Disord*. 2020;50:3450–61. 10.1007/s10803-020-04441-3. [PubMed: 32240486]
79. McBain RK, Cantor JH, Kofner A, Stein BD, Yu H. Brief report: Medicaid expansion and growth in the workforce for autism spectrum disorder. *J Autism Dev Disord*. 2021;1–9. 10.1007/s10803-021-05044-2.
80. Brookman-Frazee L, Baker-Ericzén M, Stahmer A, Mandell D, Haine RA, Hough RL. Involvement of youths with autism spectrum disorders or intellectual disabilities in multiple public service systems. *J Ment Health Res Intellect Disabil*. 2009;2:201–19. 10.1080/19315860902741542. [PubMed: 19809531]
81. National Association for Dual Diagnosis. 2020. <https://www.thenadd.org>. Accessed 18 Apr 2021.
82. Brownson RC, Kumanyika SK, Kreuter MW, Haire-Joshu D. Implementation science should give higher priority to health equity. *Implementation Sci*. 2021;16:1–16. 10.1186/s13012-021-01097-0.
83. Jones DR, Mandell DS. To address racial disparities in autism research, we must think globally, act locally. *Autism*. 2020;24:1587–9. 10.1177/1362361320948313. [PubMed: 32998555]
84. Jones DR, Nicolaidis C, Ellwood LJ, Garcia A, Johnson KR, Lopez K, et al. An expert discussion on structural racism in autism research and practice. *Autism Adulthood*. 2020;2:273–81. 10.1089/aut.2020.29015.dj.
85. Giwa Onaiwu M “They don’t know, don’t show, or don’t care”: autism’s white privilege problem. *Autism Adulthood*. 2020;2:270–2. 10.1089/aut.2020.0077.
86. Shelton RC, Adsul P, Oh A. Recommendations for addressing structural racism in implementation science: a call to the field. *Ethn Dis*. 2021;31:357–64. 10.18865/ed.31.S1.357. [PubMed: 34045837]

Service settings where mental health services may be provided to autistic individuals across the lifespan (based on [55] and [80]). Bolded settings have been the focus of recent research; italicized settings have been understudied

Table 1

Mental health service settings	Childhood (5–15 years)	Young adulthood (16–24 years)	Adulthood (25 years+)
Schools (elementary, secondary)	X	X	
Schools (postsecondary)		X	X
Outpatient	X	X	X
<i>Intensive Outpatient/Partial Hospitalization</i>	X	X	X
<i>Community Emergency Response</i>	X	X	X
<i>Emergency Departments/Crisis Stabilization Units</i>	X	X	X
<i>Inpatient</i>	X	X	X
<i>Primary Care</i>	X	X	X
<i>Alcohol and Drug Services</i>	X	X	X
<i>Juvenile Justice System</i>	X	X	
<i>Criminal Justice System</i>			X