

# Real-time conformational changes in LacY

Irina Smirnova<sup>a</sup>, Vladimir Kasho<sup>a</sup>, and H. Ronald Kaback<sup>a,b,c,1</sup>

Departments of <sup>a</sup>Physiology and <sup>b</sup>Microbiology, Immunology, and Molecular Genetics, and <sup>c</sup>Molecular Biology Institute, University of California, Los Angeles, CA 90095-7327

Contributed by H. Ronald Kaback, May 7, 2014 (sent for review May 1, 2014)

**Galactoside/H<sup>+</sup> symport across the cytoplasmic membrane of *Escherichia coli* is catalyzed by lactose permease (LacY), which uses an alternating access mechanism with opening and closing of deep cavities on the periplasmic and cytoplasmic sides. In this study, conformational changes in LacY initiated by galactoside binding were monitored in real time by Trp quenching/unquenching of bimane, a small fluorophore covalently attached to the protein. Rates of change in bimane fluorescence on either side of LacY were measured by stopped flow with LacY in detergent or in proteoliposomes and were compared with rates of galactoside binding. With LacY in proteoliposomes, the periplasmic cavity is tightly sealed and the substrate-binding rate is limited by the rate of opening of this cavity. Rates of opening, measured as unquenching of bimane fluorescence, are 20–30 s<sup>-1</sup>, independent of sugar concentration and essentially the same in detergent or in proteoliposomes. On the cytoplasmic side of LacY in proteoliposomes, slow bimane quenching (i.e., closing of the cavity) is observed at a rate that is also independent of sugar concentration and similar to the rate of sugar binding from the periplasmic side. Therefore, opening of the periplasmic cavity not only limits access of sugar to the binding site of LacY but also controls the rate of closing of the cytoplasmic cavity.**

membrane transport proteins | major facilitator superfamily

**A**lthough documented extensively with the lactose permease from *Escherichia coli* (LacY) only (reviewed in refs. 1, 2), alternating access is now generally accepted as the overall mechanism of transport for many membrane transport proteins. LacY, a member of the major facilitated superfamily (3), catalyzes the coupled stoichiometric transport of an H<sup>+</sup> and a galactopyranoside (galactoside/H<sup>+</sup> symport) across the cytoplasmic membrane (reviewed in ref. 4). With an abundance of biochemical, spectroscopic, and crystallographic data regarding structure and function, LacY is arguably the most extensively studied symport protein at the present time (reviewed in ref. 5).

LacY is organized into two pseudosymmetrical bundles, each containing six transmembrane helices, most of which are irregular, with the N and C termini on the cytoplasmic side of the membrane. WT LacY and a conformationally restricted mutant exhibit an inward-facing conformation with a tightly sealed periplasmic side and a water-filled cavity open to the cytoplasm (6–9), which is the conformation present in the membrane in the absence of sugar (10). Another conformation has been observed recently with a double-Trp mutant (11) that exhibits an occluded galactoside molecule with a narrow opening on the periplasmic side and a tightly sealed cytoplasmic aspect (12) (Fig. 14).

LacY is highly dynamic and exhibits multiple conformations at any given time, and galactoside binding triggers a shift between conformers (1). Thus, measurement of interspin distances with nitroxide-labeled Cys pairs in LacY reveals that sugar binding induces a decrease in distances on the cytoplasmic side and a corresponding increase in distances on the periplasmic side (13, 14). Site-directed alkylation of single Cys LacY mutants in either right-side-out membrane vesicles (15, 16) or dodecyl- $\beta$ -D-maltopyranoside (DDM) micelles (10), as well as single-molecule fluorescence (17) and thiol cross-linking (18), also indicates that sugar binding increases the probability of opening on the periplasmic side and closing on the cytoplasmic side.

Recently, conformational changes in LacY have been studied dynamically by site-directed Trp fluorescence quenching by a protonated His or Lys residue (19). Sugar binding leads to unquenching of Trp fluorescence in periplasmic LacY mutants N245W (helix VII) or F378W (helix XII) due to increased distance from native quenchers His35 (helix I) or Lys42 (helix II), respectively, findings consistent with opening of the periplasmic cavity. Rates of unquenching of Trp measured in DDM by stopped flow for both mutants reveal rapid opening of the periplasmic cavity. In contrast, on the cytoplasmic side, slow quenching of Trp140 (helix V) by His334 (helix X) is observed with double-mutant F140W/F334H, which indicates closing of the cytoplasmic cavity.

In proteoliposomes, sugar-binding rates measured directly for mutant N245W with 4-nitrophenyl- $\alpha$ -D-galactopyranoside (NPG) by Trp151→NPG FRET are independent of sugar concentration and very similar (56 s<sup>-1</sup>) to rates of opening of the periplasmic cavity measured in DDM micelles (50–100 s<sup>-1</sup>) by Trp unquenching as described above. Thus, opening of the periplasmic cavity is limiting for access to the galactoside-binding site with LacY reconstituted into proteoliposomes (20). However, utilization of Trp fluorescence quenching to measure rates of conformational change in proteoliposomes is challenging because of low signal amplitude. Therefore, the small fluorescence probe bimane, which is comparable in size to Trp, was selected to increase the signal-to-noise ratio for measurements in proteoliposomes. Labeling of Cys residues with (2-pyridyl)dithiobimane (PDT-bimane) generates a fluorescence probe with a short disulfide linker that is cleavable by thiol reagents, such as Tris(2-carboxyethyl)phosphine (TCEP), and reports short-range interactions in proteins by Trp-induced bimane quenching (21, 22). The efficiency of bimane quenching is dependent upon proximity to Trp as well as other factors, such as stereochemistry and local environment. The magnitude of bimane quenching by Trp in membrane proteins is typically a 10–50% change in the emission spectra, as reported with visual rhodopsin (23, 24), the cyclic nucleotide-gated ion channel (25), or the  $\beta$ -adrenoreceptor (26).

## Significance


**The lactose permease from *Escherichia coli* (LacY), a model for the major facilitator superfamily, catalyzes the symport of a galactopyranoside and an H<sup>+</sup> across the membrane by a mechanism in which the sugar-binding site in the middle of the protein becomes alternately accessible to either side of the membrane. The global conformational change is dissected into events that occur on the cytoplasmic and periplasmic aspects of LacY. Rates of individual steps are measured directly during opening or closing of periplasmic or cytoplasmic cavities by utilizing changes in Trp-bimane fluorescence with LacY in a phospholipid membrane. The findings provide a better understanding of the alternating access mechanism.**

Author contributions: I.S., V.K., and H.R.K. designed research; I.S. and V.K. performed research; I.S. and V.K. contributed new reagents/analytic tools; I.S., V.K., and H.R.K. analyzed data; and I.S., V.K., and H.R.K. wrote the paper.

The authors declare no conflict of interest.

<sup>1</sup>To whom correspondence should be addressed. E-mail: rkaback@mednet.ucla.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1408374111/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1408374111/-DCSupplemental).



**Fig. 1.** Detection of conformational changes in LacY by Trp quenching of bimane fluorescence. (A) Side view of LacY with cytoplasmic- or periplasmic-open cavities [Protein Data Bank (PDB) ID code 2CFQ (Left) and PDB ID code 4OAA (Right)]. Transmembrane helices are rainbow-colored from blue (helix I) to red (helix XII), and a bound galactoside (orange spheres) is shown at the apex of the cavities. Pairs of Cys-Trp replacements introduced individually on opposite sides of the periplasmic or cytoplasmic cavities are shown with the indicated positions of C $\alpha$  atoms of Trp (pink spheres) and PDT-bimane-labeled Cys (green spheres) residues. Blue arrows show the direction of movement. (B) Quenching of bimane fluorescence on the cytoplasmic side of LacY observed with bimane-labeled N137C/Q340W mutant. a.u., arbitrary units. (C) Unquenching of bimane fluorescence on the periplasmic side of LacY observed with bimane-labeled I32C/N245W mutant. Blue and red lines represent bimane emission spectra before and after addition of 10 mM TDG, respectively. Spectra were recorded with excitation at 380 nm with 0.3  $\mu$ M protein in 50 mM NaP/0.02% DDM (pH 7.5).


In this paper, mutants were constructed with Trp-Cys pairs in regions of LacY where movements are predicted due to conformational changes. Mutant proteins were then purified, labeled with PDT-bimane, and examined for Trp quenching of bimane fluorescence. The majority of the mutants exhibit phenomena that are entirely consistent with the alternating access mechanism. Two PDT-bimane-labeled mutants that have sugar-binding kinetics typical of WT LacY were used for stopped-flow measurements of rates of conformational change on either the periplasmic or cytoplasmic side of LacY reconstituted into proteoliposomes. The findings demonstrate that the rate of opening of the periplasmic cavity is limiting for sugar binding to reconstituted LacY and also defines the rate of closing of the cytoplasmic cavity.

## Results


**Effect of Galactoside Binding on Bimane Fluorescence.** Closing or opening of cytoplasmic or periplasmic cavities, respectively, in LacY was examined initially by galactoside-induced Trp-bimane quenching or unquenching in steady-state fluorescence experiments (21) with various PDT-bimane-labeled mutants containing Cys-Trp pairs. Cys137 (helix IV) and Trp340 (helix X) were placed on opposing sides of the cytoplasmic cavity in N137C/Q340W LacY (Fig. 1A). The PDT-bimane-labeled mutant exhibits a decrease in fluorescence upon binding of  $\beta$ -D-galactopyranosyl-1-thio- $\beta$ -D-galactopyranoside (TDG), indicating closing of the cavity (Fig. 1B). In contrast, on the periplasmic aspect, PDT-bimane-labeled mutant I32C/N245W exhibits an increase in fluorescence upon TDG binding (Fig. 1C), indicating opening of the periplasmic cavity and increasing distance between Cys32 (helix I) and Trp245 (helix VII). With both paired mutants, the change in bimane fluorescence is not associated with a spectral shift of the emission maximum, suggesting that there is no change in polarity around fluorophore. Similar effects of TDG binding, quenching on the cytoplasmic side and unquenching on the periplasmic side, are seen with several additional cytoplasmic and periplasmic bimane-Trp pairs (Fig. S1). The observations confirm and extend previous findings with Trp as a fluorescent probe as described above (19).

The change in bimane fluorescence with each mutant does not exceed 40%, and the relatively small effects are most likely associated with high flexibility of LacY and the presence of multiple conformers (13). The extent of bimane quenching by an engineered Trp residue also depends on possible background quenching by native amino acyl side chains (mostly Trp and Tyr). Therefore, TCEP reduction experiments were carried out in the absence of sugar to test the proximity between bimane and quenchers in each mutant. TCEP reduction of the S-S bond between bimane and Cys releases the fluorophore and markedly increases fluorescence if a quencher is in close proximity to the attached bimane (21). Indeed, when a PDT-bimane-labeled Cys residue is on the periplasmic side of LacY in the absence of the Trp replacement, TCEP reduction does not significantly alter bimane emission (Fig. S2A and C). However, introduction of Trp on the opposing six-helix bundle results in a dramatic increase in bimane fluorescence after TCEP reduction, indicating that the engineered Trp residue is primarily responsible for bimane quenching (compare Fig. S2A and B, C and D, and E and F) and consistent with tight packing on the periplasmic side. On the cytoplasmic side, an introduced Trp residue has little or no influence on bimane fluorescence in Cys-Trp pairs (compare Fig. S2G and H, I and J, and K and L). Thus, the cytoplasmic cavity is open, and distances between Trp and bimane are too great to cause quenching of the latter (21). Notably, several Cys mutants labeled with bimane and devoid of introduced Trp exhibit significant bimane quenching by native side chains (Fig. S2E, I, and K). In these mutants, the conformational changes triggered by galactoside binding yield relatively small quenching/unquenching of bimane fluorescence by the Trp residue inserted.

**Sugar Binding to PDT-Bimane-Labeled Mutants.** Galactoside binding and conformational rearrangements are interrelated in LacY; thus, it is important that mutants constructed for studying conformational changes exhibit sugar-binding kinetics similar to those of WT LacY. Therefore, sugar binding by bimane-labeled mutants was examined with NPG, a galactoside that participates in Trp151 $\rightarrow$ NPG FRET (27) (Fig. S3A). In addition, NPG triggers bimane quenching/unquenching by an engineered Trp residue (Fig. S3B), as observed with other galactosides (Fig. 1B and C and Fig. S3C). Moreover, the kinetic parameters of NPG binding



**Fig. 2.** Sugar binding to bimane-labeled LacY mutants. NPG-binding affinity was measured with periplasmic mutant I32C/N245W (A) and cytoplasmic mutant N137C/Q340W (B) in displacement experiments with proteins dissolved in DDM (O) or reconstituted into proteoliposomes (PL;  $\square$ ). Stopped-flow traces were recorded after mixing a saturating concentration of TDG (15 mM) with protein preincubated with the indicated concentrations of NPG (Fig. S5). The amplitude of Trp fluorescence change is expressed as a percentage of the final fluorescence level of the individual stopped-flow trace.  $K_d$  values estimated from hyperbolic fits of concentration dependencies of displacement amplitudes are given. Average values of  $k_{off}$  calculated with mutant I32C/N245W from individual experiments at nine NPG concentrations are  $17 \pm 1 \text{ s}^{-1}$  and  $25 \pm 2 \text{ s}^{-1}$  in DDM and PL, respectively. For mutant N137C/Q340W,  $k_{off}$  values calculated from experiments at eight NPG concentrations are  $50 \pm 3 \text{ s}^{-1}$  and  $69 \pm 15 \text{ s}^{-1}$  in DDM and PL, respectively.



**Fig. 3.** Rates of NPG binding measured directly for bimane-labeled mutants of LacY. (A) Stopped-flow traces of Trp fluorescence changes are recorded after mixing of NPG with periplasmic mutant I32C/N245W solubilized in DDM (Lower) or reconstituted into PL (Upper), with single-exponential fits shown as blue and brown lines, respectively. Micromolar concentrations of NPG are indicated. (B) Stopped-flow traces recorded after mixing of indicated concentrations of NPG with cytoplasmic mutant N137C/Q340W reconstituted into PL. Single-exponential fits are shown as brown lines. (C) Stopped-flow traces recorded after mixing of indicated concentrations of NPG with mutant N137C/Q340W in DDM micelles (Lower) or after dissolving the PL in DDM. Single-exponential fits are shown as solid lines. Note the different time scales in B and C. Kinetic parameters of sugar binding estimated from concentration dependences are presented in Figs. 6A and 7A.


(20) are not altered by replacement of highly reactive Cys148 with Met (C148M) (Fig. S4), which is important because the C148M replacement was included to prevent unwanted labeling in the mutants.

Both bimane-labeled Cys-Trp mutants shown in Fig. 1 display NPG-binding affinities similar to WT LacY, as measured by stopped flow in displacement experiments allowing estimation of the displacement or “off rate” ( $k_{\text{off}}$ ) and  $K_d$  for sugar binding (Fig. 2).  $K_d$  values for bimane-labeled I32C/N245W and N137C/Q340W in DDM micelles are 9 and 37  $\mu\text{M}$ , respectively, and 6 and 17  $\mu\text{M}$ , respectively, in proteoliposomes, whereas  $K_d$  values for WT LacY are 28  $\mu\text{M}$  in DDM and 11  $\mu\text{M}$  in proteoliposomes (Fig. S44). NPG  $k_{\text{off}}$  values are also comparable for the mutants and WT LacY when measured either in DDM or in reconstituted proteoliposomes (Fig. S5). Furthermore, as described for WT LacY (20), NPG-binding rates measured directly with the PDT-bimane-labeled mutant I32C/N245W or N137C/Q340W in DDM micelles increase with NPG concentration (Fig. 3, blue lines), whereas in proteoliposomes, binding rates are independent of NPG concentration (Fig. 3A and B, brown lines). Therefore, both mutants exhibit sugar-binding kinetics similar to those of WT LacY, indicating that after bimane labeling of Cys-Trp mutants, the LacY molecules remain reasonably undisturbed.


**Rates of Opening and Closing of the Periplasmic Cavity.** Use of NPG allows simultaneous determination of sugar-binding rates and rates of conformational change by measuring Trp151→NPG FRET and bimane fluorescence changes, respectively. Stopped-flow traces recorded after mixing NPG with bimane-labeled I32C/N245W mutant solubilized in DDM exhibit a rapid decrease in Trp fluorescence [Fig. 4A, green trace; observed rate ( $k_{\text{obs}}$ ) = 112  $\text{s}^{-1}$ ] and a much slower increase in bimane fluorescence (Fig. 4A, gray trace;  $k_{\text{obs}}$  = 26  $\text{s}^{-1}$ ). The data indicate that rapid saturation of the sugar-binding site accessible from the cytoplasmic side ( $K_d$  = 9  $\mu\text{M}$ ; Fig. 2A) triggers a slow shift to a conformer with an open periplasmic cavity, which is detected by bimane unquenching.

Remarkably, a similar experiment with the same mutant reconstituted into proteoliposomes (Fig. 4B) shows practically identical rates of Trp and bimane fluorescence changes ( $k_{\text{obs}}$  = 28  $\pm$  7  $\text{s}^{-1}$ ), indicating that binding of sugar occurs only after the periplasmic cavity opens. Approximately the same rate of unquenching is observed when the conformational change is detected by bimane fluorescence specifically (excitation at 380 nm). Binding of NPG, lactose, or TDG either in DDM micelles or after reconstitution into proteoliposomes results in slow unquenching of bimane fluorescence with  $k_{\text{obs}}$  = 33  $\pm$  11  $\text{s}^{-1}$  (Fig. 4C and D, traces 1, 2, and 3). However, sucrose, which is not a substrate of LacY, has no effect on bimane fluorescence (Fig. 4C, trace 4). Notably, when the mutant is preincubated with NPG and then rapidly diluted into buffer, slow bimane quenching is observed ( $k_{\text{obs}}$  = 27  $\pm$  3  $\text{s}^{-1}$  in both DDM and proteoliposomes) as a result of closing of the periplasmic cavity after sugar release (Fig. 4C, trace 5, and Fig. 4D, trace 4).

Another bimane-labeled periplasmic mutant, K42C/F378W, also exhibits a slow rate of bimane unquenching after binding TDG in DDM micelles (Fig. S64, trace 3;  $k_{\text{obs}}$  = 38  $\text{s}^{-1}$ ), although the small amplitude of the effect does not allow accurate



**Fig. 4.** Rates of opening and closing of the periplasmic cavity measured with bimane-labeled I32C/N245W mutant. (A) Comparison of the NPG-binding rate (Trp151→NPG FRET) with the rate of bimane unquenching for LacY in DDM micelles. Stopped-flow traces after mixing protein with 0.2 mM NPG were recorded simultaneously with excitation at 295 nm, with collection of emitted light from bimane (trace 1, gray) and Trp (trace 2, green) using two photomultipliers. (B) Same approach as described in A was used for protein reconstituted into proteoliposomes. Traces 1 and 2 show fluorescence change of bimane and Trp, respectively, after mixing proteoliposomes with 0.2 mM NPG. (C) Stopped-flow traces of bimane fluorescence recorded with excitation at 380 nm in DDM. Traces 1, 2, 3, and 4 represent mixing protein with 0.2 mM NPG, 20 mM lactose, 10 mM TDG, and 10 mM sucrose, respectively. Trace 5 was recorded after mixing 25  $\mu\text{L}$  of protein containing 40  $\mu\text{M}$  NPG with 250  $\mu\text{L}$  of buffer. (D) Stopped-flow traces of bimane fluorescence recorded with excitation at 380 nm for protein reconstituted into proteoliposomes. Traces 1, 2, and 3 show the effect of 0.2 mM NPG, 20 mM lactose, and 10 mM TDG, respectively. Trace 4 was recorded after 10-fold dilution of protein premixed with 30  $\mu\text{M}$  NPG. Single-exponential fits are shown as blue and brown lines for experiments with protein in DDM micelles or reconstituted into proteoliposomes, respectively. Estimated rates are given in the main text.



**Fig. 5.** Rates of opening and closing of the cytoplasmic cavity measured in DDM with bimane-labeled mutant N137C/Q340W. All traces were recorded with excitation at 295 nm, and emitted light was collected as described in Fig. 4A using two photomultipliers for simultaneous detection of Trp151→NPG FRET (NPG binding or release) and bimane fluorescence (closing or opening of the cytoplasmic cavity). (A) Mixing protein with 0.2 mM NPG results in quenching of bimane fluorescence (Lower, gray trace) and a decrease of Trp fluorescence (Lower, green trace), with the initial part of the trace (0.1-s scale, green arrow) expanded (Upper). A single-exponential fit of bimane fluorescence change (Lower, blue line) allows estimation of the rate of cavity closure ( $k_{obs} = 8.1 \text{ s}^{-1}$ ). The decrease in Trp fluorescence is fitted with a double-exponential equation (red line), which allowed estimation of the NPG-binding rate ( $k_{obs} = 94 \text{ s}^{-1}$ ), followed by a much smaller decrease in amplitude with  $k_{obs} = 7.9 \text{ s}^{-1}$ . (B) Dilution of 25  $\mu\text{L}$  of protein containing 0.2 mM NPG by mixing with 250  $\mu\text{L}$  of buffer results in unquenching of bimane fluorescence (Lower, gray trace) and an increase in Trp fluorescence (Lower, green trace), with the initial portion of the trace (0.1-s scale, green arrow) expanded (Upper). A single-exponential fit of the bimane fluorescence change (Lower, blue line) allows estimation of the rate of cavity opening ( $k_{obs} = 6.2 \text{ s}^{-1}$ ). The increase in Trp fluorescence is fitted with a double-exponential equation (red line), which allowed an estimate of the rate of NPG release ( $k_{obs} = 35 \text{ s}^{-1}$ ), followed by a much smaller increase in amplitude with  $k_{obs} = 6.0 \text{ s}^{-1}$ .

measurement when the mutant protein is reconstituted into proteoliposomes (Fig. S6B, trace 3).


**Rates of Opening and Closing of the Cytoplasmic Cavity.** Simultaneous measurements of the rate of NPG binding and the rate of conformational change were also carried out with PDT-bimane-labeled N137C/Q340W mutant (Fig. 1). Stopped-flow traces recorded after mixing of NPG with the mutant in DDM micelles show a decrease in Trp (Fig. 5A, Trp151→NPG FRET, green trace), as well as bimane fluorescence (Fig. 5A, gray trace), consistent with rapid sugar binding from the cytoplasmic side and slower closure of the cytoplasmic cavity. A double-exponential fit of the Trp fluorescence change (Fig. 5A, red line) allows determination of a major process with large amplitude and a fast rate (Fig. 5A;  $k_{obs} = 94 \text{ s}^{-1}$ ) and a much slower process (Fig. 5A;  $k_{obs} = 7.9 \text{ s}^{-1}$ ) with small amplitude. Remarkably, the rate of the slower component is identical to the rate of conformational change detected by bimane quenching (Fig. 5A, blue line;  $k_{obs} = 8.1 \text{ s}^{-1}$ ). Therefore, the slower rate detected by Trp fluorescence likely reflects the same conformational change initiated by sugar binding as reported by bimane quenching.

When the same mutant is preincubated with 0.2 mM NPG and then diluted into solution devoid of sugar, the opposite process is observed: release of NPG and opening of the cytoplasmic cavity

(Fig. 5B). Thus, simultaneously recorded stopped-flow traces demonstrate an increase in both Trp fluorescence (Fig. 5B, Trp151→NPG FRET, green trace) and bimane fluorescence (Fig. 5B, gray trace). A double-exponential fit of the change in Trp fluorescence (Fig. 5B, red line) resolves a major process with higher amplitude and a faster rate (Fig. 5B;  $k_{obs} = 35 \text{ s}^{-1}$ ) and a slower process (Fig. 5B;  $k_{obs} = 6.0 \text{ s}^{-1}$ ) with lower amplitude. A single-exponential fit of the unquenching of bimane fluorescence (Fig. 5B, blue line;  $k_{obs} = 6.2 \text{ s}^{-1}$ ) demonstrates practically the same rate as the slower component observed by Trp fluorescence. Rapid dissociation of sugar occurs via an open periplasmic cavity and leads to a slower conformational shift toward the structure with an open cytoplasmic cavity.

Comparison of the rates of bimane quenching and unquenching (Fig. 5A and B, Lower) indicates that the cytoplasmic cavity opens and closes with the same rates. Similar rates of conformational change ( $k_{obs} = 3.5\text{--}4.2 \text{ s}^{-1}$ ) were determined by bimane fluorescence specifically (excitation at 380 nm) for mutant N137C/Q340W in DDM micelles (Fig. S7A and B) or reconstituted into proteoliposomes (Fig. S7C).

**Rates of Sugar Binding and Conformational Change.** Concentration dependencies of sugar-binding rates and rates of conformational change measured with bimane-labeled mutants are presented in Fig. 6 (I32C/N245W) and Fig. 7 (N137C/Q340W). Binding of NPG to both mutants in DDM micelles is remarkably similar to that observed with WT LacY [Fig. S4B, blue lines, “on rate” ( $k_{on}$ ) =  $0.2 \mu\text{M}^{-1}\cdot\text{s}^{-1}$ ], exhibiting linear dependence of observed rates on sugar concentration with a  $k_{on}$  estimated as  $0.3\text{--}0.4 \mu\text{M}^{-1}\cdot\text{s}^{-1}$  (Figs. 6A and 7A, blue lines). After reconstitution into proteoliposomes, both mutants display sugar-binding rates that are independent of NPG concentration with  $k_{obs} = 32 \pm 5 \text{ s}^{-1}$  for periplasmic mutant I32C/N245W and  $k_{obs} = 6 \pm 1 \text{ s}^{-1}$  for



**Fig. 6.** Rates of sugar binding and opening of the periplasmic cavity. The data shown were obtained for bimane-labeled I32C/N245W LacY from the stopped-flow measurements described in Figs. 3A and 4 and Fig. S6. (A) Concentration dependencies of NPG-binding rates ( $k_{obs}$ ) were measured by Trp151→NPG FRET with protein solubilized in DDM (blue ●), reconstituted into PL (brown ■), or after dissolving the PL in DDM (◇). Displacement rates ( $k_{off}$ ) for protein in DDM micelles (○) or PL (□) are shown. Kinetic parameters for NPG binding in DDM are  $k_{on} = 0.4 \mu\text{M}^{-1}\cdot\text{s}^{-1}$  and  $k_{off} = 30 \text{ s}^{-1}$ . Reconstituted protein binds sugar with  $k_{obs} = 32 \pm 5 \text{ s}^{-1}$ . (B) Rates of change in bimane fluorescence measured with protein solubilized in DDM are plotted against NPG concentration (○ and □ correspond to excitation at 295 and 380 nm, respectively) and TDG or lactose (▽ and △, respectively, excitation at 380 nm). The pink ■ corresponds to the rate of bimane quenching after dilution of protein preincubated with NPG. The estimated rate of opening/closing of the periplasmic cavity in DDM is  $27 \pm 5 \text{ s}^{-1}$ . (C) Rates of bimane fluorescence unquenching for protein reconstituted into PL are plotted vs. the concentration of NPG (○ and □ correspond to excitation at 295 and 380 nm, respectively) and TDG, or lactose (▽ and △, respectively, excitation at 380 nm). The green ■ corresponds to the rate of bimane quenching after dilution of protein preincubated with NPG. The blue ◆ corresponds to the rate of bimane unquenching after dissolving the PL in DDM. The estimated rate of opening/closing of the periplasmic cavity for the reconstituted protein is  $23 \pm 7 \text{ s}^{-1}$ .

cytoplasmic mutant N137C/Q340W (Figs. 6A and 7A, brown lines), which are comparable to the rate measured for WT LacY (Fig. S4B, brown line,  $k_{\text{obs}} = 21 \pm 4 \text{ s}^{-1}$ ).

Rates of conformational change on the periplasmic or cytoplasmic side of bimane-labeled mutants as a function of sugar concentration are presented in Figs. 6B and C and 7B and C. With periplasmic mutant I32C/N245W, bimane quenching/unquenching rates in DDM display no dependence on sugar concentration and are similar for NPG, TDG, and lactose, with an estimated  $k_{\text{obs}} = 27 \pm 5 \text{ s}^{-1}$  (Fig. 6B). The same rate is observed for this mutant after reconstitution into proteoliposomes ( $k_{\text{obs}} = 23 \pm 7 \text{ s}^{-1}$ ) with the three galactosides (Fig. 6C). Thus, LacY embedded in a phospholipid membrane exhibits the same rates of opening/closing of the periplasmic cavity as observed in DDM. Moreover, the rate of conformational change on the periplasmic side is indistinguishable from the rate of sugar binding to the LacY reconstituted into proteoliposomes (compare brown line in Fig. 6A with data in Fig. 6B and C), indicating that opening of the periplasmic cavity limits the rate of sugar binding to reconstituted LacY.

With the cytoplasmic mutant N137C/Q340W, rates of conformational change measured in DDM and in proteoliposomes are similar ( $k_{\text{obs}} = 4.2\text{--}4.4 \text{ s}^{-1}$ ) and independent of sugar concentration (Fig. 7B and C). The observed rates are similar to sugar-binding rates measured directly by Trp151→NPG FRET with the mutant reconstituted into proteoliposomes (Fig. 7A, brown line). The slow rate of sugar binding with the reconstituted mutant N137C/Q340W is likely due to slow opening of the periplasmic cavity. Hence, opening and closing of the cytoplasmic cavity occurs with similar slow rates in DDM and proteoliposomes.

Slow opening of the periplasmic cavity is also observed with another cytoplasmic Cys-Trp mutant, F140W/V343C, labeled with bimane (Fig. S8). Fast NPG-binding rates are observed in DDM that are linearly dependent on ligand concentration ( $k_{\text{on}} = 0.8 \mu\text{M}^{-1}\text{s}^{-1}$ ), with slow binding in proteoliposomes ( $k_{\text{obs}} = 6 \text{ s}^{-1}$  for all tested NPG concentrations). The conformational change reported by bimane quenching is fast in DDM (maximum  $k_{\text{obs}} = 200 \text{ s}^{-1}$ ) and slow in proteoliposomes ( $k_{\text{obs}} = 5 \text{ s}^{-1}$ ). Clearly, with the mutant reconstituted into proteoliposomes, the periplasmic

cavity opens slowly. This slow process limits the rate of sugar binding, which subsequently defines the rate of closing on the cytoplasmic side as observed with bimane quenching (compare brown lines in Fig. S8D and F).

## Discussion


Distance-dependent quenching of bimane fluorescence by Trp is used here to study global conformational changes in LacY resulting from sugar binding to the symporter solubilized in DDM or reconstituted into proteoliposomes. Sugar-induced opening or closing of the periplasmic and cytoplasmic cavities, respectively, is confirmed by bimane fluorescence unquenching and quenching, which provides a stronger signal than the previous method (20), and the findings provide further support for the alternating access mechanism (reviewed in refs. 1 and 2).

Notably, TCEP reduction confirms that without bound sugar, LacY exists mainly in an inward-facing conformation closed on the periplasmic side, a conclusion derived from the spectra of PDT-bimane-labeled samples before and after reduction, which reveal that Trp residues effectively quench bimane in periplasmic pairs but do not affect fluorescence in cytoplasmic pairs. Also, TCEP reduction demonstrates that in several mutants, background quenching of bimane occurs without the introduced Trp, which results in an apparent decrease in Trp quenching with these engineered Cys-Trp pairs. Nevertheless, Trp-induced bimane fluorescence changes in response to sugar binding are suitable for stopped-flow measurements of conformational changes in LacY in DDM micelles or in reconstituted proteoliposomes.

Detailed analyses of sugar-binding kinetics led to selection of two bimane-labeled mutants, one on each surface of the LacY molecule, that exhibit rates of sugar binding very similar to that of WT LacY. Therefore, rates of opening and closing of the periplasmic and cytoplasmic cavities, respectively, were measured in real time with mutants that dynamically resemble the native transporter. One major conclusion from the studies is that there is a direct correlation between the rate of opening of the periplasmic cavity and the rate of sugar binding to LacY inserted into the membrane.

Furthermore, rates of sugar binding and Trp-induced bimane quenching on both the periplasmic and cytoplasmic sides of LacY allow insight into the individual steps involved in the alternating access mechanism. LacY is inserted into proteoliposomes with the same orientation as in the native bacterial membrane, with a sealed periplasmic aspect facing the external milieu (20, 28, 29). Therefore, sugar binding to LacY in proteoliposomes occurs only after spontaneous opening of the outward-facing cavity, and the rate of sugar binding does not exhibit dependence on sugar concentration.

However, once the periplasmic cavity is open, allowing diffusion-limited access of substrate to the binding site, the rate of binding becomes directly dependent upon galactoside concentration. The  $k_{\text{off}}$  and affinity ( $K_d$ ) measurements obtained from NPG displacement experiments (Fig. 2 and Fig. S5) allow estimation of the sugar  $k_{\text{on}}$  ( $k_{\text{on}} = k_{\text{off}}/K_d$ ) when the sugar-binding site is readily accessible, and they are determined to be 4.1, 4.4, and  $10 \mu\text{M}^{-1}\text{s}^{-1}$  for reconstituted N137C/Q340W, I32C/N245W, and WT LacY, respectively. The observed binding rates ( $k_{\text{obs}} = k_{\text{off}} + k_{\text{on}}[\text{NPG}]$ ) calculated for  $200 \mu\text{M}$  NPG, for example, are from 900 to  $2,100 \text{ s}^{-1}$ , which is much faster than the rates of opening of the periplasmic cavity ( $6\text{--}30 \text{ s}^{-1}$ ), as measured by bimane unquenching on the periplasmic side of LacY. The rate of closing of the periplasmic cavity is similar to the opening rate, as measured by dilution experiments either in DDM (Fig. 4C, trace 5) or in proteoliposomes (Fig. 4D, trace 4). Notably, the rate of conformational change on the periplasmic side is independent of the nature of galactoside, as shown with NPG, TDG, or lactose (Fig. 6B and C). Moreover, rates of opening of the periplasmic cavity are remarkably similar to turnover numbers for WT LacY in right-side-out membrane vesicles or in



**Fig. 7.** Rates of sugar binding and closing of the cytoplasmic cavity. The data shown were obtained for bimane-labeled N137C/Q340W LacY from the stopped-flow measurements described in Figs. 3B and C and 5 and Fig. S7. (A) Concentration dependence of NPG-binding rates ( $k_{\text{obs}}$ ) measured by Trp151→NPG FRET with protein solubilized in DDM (blue ●), reconstituted into PL (brown ■), or after dissolving the PL in DDM (◇). The  $k_{\text{off}}$  values for protein in DDM micelles (○) or PL (□) are shown. Kinetic parameters for NPG binding in DDM are  $k_{\text{on}} = 0.3 \mu\text{M}^{-1}\text{s}^{-1}$  and  $k_{\text{off}} = 43 \text{ s}^{-1}$ . Reconstituted protein binds sugar with  $k_{\text{obs}} = 6.2 \pm 1.1 \text{ s}^{-1}$ . (B) Rates of bimane fluorescence change measured in DDM after mixing protein with NPG (○ and □ with excitation at 295 nm and 380 nm, respectively) or after dilution of protein preincubated with NPG (pink ● and ■ for measurements with excitation at 295 and 380 nm, respectively). The estimated rate of opening/closing of the cytoplasmic cavity in DDM is  $4.4 \pm 2.0 \text{ s}^{-1}$ . (C) Rates of bimane quenching measured after mixing of NPG with protein reconstituted into PL (□, excitation at 380 nm) or with the same PL dissolved in DDM (blue ◆). The estimated rate of cytoplasmic cavity closing for the reconstituted protein is  $4.2 \pm 1.3 \text{ s}^{-1}$ .

reconstituted proteoliposomes with respect to uphill lactose/ $H^+$  symport [i.e., active lactose transport ( $16\text{--}21\text{ s}^{-1}$ )], downhill lactose/ $H^+$  influx ( $8\text{--}16\text{ s}^{-1}$ ), or downhill lactose/ $H^+$  efflux ( $6\text{--}9\text{ s}^{-1}$ ) (30).

Another important conclusion from these studies is that closing of the cytoplasmic cavity in WT LacY is reciprocally related to opening of the periplasmic cavity, as observed with two cytoplasmic mutants reconstituted into proteoliposomes (brown lines in Fig. 7A and C and Fig. S8D and F). Very similar rates are observed for bimane quenching on the cytoplasmic side ( $4\text{--}5\text{ s}^{-1}$ ) and for NPG binding (Trp151 $\rightarrow$ NPG FRET) from outside, the latter of which is limited by the rate of opening of the periplasmic cavity ( $6\text{ s}^{-1}$ ) in both cytoplasmic mutants.

Trp-induced bimane quenching/unquenching, coupled with direct measurements of galactoside binding, provides a powerful tool to study the alternating access mechanism of transport. Rates of global conformational change triggered by sugar binding are dissected into events that occur on the cytoplasmic and periplasmic aspects of LacY in a phospholipid membrane. Clearly, the conformational rearrangements in LacY are initiated by spontaneous opening of the periplasmic cavity, which occurs slowly, thereby limiting the rate of sugar binding; it is possible that this step is also the limiting step in the overall transport process.

## Methods

Construction of mutants, purification of LacY, reconstitution into proteoliposomes, and materials used in this study are described in *SI Methods*.

**PDT-Bimane Labeling.** Labeling of mutants with PDT-bimane was carried out at  $100\text{ }\mu\text{M}$  protein and  $100\text{ }\mu\text{M}$  PDT-bimane in  $50\text{ mM}$  sodium phosphate ( $\text{NaP}$ )/ $0.02\%$  DDM (pH 7.5) for 10 min at room temperature, followed by washing twice with the same buffer without bimane using an Amicon Ultra concentrator with  $50\text{-kDa}$  cutoff (Millipore). TDG ( $30\text{ mM}$ ) was added to the medium for labeling of Cys introduced on the periplasmic side to protect native Cys148 from labeling, and also to increase the probability of opening

the periplasmic cavity. Cys replacements on the cytoplasmic side are readily accessible without bound sugar; therefore, Cys148 was replaced with Met, and labeling was carried out without TDG.

TCEP reduction of PDT-bimane-labeled mutants was carried out at  $10\text{ }\mu\text{M}$  labeled protein in the presence of  $10\text{ mM}$  TCEP (control samples did not contain TCEP) in  $50\text{ mM}$   $\text{NaP}$ / $0.02\%$  DDM (pH 7.5) for 20 min at room temperature. Samples were diluted 25-fold in  $50\text{ mM}$  Na acetate/ $0.02\%$  DDM (pH 4.7) for bimane fluorescence measurements.

**Fluorescence Measurements.** Steady-state fluorescence spectra were measured at room temperature on a SPEX Fluorolog 3 spectrofluorometer (Horiba Scientific) as described (19) with excitation at  $295\text{ nm}$  (for Trp and bimane) and  $380\text{ nm}$  (for bimane). Stopped-flow measurements were performed at  $25\text{ }^\circ\text{C}$  on an SFM-300 rapid kinetic system equipped with a TC-50/10 cuvette (dead time =  $1.2\text{ ms}$ ) and a MOS-450 spectrofluorometer (both from Bio-Logic USA). Excitation was at  $295$  or  $380\text{ nm}$ , with emission interference filters (Edmund Optics) at  $340\text{ nm}$  (for Trp) or  $447\text{ nm}$  (for bimane). The final concentration of protein after mixing was  $0.5\text{--}2\text{ }\mu\text{M}$ . The concentration of TDG in displacement experiments was  $15\text{ mM}$ . Measurements with purified protein solubilized in DDM were done in  $50\text{ mM}$   $\text{NaP}$ / $0.02\%$  DDM (pH 7.5). Experiments with proteoliposomes were carried out in  $50\text{ mM}$   $\text{NaP}$  (pH 7.5). To dissolve proteoliposomes, DDM was added to a final concentration of  $0.3\%$ , and after 10 min, the samples were used in stopped-flow experiments. Typically,  $10\text{--}30$  traces were recorded for each data point, averaged, and fitted with an exponential equation using the built-in Bio-Kine32 software package (Bio-Logic USA) or Sigmaplot 10 (Systat Software, Inc.). All given concentrations were final after mixing unless stated otherwise. For dilution experiments, PDT-bimane-labeled protein, dissolved in DDM or reconstituted into proteoliposomes, was preincubated with NPG at a concentration fivefold higher than the  $K_d$  (measured under identical conditions) and mixed by stopped flow with a 10-fold greater volume of buffer.

**ACKNOWLEDGMENTS.** We thank Junichi Sugihara for his skillful help in preparation of mutants. This work was supported by National Institutes of Health Grants DK51131, DK069463, and GM073210 and National Science Foundation Grant MCB-1129551 (to H.R.K.).

- Smirnova I, Kasho V, Kaback HR (2011) Lactose permease and the alternating access mechanism. *Biochemistry* 50(45):9684–9693.
- Kaback HR, Smirnova I, Kasho V, Nie Y, Zhou Y (2011) The alternating access transport mechanism in LacY. *J Membr Biol* 239(1–2):85–93.
- Saier MH, Jr. (2000) Families of transmembrane sugar transport proteins. *Mol Microbiol* 35(4):699–710.
- Guan L, Kaback HR (2006) Lessons from lactose permease. *Annu Rev Biophys Biomol Struct* 35:67–91.
- Madej MG, Kaback HR (2014) The life and times of lac permease: Crystals ain't enough, but they certainly do help. *Membrane Transporter Function: To Structure and Beyond*, eds Ziegler C, Kraemer R (Springer, Berlin), pp 121–158.
- Abramson J, et al. (2003) Structure and mechanism of the lactose permease of *Escherichia coli*. *Science* 301(5633):610–615.
- Mirza O, Guan L, Verner G, Iwata S, Kaback HR (2006) Structural evidence for induced fit and a mechanism for sugar/ $H^+$  symport in LacY. *EMBO J* 25(6):1177–1183.
- Guan L, Mirza O, Verner G, Iwata S, Kaback HR (2007) Structural determination of wild-type lactose permease. *Proc Natl Acad Sci USA* 104(39):15294–15298.
- Chaptal V, et al. (2011) Crystal structure of lactose permease in complex with an affinity inactivator yields unique insight into sugar recognition. *Proc Natl Acad Sci USA* 108(23):9361–9366.
- Nie Y, Kaback HR (2010) Sugar binding induces the same global conformational change in purified LacY as in the native bacterial membrane. *Proc Natl Acad Sci USA* 107(21):9903–9908.
- Smirnova I, Kasho V, Sugihara J, Kaback HR (2013) Trp replacements for tightly interacting Gly-Gly pairs in LacY stabilize an outward-facing conformation. *Proc Natl Acad Sci USA* 110(22):8876–8881.
- Kumar H, et al. (2014) Structure of sugar-bound LacY. *Proc Natl Acad Sci USA* 111(5):1784–1788.
- Smirnova I, et al. (2007) Sugar binding induces an outward facing conformation of LacY. *Proc Natl Acad Sci USA* 104(42):16504–16509.
- Madej MG, Soro SN, Kaback HR (2012) Apo-intermediate in the transport cycle of lactose permease (LacY). *Proc Natl Acad Sci USA* 109(44):E2970–E2978.
- Kaback HR, et al. (2007) Site-directed alkylation and the alternating access model for LacY. *Proc Natl Acad Sci USA* 104(2):491–494.
- Nie Y, Ermolova N, Kaback HR (2007) Site-directed alkylation of LacY: Effect of the proton electrochemical gradient. *J Mol Biol* 374(2):356–364.
- Majumdar DS, et al. (2007) Single-molecule FRET reveals sugar-induced conformational dynamics in LacY. *Proc Natl Acad Sci USA* 104(31):12640–12645.
- Zhou Y, Guan L, Freitas JA, Kaback HR (2008) Opening and closing of the periplasmic gate in lactose permease. *Proc Natl Acad Sci USA* 105(10):3774–3778.
- Smirnova I, Kasho V, Sugihara J, Kaback HR (2009) Probing of the rates of alternating access in LacY with Trp fluorescence. *Proc Natl Acad Sci USA* 106(51):21561–21566.
- Smirnova I, Kasho V, Sugihara J, Kaback HR (2011) Opening the periplasmic cavity in lactose permease is the limiting step for sugar binding. *Proc Natl Acad Sci USA* 108(37):15147–15151.
- Mansoor SE, Farrens DL (2004) High-throughput protein structural analysis using site-directed fluorescence labeling and the bimane derivative (2-pyridyl)dithiobimane. *Biochemistry* 43(29):9426–9438.
- Mansoor SE, Dewitt MA, Farrens DL (2010) Distance mapping in proteins using fluorescence spectroscopy: The tryptophan-induced quenching (TriQ) method. *Biochemistry* 49(45):9722–9731.
- Janz JM, Farrens DL (2004) Rhodopsin activation exposes a key hydrophobic binding site for the transducin alpha-subunit C terminus. *J Biol Chem* 279(28):29767–29773.
- Tsukamoto H, Farrens DL, Koyanagi M, Terakita A (2009) The magnitude of the light-induced conformational change in different rhodopsins correlates with their ability to activate G proteins. *J Biol Chem* 284(31):20676–20683.
- Islas LD, Zagotta WN (2006) Short-range molecular rearrangements in ion channels detected by tryptophan quenching of bimane fluorescence. *J Gen Physiol* 128(3):337–346.
- Yao X, et al. (2006) Coupling ligand structure to specific conformational switches in the beta2-adrenoceptor. *Nat Chem Biol* 2(8):417–422.
- Smirnova IN, Kasho VN, Kaback HR (2006) Direct sugar binding to LacY measured by resonance energy transfer. *Biochemistry* 45(51):15279–15287.
- Herzlinger D, Viitanen P, Carrasco N, Kaback HR (1984) Monoclonal antibodies against the lac carrier protein from *Escherichia coli*. 2. Binding studies with membrane vesicles and proteoliposomes reconstituted with purified lac carrier protein. *Biochemistry* 23(16):3688–3693.
- Sun J, Wu J, Carrasco N, Kaback HR (1996) Identification of the epitope for monoclonal antibody 4B1 which uncouples lactose and proton translocation in the lactose permease of *Escherichia coli*. *Biochemistry* 35(3):990–998.
- Viitanen P, Garcia ML, Kaback HR (1984) Purified reconstituted lac carrier protein from *Escherichia coli* is fully functional. *Proc Natl Acad Sci USA* 81(6):1629–1633.

# Supporting Information

Smirnova et al. 10.1073/pnas.1408374111


## SI Methods

**Materials.** Oligonucleotides were synthesized by Integrated DNA Technologies, Inc., and restriction enzymes were purchased from New England Biolabs. The QuikChange II kit was purchased from Stratagene. The  $\beta$ -D-galactopyranosyl-1-thio- $\beta$ -D-galactopyranoside was obtained from Carbosynth Limited, and 4-nitrophenyl- $\alpha$ -D-galactopyranoside, D-lactose, and sucrose were from Sigma. Talon superflow resin was purchased from BD Clontech. Dodecyl- $\beta$ -D-maltopyranoside (DDM) and octyl- $\beta$ -D-glucoside (OG) were from Affymetrix. Synthetic phospholipids 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphoethanolamine (POPE) and 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phospho-(1'-*rac*-glycerol) (POPG) were from Avanti Polar Lipids. The (2-pyridyl)dithiobimane was purchased from Toronto Research Biochemicals, and Tris(2-carboxyethyl) phosphine was obtained from Pierce. All other materials were of reagent grade obtained from commercial sources.

**Construction of Mutants, Purification of Lactose Permease from *Escherichia coli*, and Reconstitution into Proteoliposomes.** Construction of mutants, expression in *Escherichia coli*, and purification


of lactose permease from *Escherichia coli* (LacY) were performed as described (1). Periplasmic Cys-Trp pairs were introduced on the WT LacY background, which contains eight native Cys and six native Trp residues. Mutants with cytoplasmic Cys-Trp pairs contained an additional Cys148 with Met (C148M) mutation. All mutants contained a C-terminal 6His-tag that was used for affinity purification with a Talon resin. Purified proteins (10–15 mg/mL) in 50 mM sodium phosphate [ $\text{NaP}_i$ /0.02% DDM (pH 7.5)] were frozen in liquid nitrogen and stored at  $-80^\circ\text{C}$  until use. Reconstitution into proteoliposomes was carried out with synthetic phospholipids (POPE/POPG ratio of 3:1) using the dilution method (2). Briefly, purified LacY in 0.02% DDM solution was mixed with phospholipids dissolved in 1.2% OG to yield a lipid-to-protein ratio of 5 (wt/wt). The mixture was kept on ice for 20 min and then quickly diluted 50-fold in 50 mM  $\text{NaP}_i$  buffer (pH 7.5). Proteoliposomes were harvested by centrifugation for 1 h at  $100,000 \times g$ , suspended in the same buffer, and subjected to two cycles of freeze-thaw/sonication before use.

1. Smirnova I, et al. (2007) Sugar binding induces an outward facing conformation of LacY. *Proc Natl Acad Sci USA* 104(42):16504–16509.
2. Viitanen P, Newman MJ, Foster DL, Wilson TH, Kaback HR (1986) Purification, reconstitution, and characterization of the lac permease of *Escherichia coli*. *Methods Enzymol* 125:429–452.
3. Mansoor SE, Farrens DL (2004) High-throughput protein structural analysis using site-directed fluorescence labeling and the bimeane derivative (2-pyridyl)dithiobimane. *Biochemistry* 43(29):9426–9438.
4. Smirnova IN, Kasho VN, Kaback HR (2006) Direct sugar binding to LacY measured by resonance energy transfer. *Biochemistry* 45(51):15279–15287.
5. Smirnova I, Kasho V, Sugihara J, Kaback HR (2011) Opening the periplasmic cavity in lactose permease is the limiting step for sugar binding. *Proc Natl Acad Sci USA* 108(37):15147–15151.
6. Fersht A (1999) *Structure and Mechanism in Protein Science: A Guide to Enzyme Catalysis and Protein Folding* (Freeman, New York).
7. Eccleston JF, Hutchinson JP, White HD (2001) Stopped-flow techniques. *Protein–Ligand Interactions: Structure and Spectroscopy*, eds Harding SE, Chowdhry BZ (Oxford Univ Press, New York).




**Fig. S1.** Conformational changes on the cytoplasmic and periplasmic sides of bimane-labeled Cys-Trp mutants detected by Trp-induced bimane quenching. Cys and Trp replacements are shown as green and pink spheres, respectively, indicating the positions of  $C_{\alpha}$  atoms on the backbone of the inward-facing structure of LacY (Protein Data Bank ID code 2CFQ). N-terminal and C-terminal six-helix bundles are marked with the letters N and C, respectively. Pairs of Cys-Trp replacements were introduced on opposing sides of the periplasmic or cytoplasmic cavities. Cytoplasmic Cys-Trp pairs are viewed from the cytoplasmic side (*Upper*, with the cavity open to the cytoplasm), and periplasmic pairs are shown from the periplasmic side (*Lower*). The intensity of bimane fluorescence of each (2-pyridyl)dithiobimane-labeled mutant (Trp-bimane) was measured before and after addition of 10 mM  $\beta$ -D-galactopyranosyl-1-thio- $\beta$ -D-galactopyranoside (TDG), and is shown as blue and red lines, respectively. Addition of nongalactosidic sugars had no effect on bimane fluorescence. Emission spectra were recorded with excitation at 380 nm with 0.3–0.5  $\mu$ M protein in 50 mM NaP<sub>i</sub>/0.02% DDM (pH 7.5). (*Upper*) Quenching of bimane after TDG binding to mutants with Trp-bimane pairs on the cytoplasmic side, indicating closing of the cavity, is demonstrated. Two mutants (F140W/V343C and F140W/V331C) were labeled with bimane on the C-terminal six-helix bundle; the other two mutants (R134C/V343W and R135C/V343W) were labeled with bimane on the N-terminal six-helix bundle. (*Lower*) Increase in bimane fluorescence after TDG binding to mutants with Trp-bimane pairs on the periplasmic side, indicating opening of the cavity with unquenching of bimane, is demonstrated. With mutant K42C/F378W, bimane is attached to the N-terminal six-helix bundle. With the other three mutants (F29W/G262C, S43W/N371C, and S43W/M372C), bimane is attached to the C-terminal six-helix bundle. a.u., arbitrary units.







**Fig. S2.** Proximity between Trp and bimane in PDT-bimane-labeled Cys-Trp mutants. The fluorescence intensity of bimane attached to LacY depends on the proximity of the Trp quencher. Reduction of the disulfide bond between bimane and a given Cys residue by Tris(2-carboxyethyl)phosphine (TCEP; scheme at the top) releases the fluorophore and removes the effect of the neighboring quencher(s). The bimane released has high fluorescence at acidic pH (3). The fluorescence of the bimane released remains unchanged only if no quencher is present in close proximity. Periplasmic (*Left; A–F*) and cytoplasmic (*Right; G–L*) Cys mutants (with or without given Trp replacements as indicated) were labeled with PDT-bimane and then treated with TCEP. The fluorescence intensities of bimane before and after TCEP reduction are shown as blue and pink emission spectra, respectively, measured with excitation at 380 nm with 0.4 mM protein in 50 mM sodium acetate/0.02% DDM (pH 4.7). Spectra were normalized to the maximum intensity of bimane-SH after TCEP treatment.



**Fig. S3.** Fluorescence of PDT-bimane-labeled LacY allows detection of sugar binding and/or conformational change. The effect of different galactosidic sugars on the fluorescence of periplasmic mutant I32C/N245W is demonstrated with 4-nitrophenyl- $\alpha$ -D-galactopyranoside (NPG) (*A* and *B*) or TDG (*C*). Addition of NPG, which is the acceptor in Trp151→NPG FRET (4), results in changes in both Trp and bimane fluorescence that allow detection of sugar binding and conformational change independently by using different excitation wavelengths: 295 nm for Trp fluorescence (*A*) and 380 nm for bimane fluorescence (*B*). (*C*) Galactosides that do not participate in FRET (TDG, lactose, or melibiose) were used to detect conformational changes due to sugar binding, which are shown as unquenching of bimane fluorescence in the periplasmic mutant. Note that essentially no change in Trp emission is observed with excitation at 295 nm after TDG addition. In the displacement experiments, excess TDG was used to displace NPG from the sugar-binding site, resulting in an increase in Trp151 emission (*A*, red arrow).




**Fig. 54.** Sugar binding to WT LacY solubilized in DDM or reconstituted into proteoliposomes (PL) measured by stopped flow using Trp151→NPG FRET. Stopped-flow traces of the Trp fluorescence change were recorded with excitation at 295 nm and fitted with a single-exponential equation as described (4, 5). Sugar binding was measured either directly by mixing protein with NPG or, in displacement experiments, by mixing a saturating concentration of TDG with LacY preincubated with NPG. The dependence of directly measured observed rates ( $k_{\text{obs}}$ ) on sugar concentration allowed estimations of the binding rate constant, "on rate" ( $k_{\text{on}}$ ), and dissociation rate constant or "off rate" ( $k_{\text{off}}$ ) ( $k_{\text{obs}} = k_{\text{off}} + k_{\text{on}} [\text{NPG}]$ ). In displacement experiments, mixing a high concentration of fluorescently silent ligand TDG with LacY containing bound NPG results in release of NPG that cannot reassociate with protein because of the saturating concentration of TDG. Therefore, the rate of Trp fluorescence increase (Trp151-NPG FRET) is a true  $k_{\text{off}}$  (6, 7), which does not depend on NPG concentration. (A) NPG-binding affinity determined from the concentration dependence of the amplitude of the Trp fluorescence changes in displacement experiments (at various concentrations of NPG and 15 mM TDG) with LacY in detergent (○) or reconstituted into PL (□). Estimated kinetic parameters are  $k_{\text{off}} = 41 \pm 1 \text{ s}^{-1}$  and  $K_d = 28 \pm 2 \mu\text{M}$  in DDM and  $k_{\text{off}} = 111 \pm 9 \text{ s}^{-1}$  and  $K_d = 11 \pm 2 \mu\text{M}$  in PL. (B) Rates of NPG binding directly by stopped-flow in DDM (●), reconstituted into PL (■), and after dissolving the PL in DDM (◇). The concentration dependence of NPG-binding rates measured with the C148M mutant (▲) is indistinguishable from that determined for WT LacY. Open symbols on the y axis represent rates of NPG displacement by TDG ( $k_{\text{off}}$ ) for WT LacY in DDM and PL (○ and □, respectively) or C148M in DDM (△). The concentration dependencies of the  $k_{\text{obs}}$  for NPG-binding yield estimates of  $k_{\text{on}}$  in DDM of  $0.2 \mu\text{M}^{-1} \text{ s}^{-1}$  for both WT LacY and the C148M mutant (blue lines). For WT LacY reconstituted into PL,  $k_{\text{obs}} = 21 \pm 4 \text{ s}^{-1}$ , as measured at NPG concentrations up to 0.6 mM (brown line).



**Fig. 55.** Rates of displacement of bound NPG by excess of TDG measured with bimane-labeled periplasmic mutant I32C/N245W (A) or cytoplasmic mutant N137C/Q340W (B). Stopped-flow traces of Trp fluorescence change (Trp151→NPG FRET) were recorded after mixing 15 mM TDG with a given mutant preincubated with the indicated concentrations of NPG. The  $k_{\text{off}}$  values and amplitudes of the Trp fluorescence increase were estimated from single-exponential fits to the data obtained in DDM or PL (blue and brown lines, respectively) and presented in Fig. 2.





**Fig. 58.** Rates of sugar binding and closing of the cytoplasmic cavity with PDT-bimane-labeled F140W/V343C LacY. (A) Stopped-flow traces of Trp fluorescence changes after mixing given concentrations of NPG with LacY in DDM micelles [Trp151→NPG FRET, excitation (exc.) at 295 nm]. (B) Stopped-flow traces of bimane fluorescence quenching (excitation at 380 nm) caused by closing the cytoplasmic cavity after mixing given concentrations of NPG or TDG with the protein in DDM micelles. (C) Stopped-flow traces recorded with protein reconstituted into PL show bimane fluorescence quenching after mixing protein with 100  $\mu\text{M}$  NPG (Upper; green trace, excitation at 380 nm) and the decrease in Trp fluorescence (Trp151→NPG FRET, excitation at 295 nm) caused by binding of given concentrations of NPG (Lower; four traces). Note the longer time scale in C relative to A and B. Traces were fitted with a single-exponential equation (black lines), allowing estimation of  $k_{\text{obs}}$ . (D) Concentration dependence of NPG-binding rates measured by Trp151→NPG FRET with protein in DDM as shown in A (●), reconstituted into proteoliposomes as shown in C (■), or after dissolving the PL in DDM (◇). Displacement rates measured with protein in DDM (○,  $k_{\text{off}} = 34 \text{ s}^{-1}$ ) or PL (□,  $k_{\text{off}} = 48 \text{ s}^{-1}$ ) are shown. Kinetic parameters for NPG binding in DDM are  $k_{\text{on}} = 0.8 \pm 0.1 \mu\text{M}^{-1}\text{s}^{-1}$ ,  $k_{\text{off}} = 28 \pm 15 \text{ s}^{-1}$ , and  $K_{\text{d}} = 33 \pm 17 \mu\text{M}$ . Reconstituted protein binds sugar with  $k_{\text{obs}} = 6.1 \pm 0.3 \text{ s}^{-1}$ . (E) Rates of bimane fluorescence quenching measured with protein in DDM, as shown in B, are plotted against sugar concentration and shown for NPG (○) and TDG (▽). The maximum rate of bimane quenching estimated from a hyperbolic fit (blue line) is  $196 \pm 18 \text{ s}^{-1}$ . (F) Rates of bimane fluorescence quenching measured with protein reconstituted into PL are plotted against NPG concentrations (□). The estimated rate of closing of the cytoplasmic cavity with reconstituted protein is  $5.2 \pm 1.3 \text{ s}^{-1}$ .