

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

Please note: Images will appear in color online but will be printed in black and white.

ArticleTitle	Understanding a migratory species in a changing world: climatic effects and demographic declines in the western monarch revealed by four decades of intensive monitoring	
Article Sub-Title		
Article CopyRight	Springer-Verlag Berlin Heidelberg (This will be the copyright line in the final PDF)	
Journal Name	Oecologia	
Corresponding Author	Family Name	Espeset
	Particle	
	Given Name	Anne E.
	Suffix	
	Division	Program in Ecology, Evolution, and Conservation Biology, Department of Biology
	Organization	University of Nevada
	Address	Reno, NV, USA
	Division	
	Organization	
	Address	1664 North Virginia Street, 89509, Reno, NV, USA
	Email	aespeset@nevada.unr.edu
Author	Family Name	Harrison
	Particle	
	Given Name	Joshua G.
	Suffix	
	Division	Program in Ecology, Evolution, and Conservation Biology, Department of Biology
	Organization	University of Nevada
	Address	Reno, NV, USA
	Email	
Author	Family Name	Shapiro
	Particle	
	Given Name	Arthur M.
	Suffix	
	Division	Center for Population Biology
	Organization	University of California
	Address	Davis, CA, USA
	Email	
Author	Family Name	Nice
	Particle	
	Given Name	Chris C.
	Suffix	
	Division	Department of Biology

	Organization	Texas State University
	Address	San Marcos, TX, USA
	Email	
Author	Family Name	Thorne
	Particle	
	Given Name	James H.
	Suffix	
	Division	Department of Environmental Science and Policy
	Organization	University of California
	Address	Davis, CA, USA
	Email	
Author	Family Name	Waetjen
	Particle	
	Given Name	David P.
	Suffix	
	Division	Department of Environmental Science and Policy
	Organization	University of California
	Address	Davis, CA, USA
	Email	
Author	Family Name	Fordyce
	Particle	
	Given Name	James A.
	Suffix	
	Division	Ecology and Evolutionary Biology
	Organization	University of Tennessee
	Address	37996, Knoxville, TN, USA
	Email	
Author	Family Name	Forister
	Particle	
	Given Name	Matthew L.
	Suffix	
	Division	Program in Ecology, Evolution, and Conservation Biology, Department of Biology
	Organization	University of Nevada
	Address	Reno, NV, USA
	Email	
Schedule	Received	9 September 2015
	Revised	
	Accepted	29 February 2016
Abstract	<p>Migratory animals pose unique challenges for conservation biologists, and we have much to learn about how migratory species respond to drivers of global change. Research has cast doubt on the stability of the eastern monarch butterfly (<i>Danaus plexippus</i>) population in North America, but the western monarchs have not been as intensively examined. Using a Bayesian hierarchical model, sightings of western monarchs over approximately 40 years were investigated using summer flight records from ten sites along an elevational transect in Northern California. Multiple weather variables were examined, including local and regional temperature and precipitation. Population trends from the ten focal sites and a subset of western overwintering sites were compared to summer and overwintering data from the eastern migration.</p>	

Records showed western overwintering grounds and western breeding grounds had negative trends over time, with declines concentrated early in the breeding season, which were potentially more severe than in the eastern population. Temporal variation in the western monarch also appears to be largely independent of (uncorrelated with) the dynamics in the east. For our focal sites, warmer temperatures had positive effects during winter and spring, and precipitation had a positive effect during spring. These climatic associations add to our understanding of biotic-abiotic interactions in a migratory butterfly, but shifting climatic conditions do not explain the overall, long-term, negative population trajectory observed in our data.

Keywords (separated by '-') Climate change - *Danaus plexippus* - Hierarchical model - Monarch butterfly - Western population

Footnote Information Communicated by Klaus Fischer.

Electronic supplementary material The online version of this article (doi:10.1007/s00442-016-3600-y) contains supplementary material, which is available to authorized users.

2 **Understanding a migratory species in a changing world: climatic**
3 **effects and demographic declines in the western monarch revealed**
4 **by four decades of intensive monitoring**

5 Anne E. Espeset^{1,6} · Joshua G. Harrison¹ · Arthur M. Shapiro² · Chris C. Nice³ ·
6 James H. Thorne⁴ · David P. Waetjen⁴ · James A. Fordyce⁵ · Matthew L. Forister¹

7 Received: 9 September 2015 / Accepted: 29 February 2016
8 © Springer-Verlag Berlin Heidelberg 2016

9 **Abstract** Migratory animals pose unique challenges
10 for conservation biologists, and we have much to learn
11 about how migratory species respond to drivers of global
12 change. Research has cast doubt on the stability of the east-
13 ern monarch butterfly (*Danaus plexippus*) population in
14 North America, but the western monarchs have not been
15 as intensively examined. Using a Bayesian hierarchical
16 model, sightings of western monarchs over approximately
17 40 years were investigated using summer flight records
18 from ten sites along an elevational transect in Northern
19 California. Multiple weather variables were examined,
20 including local and regional temperature and precipitation.
21 Population trends from the ten focal sites and a subset of
22 western overwintering sites were compared to summer and

overwintering data from the eastern migration. Records 23
showed western overwintering grounds and western breed- 24
ing grounds had negative trends over time, with declines 25
concentrated early in the breeding season, which were 26
potentially more severe than in the eastern population. 27
Temporal variation in the western monarch also appears to 28
be largely independent of (uncorrelated with) the dynam- 29
ics in the east. For our focal sites, warmer temperatures had 30
positive effects during winter and spring, and precipitation 31
had a positive effect during spring. These climatic associa- 32
tions add to our understanding of biotic-abiotic interactions 33
in a migratory butterfly, but shifting climatic conditions do 34
not explain the overall, long-term, negative population tra- 35
jectory observed in our data. 36

Keywords Climate change · *Danaus plexippus* · 37
Hierarchical model · Monarch butterfly · Western 38
population 39

Introduction 40

Issues of spatial and temporal scale have always been 41
among the greatest challenges that face ecologists wishing 42
to extrapolate beyond single species and local conditions 43
(McGill 2010; Chave 2013). These concerns have been 44
brought to the fore by recent decades of anthropogenic 45
influence on the environment, as the public looks to ecolo- 46
gists for predictions regarding changes in regional or con- 47
tinental floras and faunas (Morissette et al. 2008; Tyliana- 48
kis et al. 2008). An important advance in the process has 49
involved meta-analyses that allow global phenomena to be 50
perceived through the accumulation of smaller-scale case 51
studies (Parmesan 2006; Wu et al. 2011; Mantyka-Pringle 52
et al. 2012). A further key contribution has come from the 53

A1 Communicated by Klaus Fischer.

A2 **Electronic supplementary material** The online version of this
A3 article (doi:10.1007/s00442-016-3600-y) contains supplementary
A4 material, which is available to authorized users.

A5 Anne E. Espeset
A6 aespeset@nevada.unr.edu

A7 ¹ Program in Ecology, Evolution, and Conservation Biology,
A8 Department of Biology, University of Nevada, Reno, NV,
A9 USA

A10 ² Center for Population Biology, University of California,
A11 Davis, CA, USA

A12 ³ Department of Biology, Texas State University, San Marcos,
A13 TX, USA

A14 ⁴ Department of Environmental Science and Policy, University
A15 of California, Davis, CA, USA

A16 ⁵ Ecology and Evolutionary Biology, University of Tennessee,
A17 Knoxville, TN 37996, USA

A18 ⁶ 1664 North Virginia Street, Reno, NV 89509, USA



54 development of hierarchical models that can effectively
55 estimate parameters (such as the influence of weather)
56 across large numbers of species and locations (Royle and
57 Dorazio 2008; Ponciano et al. 2009; Congdon 2014; Nice
58 et al. 2014). A logical implementation of such models
59 involves species that utilize large regions, and migratory
60 species are of particular interest because of complex life
61 cycles that integrate climatic variation across heterogene-
62 ous local climates (Zipkin et al. 2012).

63 A prominent migratory animal in North America is the
64 monarch butterfly (*Danaus plexippus*). This species has
65 two independent migratory populations or subpopulations
66 that together traverse much of the continent (Brower 1995;
67 Brower and Malcolm 1991). In general, monarchs east of
68 the Rocky Mountains migrate to Mexico while monarchs
69 west of the Rocky Mountains migrate to locations along the
70 Pacific coast of California (Urquhart and Urquhart 1977),
71 aggregating in groves of Monterey pine (*Pinus radiata*),
72 Monterey cypress (*Cupressus macrocarpa*), and blue gum
73 (*Eucalyptus globulus*) (Weiss et al. 1991). Genetic stud-
74 ies have suggested that these populations are not distinct
75 (Brower and Malcolm 1991; Lyons et al. 2012; Zhan et al.
76 2014), and some western monarchs potentially overwinter
77 in Mexico (Dingle et al. 2005; Morris et al. 2015). Both
78 populations depend on host plants in the genus *Asclepias*,
79 the milkweeds. These plants are ruderal in nature, and have
80 experienced declines in recent years in some areas, poten-
81 tially in association with increased herbicide use on agri-
82 cultural lands (Hartzler 2010; Pleasants and Oberhauser
83 2012; Zalucki and Lammers 2010). For the overwintering
84 monarchs in Mexico, severe weather and forest degradation
85 are further stressors that compound habitat and host loss on
86 breeding grounds in the USA (Brower et al. 2012; Flock-
87 hart et al. 2015). Despite apparent stressors and declines in
88 monarchs at their overwintering grounds (Vidal and Ren-
89 don-Salinas 2014; Saenz-Romero et al. 2012; Brower et al.
90 2002), numbers have not declined at some of the fall stop-
91 over sites in the Eastern USA (e.g., Davis 2012) or summer
92 breeding grounds (Ries et al. 2015). Also, weather has not
93 been considered to have a significant effect on the eastern
94 Monarch population during spring and summer (Zalucki
95 et al. 2015). In contrast to the many detailed studies pub-
96 lished on the eastern monarchs (e.g., Oberhauser and Peter-
97 son 2003; Batalden et al. 2007; Brindza et al. 2008; Davis
98 and Dyer 2015; Oberhauser et al. 2015), the western migra-
99 tion has received less attention (Koenig 2006).

100 Here we utilize a single-observer dataset on monarch
101 populations at ten locations throughout the breeding range
102 across northern California (Fig. 1). To our knowledge,
103 this is the longest and most temporally intensive dataset
104 on western monarchs, and consists of biweekly observa-
105 tions during monarch flights for between 27 and 42 years,
106 depending on the site. In addition to these biweekly data,

107 we use publically available numbers characterizing adult
108 densities at coastal overwintering locations, as well as data
109 describing abundances of eastern monarchs to compare
110 western and eastern population dynamics [Shapiro 2014;
111 The Xerces Society 2015; North American Butterfly Asso-
112 ciation (NABA) 2015; Monarch Net 2015]. Through exam-
113 ination of these data we address the following questions:
114 have monarch observations per year changed over time
115 (between years and within years) at our ten focal sites?
116 And, to what extent can fluctuations in observed monarchs
117 per year at focal sites be predicted by local and regional
118 weather variables, both at the summer sites and at the over-
119 wintering grounds? With respect to these questions, we pre-
120 dict monarch numbers to be declining, as previous studies
121 have shown negative trends over time for most butterflies
122 in the region, especially at low-elevation sites (Forister
123 et al. 2011; Harrison et al. 2015). We also hypothesize that
124 warming conditions will have had a negative influence on
125 the population, as has been observed for other butterflies in
126 northern California (Casner et al. 2014). We also ask, as an
127 issue of secondary interest, if dynamics at our transect sites
128 (on the western summer grounds) are similar to patterns
129 observed at the California overwintering sites. This ques-
130 tion is motivated in part by previous work by Stevens and
131 Frey (2010) who suggested a positive association between
132 breeding season precipitation in our study area and inter-
133 annual variation in monarch counts at the coastal overwin-
134 tering areas. Finally, in order to place the western popula-
135 tions in the larger, continental context for monitoring and
136 management of this migratory species, we compare tem-
137 poral dynamics among the following datasets: the western
138 summer grounds (our ten focal sites), the western overwin-
139 tering grounds, the eastern summer grounds, and the east-
140 ern (Mexican) overwintering grounds (Shapiro 2014; the
141 Xerces Society 2015; NABA 2015; Monarch Net 2015).

142 Materials and methods

143 Data collection and sampling locations

144 Data were recorded from 1972 up to and including 2014
145 at ten locations in Northern California by A. M. S. (Shap-
146 iredo 2014). These locations describe an elevational transect
147 starting at sea level and extending up over the crest of the
148 Sierra Nevada mountains at 2775 m, and down the eastern
149 slope to Sierra Valley. Sites encompass an array of habitat
150 types, from saltwater marsh to sub-alpine barrens. Each site
151 was visited every 2 weeks and the presence or absence of
152 monarchs was noted (henceforth presences are referred to
153 as “day positives”, as in other publications from these data,
154 e.g., Forister et al. 2010). Surveys were conducted via the
155 Pollard walk method (Pollard 1977) on days suitable for




Fig. 1c **a** Map of California in the Western USA, including focal sites where observations of adults during the summer flight season were recorded [Suisun Marsh (A), Gates Canyon (B), West Sacramento (C), North Sacramento (D), Rancho Cordova (E), Washington (F), Lang Crossing (G), Castle Peak (H), Donner Pass (I), and Sierra Valley (J)]. *Large, open circles* along the coast are overwintering locations (see main text for details) from which abundance data were collated for use in analyses; *solid dots* are overwintering locations from which weather data were gathered for use in a subset of climatic analyses. **b** Map of North America showing eastern regions represented by count data from the summer flight season, as follows: North Central (1), (2) North East (2), Mid Central (3), Mid East (4), and South (5). **c** Diagram of datasets analyzed, as follows: “day positives” (I; counts of days on which adult monarchs were observed per year) at ten sites across the western breeding grounds; counts of adults at five of the low-elevation western breeding sites (II); counts of overwintering adults from a subset of western overwintering sites (III; circled on map; summary data from the eastern migration including summer population indices and hectares occupied by overwintering adults in Mexico (IV); climatic data from each of the ten focal western breeding sites (V); regional climatic conditions (VI; MEI 1 and MEI 2); climatic conditions at the Pacific overwintering sites (VII, VIII; treated separately, as explained in main text). Our focal dataset (day positives) is highlighted with a *gray background* and connected to climatic datasets (V–VIII) by *single-headed arrows* to represent multiple regressions. Relationships among monarch datasets (I–IV) were explored with correlations, indicated by *double-headed arrows* (for simplicity, not all connections are drawn). Illustration of adult monarch butterfly, *Danaus plexippus*, by Anne Espeset

156 butterfly flight (sunny days with little wind). Abundance
 157 data (counts of adult monarchs) are also available from five
 158 of our ten sites, and from a subset of years (1999–2012).
 159 The abundance data have been used previously to show that
 160 day positives are suitable proxies for monarch abundance
 161 (Casner et al. 2014), and we present limited analyses of the
 162 abundance data here [Electronic Supplementary Material
 163 (ESM) 1].

164 Count data from California overwintering locations were
 165 obtained from the Xerces Society Thanksgiving Count
 166 Database (The Xerces Society 2015). Data were collected
 167 from 1997 to 2014 by volunteers. Data from six overwin-
 168 tering counties were gathered for use in the present analy-
 169 ses: Marin, Monterey, San Luis Obispo, San Diego, Santa
 170 Barbara, and Santa Cruz. The overwintering data contains
 171 gaps (years without observations), but counties were cho-
 172 sen that had enough individual sites within them to provide
 173 coverage of the greatest number of years (1997–2014). All
 174 sightings within a county were averaged per year, giving
 175 a mean count/site per year. Data for the eastern monarch
 176 population were obtained from Monarch Net (Monarch

Net 2015). These data consist of NABA Fourth of July
 177 counts from 1990 to 2009 (NABA 2015). Counts span five
 178 large geographical regions (north east, north central, mid
 179 east, mid central, and south; Fig. 1) and were collected
 180 by volunteers. Hectare overwintering data from Mexico
 181 were obtained from Monarch Butterfly Biosphere Reserve
 182 from 1994 to 2003 and World Wildlife Fund-Telcel Alli-
 183 ance from 2004 to 2015; the compiled data were accessed
 184 through The Xerces Society (2015). 185

Weather data 186

Weather specific to our ten transect sites was obtained from
 187 the PRISM working group (PRISM Climate Group 2015). **AQI** 8
 188 These data are interpolated from neighboring weather sta-
 189 tions incorporating local differences in topography, thus
 190 they potentially provide a more robust estimate of site-
 191 specific weather than raw data from the closest weather
 192 stations (Daly et al. 2008), which are subject to error and
 193 missing values. Data were grouped seasonally and reflect
 194 the water year, such that the 1980 water year, for exam-
 195 ple, starts with the fall of 1979. Specifically, “fall” is
 196 the previous year’s September, October, and November; “win-
 197 ter” consists of the previous year’s December and the focal
 198 year’s January and February; “spring” is March up to and
 199 including May; and “summer” is June up to and including
 200 August. For each season, average daily temperature and
 201

202 total precipitation were calculated (precipitation included
203 snow for the high-elevation sites). Winter average tempera-
204 ture and precipitation data from PRISM were also obtained
205 for each California overwintering location within the fol-
206 lowing eight counties: Alameda, Marin, Mendocino, Mon-
207 terey, San Diego, San Luis Obispo, Sonoma, and Ventura.
208 These counties were chosen to ensure that sampling encap-
209 sulated a wide range of overwintering conditions along
210 the coast. In all cases (for focal and overwintering sites),
211 PRISM data were taken from the latitude and longitude
212 centroid of each site (using the default setting of 4-km²
213 cells).

214 Because of the migratory nature of the monarch, we
215 were interested in the possibility that regional weather vari-
216 ables could provide an informative contrast to the local data
217 generated by PRISM. The multivariate El Niño Southern
218 Oscillation index (MEI) was used to explore the impact of
219 regional weather drivers (Wolter and Timlin 1993). The El
220 Niño Southern Oscillation index (ENSO) is associated with
221 unusual precipitation patterns throughout northern Califor-
222 nia (Schonher and Nicholson 1989), and MEI is the first
223 principal component extracted from the analysis of six vari-
224 ables that together provide an index of the intensity of the
225 ENSO for a given month. In order to reduce the complex-
226 ity of the data to a manageable form, an additional prin-
227 ciple components analysis on the MEI values across all
228 12 months was performed, and the first two components
229 (MEI 1 and MEI 2) were extracted for analyses of monarch
230 dynamics.

231 Overview of statistical methods

232 Analyses consisted of multiple, distinct models and com-
233 binations of data sources (Fig. 1c). The focus of analyses
234 is data from our ten sites across the breeding grounds,
235 because it is from those sites that we have the most reliable,
236 temporally consistent data.

- 237 1. First, a hierarchical Bayesian model was used to study
238 the effects of local and regional weather variables on
239 monarch observations at focal sites across the sum-
240 mer breeding grounds (“local” weather for these mod-
241 els refers to PRISM data from the focal sites, while
242 “regional” refers to MEI, as described above). This
243 model included year as a predictor variable, and thus
244 produced estimates of change in monarch observations
245 across the breeding sites through time. As a comple-
246 mentary analysis, individual count data (available for
247 a subset of years and sites) were used to ask if any
248 changes across years have been localized to particular
249 times of the year (more details below).
- 250 2. Next, a path analysis was used to place effects of
251 weather and year effects (change over time in monarch

observations) into a context that allows for the direct
quantification of shifting climatic conditions on butter-
fly observations.

3. The approach in the first step (a hierarchical Bayes-
ian model) was repeated, but investigated the effects
of climate at the Pacific overwintering sites on mon-
arch observations across the summer breeding grounds
(in other words, investigating potential connections
between conditions experienced during overwintering
and observations made the following flight season).
4. Finally, simple correlations were used to investigate
associations between all monarch datasets: our focal
sites (observations during the breeding season), west-
ern overwintering sites, eastern regions, and Mexican
overwintering sites. Because this step involved a large
number of comparisons, we focus primarily on overall
patterns rather than significance testing of individual
correlations.

Hierarchical models

The impact of weather variables on monarch day posi-
tives was explored using a hierarchical Bayesian modeling
approach. This analytical method is described in detail
elsewhere (see Nice et al. 2014; Harrison et al. 2015).
Briefly, the model estimates posterior probability distribu-
tions (PPDs) for partial regression coefficients associated
with model terms at multiple hierarchical levels, in this
case site and transect wide. Information from each site is
used to inform transect-wide estimates and vice versa. A
binomial response consisting of day positives and num-
ber of visits for a given year (i.e., the proportion of posi-
tive visits in a year) and site was modeled, thus account-
ing for variation in sampling effort among years. Model terms
included site-specific seasonal average temperatures and
total precipitation (summed over season) and our indices of
MEI (as described above). Year was included in the model
to quantify inter-annual population trends not directly
associated with fluctuating climatic conditions. All predic-
tor variables were converted to standardized *z*-scores prior
to modeling. PPDs for each model term were estimated
using the JAGS sampler [version 3.4.0 (Plummer 2013)],
a Markov chain Monte Carlo (MCMC) algorithm, imple-
mented in R (R Core Team 2014) using the rjags package
[version 3-15 (Plummer 2015)]. The model was run using
two search chains and uninformative priors and hyperpriors
for 500,000 iterations of the sampling algorithm. To gauge
model performance, effective sample sizes (ESS) were cal-
culated for each parameter estimate, and trace plots of esti-
mates against iterations were examined to evaluate mixing.

As described above, this modeling approach was used
for different sets of weather variables (Fig. 1c). First,
regional and local (associated with each focal site) weather

303 variables (MEI 1, MEI 2, temperature, and precipitation)
 304 were used to predict monarch observations at the focal sites
 305 across the breeding grounds. Second, the impact of over-
 306 wintering weather conditions on subsequent-year observa-
 307 tions was examined across the breeding grounds. Because
 308 of the large number of overwintering weather variables and
 309 the large number of relationships (associated with eight
 310 overwintering counties potentially affecting observations
 311 at ten breeding sites), two analyses were run with overwin-
 312 tering weather variables: first with all of the overwintering
 313 temperature data, and second with all of the overwintering
 314 precipitation data (analyses were also run with all weather
 315 data combined, which did not qualitatively alter results
 316 but did broaden credible intervals and lower precision as a
 317 result of reduced power). Year was included as a covariate
 318 in all models.

319 Climatic trends and path analyses

320 The analyses described in the previous section addressed
 321 the impact of climatic variables (local, regional and at the
 322 overwintering sites) on monarch observations at the focal
 323 breeding ground sites. To explicitly examine the impact of
 324 climate change on monarch day positives, a path analysis
 325 was used to compare the direct effect of year on monarchs
 326 with the indirect effect of year as mediated by weather. The
 327 path analysis was built using a suite of hierarchical Bayesian
 328 regression models that together characterized the path
 329 models. Specifically, we separately modeled the effect of
 330 year on each endogenous climate variable (assuming a
 331 normally distributed response variable, as opposed to the
 332 binomially distributed response variable described above).
 333 Path coefficient estimates describing the effect of climate
 334 variables on day positives were taken from the hierarchical
 335 multiple regression model described above which included
 336 all climate variables as predictors. PPDs of all models
 337 were characterized using two search chains each of 25,000
 338 MCMC iterations. The mean and 95 % credible intervals
 339 from these distributions were used to characterize associ-
 340 ated path coefficients. Indirect effects of year as mediated
 341 by a given climate variable were calculated by taking the
 342 product of the appropriate path coefficients. In order to
 343 incorporate uncertainty from parameter estimates, those
 344 products were generated from 50,000 samples from each
 345 of the PPDs for the two path coefficients involved in each
 346 comparison (means and 95 % credible intervals were then
 347 retained from the distribution of products).

348 Localization of temporal trends within years

349 The analyses described above included year as a predictor
 350 variable and thus estimated changes in monarch observa-
 351 tions across the decades encompassed by our study. It is

also of interest to ask if any demographic trends over the
 years were focused on any particular time during the breed-
 ing season, which could be informative with respect to
 causes of population trends (e.g., associated with the over-
 wintering generation or with late-season immigrants from
 more distant breeding areas). To address this, counts of
 individuals were utilized from five of our focal sites (where
 such data are available), for 1999 to 2013, in a sliding win-
 dow regression analysis. Specifically, counts of monarchs
 were calculated from 50-day windows organized by ordinal
 dates, counting from the first of the year. For each of those
 windows, the total count was regressed against years and
 the beta coefficient (slope of count vs. years) was saved and
 examined for intra-annual patterns in inter-annual trends.

Comparisons between western and eastern populations

To examine the relationship between eastern and western
 monarch populations, Spearman's rank correlation coef-
 ficients were calculated using data from 1997 to 2014 for
 eastern monarchs obtained from NABA through Monarch
 Net, overwintering data for eastern (at Mexican sites) and
 western monarchs (at California sites) obtained through the
 Xerces society, and day positives per year for each of our
 transect locations. Eastern regional count data were only
 available from 1997 onwards, therefore only those Cali-
 fornia overwintering locations with high abundances and
 complete records since 1997 were used for this analysis.
 California overwintering data were obtained from counts
 conducted in six counties spanning the mid to southern
 coast of California (Fig. 1). As with hierarchical Bayesian
 models, these analyses were performed using R (version
 3.1.2, R Core Team 2014).

Results

Annual trends, climatic impacts, and the effect of a changing climate

The hierarchical Bayesian approach successfully character-
 ized PPDs for partial regression coefficients for all models
 in which weather variables and year predicted monarch
 observations (day positives). Visual inspection of trace
 plots confirmed adequate mixing of model chains. ESS
 necessarily varied between parameter estimates, but were
 always greater than 1000. A decline in monarch observa-
 tions is evident in both raw day positives (Fig. 2a), and in
 the year coefficient estimated across sites from the hier-
 archical model (Fig. 2b). The point estimate for the year
 coefficient as a log odds ratio across all sites was -0.6
 (Fig. 2a). The exponential transformation of that coefficient
 (from log odds ratio to odds ratio) is 0.55, which means




Fig. 2 **a** Temporal trends of monarch fractional day positives (*FDPs*) from 1972 to 2014. These values (*y*-axis) correspond to the fraction of days during the year in which a monarch adult was seen (out of the total number of visits in that year) at the focal sites. Letters (labeling the observations for each year) correspond to the site labels, as in Fig. 1. Fitted lines from simple linear regressions are shown for visualization (for each site separately as *gray lines*, and for all of the sites as the *darker line*). **b** Posterior probability distributions (PPDs) for the coefficient associated with year from a hierarchical Bayesian model predicting monarch observations across the ten focal sites. PPDs are shown in *light gray* for each site, and across sites in *black*; similarly, *tick marks* at the bottom of the graph show the mean estimate for each curve. *Dashed lines* indicate 95 % credible intervals for the PPD across sites. **c** Beta coefficients from moving window analysis of monarch abundance at five summer breeding sites where data on counts of individuals were available: *each point* indicates the relationship between abundance and year for monarch counts in 50-day windows (the *x*-axis is the midpoint of those windows in days from the start of the year). *Increasingly negative values* indicate more severe declines, with regressions significant at $P < 0.05$ shown in *black*. *Dotted horizontal line* at zero shown for reference

declining observations of monarchs are not spread equally across the breeding season, but appear to be localized earlier in the season (Fig. 2c). It is important to note that local breeding at the low-elevation sites (Fig. 1c, sites A–E) has not been observed before May (A. M. S., personal observation), and in some years local breeding never happens. Thus the reduced early spring numbers (Fig. 2c) likely involve a reduction in immigration from coastal overwintering sites. Reduced numbers of individuals observed at a particular time of the year could also be a result of phenological shifts, but monarchs at the five low-elevation sites have not been appearing earlier or later in the spring ($F_{1,13} = 0.06$; $P = 0.81$) nor has phenology shifted at the end of the season ($F_{1,13} = 0.41$; $P = 0.53$).


Warmer temperatures in winter and spring (Fig. 3; for results from all weather variables, see Table 1 and ESM 2–4) were positively and consistently associated with monarch sightings at our ten focal sites. The standardized beta coefficients for winter and spring temperatures estimated across all breeding sites were 0.23 and 0.26, respectively. Spring precipitation had a positive effect across sites, while summer precipitation had a more heterogeneous effect across sites (Fig. 3). The impact of overwintering conditions on monarch observations at our focal sites the following summer was also examined; complex and site-specific relationships were revealed (ESM 5). In particular, both temperature and precipitation have strong and significant effects at a subset of the overwintering sites, but the direction of the effect (from positive to negative) varies. A path analysis allowed us to address the potential influence of climate change on monarch populations through the examination of the indirect effect of year as mediated by a given weather variable. The direct effect of year was much greater than the indirect effect of year as mediated by any one weather variable (Fig. 4).

Relationship between eastern and western monarch populations

In the west, overwintering population size and day positives at the transect sites (in the subsequent year) were positively correlated (Fig. 5). Monarch overwintering counts and day positives along the transect were all negatively correlated with year, consistent with the declining annual trend reported from analyses above. Inspection of abundance data from the western overwintering sites (ESM 6) confirms the downward trajectories during the years studied. In contrast, observations of eastern breeding locations tended to be weakly or positively related to year, although a decline is evident at the Mexican overwintering grounds (negatively associated with year). Eastern and western populations show no significant correlations (neither positive or negative; all P -values were > 0.05), with one anomalous exception (Gates Canyon and

399 that the odds ratio of observing a monarch has decreased by
400 that factor (0.55) for every year. A decline can also be seen
401 in plots of adult counts from our focal sites for the subset
402 of recent years in which counts are available (ESM 1). The

Fig. 3 PPDs for a subset of weather variables predicting monarch observations from hierarchical Bayesian models [for results from all weather variables, see Table 1 and Electronic Supplementary Material (ESM) 2–4]. PPDs are shown in *light gray* for each site, and across sites in *black*; similarly, *tick marks* at the bottom of the graph show the mean estimate for each curve. *Insets* The probability that each site has a non-zero (either positive or negative) coefficient is shown (calculated as the proportion of the corresponding PPD greater than or less than zero). The *bars* indicating probability are labeled by site (A–J; see Fig. 1), and for the whole model. *Bars* above the *horizontal line* (at zero) are for coefficients with mean positive values, while *bars* below the *horizontal line* are for negative coefficients (the *top three panels* are dominated by positive coefficients, while the *bottom panel* includes a mix of positive and negative coefficients)



454 the mid-central eastern region), suggesting in general that
 455 an abundant year for one region is not necessarily an abundant
 456 year for the other (Fig. 5); for further details, see ESM
 457 7 for a comparison of distributions of correlation coefficients
 458 within and among geographic regions.

459 **Discussion**

460 Monarch butterflies have received a great deal of attention
 461 as one of the most conspicuous migratory species in North
 462 America, but most research has focused on the larger,

eastern migration. In this study, we found that the western
 migration of *Danaus plexippus* shows evidence of decline
 which is apparent in raw day positive data (Fig. 2a), abundance
 data from our focal sites (ESM 1), and abundance data from the
 Pacific coast overwintering sites (ESM 6). Moreover, the declines
 at the western breeding sites are concentrated early in the season,
 while abundances of adults have not been as reduced closer to the
 end of the breeding season (Fig. 2c). When considering this pattern
 in light of the associated decline in overwintering population size,
 it suggests that mortality could be increasing either during or
 immediately after overwintering. Alternatively,

463
464
465
466
467
468
469
470
471
472
473
474

Table 1 Standardized regression coefficients from hierarchical Bayesian models relating temperature, precipitation, and multivariate El Niño Southern Oscillation index (*MEI*) values (El Niño Southern Oscillation index indices) to monarch observations at the ten focal sites (coefficients are in log-linear units from binomial regressions)

	Temperature				Precipitation				MEI	
	Winter	Spring	Summer	Fall	Winter	Spring	Summer	Fall	1	2
Suisun Marsh	0.26 ^a	0.21 ^a	-0.08	0.04	-0.14	0.26 ^a	0.17	0.05	0.32	0.0029
Gates Canyon	0.30 ^a	0.30 ^a	-0.01	0.09	-0.05	0.17	-0.23	0.03	0.36	-0.052
West Sacramento	0.25 ^a	0.32 ^a	-0.01	0.09	-0.06	0.14	-0.02	0.12	0.37	-0.017
North Sacramento	0.24 ^a	0.27 ^a	-0.08	0.11	-0.05	0.16	-0.16	0.14	0.28	0.11
Rancho Cordova	0.27 ^a	0.27 ^a	-0.01	0.12	-0.03	0.23 ^a	0.09	0.06	0.35	-0.034
Washington	0.25 ^a	0.25 ^a	-0.09	0.11	-0.07	0.15	0.06	0.06	0.33	0.0041
Lang Crossing	0.14	0.28 ^a	-0.05	0.13 [†]	-0.08	0.14	-0.11	0.07	0.37	0.023
Donner Pass	0.17	0.22	0.06	0.07	-0.09	0.1	-0.2	0.05	0.34	-0.17
Castle Peak	0.24	0.26 ^a	0.05	0.12	-0.08	0.16	0.12	0.07	0.35	-0.027
Sierra Valley	0.23 ^a	0.24 ^a	0.05	0.06	-0.05	0.12	0.04	0.05	0.32	-0.13
Across sites ^b	0.23 ^a	0.26 ^a	-0.02	0.09	-0.07	0.16 ^a	-0.02	0.07	0.34	-0.028

Numbers correspond to posterior probability distributions visualized in Fig. 3, and Electronic Supplementary Material (ESM) 2–4

^a Coefficients with 95 % credible intervals that do not overlap zero

^b Coefficients estimated across all sites

perhaps fewer butterflies are able to successfully migrate in the fall from their summer breeding grounds back to their overwintering colonies. Regardless, subsequent generations during the summer are able to at least partially rebound.

The climatic models we developed successfully predicted monarch observations. Warmer springs and winters, for example, have pronounced and positive effects on the frequency of monarch observations during the summer flight (Fig. 3). These effects of weather could be the result of positive associations with overwinter survival and reproduction of the first summer generation, although it is important to remember that these effects are of much smaller magnitude compared to the direct, negative association with years (Fig. 4). The positive effects of temperature could also be mediated through increased nectar and host plant growth, although we can only pose these possibilities as hypotheses at this time. The positive effect of precipitation is consistent with previous climatic modeling for the western monarch that suggested water as a limiting factor (Stevens and Frey 2010).

Although our analyses revealed effects of weather on monarch observations, it is clear that shifting climatic conditions are not the major factor in the observed linear population declines. This result can be seen both in the lack of directional change in most weather variables studied, and in the very small indirect effects of year as mediated through weather (Fig. 4). Summer temperatures are an exception as they are rising across our focal sites, but they do not appear to have an effect on monarch observations (Fig. 4; ESM 3).

The large, negative, direct effect of year suggests other (non-climatic) drivers of decline. In studies on eastern monarchs, it has been posited that dwindling host plant

populations (Flockhart et al. 2015; Pleasants and Oberhauser 2012; Zalucki and Lammers 2010; Brower et al. 2006), the use of insecticides (Krischik et al. 2015; Pecenka and Lundgren 2015), and overwintering habitat destruction (Vidal and Rendon-Salinas 2014; Saenz-Romero et al. 2012; Brower et al. 2002) are primary drivers of decline. To our knowledge, a decline in milkweed abundance has not been reported within the range of the western monarch. The intra-annual pattern of declines (Fig. 2c) is relevant to this issue, as it suggests a reduced number of early spring immigrants over the years. If host plants throughout the breeding season were limiting, we might expect declines to be spread more evenly throughout the year or even concentrated towards the end of the season. Given the likely link between overwintering declines (ESM 6) and reduced immigration to the breeding grounds, habitat loss along the California coast is potentially important (Jepsen and Black 2015), and we can suggest that overwintering sites would be the logical next step for focused investigation. The need to better understand overwintering sites is also highlighted by the heterogeneous weather effects that we observed along the California coast, potentially mediated by variation in habitat and micro-climate. In general, a large number of butterflies are known to be declining in Northern California (Forister et al. 2010, 2011), for which causes are likely multifarious, though a combined effect of land use change and warming conditions has been implicated for many species (Casner et al. 2014). Finally, we compared population dynamics among western summer grounds, western overwintering sites, and the eastern migration, both breeding and overwintering sites (Fig. 5). The lack of correlation between western and eastern observations is consistent with previous studies (Frey and Schaffer 2004;




Fig. 4 Path diagram illustrating direct and indirect effects of year and weather variables on monarch sightings (day positives), as well as effects of year on weather variables. Values next to each path are means from Bayesian PPDs, and 95 % credible intervals (from the same analyses reported in Fig. 3: ESM 2–4). Indirect effects of year on day positives mediated through weather variables were all small

in comparison to direct coefficients, and are shown as *italicized gray font* above the names of the weather coefficients. Asterisks indicate coefficients whose 95 % credible intervals do not overlap zero. Lines representing negative relationships end in *circles*. For abbreviations, see Figs. 2 and 3


541 Stevens and Frey 2010) and suggests that the two popula- 553
 542 tions are fluctuating independently. 554

543 In conclusion, we have successfully modeled effects 555
 544 of weather on monarch observations along an elevational 556
 545 transect encompassing a portion of the summer breeding 557
 546 ground of the western subpopulation of the monarch butterfly. 558
 547 Although we were able to detect climatic effects, there 559
 548 is clearly more to be learned with respect to biotic-abiotic 560
 549 interactions playing out across the geographic extent of the 561
 550 range of the western monarch. This is particularly appar- 562
 551 ent in the variety of weather effects observed at the coastal 563
 552 overwintering grounds. In contrast to the complexity of 564

553 weather, a negative, annual trend in monarch observa- 554
 555 tions was readily detected across all of the focal sites. The 556
 557 annual trend is not explained by shifting climatic condi- 558
 559 tions, which have been implicated in the declines of other 560
 561 butterflies in the region (Casner et al. 2014). Furthermore, 562
 563 the decline in observations is correlated with decreasing 564
 564 numbers at the overwintering sites that we studied along 565
 565 the Pacific coast. 566

561 An important caveat to these results is the fact that we 562
 562 have focused our analyses on observations of adults during 563
 563 the breeding season from ten sites that encompass a broad 564
 564 elevational transect, but a narrow portion of the breeding 565
 565 season. 566

Fig. 5 Correlations among our focal sites (“Transect sites” across the western breeding grounds), western overwintering sites averaged by county, summer observations from the eastern region, and areal extent of Mexican overwintering grounds (for all sets of data, years covered were 1997–2014). Colors indicate the magnitude of Spearman’s rank correlation coefficients, and year was included (at the bottom of the figure) for quantification of temporal trends. Rank correlations were used because of the heterogeneous nature of the data (including fractional day positives for our focal sites, counts from the overwintering sites, and population indices from the eastern sites). Black boxes outline western transect sites vs. western overwintering sites (upper left), and western transect sites vs. eastern summer sites (lower right). To compare patterns among regions, histograms of correlations can be found in ESM 7



565 grounds. It is possible that the declines we have observed
 566 are the consequence of a shift in migration behavior rather
 567 than a demographic effect, although we have no particu-
 568 lar reason (anecdotal or otherwise) to expect such a geo-
 569 graphic shift, especially since our ten sites are not marginal
 570 to the breeding range. Indeed, analyses by Stevens and Frey
 571 (2010) place our transect within the geographical region
 572 best suited for monarch breeding, as determined by both
 573 thermal conditions, and host plant availability. The corre-
 574 lations observed between our focal sites and the western
 575 overwintering sites (Fig. 5) also suggest general declines
 576 rather than any localized shift in migration patterns.
 577 Finally, it is interesting to note that dynamics of the sites
 578 that we have studied (both the summer sites and overwin-
 579 tering sites) appear to be uncorrelated with dynamics in the
 580 eastern subpopulation. Thus, even though genetic differen-
 581 tiation between the western and eastern subpopulations has
 582 not been detected, from a conservation and management
 583 perspective they should be considered different entities.

584 **Acknowledgments** We thank the citizen science participants of
 585 the Western Monarch Thanksgiving Count and the Fourth of July
 586 Butterfly Counts, as well as two anonymous reviewers of an early
 587 draft. Much appreciation goes to the Xerces Society and the North
 588 American Butterfly Association, respectively associated with the

aforementioned counts, for making the data publicly available and
 providing them online through Monarch Net. This research was
 supported in part by the Trevor James McMinn professorship to
 M. L. F. and by the National Science Foundation (DEB-1050726, and
 DEB-1145609).

Author contribution statement A. E. E. and M. L. F. conceived the
 idea. A. M. S. collected the data. C. C. N., A. E. E., J. G. H., M. L. F.,
 and J. A. F. developed the statistical models; J. H. T. and D. P. W.
 managed and archived the data; A. E. E., J. G. H., and M. L. F. wrote
 the manuscript, while A. M. S., C. C. N., J. H. T., D. P. W., and J. A. F.
 reviewed the manuscript before submission.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict
 of interest.

References

Batalden RV, Oberhauser K, Peterson AT (2007) Ecological niches
 in sequential generations of eastern North American monarch
 butterflies (Lepidoptera: Danaidae): the ecology of migra-
 tion and likely climate change implications. *Environ Entomol*
 36:1365–1373
 Brindza LJ, Brower LP, Davis AK, Van Hook T (2008) Comparative
 success of monarch butterfly migration to overwintering sites in

589
590
591
592
593

594
595
596
597
598
599

AQ3
600

601
602

AQ4
3

604
605
606
607
608
609
610

- 611 Mexico from inland and coastal sites in Virginia. *J Lepid Soc* 62:189–200
- 612 Brower LP (1995) Understanding and misunderstanding the migration of the monarch butterfly (Nymphalidae) in North America: 1857–1995. *J Lepid Soc* 49:304–385
- 613 Brower AVZ, Boyce TM (1991) Mitochondrial-DNA variation in monarch butterflies. *Evolution* 45:1281–1286
- 614 Brower LP, Malcolm SB (1991) Animal migrations—endangered phenomena. *Am Zool* 31:265–276
- 615 Brower LP, Castilleja G, Peralta A, Lopez-Garcia J, Bojorquez-Tapia L, Diaz S, Melgarejo D, Missrie M (2002) Quantitative changes in forest quality in a principal overwintering area of the monarch butterfly in Mexico, 1971–1999. *Conserv Biol* 16:346–359
- 616 Brower LP, Fink LS, Walford P (2006) Fueling the fall migration of the monarch butterfly. *Integr Comp Biol* 46:1123–1142
- 617 Brower LP, Taylor OR, Williams EH, Slayback DA, Zubieta RR, Ramirez MI (2012) Decline of monarch butterflies overwintering in Mexico: is the migratory phenomenon at risk? *Insect Conserv Diversity* 5:95–100
- 618 Casner KL, Forister ML, O'Brien JM, Thorne J, Waetjen D, Shapiro AM (2014) Contribution of urban expansion and a changing climate to decline of a butterfly fauna. *Conserv Biol* 28:773–782
- 619 Chave J (2013) The problem of pattern and scale in ecology: what have we learned in 20 years? *Ecol Lett* 16:4–16
- 620 Congdon P (2014) Applied Bayesian modeling, vol. 595. Wiley
- 621 Daly C, Halbleib M, Smith JI, Gibson WP, Doggett MK, Taylor GH, Curtis J, Pasteris PP (2008) Physiographically sensitive mapping of climatological temperature and precipitation across the conterminous United States. *Int J Climatol* 28:2031–2064
- 622 Davis AK (2012) Are migratory monarchs really declining in eastern North America? Examining evidence from two fall census programs. *Insect Conserv Divers* 5:101–105
- 623 Davis, AK, Dyer LA (2015) Long-term trends in eastern North American monarch butterflies: a collection of studies focusing on spring, summer, and fall dynamics. *Ann Entomol Soc Am* 1–3
- 624 Dingle H, Zalucki MP, Rochester WA, Armijo-Prewitt T (2005) Distribution of the monarch butterfly, *Danaus plexippus* (L.) (Lepidoptera: Nymphalidae), in western North America. *Biol J Linn Soc Lond* 85:491–500
- 625 Flockhart DTT, Pichancourt J-B, Norris DR, Martin TG (2015) Unravelling the annual cycle in a migratory animal: breeding-season habitat loss drives population declines of monarch butterflies. *J Anim Ecol* 84:155–165
- 626 Forister ML, McCall AC, Sanders NJ, Fordyce JA, Thorne JH, O'Brien J, Waetjen DP, Shapiro AM (2010) Compounded effects of climate change and habitat alteration shift patterns of butterfly diversity. *Proc Natl Acad Sci USA* 107:2088–2092
- 627 Forister ML, Jahner JP, Casner KL, Wilson JS, Shapiro AM (2011) The race is not to the swift: long-term data reveal pervasive declines in California's low-elevation butterfly fauna. *Ecology* 92:2222–2235
- 628 Frey DF, Schaffer A (2004) Spatial and temporal patterns of monarch overwintering abundance in Western North America. *Monarch butterfly biology and conservation*. Cornell University Press, Ithaca, NY, pp 167–176
- 629 Harrison JG, Shapiro AM, Espeset AE, Nice CC, Jahner JP, Forister ML (2015) Species with more volatile population dynamics are differentially impacted by weather. *Biol Lett* 11:5
- 630 Hartzler RG (2010) Reduction in common milkweed (*Asclepias syriaca*) occurrence in Iowa cropland from 1999 to 2009. *Crop Prot* 29:1542–1544
- 631 Jepsen S, Black SH (2015) Understanding and conserving the Western North American monarch population. In: Oberhauser KR, Altizer NS (eds) *Monarchs in a changing world. Biology and conservation of an iconic butterfly*, chapter 11. Cornell University Press, Ithaca
- 632 Koenig WD (2006a) Spatial synchrony of monarch butterflies. *Am Midl Nat* 155:39–49
- 633 Krischik V, Rogers M, Gupta G, Varshney A (2015) Soil-applied imidacloprid translocates to ornamental flowers and reduces survival of adult *Coleomegilla maculata*, *Harmonia axyridis*, and *Hippodamia convergens* lady beetles, and larval *Danaus plexippus* and *Vanessa cardui* butterflies. *PLoS One* 10:22
- 634 Lyons JI, Pierce AA, Barribeau SM, Sternberg ED, Mongue AJ, de Roode JC (2012) Lack of genetic differentiation between monarch butterflies with divergent migration destinations. *Mol Ecol* 21:3433–3444
- 635 Mantyka-Pringle CS, Martin TG, Rhodes JR (2012) Interactions between climate and habitat loss effects on biodiversity: a systematic review and meta-analysis. *Global Change Biol* 18:1239–1252
- 636 McGill BJ (2010) Matters of scale. *Science* 328:575–576
- 637 Monarch Net (2015) Monarch abundance data. <http://monarchnet.uga.edu/>
- 638 Morissette JT, Richardson AD, Knapp AK, Fisher JI, Graham EA, Abatzoglou J, Wilson BE, Breshears DD, Henebry GM, Hanes JM, Liang L (2008) Tracking the rhythm of the seasons in the face of global change: phenological research in the 21st century. *Front Ecol Environ* 7:253–260
- 639 Morris GM, Kline C, Morris SM (2015) Status of *Danaus plexippus* population in Arizona. *J Lepid Soc* 69:91–107
- 640 Nice CC, Forister ML, Gompert Z, Fordyce JA, Shapiro AM (2014) A hierarchical perspective on the diversity of butterfly. Species' responses to weather in the Sierra Nevada Mountains. *Ecology* 95:2155–2168
- 641 North American Butterfly Association (NABA) (2015) NABA butterfly counts from 1990 to 2009
- 642 Oberhauser K, Peterson AT (2003) Modeling current and future potential wintering distributions of eastern North American monarch butterflies. *Proc Natl Acad Sci USA* 100:14063–14068
- 643 Oberhauser KS, Nail KR, Altizer SM (2015) Monarchs in a changing world: biology and conservation of an iconic butterfly. *Comstock, Ithaca*
- 644 Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annu Rev Ecol Evol Syst* 37:637–669
- 645 Pecenka JR, Lundgren JG (2015) Non-target effects of clothianidin on monarch butterflies. *Nat Sci* 102:19
- 646 Pleasants JM, Oberhauser KS (2012) Milkweed loss in agricultural fields because of herbicide use: effect on the monarch butterfly population. *Insect Conserv Divers* 6:135–144
- 647 Plummer M (2013) JAGS: a program for analysis of Bayesian graphical models using Gibbs sampling. Version 3:4
- 648 Plummer M (2015) Package rjags. Version 3-15
- 649 Pollard E (1977) Method for assessing changes in abundance of butterflies. *Biol Conserv* 12:115–134
- 650 Ponciano JM, Taper ML, Dennis B, Lele SR (2009) Hierarchical models in ecology: confidence intervals, hypothesis testing, and model selection using data cloning. *Ecology* 90:356–362
- 651 PRISM Climate Group (2015) Weather data. Oregon State University. <http://prism.oregonstate.edu/>
- 652 Ries L, Taron DJ, Rendón-Salinas E (2015) The disconnect between summer and winter monarch trends for the Eastern migratory population: possible links to differing drivers. *Ann Entomol Soc Am* 1–9
- 653 Royle JA, Dorazio RM (2008) Hierarchical modeling and inference in ecology: the analysis of data from populations, metapopulations and communities. Academic Press, Boston
- 654 R Core Team (2014) R: a language and environment for statistical computing. Version 3.1.2. R Foundation for Statistical computing, Vienna
- 655 Saenz-Romero C, Rehfeldt GE, Duval P, Lindig-Cisneros RA (2012) *Abies religiosa* habitat prediction in climatic change scenarios



- 743 and implications for monarch butterfly conservation in Mexico. 768
 744 For Ecol Manage 275:98–106 769
- 745 Schonher T, Nicholson SE (1989) The relationship between California 770
 746 rainfall and ENSO events. J Clim 2:1258–1269 771
- 747 Shapiro AM (2014) Species presence by site. Art Shapiro's butterfly 772
 748 site. <http://butterfly.ucdavis.edu/> 773
- 749 Stevens SR, Frey DF (2010) Host plant pattern and variation in climate 774
 750 predict the location of natal grounds for migratory monarch 775
 751 butterflies in western North America. J Insect Conserv 14:731–744 776
- 752 The Xerces Society (2015) Western Monarch Thanksgiving Count 777
 753 from 1997 to 2014. [http://www.xerces.org/wp-content/](http://www.xerces.org/wp-content/uploads/2011/04/WMTC-Data-1997-2014.pdf) 778
 754 [uploads/2011/04/WMTC-Data-1997-2014.pdf](http://www.xerces.org/wp-content/uploads/2011/04/WMTC-Data-1997-2014.pdf) 779
- 755 Tylianakis JM, Didham RK, Bascompte J, Wardle DA (2008) Global 780
 756 change and species interactions in terrestrial ecosystems. Ecol 781
 757 Lett 11:1351–1363 782
- 758 Urquhart FA, Urquhart NR (1977) Overwintering areas and migratory 783
 759 routes of monarch butterfly (*Danaus plexippus*, Lepidoptera 784
 760 Danaidae) in North America, with special reference to western 785
 761 population. Can Entomol 109:1583–1589 786
- 762 Vidal O, Rendon-Salinas E (2014) Dynamics and trends of overwintering 787
 763 colonies of the monarch butterfly in Mexico. Biol Conserv 180:165–175 788
- 764 Weiss SB, Rich PM, Murphy DD, Calvert WH, Ehrlich PR (1991) 789
 765 Forest canopy structure at overwintering monarch butterfly sites: 790
 766 measurements with hemispherical photography. Conserv Biol 5:165–175 791
 767 Wolter K, Timlin MS (1993) Monitoring ENSO in COADS with a 768
 769 seasonally adjusted principal component index. Proceedings of 770
 771 the 17th Climate Diagnostic, Norman, OK. pp 52–57 772
- 773 Wu Z, Dijkstra P, Koch GW, Penuelas J, Hungate BA (2011) 774
 775 Responses of terrestrial ecosystems to temperature and precipitation 776
 777 change: a meta-analysis of experimental manipulation. Global Change Biol 17:927–942 778
- 779 Zalucki MP, Lammers JH (2010) Dispersal and egg shortfall in monarch 780
 781 butterflies: what happens when the matrix is cleaned up? Ecol Entomol 35:84–91 782
- 783 Zalucki MP, Brower LP, Malcolm SB, Slager BH (2015) Estimating 784
 785 the climate signal in monarch population decline. In: Oberhauser 786
 787 KR, Altizer NS (eds) Monarchs in a changing world. Biology and 788
 789 conservation of an iconic butterfly, chapter 11. Cornell University 790
 791 Press, Ithaca 791
- 792 Zhan S, Zhang W, Niitepold K, Hsu J, Haeger JF, Zalucki MP, Altizer 793
 794 S, de Roode JC, Reppert SM, Kronforst MR (2014) The genetics 795
 796 of monarch butterfly migration and warning colouration. Nature 797
 798 514:317–321 799
- 800 Zipkin EF, Ries L, Reeves R, Regetz J, Oberhauser KS (2012) Tracking 801
 802 climate impacts on the migratory monarch butterfly. Global 803
 804 Change Biol 18:3039–3049 805

UNCORRECTED PROOF

Journal:	442
Article:	3600

Author Query Form

Please ensure you fill out your response to the queries raised below and return this form along with your corrections

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Query	Details Required	Author's Response
AQ1	PRISM Climate Group 2013 has been changed to PRISM Climate Group 2015 so that this citation matches the list.	
AQ2	Please provide a definition for the significance of dagger symbol (†) in Table 1.	
AQ3	Koenig 2006b has been deleted as it is identical to Koenig 2006a.	
AQ4	The following references have not been cited: Brower and Boyce (1991), Pecenka and Lundgren (2015). Please cite these references or delete them from the list.	