

Lawrence Berkeley National Laboratory

Recent Work

Title

CHARGE CHANGING CROSS SECTIONS FOR Pb AND Xe IONS AT VELOCITIES UP TO 4×10^9 cm/sec

Permalink

<https://escholarship.org/uc/item/09f018wf>

Authors

Alonso, J.
Gould, H.

Publication Date

1982-02-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

RECEIVED
LAWRENCE
BERKELEY LABORATORY

APR 5 1982

LIBRARY AND
DOCUMENTS SECTION

Submitted to Physical Review Letters


CHARGE CHANGING CROSS SECTIONS FOR Pb AND Xe
IONS AT VELOCITIES UP TO 4×10^9 cm/sec

Jose Alonso and Harvey Gould

February 1982

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782



LBL-14108
2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Charge changing cross sections for Pb and Xe ions at velocities up to 4×10^9 cm/sec.

Jose Alonso and Harvey Gould

Accelerator & Fusion Research Division, Building 64,
and Materials & Molecular Research Division, Building
71, Lawrence Berkeley Laboratory, University of California,
Berkeley, CA 94720

We have made a comprehensive study of charge changing cross sections in N_2 of Pb and Xe ions at velocities near 4×10^9 cm/sec. We determine the velocity, ionization state, and Z dependence of the capture and ionization cross sections and make an extensive comparison with theory.

PACS numbers 34.70+e, 34.50.Hc

Capture or loss of an electron by a fast very heavy complex ion colliding with an atom is one of the more refractory problems in atomic collisions^{1,2}. For very heavy ions at velocities $\geq 10^9$ cm/sec. only a small amount of experimental information is available^{3,4}. Also, there is a general perception that theoretical models used for lighter and simpler ions may be inadequate².

In this letter we present the first comprehensive study of charge changing collisions of very heavy ions at velocities above 10^9 cm/sec. From experimental measurements of Pb ($Z=82$) and Xe ($Z=54$) ions in N_2 at velocities (relative to the speed of light) of $\beta = v/c = 0.134$ and $\beta = 0.099$, and Xe at $\beta = 0.072$ we extract a total of 178 cross sections. We determine the velocity, ion charge, and nuclear charge dependence of these cross sections. In comparison with theory, we find several simple models which adequately describe our results.

The apparatus and experimental methods used are similar to those described in Refs.2,5. In our experiment, the incident ionization states are obtained by magnetically separating a beam of ions accelerated at the Lawrence Berkeley Laboratory's Super-HILAC. These ions then pass through a differentially pumped 24 cm long chamber containing nitrogen gas at a maximum pressure of 1.0 torr. The final ionization states are analyzed in a second electromagnet and are

detected by a position-sensitive gas-filled proportional counter. The raw data collected are the relative number of ions striking the detector at the locations corresponding to the different ionization states as a function of the pressure in the charge exchange chamber.

Cross sections (Fig. 1a,b) were fit to the data by the least squares method using a computer code developed by Betz^{4,5}. Satisfactory fits could be obtained only by including cross sections for the loss of two or more electrons. The sum of the multiple loss cross sections here averages about 15% of the one electron loss cross section. For most ionization states, cross sections were found by fitting data from 2 or 3 different incident ionization states. The variation in a cross section when fit to data from 3 different incident ionization states is the basis of our fitting error. We find a (one standard deviation) fitting error of 14% for the capture cross section and 26% for the loss cross section. These errors are shown as error bars in the lower left of Fig 1a,b. The larger error in the single electron loss cross section is consistent with the larger number of degrees of freedom available when fitting to multiple cross sections. We estimate the systematic error, principally an uncertainty in the thickness of the gas target, to be 20%.

The velocity dependence of the capture cross sections σ_c is extracted by fitting the cross sections in Fig. 1a to $\sigma_c = A\beta^n$ where A is a constant and $\beta = v/c$. Between $\beta=0.099$ and $\beta = 0.134$ we find, for Pb averaged over ionization states +34 to +44, that $n = -5.83$ (0.12); and for Xe +28 to +43, that $n = -5.8$ (0.3). From the Xe⁺²⁷ cross section measured at $\beta = 0.072$ and $\beta = 0.099$ we find $n = -5.9$.

We obtain the ionization state dependence of the capture cross section by fitting to $\sigma_c = Aq^n$ where q is the ionization state. At $\beta = 0.134$ we find for Pb, that $n = 2.9$ (0.4), and for Xe, $n = 3.1$ (0.9). At $\beta = 0.099$, we find for Pb, that $n = 3.0$ (0.3), and for Xe, $n = 3.1$ (0.7); and at $\beta = 0.072$, we find for Xe, $n = 3.3$ (0.3).

The capture cross sections in Fig. 1a exhibit at most a weak dependence on the nuclear charge Z. For $\sigma_c = AZ^n$ we find at $\beta = 0.134$, that $n = 0.2$ (0.1), and at $\beta = 0.099$, $n = -0.2$ (0.2).

Experimental cross sections, and the Oppenheimer-Brinkman-Kramer (OBK) model have been used by several authors^{1,2} to predict the velocity, ionization state, and Z dependence of the capture cross sections for high ionization states of very fast very heavy ions. Janev and Hvelplund¹ propose $\sigma_c = Aq^3$ where q is the ionization state and A a constant, while Betz² suggests $\sigma_c = AZ^0 q^2$ for $q \geq 20$ and $\beta \geq 0.03$; Our experiment is consistent with Z^0 but strongly

favors q^3 over q^2 . On the basis of the OBK model, both authors predict $\sigma_c = A\beta^{-12}$ in the high velocity limit. At $\beta = 0.134$ we find no evidence of this velocity dependence.

Simple calculations based upon semiclassical models⁶⁻¹⁰ have provided some accurate capture cross sections for lighter ions and lower velocities. However their validity for heavy ions at high velocity has been open to question². The results of two semiclassical calculations multiplied by a factor of two to roughly account for molecular nitrogen, are plotted in Fig 1a. Knudsen, Haugen and Hvelplund (KHH)⁹ use a Lenz-Jensen statistical model for the target atom; while Nikolaev¹⁰ (N) uses a treatment similar to Bohr⁶. Both models are in excellent agreement with this experiment.

We determine the velocity dependence of the loss cross sections σ_1 between $\beta = v/c = 0.099$ and $\beta = 0.134$ by fitting the cross sections in Fig. 1b to $\sigma_1 = A\beta^n$ where A is a constant. We find, for Pb averaged over ionization states +36 to +51, that $n = -0.98$ (0.2), and for Xe +27 to +42, $n = -1.56$ (0.25). These values disagree with the β^{-2} dependence predicted for very fast collisions in a Born approximation^{11,12}, and a binary encounter approximation calculation¹³.

The ionization state (q) dependence of the loss cross section σ_1 is found by fitting to $\sigma_1 = Aq^n$. From the data in Fig 1b, we find for Pb, $n = -3.8$ (0.4) at $\beta = 0.134$ and $n =$

-3.3 (0.6) at $\beta = 0.099$. For Xe we find $n = -4.4$ (0.5), $n = -3.3$ (0.4), and $n = -3.1$ (0.7) at $\beta = 0.134$, $\beta = .099$, and $\beta = 0.072$ respectively.

The Z dependence of the loss cross section is obtained by fitting to $\sigma_1 = AZ^n$. We find, for ionization states +36 to +42, that $n = 2.6$ (0.3) for Xe - Pb at $\beta = 0.134$, and $n = 1.9$ (0.2) at $\beta = 0.099$.

To predict the loss cross sections of fast heavy ions in the velocity region $\beta \geq 0.2$ for heavy ion fusion studies, Gillespie¹¹ and others¹² have used a Born approximation sum rule (BASR) and Yu¹³ has used a binary encounter approximation (BEA). For accelerator design studies, Dmitriev, Zaikov, and Tashaev¹⁴ (DZT) have developed a semi-empirical model to predict cross sections over the region $\beta > 0.325/Z$. Loss cross sections calculated at $\beta = 0.2$ by the BASR^{11,12} and BEA¹³ for Hg^{+52} , and Hg^{+34} ($Z=80$); and the BASR¹¹ for Xe^{+36} are indicated in Fig. 1b. The BASR cross sections (at $\beta = 0.2$) are at least a factor of five larger than our experimental values (at $\beta = 0.134$). The BEA calculation is within a factor of 2-3. Correction for the difference in β would make the experimental disagreement with the BASR and BEA calculations larger. The DZT semi-empirical values¹⁴ (Fig. 1b) for U^{+60} ($Z=92$) and U^{+40} at $\beta = 0.134$ and $\beta = 0.099$ are a good match to the experimental cross sections for Pb, while their values for I^{+40} ($Z=53$) and I^{+30} at $\beta = .099$ and β

=.072 underestimate our Xe cross sections by only a factor of 2.5.

Good agreement with all of our data is obtained with a semi-classical (SC) model. We start with Bohr's formula for the loss cross section, from eq. 3.1.3 of Ref.6:

$$\sigma_1 = 2\pi a_0^2 Z_T^2 (\alpha/\beta)^2 \sum_j \left(\frac{1}{I_j} - \frac{1}{2\beta^2} \right)$$

where Z_T is the effective charge of the target, I_j is the binding energy (in units of the rest mass of the electron) of the j^{th} electron in the ion, and the sum is over all electrons j for which $2\beta^2 > I_j$. However, instead of a statistical model for the electron binding energies used by Bohr and Lindhard (Ref. 8 p15), we use binding energies obtained from a relativistically corrected Hartree Fock calculation¹⁵. We neglect excitation to autoionizing or metastable states. The single electron loss cross sections in N_2 obtained from the calculation are plotted in Fig.1b. The agreement with experiment is on average within a factor of two, and better at the lower ionization states and lower Z .

The velocity dependence of the loss cross sections in the SC model is largely determined by the ratio of the projectile velocity β to the binding energy of its electrons I_j . When β is very large compared to I_j for most of the electrons in the ion, the velocity dependence can approach β^{-2} . However at velocities (where the kinetic energy is)

just above I_j of the most loosely bound electrons, the slope of σ_1 versus β can become zero or assume positive values.

The loss cross sections for partially ionized Pb and U at velocities $\beta \geq 0.2$ are required for the use of fast heavy ions as igniters for pellet fusion. The cross sections are critical in determining the (vacuum) conditions for acceleration and final focusing of the ions in the reaction vessel¹⁶⁻¹⁸, and in determining the energy deposition in the pellet. Our SC calculation of the cross section for single electron loss in N_2 for Pb^{+2} at $\beta = 0.2$ yields a value of $1.4 \times 10^{-16} \text{ cm}^2 / \text{molecule}$.

In summary, from measured charge changing cross sections of Pb and Xe in N_2 we find that in the region $\beta \approx 0.1$ the capture cross sections scale approximately as $Z^0 q^3 \beta^{-6}$ and are in good agreement with semi-classical models. We also find that the loss cross sections decrease more slowly than β^{-2} and are in disagreement with a Born approximation calculation. Agreement however is good with the DZT semiempirical model and a Bohr semi-classical calculation.

Acknowledgments

We thank Dr. Hermann Grunder and Professor Richard Marrus for their enthusiastic support; Mr. Douglas MacDonald and Ms. Jeannette Mahoney for mechanical design and fabrication; Dr. Hans Betz for use of his computer code, Ms.


Rebecca Clark for assistance in data analysis; and the operators and staff of the Super-HILAC, whose skill and dedication made this experiment possible. This work was supported by the Director, Office of Energy Research: Office of Basic Energy Sciences, Chemical Sciences Division, and Office of High Energy and Nuclear Physics, Nuclear Science Division, of U.S. Department of Energy, under Contract No. DE-AC03-76SF00098.

Footnotes

- (1) R.K. Janev, and P. Hvelplund, Comments At. Mol. Phys. 11, 75 (1981).
- (2) H.D. Betz, in "Methods of Experimental Physics: Atomic Physics, Accelerators", edited by P. Richard (Academic Press, N.Y. 1980), Vol 17, p73.
- (3) B. Franzke, IEEE Trans. Nucl. Sci. NS-28, 2116 (1981).
- (4) S. Datz, H.O. Lutz, L.B. Bridwell, C.D. Moak, H.D. Betz, and L.D. Ellsworth, Phys. Rev. A2, 430 (1970).
- (5) H.D. Betz, Rev. Mod. Phys. 44, 465 (1972), and references contained therein.
- (6) N.Bohr, K. Dan. Vidensk. Selsk, Mat.-Fys. Medd. 18, No 8 (1948).
- (7) G.I. Bell, Phys. Rev. 90, 548 (1953); R.L. Gluckstern, Phys. Rev. 98, 1817 (1955);
- (8) N.Bohr and J. Lindhard, K. Dan. Vidensk. Selsk, Mat.-Fys. Medd. 28 No 7 (1954).
- (9) H. Knudsen, H.K. Haugen and P. Hvelplund, Phys. Rev. A23, 597 (1981).
- (10) V.S. Nikolaev, Zh. Eksp. Teor. Fiz. 33, 534 (1957) [Sov. Phys.- JETP 6, 417 (1957)]. For comparison with our experiment we use $n=7$ and $q=1$.

- (11) G.H. Gillespie, See Ref.16, p45; Nucl. Instrum. Methods 176, 611 (1980)
- (12) K.T. Cheng, G. Das, Y.-K. Kim, and R.C. Raffanetti, See Ref.16, p43; G.H. Gillespie, K.T. Cheng, and Y.-K. Kim, Proc. of the Heavy Ion Fusion Workshop held at Argonne National Lab. Sept.19-26, 1978, edited by R.C. Arnold. Argonne National Lab. Pub. ANL-79-41 (1979). p175.
- (13) S. Yu, See Ref.16, p50.
- (14) I.S. Dmitriev, V.P. Zaikov, and Yu. A. Tashaev, Nucl. Instrum. Methods 164, 329 (1979).
- (15) R.D. Cowan, Phys. Rev. 163, 54 (1967); R.D. Cowan, and D.C. Griffin, J. Op. Soc. Am. 66, 1010 (1976).
- (16) Proc. of the Heavy Ion Workshop held at Brookhaven National Lab. Oct 17-21, 1977, edited by L.W. Smith, Brookhaven National Lab Report BNL 50769 (1978). p41-58, 136.
- (17) R.F. Hubbard and D.A. Tidman, Phys. Rev. Lett. 41, 866 (1978).
- (18) Report on the Workshop on Atomic and Plasma Phys. Requirements for Heavy Ion Fusion, Argonne National Lab., Dec. 13-14, 1979, Argonne National Lab. Report ANL-80-17 (1980).

Figure Caption Figs. 1a and 1b show the capture (a) and single electron loss (b) cross sections in N_2 as a function of the ionization state for Pb and Xe ions. Experiment is shown as open symbols, theory with the corresponding closed symbols and with broken and solid lines. In (a) KHH is Ref. 9, and N is Ref. 10. In (b) theory in closed symbols should be compared to experiment with corresponding open symbols. BASR (Ref. 11,12) and BEA (Ref 13) were calculated for Hg ($Z=80$) and Xe at $\beta = 0.2$ and DZT (Ref. 14) was calculated for U ($Z=92$) and I ($Z=53$) at $\beta= 0.134$ and $\beta= 0.099$. SC is semi-classical (this work and Ref.6).



XBL 822-8112

Fig. 1

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720