Lawrence Berkeley National Laboratory

Recent Work

Title

Phylotype Dynamics of Bacterial P Utilization Genes in Microbialites and Bacterioplankton of a Monomictic Endorheic Lake.

Permalink

<https://escholarship.org/uc/item/0b29w5mr>

Journal

Microbial ecology, 73(2)

ISSN

0095-3628

Authors

Valdespino-Castillo, Patricia M
Valdespino-Castillo, Patricia M
Alcántara-Hernández, Rocío J
et al.

Publication Date

2017-02-01

DOI

10.1007/s00248-016-0862-1

Peer reviewed

Phylotype Dynamics of Bacterial P Utilization Genes in Microbialites and Bacterioplankton of a Monomictic Endorheic Lake

P
a
t
r
i
c
i
a

M
.

V
a
l
d
e
s
p
i
n
o
-
C
a
s
t
i
l
l
o
1
,
2

&

R
o
c
í
o

J

.
Alcánt
ara-
Herná
ndez³
&
Martín
Merin
o-
Ibarra⁴
&
Javier
Alcoc
er⁵ &
Mirosł
av
Macek
^{5,6} &
Octavi
o A.
Moren
o-
Guillé
n^{1,7} &
Luisa
I.
Falcón
1

spaceAbstract Microbes can modulate ecosystem function since they harbor a vast genetic potential for biogeochemical cycling. The spatial and temporal dynamics of this genetic diversity should be acknowledged to establish a link between ecosystem function and community structure. In this study, we analyzed the genetic diversity of bacterial phosphorus utilization genes in two microbial assemblages, microbialites and bacterioplankton of Lake Alchichica, a semiclosed (i.e., endorheic) system with marked seasonality that varies in nutrient conditions, temperature, dissolved oxygen, and water column stability. We focused on dissolved organic phosphorus (DOP) utilization gene dynamics during contrasting mixing and stratification periods. Bacterial alkaline phosphatases (*phoX* and *phoD*) and alkaline beta-propeller phytases (*bpp*) were surveyed. DOP utilization genes showed different dynamics evidenced by a marked change within an intra-annual period and a differential circadian pattern of expression. Although Lake Alchichica is a semiclosed system, this dynamic turnover of phylotypes (from lake circulation to stratification) points to a different potential of DOP utilization by the microbial communities within periods. DOP utilization gene dynamics was different among genetic markers and among assemblages (microbialite vs. bacterioplankton). As estimated by the system's P mass balance, P inputs and outputs were

similar in magnitude (difference was <10 %). A theoretical estimation of water column P monoesters was used to calculate the potential P fraction that can be remineralized on an annual basis. Overall, bacterial groups including Proteobacteria (Alpha and Gamma) and Bacteroidetes seem to be key participants in DOP utilization responses.

Keywords Extracellular enzymes · DOP utilization · Phytase · P turnover · Phylotype seasonality · Microbial functional diversity space

Luisa I. Falcón
falcon@ecologia.unam.mx

¹ Laboratorio de Ecología Bacteriana, Instituto de Ecología, Universidad Nacional Autónoma de México, 04510 Mexico City, Mexico

² Secretaría de Ciencia, Tecnología e Innovación del Distrito Federal-Centro Latino-Americano de Física, Olivo 39, Col. Florida, 01030 Mexico City, Mexico

³ Instituto de Geología, Universidad Nacional Autónoma de México, 04510 Mexico City, Mexico

⁴ Unidad Académica de Ecología y Biodiversidad Acuática, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, 04510 Mexico City, Mexico

⁵ Proyecto de Investigación en Limnología Tropical, FES Iztacala, UNAM, Av. de los Barrios No. 1, Los Reyes Iztacala, 54090 Tlalnepantla, State of Mexico, Mexico

⁶ Biology Centre v. v. i., Institute of Hydrobiology, Academy of Sciences of the Czech Republic, 37001 České Budějovice, Czech Republic

⁷ Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, 04510 Mexico City, Mexico

Introduction

D
i
s
s
o
l
v
e
d

o

r
g
a
n
i
c

p
h
o
s
p
h
o
r
u
s

(
D
O
P
)

i
s

a
n

a
b
u
n
d
a
n
t

p
h
o
s
-

p
h
o
r
u
s

(
P
)

able 1Phylotype assignment of DOP utilization genes *phoD*, *phoX*, and *bpp*; richness and diversity estimators; as well as coverage for each group (microbials and bacterioplankton) and period (circulation and stratification) are shown

<i>phoX</i> (sequences <i>opp</i> (sequences obtained = 58)			
SCoverage (%)	OTUs	Chao	Shannon (<i>H'</i>) Coverage (%)
2 88	15	33	2.3527 5 74

f
r
a
c
t
i
o
n

i
n

a
q
u
a
t
i
c

s
y
s
t
e
m
s
,

r
e
p
r
e
s
e
n
t
i
n
g

u
p

t
o

8
0

%

o

f

t
h
e

t
o
t
a
l

d
i
s
s
o
l
v
e
d

p
h
o
s
p
h
o
r
u
s

i
n

t
h
e

w
a
t
e
r

c
o
l
u
m
n

o
f

o
l
i
g
o
t
r
o
p
h
i
c

s
y
s
t
e
m
s

[
[1](#)
,

[2](#)
]
.

Y
e
t

D
O
P

i
s

t
h
e

l
e
s
s

k
n
o
w
n

P

f
r
a
c
t
i
o
n

[
[3](#)
]
.

O
r
g
a
n
i
c

p
h
o
s
p
h
o
r
u
s

t
r
a
n
s
f
o
r
m
a
t
i
o
n
s

o
c
c

u	f
r	a
	c
e	e
f	
f	w
i	a
-	t
	e
c	r
i	
e	r
n	a
t	p
l	i
y	d
i	P
n	-
	t
n	u
a	r
t	n
u	o
r	v
a	e
l	r
s	r
y	a
s	t
t	e
e	s
m	
s	[
,	2
	,
a	
s	4
	]
r	.
e	
v	W
e	e
a	
l	k
e	n
d	o
	w
b	
y	s
	o
s	
u	f
r	a

r	i
	n
t	g
h	
a	d
t	i
	s
m	s
i	o
c	l
r	v
o	e
b	d
e	
s	o
	r
a	g
r	a
e	n
	i
k	c
e	
y	n
	u
c	t
o	r
m	i
p	e
o	n
n	t
e	
n	f
t	r
s	a
	c
i	t
n	i
v	o
o	n
l	s
v	
e	[
d	5
	,
i	
n	6
	]
r	;
e	
u	t
t	h
i	e
l	r
i	e
z	f

o	a
r	t
e	i
,	o
	n
D	s
O	
P	f
	o
u	r
t	
i	c
l	a
i	r
z	b
a	o
t	n
i	
o	u
n	p
	t
b	a
y	k
	e
m	
i	a
c	n
r	d
o	
b	e
e	x
s	p
	o
h	r
a	t
s	a
	t
s	i
i	o
g	n
n	
i	[
f	Z
i	]
c	.
a	
n	A
t	
	r
i	e
m	c
p	o
l	g
i	n
c	i

zed
D
O
P
t
r
a
n
s
f
o
r
m
a
t
i
o
n
p
r
o
c
e
s
s
i
s
t
h
e
s
y
n
t
h
e
s
i
s
o
f
e
x
t
r

a
c
e
l
l
u
l
a
r
p
h
o
s
p
h
a
t
a
s
e
s
t
h
a
t
b
e
c
o
m
e
a
c
t
i
v
e
t
o
f
a
c
e
P
s
t
a

r	e
v	r
a	s
t	
i	a
o	s
n	
	t
i	h
n	e
t	m
h	a
e	i
	n
w	
a	s
t	u
e	b
r	s
	t
c	r
o	a
l	t
u	e
m	s
n	
	[
w	8
i	–
t	1
h	1
	]
p	.
m	A
o	n
n	o
o	t
e	h
s	e
t	r
e	
r	i
s	n
	t
a	e
n	r
d	e
	s
d	t
i	i
e	n
s	g
t	

e	p
x	h
a	y
m	t
p	a
l	t
e	e
f	(
o	m
r	y
o	o
r	-
g	i
a	n
n	o
i	s
c	i
p	t
h	o
o	l
s	h
p	e
h	x
o	a
r	k
u	i
s	s
u	p
t	h
i	o
l	s
i	p
z	h
a	a
t	t
i	e
o	o
n	r
c	I
o	P
n	6
s	)
i	d
s	e
t	g
s	r
o	a
f	d
	a

t	t
i	h
o	e
n	
.	t
	o
p	t
h	a
y	l
t	
a	p
t	h
e	o
	s
r	p
e	h
p	o
r	r
e	u
-	s
s	f
e	o
n	u
t	n
s	d
a	i
p	n
p	
r	n
o	a
x	t
i	u
m	r
a	e
t	
e	[
l	1
y	2
	]
7	.
5	
	T
t	h
o	i
	s
8	
0	m
	o
%	l
	e
o	c
f	u
	l

e
,
a
b
u
n
d
a
n
t

i
n

v
e
g
e
t
a
l

t
i
s
s
u
e
s
,
a
c
c
u
m
u
l
a
t
e
s

i
n

t
h
e

s
o
i
l

b
u
t

i
s

a

m
i
n
o
r

c
o
m
p
o
n
e
n
t

o
f

m
a
r
i
n
e

s
e
d
i
m
e
n
t
s
,

i
n
d
i
c
a
t
i

n	o
g	l
	u
i	m
t	n
s	
	[
e	1
f	3
f	]
i	.
c	
i	R
e	e
n	c
t	e
	n
d	t
e	l
g	y
r	,
a	
d	a
a	
t	b
i	a
o	c
n	t
/	e
u	r
t	i
i	a
l	l
i	
z	a
a	l
t	k
i	a
o	l
n	i
	n
i	e
n	
	p
t	h
h	y
e	t
	a
w	s
a	e
t	
e	(
r	b
	e
c	-

t
a
-
p
r
o
p
e
l
l
e
r

p
h
y
t
a
s
e
,

b
p
p
)

w
a
s

i
d
e
n
t
i
f
i
e
d

i
n

a

v
a
r
i
e
t
y

o
f

b
i
o
-

l
o
g
i
c
a
l

s
y
s
t
e
m
s
,

r
e
v
e
a
l
i
n
g

i
t
s

v
a
s
t

p
o
t
e
n
t
i
a
l

for
o
r

p
h
y
t
a
t
e

d
e
g
r
a
-

d
a
t
i
o
n

[
[1](#)
[4](#)
,

[1](#)
[5](#)
]
.

P
h
y
t
a
s
e
s

h
y
d
r
o
l
y
z
e

p
h
y
t
a
t
e

i
n
t
o

l
e
s
s

p
h
o
s
-

p
h
o
r
y
l
a
t
e
d

m
y
o
-
i
n
o
s
i
t
o
l

r
e
s
i
d
u
e

s
o
r
e
l
e
a
s
e
s
m
e
t
a
l
s
c
h
e
l
a
t
e
d
b
y
p
h
y
t
a
t
e
(
e
.
g
.
,
F
e
+
3
,
Z
n
+
2
a
n
d
i
n
o
r
g
a
n
i
c
p
h
o
s
p
h
a
t
e
s
[
[1](#)
[6](#)
],
,
a
n
d
t
h
e
i
r
a
c
t
i
v
i
t
y
a
l
s

a	m
r	s
t	
i	t
n	h
g	a
	t
t	
o	h
	a
u	r
n	b
d	o
e	r
r	
s	P
t	
a	t
n	r
d	a
	n
t	s
h	f
e	o
	r
i	m
d	a
e	t
n	i
-	o
	n
t	
i	g
t	e
y	n
	e
o	s
f	
	[
t	3
h	,
e	
	1
m	8
i	]
c	,
r	
o	t
o	h
r	e
g	
a	f
n	a
i	c
s	t

o	f
r	
s	P
t	t
h	r
a	a
t	n
	s
d	f
r	o
i	r
v	m
e	a
	t
t	i
h	o
e	n
p	g
r	e
e	n
s	e
e	s
n	/
c	p
e	r
,	o
	t
d	e
i	i
v	n
e	s
r	
s	[
i	<u>1</u>
t	<u>0</u>
y	,
,	
	<u>1</u>
o	<u>1</u>
r	,
e	<u>1</u>
x	<u>8</u>
p	,
r	
e	<u>1</u>
s	<u>9</u>
s	]
i	,
o	
n	a
	s
o	

w
e
l
l

a
s

t
h
e
i
r

s
p
a
t
i
a
l

a
n
d

t
e
m
p
o
r
a
l

r
e
p
l
a
c
e
m
e
n
t

o
r

t
u
r
n
o

v
e
r

[
[4](#)
,

[2](#)
[0](#)
]
.

R
e
m
i
n
e
r
a
l
i
z
a
t
i
o
n

p
a
t
t
e
r
n
s

a
n
d

r
a
t
e
s

m
a
y

v
a

r	b
y	u
	t
d	
u	t
e	h
	e
t	
o	p
	a
s	t
h	t
i	e
f	r
t	n
s	s
i	a
n	n
	d
b	
a	m
c	e
t	c
e	h
r	a
i	n
a	i
l	s
	m
a	s
s	
s	o
e	f
m	
b	f
l	u
a	n
g	c
e	t
s	i
	o
[	n
4	a
,	l
2	g
1	e
,	n
	e
2	
2	t
]	u
	r
,	n

o	n
v	s
e	f
r	o
	r
s	m
t	a
i	t
l	i
l	o
	n
r	
e	p
m	o
a	t
i	e
n	n
	t
u	i
n	a
c	l
l	
e	i
a	n
r	
.	m
H	i
e	c
r	r
e	o
,	b
	i
w	a
e	l
	i
s	t
t	e
u	s
d	
i	a
e	n
d	d
t	b
h	a
e	c
	t
D	e
O	r
P	i
	o
t	p
r	l
a	a
	n

k	r
t	s
o	
n	h
	a
o	v
f	e
L	s
a	h
k	o
e	w
	n
A	
l	t
c	h
h	a
i	t
c	
h	a
i	l
c	k
a	a
.	l
	i
R	n
e	e
c	
e	p
n	h
t	o
	s
s	p
u	h
r	a
v	t
e	a
y	s
s	e
	s
i	
n	(
	A
n	P
a	)
t	
u	<i>p</i>
r	<i>h</i>
a	<i>o</i>
l	<i>D</i>
w	a
a	n
t	d
e	

p
h
o
X

a
r
e

m
o
r
e

a
b
u
n
d
a
n
t

t
h
a
n

p
h
o
A
,

t
h
e

c
l
a
s
s
i
c
a
l

m
o
d
e
l

t

o

s
t
u
d
y

a
l
k
a
l
i
n
e

p
h
o
s
p
h
a
t
a
s
e
s

[
[1](#)
[0](#)
,

[1](#)
[1](#)
]

.

H
e
r
e
,

t
h
e
s
e

t
w
o

b
a
c
t
e
r
i
a
l

a
l
k
a
l
i
n
e

p
h
o
s
p
h
a
t
a
s
e
s
,

p
h
o
X

a
n
d

p
h
o
D
,

a
n
d

a
l

k
a
l
i
n
e

b
e
t
a
-
p
r
o
p
e
l
l
e
r

p
h
y
t
a
s
e
s
,

b
p
p
,

w
e
r
e

s
u
r
v
e
y
e
d

i
n

e

n
v
i
r
o
n
m
e
n
t
a
l

D
N
A

a
n
d

R
N
A

(
T
a
b
l
e

[1](#)
)
.

S
i
n
c
e

s
h
i
f
t
s

i
n

t
h
e

b
a
c
t
e
r
i
a
l

a
s
s
e
m
b
l
a
g
e
s

s
t
r
u
c
t
u
r
e

c
o
u
l
d

a
f
f
e
c
t

r
e
m
i
n
e
r
a
l

l
z
a
t
i
o
n

p
a
t
t
e
r
n
s

[
[4](#)
,

[2](#)
[1](#)
]
,

t
h
e

d
y
n
a
m
i
c
s

o
f

t
h
e
s
e

c
a
l
c
i
u
m

-
d
e
p
e
n
d
e
n
t

m
e
t
a
l
l
o
e
n
z
y
m
e
s

w
a
s

f
o
l
l
o
w
e
d

d
u
r
i
n
g

l
a
k
e

c
i
r
c

u
l
a
t
i
o
n

a
n
d

s
t
r
a
t
i
f
i
c
a
t
i
o
n

p
e
r
i
o
d
s
,

w
h
i
c
h

o
f
f
e
r

c
o
n
t
r
a
s

t
i
n
g

p
h
y
s
i
c
o
c
h
e
m
i
c
a
l

c
o
n
d
i
t
i
o
n
s

w
i
t
h
i
n

t
h
e

s
y
s
t
e
m
.

A

p
h

o
s
p
h
o
r
u
s

m
a
s
s

b
a
l
a
n
c
e

w
a
s

p
e
r
f
o
r
m
e
d

t
o

f
r
a
m
e

P

s
t
a
t
u
s

a

n
d

u
t
i
l
i
z
a
t
i
o
n

i
n

L
a
k
e

A
l
c
h
i
c
h
i
c
a
.
space

Methods

Study Area

Alchichica is a maar lake located in the easternmost region of the Trans-Volcanic Belt, Puebla, Mexico (19° 24' N, 97° 24' W; 2,340 m a.s.l.). It is a saline and alkaline lake (salinity 8.5 g l⁻¹, pH 9.5) that exhibits a particular chemical signature where sodium is abundant [23–25], calcium concentration is particularly low (~0.3 mM), and Mg/Ca ratio is high. Alchichica's water column production seems to be limited mainly by nitrogen, although phosphorus and silica may alter- natively limit phytoplankton productivity [26, 27]. Lake Alchichica is warm-monomictic [28, 29]. Stratification period occurs from April to December and circulation, in the dry winter, from January to March; the water column (Z_{mean} = 40.9 m and Z_{max} = 62 m) mixes completely by the end of January or early February. Stratification and circulation pe- riods offer contrasting (intra-annual) physicochemical and nutrimental conditions [19] within the system (Fig. 1).

Two different microbial assemblages characterize Alchichica's microbiota: bacterioplankton and a littoral ring of microbialites. Macek et al. [30] found that picoplankton dynamics follow a regular pattern, linked to the hydrological cycle of this system. Nutrient availability relates to the tempo-ral pattern and the vertical zonation of the bacterioplankton components [31]. Two microbialite morphotypes can be found in Lake Alchichica (columnar and spongy morphology); their microbial composition shows high prokaryote diversity [32]. Microbialites have shown relevant activity related to the N cycle, such as high N_2 fixation rates [33, 34] as well as potential for DOP utilization through bacterial alkaline phosphatases [19].

Environmental Characterization

Water samples from the littoral zone, central station of the lake with depths of 5 m (surface mixed layer), 25 m (metalimnion), and 61 m (1 m above the bottom)—detailed description of the thermal structure of Alchichica can be consulted in Macek et al. [30]—and from nearby wells were taken in 2011: during lake circulation (February) and the well-established stratification (August). Physicochemical characterization of the water column and the littoral zone (water surrounding the microbialites) included temperature, dissolved oxygen (DO), pH, and conductivity of the water column and of the water surrounding the microbialites, recorded with a YSI 6600 multi-parameter probe. Water samples for nutrient analysis included determinations of soluble reactive phosphorus (SRP), NO_3^- , NO_2^- , dissolved inorganic nitrogen (DIN), soluble

spacein polypropylene containers. Samples were analyzed (within 24 to 48 h) with a segmented flow Autoanalyzer (Skalar San-Plus) following the standard methods adapted by Grasshoff et al. [35] and the circuits indicated by Kirkwood [36]. Total P and N were analyzed following the methods proposed by Valderrama [37], and total particulate P (TPP) was calculated as the difference between total P – SRP (this represents the minimum benchmark of organic phosphorus since SRP contains some organic species).

Sampling

Microbialites and bacterioplankton for total DNA and RNA surveys were collected in both periods (circulation and stratification). The two morphologically different microbialites (spongy and columnar) were sampled in six stations ($n = 6$ for each morphotype, at a depth of ~ 0.30 m) along the lakes littoral zone. Subsamples of ~ 10 g were excised and kept frozen ($-20^\circ C$) until DNA extraction. After

extraction, DNA from all six stations was mixed to obtain a single pooled DNA sample for each microbialite morphotype per period.

At the central station of the lake, water column samples of 0.5 l (triplicates) were filtered through polycarbonate membranes (pore size $0.22\ \mu m$, Osmonics; Poretics Corp.) taken at the following water depths: 5 m (surface), 25 m (metalimnion), and 61 m (bottom). Filters were then collected and kept frozen ($-20^\circ C$) in DNA-free 2.0 ml tubes, until analysis.

Collection of samples for total RNA extraction was performed in a diel sampling scheme. Microbialite sections from all sampling stations (six microbialite samples) were incubated at on-site conditions (light and temperature). Subsamples were taken (in RNase-free surfaces) at 0600, 1200, 1800, and 2400 hours; placed in liquid nitrogen; and kept frozen at $-80^\circ C$ until analysis.

Nucleic Acid Extraction

DNA extraction was performed using the method of Zhou et al. [38] modified by Centeno et al. [32] in which samples of approximately 5 g of each microbialite were pulverized in liquid nitrogen with a buffer solution (100 mM Tris-HCl, 20 mM NaCl, 100 mM EDTA, pH 8) and 0.06 V of cetyl trimethylammonium bromide (CTAB). A total volume of

0.5 l of water was filtered in $0.22\text{-}\mu m$ membranes (Durapore, Millipore) to obtain total DNA from bacterioplankton samples. A volume of 3 ml of the extraction buffer was added and the mixtures were then incubated with lysozyme (30 mg ml^{-1}) (Sigma Aldrich, USA) at $37^\circ C$ for

30 min. A second incubation followed, adding proteinase K (10 mg ml^{-1} , Sigma Aldrich, USA) and 0.1 V of sodium dodecyl sulfate (SDS) at $55^\circ C$, overnight. Incubations were then centrifuged (20 min, $1800g$). The aqueous phase of each

Fig. 1 a–c Physicochemical conditions associated to microbialites (littoral zone) and bacterioplankton (water column) during contrasting lake periods: stratification (summer) and circulation (winter). Profiles (on top) show water column conditions and a table (below) shows littoral zone ambient conditions

spacea. Lake circulation

frozen in liquid nitrogen in a circadian sampling scheme (0600, 1200, 1800, and 2400 hours). Total RNA extraction was performed using the RNA PowerSoil® Total RNA Isolation Kit (Mo Bio Laboratories, Carlsbad, CA) following the protocol of the manufacturer. RNA was then purified with the RNeasy Mini Kit (Qiagen, Venlo, Netherlands) including a step with DNaseI (Qiagen) to avoid DNA contamination. Immediately after RNA purification, primer-conducted complementary DNA (cDNA) synthesis was performed using the Reverse Transcription System (Promega, WI, USA) with

space

space 0.5 µg of each primer (Invitrogen). Reactions of 20 µl were incubated in the following sequence: 10 min at room temperature, 15 min at 42 °C, and finally 5 min at 95 °C to inactivate reverse transcriptase. cDNA obtained was kept frozen (at –20 °C) until analysis. Samples were verified for DNA contamination after DNase incubation with PCR before reverse transcription.

Amplification of DOP Utilization Genes

Partial sequences of alkaline phosphatases (*phoX* and *phoD*) and beta-propeller phytase (*bpp*) were amplified using reported degenerate primers [10, 15, 39]. PCR-specific conditions, primers used, and partial sequence length are shown in Table 2 (Online Resource 1). In a general scheme, PCRs were performed using ~10 ng of DNA and cDNA in a final reaction mixture of 25 µl, containing 2.5 µl of 10× ViBuffer A (Vivantis, Oceanside, USA), a specific MgCl₂ concentration (Table 2, Online Resource 1), 0.2 mM of each deoxynucleotide triphosphate, 0.5 µg µl⁻¹ bovine serum albumin (BSA, Biolabs, New England), and 0.5 U of Taq DNA Polymerase (Vivantis, Oceanside, USA). PCR products obtained from five reactions were pooled and then gel purified using QIAquick spin columns (Qiagen). DNA and cDNA amplicons were used to build clone libraries for each season and sample. Amplified fragments were ligated to pCR®2.1 vector following the manufacturer's suggestions (Original TA Cloning Kit, Invitrogen, Carlsbad, CA, USA); *Escherichia coli* TOP-10 chemically competent cells were transformed and selected with ampicillin (100 µg ml⁻¹) and X-Gal (80 µg ml⁻¹); M13 primers were used to screen the fragments (~100 transformant clones were picked for each DNA sample, and transcript libraries had a lower efficiency, with ~50 clones per sample).

Sequence Analyses

To explore diversity and dynamics of the genetic markers surveyed, operational taxonomic units (OTUs) were defined and considered as P-gene phylotypes. Phylotype assignment was done using cutoff values of 0.25 and 0.05 (see Table 1), following the similitude cutoff of 0.25 has been defined in

spacesilico for *phoD* [40], and cutoff value of 0.05 suggested for the analysis of functional genes [41]. Although both cutoff values (0.25 and 0.05) were used for OTU assignment (Table 1), a theoretical analysis of *phoD* sequences indicates a specific sequence cutoff of 0.25 [40], and the resulting phylotypes were used further to compare phylotypes among groups and periods (Fig. 3). In contrast, while no specific cutoff values have been determined for *phoX* and *bpp* (perhaps due to the complexity for the reconstruction of the evolution of these proteins), a cutoff of 0.05, suggested for functional genes [41], was used to examine phylotype dynamics of *phoX* and *bpp* (Fig. 3).

Using the predicted amino acid sequences, alignments of each marker (using the Multiple Sequence Alignment software MUSCLE) and reference sequences of phosphatases (*phoX* and *phoD*) and *bpp* phytases were analyzed and trees were constructed with neighbor-joining and maximum likelihood algorithms (PhyML 3.0) [42]. Functions Bdist.seqs^ and Bcluster^ (using the furthest neighbor algorithm) of the platform for bioinformatic analyses MOTHUR v.1.33.3 [43] were used to group sequences into OTUs and to explore diversity within each marker (nonparametric richness estimator Chao, Shannon diversity index, as well as coverage, are reported in Table 1). The exploration of the sequence sets of each microbial consortia and season (stratification or circulation) was used to construct Venn diagrams showing the overlap and turnover of the markers explored.

Lake Alchichica AP (*phoX* and *phoD* partial gene-nucleotide sequences) from microbialites and bacterioplankton can be consulted in GenBank under the accession numbers KF891484–KF891515, KF891517–KF891828, and KF891830–KF891882. PhoD-like transcripts (nucleotide and predicted amino acid sequences) were identified using the non-redundant (NR) database of NCBI BLAST. Sequences obtained have been submitted to the GenBank database under accession numbers KT763020–KT763031. Sequences shorter than 200 nucleotides are compiled in fasta archives (Online Resources 2, 3, and 4). Their affiliation and general features were explored through searches that correspond to the particular features of each set of sequences (see transcript sequences analyses in Online Resource 1).

space

Table 2 Online Resource 1. Primers and PCR amplification conditions for each marker

Marker	[Primer] (mM)	MgCl ₂ (mM)	PCR conditions
phoX	0.4	1.4	Denaturalization (95 °C for 5 min) 35 cycles (94 °C for 30 s, 52 °C for 30 s, and 72 °C for 1 min); spacefinal extension (72 °C for 10 min)
phoD	2	2	Denaturalization (94 °C for 3 min) 35 cycles (94 °C for 1 min, 57 °C for 1 min, and 72 °C for 2 min); spacefinal
BPP	1	2	Denaturalization (95 °C for 4 min), 8 cycles (95 °C for 30 s, 57 °C for 30 s, diminishing 1 °C each cycle and 72 °C for 30 s); 27 cycles (95 °C for 30 s, 48 °C for 30 s, and 72 °C for 5 min)

space
[39]

[15]

space

Database Searches and Structural Determination of PhoX and BPP transcripts

PhoX transcripts and BPP sequences were explored manually (using and modifying the default search parameters of HMMER, nBLAST, and pBLAST algorithms) [44–46] to confirm the identity of the aforementioned sequences. A detailed description of the strategies followed to perform these analyses can be consulted in Online Resource 1. For sequences less than 18 amino acids in length (some phoX transcripts), we decided to analyze the linear predisposition of the amino acids in the sequence to form regular secondary structures [47]. Due to sequence length limitations, we considered the overall hydrophobicity, using the scale of Kyte and Doolittle [48] of a linear string of amino acids [49–51].

System's Water Balance

The water balance of Lake Alchichica was modeled by García Martínez [52] using a continuity algorithm. Alchichica water balance was estimated in an annual basis using primary and secondary data [53]. Primary data were obtained through direct measurements (e.g., wells), while secondary data were obtained from reported literature and databases. The water balance considered the following climatic data: temperature, precipitation, evaporation, real (ETR) and potential (ETP) evapotranspiration, as well as the surface area of the lake, altitude, and static level [52].

Phosphorus Mass Balance

A stationary state was assumed ($\Delta P/\Delta T = 0$) because Lake Alchichica's trophic status has exhibited relatively low variation in SRP, TP, or SiRP concentrations in recent decades [54–56], and a whole system P balance was

performed. Data for the estimation of the phosphorus mass balance included primary and secondary data. Phosphorus balance is defined by the equation provided below as the difference between P inputs and outputs in a time basis to estimate the difference in magnitude between in- and outflows [57].

cycles (94 °C for 30 s, 52 °C for 30 s, and 72 °C for 1 min);

$$\Delta M \delta P \Delta t \frac{1}{4} \sum \delta P \text{ inputs} - P \text{ outputs}$$

$$\pm P \text{ Net Internal Processes } \delta P_{\text{NIPP}} \delta P$$

where $\Delta M(P)/\Delta t$ is the mass change of P within the system and was evaluated numerically from the volume-weighted concentrations and volume data of the lake. P inputs were computed and estimated for each water source or outflow (Table 6).

Some potential P sources were not considered for the P mass balance due to the following reasons: (1) local sewage inflows or water withdrawal was considered negligible (there is no information or evidence about their existence), (2) net P internal flow

(no studies on P capture and release have been reported for the lake), and (3) P precipitation in OH-apatite was not considered since this mineral is unsaturated in Alchichica water column (OH-apatite saturation index = −2.97) [25]. After these considerations, Lake Alchichica's P balance (Table 6) was constrained to the equation:

P Precipitation $\pm$ P Atmospheric deposition

$\pm$ P Basin runoff $\pm$ P groundwater

$\frac{1}{4}$ P Sedimentation

δP

To estimate the ratio of total P mass/P sedimentation flux (Table 6), lake's total P mass was calculated using the concentration on TP in each layer of the lake times the volume of each layer.

Results

Genetic Diversity of Bacterial P Utilization Genes

Microbialite and bacterioplankton DOP utilization genes showed affiliation with Actinobacteria (*phoD*), Alphaproteobacteria (*phoX*, *phoD*), Betaproteobacteria (*phoD*), Gammaproteobacteria (*phoD* and *bpp*), and Bacteroidetes (*bpp*).

Partial phytase sequences recovered from environmental DNA of Lake Alchichica's bacterioplankton and microbialites (~200 bp) were similar to *bpp* phytases from other environmental studies [15] and affiliated with annotated genomic sequences (3-Phytase) of Bacteroidetes (Flavobacteriales), Beta, and Gammaproteobacteria (Fig. 2). Phytase (*bpp*) predicted amino acid sequences corresponded to protein domain 3- Phytase BPP (Pfam PF02333, InterPro IPR003431).

Alchichica's *phoX* partial sequences (~650 bp) showed phylogenetic affiliation with Alphaproteobacteria and Actinobacteria from aquatic environmental samples of different trophic states [10, 19]. Amino acid predicted sequences corresponded to protein families COG3211: PhoX, Pfam 05787 (residues E273 and E873, Protein Data Bank 3ZWU_A), and to the Domain of Unknown Function (DUF) DUF839. This domain has been directly associated to a large family of proteins that are able to metabolize phosphorus (Interpro Database) [59], while in the Pfam Database [60], DUF839 is considered a featured domain of the proteins which are included as part of the family with the PhoX domain [61].

Lake Alchichica partial *phoD* sequences (~350 bp) affiliated with Alpha, Beta, and Gammaproteobacteria from soil studies [39, 40]. Most (96 %) of the *phoD* amino acid predicted sequences contained the calcium binding site (residues N215, N216; 2YEQ; of the Protein Data Bank).

space

spaceFig. 2 Partial *bpp* sequences maximum likelihood topology. Alchichica Lake sequences (spongy and columnar microbialites and bacterioplankton); environmental studies *bpp* sequences are shown in black branches and genomic reference sequences are shown in cyan

spacecolor. Branches show bootstrap testing over 1000 replicates (only bootstrap values >50 are shown). Colored dots over clades show the affiliation of BPP transcript sequences (blue for transcripts of 24 h and green for 18 h)

space

space

space

space

space

Stratification.

spaceStratification. microbialites (7 phylotypes)

0

2

phosphatase *phoX*
Venn Diagram at distance 0.05

BPP phytase
Venn Diagram at distance 0.05

phosphatase *phoD*
Venn Diagram at distance 0.25

Fig. 3 Venn diagrams showing the distribution of phylotypes of alkaline phosphatases *phoX* and *phoD*, and beta-propeller phytase *bpp* in microbialites and bacterioplankton of Lake Alchichica. The cutoff value

for phylotype determination is shown, as well as the total number of phylotypes (in parenthesis) for each marker

space

Phylotype Distribution Between Microbial Assemblages (Microbialites and Bacterioplankton)

Alkaline phosphatases (*phoX*) exhibited the largest number of phylotypes (27), followed by *phoD* (23) and *bpp* (18).

Diversity indexes indicate that overall, the three studied markers were more diverse in microbialites than in bacterioplankton (Table 1). Analysis of alkaline phosphatases *phoX* and *phoD* (cutoffs = 0.05 and 0.25, see Table 1) showed that some phylotypes are shared between microbialites and bacterioplankton and between lake circulation periods (stratification and circulation), while other phylotypes are unique (Fig. 3). The proportion of unique sequences was higher for *bpp* (72.4 %) followed by *phoX* (62.5 %) and *phoD* (52.2 %). Microbialites shared more *phoX* phylotypes within seasons than bacterioplankton. The proportion of *phoD* phylotypes shared between periods was the same in microbialites and bacterioplankton although microbialites exhibited a larger number of shared *phoD* phylotypes between periods (Table 3). Unlike alkaline

phosphatases (that were present in microbialites and bacterioplankton during lake's circulation and stratification), *bpp* phytases were not found in the bacterioplankton during lake's circulation. Microbialites, on the other hand, contained four times more *bpp* phylotypes in circulation than during stratification. No phylotypes were common to all the *bpp* groups (Fig. 3).

The groups of sequences of *phoX*, *phoD*, and *bpp* showed a distinctive distribution pattern. The type of assemblage (microbialite or bacterioplankton) was more relevant than seasonality in the distribution of *phoX*. In contrast, seasonality distinguished *phoD* and *bpp* phylotype distribution. Bacterioplankton in circulation for *phoD* and microbialites in circulation for *bpp* were the groups that showed the highest number of unique phylotypes (Fig. 3).

A differential distribution of markers was explored by calculating the ratio of unshared phylotypes: shared phylotypes among assemblages (Table 4). Ratios indicate that exclusivity of DOP utilization phylotypes is favored in microbialites compared to bacterioplankton. A differential pattern was observed where *phoD* showed the lowest values for both assemblages, and *phoX* the highest

Transcription Patterns of *phoX*, *phoD*, and *bpp* Following a Diel Cycle

Alkaline phosphatases and phytases showed different expression patterns (Fig. 4). *PhoX* transcripts exhibited a diurnal expression pattern, 40 % of the transcripts was recovered at sunrise (06:00 h) and 60 % at noon (1200 hours). *PhoD* transcripts showed a broad pattern of expression throughout the diel cycle, where 42 % of the transcripts corresponded to 0600 hours and 50 % to 1200 hours, although 8 % was recovered at midnight (none for 1800 hours). *BPP* phytase

transcripts showed an afternoon pattern of expression where most of the transcripts were recovered at midnight (70 %) and 30 % at 1800 hours (Table 5, Online Resource 1).

Alkaline phosphatase *PhoD* transcripts were very similar in length (approximately 369 bp) to the ones recovered from total DNA. The closest affiliation of *PhoD* transcripts corresponded to alkaline phosphatase D (*Bacillus subtilis* type), two of the *PhoD* transcripts affiliated to precursor sequences of alkaline phosphatase D of *Pseudomonas stutzeri* (Gammaproteobacteria) and *Bhadrirhizobium* sp (Alphaproteobacteria); both are bacterial genus known to be involved in denitrification and nitrogen fixation, respectively (Table 5, Online Resource 1). *PhoX* and *BPP* phytase transcripts were smaller than the genomic sequences recovered from total DNA (transcript sequences can be consulted in Online Resource 1, also an ad hoc scrutiny of each of these sequences to verify their structural features and the presence of protein domains).

PhoX transcript sequences affiliated to three main functional categories: signaling, DNA binding, and phosphorus metabolism (see Online Resource 1); transcripts also associated to the *PhoX* domain, linker domains (like

the SH3 domains) [62–64], and to a domain that is still not functionally characterized, DUF839. By performing the hydrophobicity plot of our sequences [48], we found that some of the PhoX transcript sequences could, theoretically, form alpha-helices and loops, while the rest of them may be forming special arrangements of beta-sheets (see Online Resource 1). The PhoX domain is approximately 120 amino acids long and folds into a very particular three stranded beta-sheet followed by three helices and a proline-rich region that it is located before a membrane-interaction loop and spans approximately eight hydrophobic and polar residues [64, 65].

Transcript of BPP sequences showed affiliation (with significant scores) to sequences identified in the NR database of NCBI [45]. Transcript sequences of BPP showed domains involved in phosphorous metabolism, some specifically belonging to phosphoenolpyruvate carboxykinase (PEPCK), a protein that catalyzes the reversible decarboxylation and phosphorylation of oxaloacetate to yield phosphoenolpyruvate and carbon dioxide using ATP or GTP for the phosphoryl transfer [66]. Notably, in one of Alchichica's sequences (bppSsMcDNAOh_17), the Domain of Unknown Function DUF4397 (<http://pfam.xfam.org/family/PF05787>) was detected; DUF4397 has not been yet associated to any specific role [60].

P Mass Balance

Lake Alchichica's P mass balance is summarized in Fig. 5. Atmospheric deposition was the highest (annual basis average was 4.94 ton P/year) among P inputs to Alchichica Lake

space

Table 3 Alkaline phosphatases *phoX* and *phoD* and beta- propeller phytase *bpp* phylotype distribution among microbial assemblages

space

Marker Total phylotypes

space

Phylotypes shared between assemblages

space

Phylotypes only in bacterioplankton

Total Present in both seasons

space

Phylotypes only in microbialites

Total Present in both seasons

space

<i>phoX</i>	36	2
<i>phoD</i>	30	7
<i>bpp</i>	18	2

space

(Table 6). The sum of all inputs (precipitation, basin runoff, groundwater, and atmospheric deposition) reached 5.26 ton P/year, which was almost 10 % higher than P sedimentation (4.79 ton P/year), the main output identified. Since lake TP mass (calculated integrating TP concentrations over the water column) was 4.33 ton, we estimated the P residence time in the water column in the range of 0.82–0.90 y (i.e., 9.8–

10.8 months). P mass balance at Alchichica is further discussed in relation to the lake's physicochemical conditions during stratification and circulation periods, in the [BLake Alchichica's P Status^](#) section.

Discussion

Potential for DOP Remineralization

The potential for DOP remineralization found in Lake Alchichica's microbialites and bacterioplankton concur with those obtained by Sebastián and Gasol [67], who identified (using microautoradiography combined with in situ hybridization) a stronger response to P stimulation/starvation of Gammaproteobacteria and Bacteroidetes over other microbial groups (in bacterioplankton samples of an environment where P starvation was presumably prevalent). Jorquera et al. [68] also described the relevant abundance and diversity of *bpp* phytases in Gammaproteobacteria and in some Bacteroidetes genus such as *Flavobacterium*. Although Bacteroidetes (Flavobacteriales) has not been recognized as an abundant

space

phylum in Alchichica's microbialites [32], this group may be a key participant in DOP transformation.

PhoD and BPP transcript sequences were associated with DNA sequences from Alchichica assemblages that showed their closest affiliation with Proteobacteria and Actinobacteria (PhoD) and Gammaproteobacteria and Bacteroidetes (BPP). PhoD transcripts of 0600 hours were significantly affiliated to sequences of Gammaproteobacteria (mRNA 6h_14 and mRNA 6h_7 of Table 5; Online Resource 1; E values = e-142 and e-175).

Phylotype Dynamics and Turnover

The type of microbial assemblage defined the abundance of DOP utilization genetic markers including *phoX*, *phoD*, and *bpp* when metabolic potential for DOP utilization of bacterioplankton (of the whole water column) was compared with that of microbialites. Although a functional

parallelism was shown, results indicate that the potential of bacteria (as defined by OTU numbers) to transform DOP through AP *phoX*, *phoD*, and *bpp* phytases was higher in microbialites than in bacterioplankton (Fig. 3, Table 1). In general, the bacterioplankton of the stratification period shared more phylotypes with microbialites than the bacterioplankton of the circulation period.

Although we studied only three molecular markers associated to DOP utilization, here, we explore some ideas about the meaning of these phylotype dynamics-patterns. While microbialites showed a higher potential (higher genetic diversity) than

space

spaceTable 4 Ratios of *phoX*, *phoD*, and *bpp* unshared phylotypes of

space
Microbialites
each microbial assemblage:
Si shared phylotypes among
assemblages

space

Transcripts (PhoD)

Transcripts (PhoX)

Transcripts (BPP)

spacecirculation to stratification. The divergence of the groups *phoD*-Bacterioplankton and *bpp*-microbialites (of circulation) stands out overall, suggesting a faster dynamics of these markers, compared to *phoX*. In contrast, *phoX* dynamics did not show a marked difference among assemblages or periods. APs and phytase exhibited phylotype overlap (between assemblages and periods), as seen in the Venn diagrams (Fig. 3). These relationships suggest not only a structural connection between microbialites

and bacterioplankton assemblages but also between the littoral and the pelagic zone of the system. Unchanged (unshared) phylotypes constitute therefore the genetic reservoir (resilient stock) for DOP transformation (Table 4). Bacterioplankton seems to harbor overall a minor genetic potential than microbialites (for DOP utilization); how-

spaceFig. 4 Expression patterns of DOP utilization genes in a diel sampling scheme

bacterioplankton to remineralize DOP through APs, the difference was markedly higher in microbialites to remineralize phytate through BPP phytase. Microbial assemblages harboring these markers showed change (phylotype replacement or seasonality) in a timespan as short as the seasonal cycle of spaceever, bacterioplankton showed a more dynamic phylotype turnover (Table 4). This result may be coupled to their contrasting lifestyles: free-living vs. benthic consortia.

PhoD and BPP transcripts were obtained only during stratification, suggesting a more intense P starvation condition for the bacterial communities in this period (Table 5).

Table 5 Online Resource 1. Characteristics of the PhoD, PhoX, and BPP transcripts recovered and their closest affiliation in the general databases

Transcript	Expression pattern	Season	Length (bp)
PhoX			
alpsSsMcDNA6h_2	0600	S	369
alpsSsMcDNA6h_003	0600	S	369
C_alpsSsMcDNA6h_7	0600	S	369
alpsSsMcDNA6h_10	0600	S	369
alpsSsMcDNA6h_14	0600	S	369
alpsSsMcDNA6h_16	0600	S	369
C_alpsSsMcDNA6h_19	0600	S	363
alpsSsMcDNA12h_5	1200	S	369
alpsSsMcDNA12h_6	1200	S	369
alpsSsMcDNA12h_007	1200	S	369
alpsSsMcDNA12h_11	1200	S	369
alpsSsMcDNA12h_12	2400	S	357
PhoX			
phoXcDNA.S6h.29	0600	S	24
phoXcDNA.S6h.32	0600	S	44
phoXcDNA.S12h.20	1200	S	24
phoXcDNA.S12h.22	1200	S	58
phoXcDNA.S12h.23	1200	S	25
NCBIBPP (3-phytase)			
bppSsMcDNA18h_1	1800	S	93
bppSsMcDNA18h_3	1800	S	93
bppSsMcDNA18h_4	1800	S	93
bppSsMcDNAOh_2	1800	S	114
bppSsMcDNAOh_3	2400	S	118
bppSsMcDNAOh_6	2400	S	63
bppSsMcDNAOh_15	2400	S	604
bppSsMcDNAOh_16	2400	S	57
bppSsMcDNAOh_17	2400	S	114
bppSsMcDNAOh_18	2400	S	54

^a Sequence fragment affiliation corresponds to the results in the Protein Data Bank (PDB) and in the nonredundant database of the National Center for Biotechnology

Information (NCBI). Predicted sequences of amino acids were used as templates

as well as beta-propeller phytase *bpp* to release inorganic phosphorus from dissolved inorganic phosphorus (DOP)

space

space

Precipitation (0.03 ton P year⁻¹)

space

Atmospheric deposition (4.94 ton P year⁻¹)

showed mostly a daytime expression pattern compared to the overall evening expression of BPP. These patterns should correspond to peaks of activity of the community harboring such a metabolic potential. Although more effort should be done to understand the whole functional community, an indication of the activity of Gammaproteobacteria and Bacteroidetes in DOP utilization has been previously described in a low phosphorus environment [67]. The identification of Lake Alchichica BPP transcripts affiliated to diazotroph genus *Scytonema hofmanni* also suggests that Cyanobacteria may be an important group in phytate degradation, as reviewed by Jorquera et al. [68]. The type and frequency to which Alchichica's sequences were associated to best-hit proteins into the functional categories of

space

space

littoral-pelagic exchange?

microbialites

space

Lake TP mass (4.33 ton)
Estimated TPP (3.81 ton)
P monoesters (0.48 1.68 ton)

Table 6 P mass balance in Lake Alchichica, inputs (positive sign fluxes) and outputs (negative sign)

Fluxes of P	(P ton/year)	% of P mass	Reference
Fluxes of P	(P ton/year)	% of P mass	Reference
Unaccounted sources/sinks and/or error (0.24 ton P year ⁻¹)	0.24	5.5	[25, 52]
Groundwater input (0.24 ton P year ⁻¹)	0.24	5.5	[25, 52]
P sedimentation (4.79 ton P year ⁻¹)	-4.79	-111	[27]
Water-column P residence time (0.82 - 0.90 year)			
Lake total P mass (tons)	4.33	100	This study
Precipitation	0.03	0.8	[52]
Basin runoff	0.05	1.1	[52]
Atmospheric deposition	4.94	114	[52, 58]
Groundwater	0.24	5.5	[25, 52]
Sedimentation	-4.79	-111	[27]
System P balance	0.47		This study

space

general databases may reflect that affiliations are not completely at random, and these results are not false

positives but structurally significant sequences (see Online Resource 1). DUF839 has been previously associated to *phoX* alkaline phosphatase [10, 19]; DUFs are frequently uncharacterized protein families found in the protein databases, and they seem to play important roles in the clarification of DUFs' challenge.

of the lake, including water column P availability and dissolved oxygen and nutrients, vary importantly between circulation and stratification (Fig. 1). Although the littoral zone exhibits low availability of N and P (low P_{org} and DIN/SRP ratio <16), the littoral SRP concentration during stratification period is almost half of the SRP concentration

space

space

space

space

space

Fig. 5 Lake Alchichica phosphorus condition. The diagram shows the main phosphorus in- and outflows (from atmospheric deposition, basin runoff, groundwater, and sedimentation), lake total phosphorus (TP) mass, total particulate (TPP), P monoester fraction, and residence time

space

and in an annual basis. Boxes enclose enzymatic mechanisms of alkaline phosphatases *phoX* and *phoD*,

space

Lake Alchichica's P balance shows that allochthonous P total input was 9.8 % higher than the TP exported to the sediments (Fig. 5).

This difference (or unaccounted P) could correspond to presently unknown sources or sinks of P, as well as to errors in the present estimates, or due to intra-annual heterogeneity. In any case, this P mass balance revealed that the magnitude of P fluxes is similar (between 8 and 10 % higher) to the lake's total P mass. The average time that each P atom spends in the water column of Lake Alchichica before it sediments (i.e., the P water column-residence time) ranges between 0.82 and 0.90 year, as estimated by the P mass/P flux ratio calculated using either total inputs or total outputs.

Temporal heterogeneity is a factor that should be considered in this exploration, although at the moment, a dynamic water balance in Lake Alchichica is constrained due to the uncertainty on the lake's groundwater flows and particularly on their variability. During winter, dry and dusty conditions of the Mexican highlands favor higher atmospheric P deposition rates than during the rest of the year (see seasonal atmospheric dust deposition in Oseguera et al. [58]). On an annual basis, the circulation bloom of big-size diatoms [27] is assumed to be the main driver of organic matter sedimentation.

Alchichica Lake's TPP accounted for 3.67 and 3.94 tons of organic P in the water column during circulation and stratification, respectively. According to novel ^{31}P NMR spectroscopy P fractioning studies, P monoesters would account for 11 to 44 % of TPP (in a productive system) [70]; using this range (in the absence of any similar measurements for Alchichica), the P monoesters fraction of the lake would range from 0.42 to

1.68 tons, a considerable size for a pool that would have to be remineralized by microorganisms [71] fitted with P extracellular enzymes (i.e., *phoD*, *phoX*, *phoA*, etc). Nevertheless, understanding the metabolic strategies to utilize this P fraction likely requires multiple approaches, from microscopic to community-scale research strategies. It is also important to assess the exchange of P, organic matter, and other nutrients from the littoral zone (where there are areas with high coverage of macrophytes, epibenthic diatoms, and living microbialites) to the pelagic zone, as outlined by Ardiles et al. [27], who also reported benthic littoral phytoplankton species (pennate diatoms) in the central and deep areas of the lake. Finally, it is also remarkable that, likely because of the high P output flow associated to sedimentation, there is no experimental evidence of net P release from the sediments.

Final Remarks

Microbialite and bacterioplankton assemblages seem to exhibit different strategies for DOP utilization (i.e., through alkaline phosphatases and phytases). A Bstock strategy^ favored in microbialites, focused on preserving a

diverse community (a higher potential to utilize DOP, e.g., maintaining

spaceproteobacterial participants) and a Bfast-change strategy^, overall observed in bacterioplankton, in which other (less abundant) phylotypes exhibit a higher turnover. Although microbialites harbor a higher genetic potential for DOP utilization (through *phoX*, *phoD*, and *bpp*), the resilience potential of each assemblage is still to be assessed and may be related to the time scale as well as to the magnitude of environmental changes. Under this scenario, we hypothesize that the water column communities may be more effective to face drastic environmental change, although their effective rate of DOP transformation may be lower (due to their lower phylotype diversity) than in their benthic counterpart.

Accounting for the area covered by microbialites in Lake Alchichica, and considering their extensive genetic potential and functional capabilities for DOP utilization, both assemblages (microbialites and bacterioplankton) may play an important role in contributing to the availability of nutrient and trace metals and in maintaining a relatively high production in a system where allochthonous nutrient input is relatively low. This Befficient^ remineralization would help explain observations that refer higher production features than expected for an oligotrophic system (with low P inputs). This evidence includes high abundance of total picoplankton relative to low trophic state systems [30] and regular *Nodularia* blooms [33, 55].

DOP utilization genes/transcripts showed different dynamics in contrasting hydrological conditions (within microbialites and bacterioplankton). Our findings indicate that microbial groups have different responses to face P stimulation/starvation. This provides dynamic capacities (system's metabolic potential) to utilize P (DOP). It is relevant to observe seasonality of microbial phosphorus utilization potential to understand transformations of other major biogeochemical cycles such as carbon and nitrogen.

Acknowledgments We thank Luis Oseguera and the Proyecto de Investigación en Limnología Tropical (FES Iztacala, UNAM) and Bernardo Valdespino for fieldwork support in Lake Alchichica. We acknowledge F. Sergio Castillo-Sandoval of the Aquatic Biogeochemistry Laboratory (ICMyL, UNAM) for the accomplishment of chemical analyses. Additionally, we thank Osiris Gaona Pineda and Antonio Cruz-Peralta for technical and laboratory assistance. Funding for this project was granted to L.I. Falcón through SEP-CONACyT No. 151796 and PAPIIT-UNAM No. IT100212-3. All sampling was done under collector permit No. PPF/DGOPA.033/2013 (LIF). P Valdespino acknowledges CONACyT for a doctoral fellowship and SECITI-CLAF for a postdoctoral fellowship.

References

1. Kolowith LC, Ingall ED, Benner R (2001) Composition and cycling of marine organic phosphorus. *Limnol Oceanogr* 46:309–320
2. Björkman KM, Karl DM (2003) Bioavailability of dissolved organic phosphorus in the euphotic zone at Station ALOHA, North Pacific Subtropical Gyre. *Limnol Oceanogr* 48(3):1049–1057
3. ^{space}Dyhrman ST, Ammerman JW, Van Mooy BAS (2007) Microbes and the marine phosphorus cycle. *Oceanography* 20:110–116
4. White AE, Watkins-Brandt KS, Engle MA, Burkhardt B, Paytan A (2012) Characterization of the rate and temperature sensitivities of bacterial remineralization of dissolved organic phosphorus compounds by natural populations. *Front Microbiol* 3:276
5. Biddanda B, Opsahl S, Benner R (1994) Plankton respiration and carbon flux through bacterioplankton on the Louisiana shelf. *Limnol Oceanogr* 39:1259–1275
6. del Giorgio PA, Cole JJ, Cimbleris A (1997) Respiration rates in bacteria exceed phytoplankton production in unproductive aquatic systems. *Nature* 385:148–151
7. Benitez-Nelson CR, Buesseler KO (1999) Variability of inorganic and organic phosphorus turnover rates in the coastal ocean. *Nature* 6727:502–505
8. Cotner JB, Wetzel RG (1992) Uptake of dissolved inorganic and organic phosphorus compounds by phytoplankton and bacterioplankton. *Limnol Oceanogr* 37(2):232–43
9. Štrojsová A, Vrba J, Nedoma J, Komárková J, Znachor P (2003) Seasonal study of extracellular phosphatase expression in the phytoplankton of a eutrophic reservoir. *Eur J Phycol* 38(4):295–306
10. Sebastian M, Ammerman JW (2009) The alkaline phosphatase PhoX is more widely distributed in marine bacteria than the classical PhoA. *ISME J* 3:563–572
11. Luo H, Benner R, Long RA, Hu J (2009) Subcellular localization of marine bacterial alkaline phosphatases. *Proc Natl Acad Sci U S A* 106:21219–21223
12. Oh BC, Chang BS, Park KH, Ha NC, Kim HK, Oh BH et al (2001) Calcium-dependent catalytic activity of a novel phytase from *Bacillus amyloliquefaciens* DS11. *Biochemistry* 40(32):9669–9676
13. Suzumura M, Kamatani A (1995) Origin and distribution of inositol hexaphosphate in estuarine and coastal sediments. *Limnol Oceanogr* 40:1254–1261
14. Lim BL, Yeung P, Cheng C, Hill JE (2007) Distribution and diversity of phytate-mineralizing bacteria. *ISME J* 1(4):321–330
15. Huang H, Shi P, Wang Y, Luo H, Shao N, Wang G et al (2009) Diversity of beta-propeller phytase genes in the intestinal contents of grass carp provides insight into the release of major phosphorus from phytate in nature. *Appl Environ Microbiol* 75(6):1508–1516
16. Cheng C, Lim BL (2006) Beta-propeller phytases in the aquatic environment. *Arch Microbiol* 185(1):1–13
17. Jacobsen T, Slotfeldt-Ellingsen D (1983) Phytic acid and metal availability: a study of Ca and Cu binding. *Cereal Chem* 60(5):392–395
18. White AE (2009) New insights into bacterial acquisition of phosphorus in the surface ocean. *PNAS* 106(50):2013–2014
19. Valdespino-Castillo PM, Alcántara-Hernández RJ, Alcocer J, Merino-Ibarra M, Macek M, Falcón LI (2014) Alkaline phosphatases in microbialites and bacterioplankton from Alchichica soda lake, Mexico. *FEMS Microbiol Ecol* 90(2):504–519
20. Giovannoni SJ, Vergin KL (2012) Seasonality in ocean microbial communities. *Science* 335:671–676
21. Martínez J, Smith DC, Steward GF, Azam F (1996) Variability in ectohydrolytic enzyme activities of pelagic marine bacteria and its significance for substrate processing in the sea. *Aquat Microb Ecol* 10:223–230. doi:10.3354/ame010223
22. Arnosti C, Durkin S, Jeffrey WH (2005) Patterns of extracellular enzyme activities among pelagic marine microbial communities: implications for cycling of dissolved organic carbon. *Aquat Microb Ecol* 38(2):135–145
23. Vilaclara G, Chávez M, Lugo A, González H, Gaytán M (1993) Comparative description basic chemistry of crater-lakes in Puebla State, México. *Verh Int Ver Theoret Angew Limnol* 25:435–440
24. Armienta MA, Vilaclara G, De la Cruz-Reyna S, Ramos S, Ceniceros N, Cruz O, Aguayo A, Arcega-Cabrera F (2008) Water ^{space}chemistry of lakes related to active and inactive Mexican volcanoes. *J Volcanol Geoth Res* 178(2):249–258
25. Kazmierczak J, Kempe S, Kremer B, Lopez-García P, Moreira D, Tavera R (2010) Hydrochemistry and microbialites of the alkaline crater lake Alchichica, Mexico. *Facies* 57:543–570
26. Ramírez-Olvera MA, Alcocer J, Merino-Ibarra M, Lugo A (2009) Nutrient limitation in a tropical saline lake: a microcosm experiment. *Hydrobiologia* 626:5–13
27. Ardiles V, Alcocer J, Vilaclara G, Oseguera LA, Velasco L (2012) Diatom fluxes in a tropical, oligotrophic lake dominated by large-sized phytoplankton. *Hydrobiologia* 679(1):77–90
28. Lugo A, Alcocer J, Sanchez M, Escobar E (1998) Littoral protozoan assemblages from two Mexican hyposaline lakes. *Hydrobiologia* 381:9–13
29. Alcocer J, Lugo A, Escobar E, Sánchez MR, Vilaclara G (2000) Water column stratification and its implications in the tropical warm monomictic lake Alchichica, Puebla, Mexico. *Verh Int Ver Theoret Angew Limnol* 27:3168–3169
30. Macek M, Alcocer J, Lugo-Vázquez A, Martínez-Pérez ME, Peralta Soriano L, Vilaclara Fatjó G (2009) Long term picoplankton dynamics in a warm-monomictic, tropical high altitude lake. *J Limnol* 68(2):183–192
31. Hernández-Avilés JS, Macek M, Alcocer J, Lopez-Trejo B, Merino-Ibarra M (2010) Prokaryotic picoplankton dynamics in a warm-monomictic saline lake: temporal and spatial variation in structure and composition. *J Plankton Res* 32:1301–1314
32. Centeno CM, Legendre P, Beltrán Y, Alcántara-Hernández RJ, Lidström UE, Ashby MN, Falcón LI (2012) Microbialite genetic diversity and composition relate to environmental variables. *FEMS Microbiol Ecol* 82(3):724–735
33. Falcón LI, Escobar-Briones E, Romero D (2002) Nitrogen fixation patterns displayed by cyanobacterial consortia in Alchichica crater-lake, Mexico. *Hydrobiologia* 467(1–3):71–78
34. Beltrán Y, Centeno CM, García-Oliva F, Legendre P, Falcón LI (2012) N₂ fixation rates and associated diversity (*nifH*) of microbialite and mat-forming consortia from different aquatic environments in Mexico. *Aquat Microb Ecol* 67(1):15–24
35. Grasshoff K, Kremling K, Ehrhardt M (1983) Methods of seawater analysis. Verlag Chemie, Weinheim
36. Kirkwood DS (1994) Sanplus segmented flow analyzer and its applications. Seawater analysis. Skalar, Amsterdam
37. Valderrama JC (1981) The simultaneous analysis of total nitrogen and total phosphorus in natural waters. *Mar Chem* 10:109–122
38. Zhou J, Bruns MA, Tiedje JM (1996) DNA recovery from soils of diverse composition. *Appl Environ Microbiol* 62:316–322
39. Sakurai M, Wasaki J, Tomizawa Y, Shinano T, Osaki M (2008) Analysis of bacterial communities on alkaline phosphatase genes in soil supplied with organic matter. *Soil Sci Plant Nutr* 54:62–71
40. Tan H, Barret M, Mooij MJ, Rice O, Morrissey JP, Dobson A et al (2013) Long-term phosphorus fertilisation increased the diversity of the total bacterial community and the *phoD* phosphorus mineraliser group in pasture soils. *Biol Fert Soils* 49(6):661–672

41. Iwar S, Chai B, Jesus EDC, Penton CR, Lee TK, Cole JR, Tiedje JM (2011) Gene-targeted metagenomics (GT Metagenomics) to explore the extensive diversity of genes of interest in microbial communities. In: De Bruijn FJ (ed) Handbook of molecular microbial ecology I: metagenomics and complementary approaches. Wiley, Hoboken, pp 235–243
42. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59(3):307–321
43. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB et al (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and space
- spaceComparing microbial communities. *Appl Environ Microbiol* 75(23):7537–7541
44. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
45. Geer LY, Marchler-Bauer A, Geer RC, Han L, He J, He S, Liu C, Shi W, Bryant SH (2010) The NCBI BioSystems database. *Nucleic Acids Res* 38(Database issue):D492–D496
46. Finn RD, Clements J, Eddy SR (2011) HMMER web server: inter-active sequence similarity searching. *Nucleic Acids Res* 39:W29–W37
47. Chou PY, Fasman GD (1974) Conformational parameters for amino acids in helical, β -sheet, and random coil regions calculated from proteins. *Biochemistry* 13(2):211–222
48. Kyte J, Doolittle RF (1982) A simple method for displaying the hydropathic character of a protein. *J Mol Biol* 157(1):105–132
49. Dworkin JE, Rose GD (1987) Hydrophobicity profiles revisited. In: Walsh KA (ed) Methods in protein sequence analysis. Humana, Clinton, pp 573–586
50. Baker D, Sali A (2001) Protein structure prediction and structural genomics. *Science* 294(5540):93–96
51. Krystek SR Jr, Metzler WJ, Novotny J (2001) Hydrophobicity profiles for protein sequence analysis. *Curr Protoc Protein Sci* 00:2.2: 2.2.1–2.2.13
52. García Martínez J (2010) Efectos climáticos sobre el agua subterránea y el lago Alchichica Puebla, México. Dissertation, Universidad Nacional Autónoma de México
53. Custodio E, Llamas MR (1983) Hidrología Subterránea Vol. II. Editorial Omega, México
54. Oliva MG, Lugo A, Alcocer J, Peralta L, Sánchez R (2001) Phytoplankton dynamics in a deep, tropical, hyposaline lake. *Hydrobiologia* 466:299–306
55. Oliva MG, Lugo A, Alcocer J, Peralta L, Oseguera LA (2009) Planktonic bloomforming *Nodularia* in the saline Lake Alchichica, Mexico. *Nat Resour Env Iss* 15(1):22, <http://digitalcommons.usu.edu/nrei/vol15/iss1/22>
56. Alcocer J, López-Anaya DP, Oseguera LA (2007) Dinámica del carbono orgánico particulado en un lago tropical profundo. In: Hernández de la Torre B, Gaxiola Castro G (ed) Carbono en ecosistemas acuáticos de México. INE, México, pp 239–247
57. Ramírez-Zierold JA, Merino-Ibarra M, Monroy-Ríos E, Olson M, Castillo FS, Gallegos ME, Vilaclara G (2010) Changing water, phosphorus and nitrogen budgets for Valle de Bravo reservoir, water supply for Mexico City Metropolitan Area. *Lake Reserv Manag* 26(1):23–34
58. Oseguera LA, Alcocer J, Vilaclara G (2011) Relative importance of dust inputs and aquatic biological production as sources of lake sediments in an oligotrophic lake in a semi-arid area. *Earth Surf Processes* 36(3):419–426
59. spaceMitchell A, Chang HY, Daugherty L, Fraser M, Hunter S, Lopez R et al (2015) The InterPro protein families database: the classification resource after 15 years. *Nucleic Acids Res* 43(Database issue): D213–D221
60. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A, Hetherington K, Holm L, Mistry J, Sonnhammer ELL, Tate J, Punta M (2014) Pfam: the protein families database. *Nucleic Acids Res* 42:D222–D230
61. Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY et al (2014) CDD: NCBI's conserved domain database. *Nucleic Acids Res* 43(Database issue):D222–D226
62. Ponting CP (1996) Novel domains in NADPH oxidase subunits, sorting nexins, and PtdIns 3-kinases: binding partners of SH3 domains? *Protein Sci* 5(11):2353–2357
63. Hiroaki H, Ago T, Ito T, Sumimoto H, Kohda D (2001) Solution structure of the PX domain, a target of the SH3 domain. *Nat Struct Biol* 8(6):526–530. doi:[10.1038/88591](https://doi.org/10.1038/88591). PMID11373621
64. Ago T, Kuribayashi F, Hiroaki H, Takeya R, Ito T, Kohda D, Sumimoto H (2003) Phosphorylation of p47phox directs phox homology domain from SH3 domain toward phosphoinositides, leading to phagocyte NADPH oxidase activation. *Proc Natl Acad Sci U S A* 100(8):4474–4479
65. Wishart MJ, Taylor GS, Dixon JE (2001) Phox lipids: revealing PX domains as phosphoinositide binding modules. *Cell* 105(7): 817–820. doi:[10.1016/S0092-8674\(01\)00414-7](https://doi.org/10.1016/S0092-8674(01)00414-7). PMID11439176
66. Márquez JA, Hasenbein S, Koch B, Fieulaine S, Nessler S, Russell RB et al (2002) Structure of the full-length HPr kinase/phosphatase from *Staphylococcus xylosus* at 1.95 Å resolution: mimicking the product/substrate of the phosphotransfer reactions. *Proc Natl Acad Sci U S A* 99(6):3458–3463
67. Sebastián M, Gasol JM (2013) Heterogeneity in the nutrient limitation of different bacterioplankton groups in the Eastern Mediterranean Sea. *ISME J* 7(8):1665–1668
68. Jorquera M, Martínez O, Maruyama F, Marschner P, de la Luz Mora M (2008) Current and future biotechnological applications of bacterial phytases and phytase-producing bacteria. *Microbes Environ* 23(3):182–191
69. Goodacre NF, Gerloff DL, Uetz P (2013) Protein domains of unknown function are essential in bacteria. *mBio* 5(1):e00744–13
70. Read EK, Ivancic M, Hanson P, Cade-Menun BJ, McMahon KD (2014) Phosphorus speciation in a eutrophic lake by ³¹P NMR spectroscopy. *Water Res* 62:229–240
71. Cotner J, Wetzel RG (1992) Uptake of dissolved inorganic and organic phosphorus compounds by phytoplankton and bacterioplankton. *Limnol Oceanogr* 37(2):232–243

ization genes *phoD*, *phoX*, and *bpp*; richness and diversity estimators; as well as coverage for each group (microbialites and bacterioplankton) and period

#	<i>phoD</i> (sequences obtained = 180)				<i>phoX</i> (sequences obtained = 192)				<i>bpp</i> (sequences obtained = 58)			
	OTUs	Chao	Shannon (<i>H'</i>)	Coverage (%)	OTUs	Chao	Shannon (<i>H'</i>)	Coverage (%)	OTUs	Chao	Shannon (<i>H'</i>)	Coverage (%)
23	58	2.51	2.51	80	16	30	2.28	88	15	33	2.3527	74
15	20	2.22	2.22	92	14	17.3	2.21	92	10	20	1.95	86
17	27.5	2.13	2.13	91	7	7	1.67	100	3	4	0.7963	71
13	20.5	1.68	1.68	92	6	6	1.46	100	3	4	0.8	71
5	8	1.56	1.56	33	7	7.25	1.15	97	-	-	-	-
5	8	1.56	1.56	33	6	6	1.12	98	-	-	-	-
9	14	1.56	1.56	83	6	16	1.04	72	3	3	0.7356	100
7	7.33	1.44	1.44	93	6	16	1.04	72	3	3	0.74	100

a. Lake circulation

b. Lake stratification

c. Littoral zone

	T (°C)	SRP (μM)	P _{org} (μM)
Circulation	16.1	0.68	2.5
Stratification	19.8	0.38	2.5

Fig. 3 Venn diagrams showing the distribution of phyl phosphatases *phoX* and *phoD*, and beta-propeller microbialites and bacterioplankton of Lake Alchichica

Table 4 Ratios of *phoX*, *phoD*, and *bpp* unshared phylotypes of each microbial assemblage: shared phylotypes among assemblages

	Ba
	Un
	phy
<i>phoX</i>	4.5
<i>phoD</i>	0.3
<i>bpp</i>	0.5

Table 3 Alkaline phosphatases *phoX* and *phoD* and beta-propeller phytase *bpp* phylotype distribution among microbial assemblages

Marker	Total phylo
<i>phoX</i>	36
<i>phoD</i>	30
<i>bpp</i>	18

Fig. 5 Lake Alchichica phosphorus condition. The diagram shows main phosphorus in- and outflows (from atmospheric deposition, runoff, groundwater, and sedimentation), lake total phosphorus mass, total particulate (TPP), P monoester fraction, and residence

Table 6 P mass balance in Lake Alchichica, inputs (positive fluxes) and outputs (negative sign)

Fluxes of P	(P ton/year)	% of P mass	Re
Lake total P mass (tons)	4.33	100	Th
Precipitation	0.03	0.8	[5]
Basin runoff	0.05	1.1	[5]
Atmospheric deposition	4.94	114	[5]
Groundwater	0.24	5.5	[2]
Sedimentation	-4.79	-111	[2]
System P balance	0.47		Th