

UC Irvine

UC Irvine Previously Published Works

Title

Adenosine A1 and Prostaglandin E Receptor 3 Receptors Mediate Global Airway Contraction after Local Epithelial Injury

Permalink

<https://escholarship.org/uc/item/0c20f2rd>

Journal

American Journal of Respiratory Cell and Molecular Biology, 48(3)

ISSN

1044-1549

Authors

Zhou, Jian

Alvarez-Elizondo, Martha B

Botvinick, Elliot

et al.

Publication Date

2013-03-01

DOI

10.1165/rcmb.2012-0174oc

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Adenosine A₁ and Prostaglandin E Receptor 3 Receptors Mediate Global Airway Contraction after Local Epithelial Injury

Jian Zhou¹, Martha B. Alvarez-Elizondo^{1,2}, Elliot Botvinick^{1,2,4,5}, and Steven C. George^{1,3,5}

¹Department of Biomedical Engineering, ²The Beckman Laser Institute and Medical Clinic, ³Department of Chemical Engineering and Materials, ⁴Department of Surgery, and ⁵The Edwards Lifesciences Center for Advanced Cardiovascular Technology, University of California, Irvine, Irvine, California

Epithelial injury and airway hyperresponsiveness are prominent features of asthma. We have previously demonstrated that laser ablation of single epithelial cells immediately induces global airway constriction through Ca²⁺-dependent smooth muscle shortening. The response is mediated by soluble mediators released from wounded single epithelial cells; however, the soluble mediators and signaling mechanisms have not been identified. In this study, we investigated the nature of the epithelial-derived soluble mediators and the associated signaling pathways that lead to the L-type voltage-dependent Ca²⁺ channel (VGCC)-mediated Ca²⁺ influx. We found that inhibition of adenosine A₁ receptors (or removal of adenosine with adenosine deaminase), cyclooxygenase (COX)-2 or prostaglandin E receptor 3 (EP₃) receptors, epidermal growth factor receptor (EGFR), or platelet-derived growth factor receptor (PDGFR) all significantly blocked Ca²⁺ oscillations in smooth muscle cells and airway contraction induced by local epithelial injury. Using selective agonists to activate the receptors in the presence and absence of selective receptor antagonists, we found that adenosine activated the signaling pathway A₁R→EGFR/PDGFR→COX-2→EP₃→VGCCs→calcium-induced calcium release, leading to intracellular Ca²⁺ oscillations in airway smooth muscle cells and airway constriction.

Keywords: ATP; epidermal growth factor receptor; platelet-derived growth factor receptor; cyclooxygenase-2; L-type voltage-dependent Ca²⁺ channels

We have recently demonstrated that laser ablation of a single epithelial cell reproducibly induces rapid and global airway constriction (1). The dynamics of the response suggested that local epithelial injury released a soluble mediator(s) that was transported to underlying smooth muscle cells by diffusion. The soluble mediator(s) evoked multiple Ca²⁺ oscillations in smooth muscle cells by stimulating L-type voltage-dependent Ca²⁺ channels (VGCCs), thus increasing intracellular Ca²⁺ levels via the calcium-induced calcium release (CICR) mechanism. In this study, we investigated the specific nature of the soluble mediator (s) and signaling pathway(s) underlying the VGCC-mediated Ca²⁺ influx.

In response to mechanical stimulation, ATP is released from airway epithelial cells and stimulates Ca²⁺ waves in the epithelium (2–4); however, the effects of local epithelial injury-induced ATP on

CLINICAL RELEVANCE

Our study provides direct evidence that local epithelial injury could contribute to airway hyperresponsiveness in subjects with asthma, and provides potentially new pharmacological targets for asthma treatment.

airway caliber have not been studied. The level of intracellular ATP is high (millimolar range) for metabolism, but is extremely low in the extracellular space, where it can function as a signaling molecule (5). Extracellular ATP activates two subtypes of purinergic receptors, P2X and P2Y, and both of them are expressed on airway epithelial and smooth muscle cells (6). The P2X receptors are ligand-gated ion channels that mediate Ca²⁺ and Na⁺ influx, and P2Y receptors are G protein-coupled receptors that regulate phospholipase C pathway, leading to inositol trisphosphate (IP₃) production and intracellular Ca²⁺ release (6). Because ATP has been shown to stimulate small airway contraction in mouse lung tissue slices by activating P2Y receptors (7), it was a likely candidate to be the soluble mediator(s) involved in local epithelial injury-induced smooth muscle contraction. However, in our previous studies, we found that inhibition of P2 purinoceptor receptors did not block Ca²⁺ signaling in smooth muscle cells and airway contraction induced by local epithelial injury. Furthermore, inhibition of P2 purinoceptor receptors or neutralizing ATP with apyrase did not block extracellular ATP-induced airway contraction. Thus, we previously ruled out the involvement of P2 purinoceptor receptors on smooth muscle cells in local epithelial injury-induced airway contraction, but we could not completely rule out ATP as a soluble mediator (1).

Exogenous ATP is rapidly converted to its metabolic products, such as adenosine monophosphate (AMP), ADP, and adenosine by ecto-apyrase enzymes expressed on the cell surface (8). Thus, ATP released from wounded single epithelial cells could potentially stimulate adenosine receptors after degradation to AMP (9) or adenosine (6, 10). Adenosine receptors are G protein-coupled receptors and have four subtypes: A₁, A_{2A}, A_{2B}, and A₃ receptors. Although the four adenosine receptors are all expressed on airway smooth muscle cells (6), adenosine is traditionally considered to indirectly induce bronchoconstriction either through activating A_{2B} or A₃ receptors on mast cells or through neural nerves via A₁ receptors (11–14). However, recent studies demonstrated that adenosine could directly activate A₁ and A_{2B} receptors to regulate adenylyl cyclase on human tracheal smooth muscle cells (15), and stimulate A₁ receptors on human bronchial smooth muscle cells to increase Ca²⁺ signaling (16). Furthermore, adenosine A₁ receptors are colocalized with epidermal growth factor receptors (EGFRs) and induce the transactivation of EGFR and their downstream pathways, such as phosphoinositide 3-kinase and Src kinase in neural cells (17).

(Received in original form May 11, 2012 and in final form October 19, 2012)

This work was supported by grant R01 HL067954 from the National Heart, Lung, and Blood Institute (S.C.G.) and by grant P41RR01192 from the National Center for Research Resources.

Correspondence and requests for reprints should be addressed to Steven C. George, M.D., Ph.D., Department of Biomedical Engineering, 2420 Engineering Hall, University of California, Irvine, CA 92697-2715. E-mail: scgeorge@uci.edu

This article has an online supplement, which is accessible from this issue's table of contents at www.atsjournals.org

Am J Respir Cell Mol Biol Vol 48, Iss. 3, pp 299–305, Mar 2013

Copyright © 2013 by the American Thoracic Society

Originally Published in Press as DOI: 10.1165/rcmb.2012-01740C on December 6, 2012

Internet address: www.atsjournals.org

Prostanoids, which are derived from arachidonic acid, include prostaglandins (PGE₂, PGD₂, and PGF₂), prostacyclins (PGI₂), and thromboxane (18). Cyclooxygenase (COX-1 and COX-2) is an enzyme that converts arachidonic acid into the prostanoids. High levels of prostanoids in bronchoalveolar lavage fluid and the increased expression of COX-2 in lung tissue have been detected in subjects with asthma (18). PGE₂ is produced by airway epithelial cells (19–21) and by smooth muscle cells (22–25), and has been demonstrated to regulate airway caliber (26–28). There are four subtypes of PGE₂ receptors: EP_{1–4}. Activation of EP₂ or EP₄ receptors increases intracellular cyclic AMP level and causes relaxation of smooth muscles (29, 30), whereas activation of EP₁ receptors increases intracellular Ca²⁺ and thus causes smooth muscle contraction (23, 31, 32). EP₃ has multiple isoforms, and its activation can stimulate smooth muscle contraction through either decreasing cyclic AMP levels or increasing intracellular Ca²⁺ (30, 33).

The goal of this study was to identify the soluble mediator(s) and signaling pathway(s) mediating the local epithelial injury-induced smooth muscle contraction in rat lung tissue slices. We hypothesized that adenosine and PGE₂ were both involved in the underlying mechanisms of local epithelial injury-induced airway contraction. Our results show that ATP is released from wounded single epithelial cells, and activates the following sequence of events: A₁R→EGFR/PDGFR→COX-2→EP₃→VGCCs→CICR, leading to the Ca²⁺ oscillations in smooth muscle cells and airway contraction.

MATERIALS AND METHODS

Materials

Fluo-4/AM, Pluronic F-127, Hanks' balanced salt solution (HBSS), Dulbecco's modified Eagle medium, and Antibiotic-Antimycotic were purchased from Invitrogen (Carlsbad, CA). AH6809, AG18, AG1478, AG1296, and 11-deoxy-16,16-dimethyl PGE₂ (11-PGE₂) were purchased from Cayman Chemical (Ann Arbor, MI). Sulfobromophthalein, ATP, adenosine 5'-[γ-thio]triphosphate tetralithium salt (ATP-γ-S), indomethacin, L-798106, and N6-cyclopentyladenosine (CPA) were purchased from Sigma-Aldrich (St. Louis, MO). Adenosine, 9-chloro-2-(2-furanyl)-[1,2,4] triazololo[1,5-c]quinazolin-5-amine (CGS15943), 1-butyl-8-(hexahydro-2,5-methanopentalen-3a(1H)-yl)-3,7-dihydro-3-(3-hydroxypropyl)-1H-purine-2,6-dione (PSB36), 8-[4-[4-(4-chlorophenyl)piperazine-1-sulfonyl]phenyl]-1-propylxanthine (PSB603), SLV320, ZM241385, and MRS1334 were purchased from Tocris Bioscience (Ellisville, MO). Supplemented HBSS (sHBSS) was made from HBSS with Ca²⁺ and Mg²⁺ supplemented with 20 mM Hepes (pH 7.4) (34).

Because the selectivity of an inhibitor is dependent on species (e.g., human versus rat) and cell type, for each chemical compound that we used in this study, we have provided detailed information for the concentration used based on the previous studies in rat (*see* Table E1 in the online supplement). Because molecular methods, such as small interfering RNA or lentivirus-based short hairpin RNA, to confirm the molecular mechanism in our current studies, introduce significant technical challenges in the lung tissue slice model, we applied multiple selective inhibitors to confirm our results.

Preparation of Lung Tissue Slices

All procedures involving animals were approved by the Institutional Animal Care and Use Committee of the University of California, Irvine, and were consistent with guidelines published by the National Institutes of Health. The preparation of rat lung tissue slices has been previously described in detail (1), and the procedure is also available in the MATERIALS AND METHODS section of the online supplement.

Measurement of Intracellular Ca²⁺ Signaling

To monitor free intracellular Ca²⁺ in both epithelial and smooth muscle cells, lung tissue slices were incubated in sHBSS with 20 μM Fluo-4/AM,

100 μM sulfobromophthalein, and 0.2% Pluronic F-127 for 1 hour at room temperature (35). Subsequently, the slices were kept in sHBSS with 100 μM sulfobromophthalein for another hour at room temperature. The slices were then transferred to a glass-bottom dish (MatTek, Ashland, MA) and held in place with a slice anchor (Warner Instruments, Hamden, CT). Confocal imaging was performed on a Zeiss 510 Meta multiphoton laser scanning microscope (LSM 510; Zeiss, Jena, Germany). Fluo-4 was excited with a 488-nm laser, and the fluorescence images (512 × 512 pixels) were collected.

Laser Ablation

The procedure for femtosecond (fs) laser ablation has been previously described in detail (1). Briefly, the laser ablation was performed on the LSM 510 with an Achroplan 40×/0.8 NA water-immersion objective. A single epithelial cell was ruptured by focusing the Mode-locked Ti:Sapphire femtosecond laser beam over a triangular region of interest (~6 μm²) that included the apical membrane of the epithelial cell. The region of interest was scanned horizontally by the femtosecond laser at 100 μs/μm. By using the "bleach control" program in the LSM 510, we were able to immediately (less than 1 second) switch between the imaging mode and the ablation mode. The femtosecond laser beam was produced from a Coherent Chameleon system (Coherent, Santa Clara, CA) with 800-nm wavelength, 140-fs pulse duration, and 80-MHz repetition rate. The average power at the sample plane was ~600 mW, the pulse energy was ~7.5 nJ per pulse, and the peak power was ~37.5 kW.

Statistical Analysis

The ratio of lumen area was defined as the minimum cross-sectional area of airways after treatment divided by initial cross-sectional area. Statistical tests of significance of the ratio of lumen area were performed with one-way ANOVA using commercial software (SPSS v. 16; SPSS, Chicago, IL), and a *P* value less than 0.05 was considered statistically significant.

RESULTS

Adenosine and A₁ Receptor Mediate the Local Epithelial Injury-Induced Airway Contraction

In our previous studies, we ruled out the involvement of P2 purinoceptor receptors on smooth muscle cells in local epithelial injury-induced airway contraction; however, we did not completely rule out ATP as a soluble mediator, because inhibition of P2 purinoceptor receptors did not block the ATP-induced airway contraction (1). One explanation for these results would be activation of adenosine receptors to stimulate airway contraction from ATP metabolites, such as AMP and adenosine. To test this possibility, we inhibited the adenosine receptors with 2 μM CGS15943, a non-selective adenosine receptor antagonist (36, 37). Laser ablation of single epithelial cells induced an increase in Ca²⁺ oscillations (*see* Figure E1A in the online supplement), and airway contraction to 70% of the original cross-sectional area (Figures 1A and 1C and Movie E1). However, inhibition of adenosine receptors with CGS15943 completely blocked Ca²⁺ oscillations in smooth muscle cells and airway contraction induced by local epithelial injury, but did not block the Ca²⁺ wave in epithelial cells (Figure 1C, Figure E1B, and Movie E2). To confirm that adenosine is the soluble mediator, we incubated lung tissue slices with 5 units/ml adenosine deaminase (ADA), an enzyme that catalyzes the deamination of adenosine (38), and found that ADA significantly blocked the local epithelial injury-induced airway contraction (Figure 1C).

Because all the four subtypes of the adenosine receptors (A₁, A_{2A}, A_{2B}, and A₃) have been demonstrated to mediate smooth muscle contraction (5, 11–16, 39–41), we next investigated the roles of these four adenosine receptors. We found that selective inhibition of the A₁ receptor with 5–50 μM PSB36 (42, 43) or 4 μM SLV320 (44) significantly blocked Ca²⁺ oscillations in smooth muscle cells and airway contraction induced by local




Figure 1. Adenosine released from a wounded single epithelial cell activates A₁ receptors to induce airway contraction. (A) Bright-field images of a small airway embedded in a lung tissue slice before and ~40 seconds after laser ablation demonstrate that damage of a single epithelial cell induces global airway contraction. Blue arrow points to the ablated epithelial cell, the apical membrane of which was ruptured by a pulsed femtosecond laser. Red dashed line outlines the lumen cross-sectional area before laser ablation. (B) Inhibition of A₁ receptors with 50 μM PSB36 blocked local epithelial injury-induced airway contraction. (C) Local epithelial injury induced airway contraction, which leads to a roughly 30% reduction of the original cross-sectional area in control airways ($n = 33$ airways from 12 rats). Statistical tests demonstrate that inhibition of adenosine receptors with 2 μM 9-chloro-2-(2-furanyl)-[1,2,4]triazolo[1,5-c]quinazolin-5-amine (CGS15943) (CGS, a nonselective adenosine receptor antagonist; $n = 7$ airways from 5 rats), inhibition of adenosine A₁ receptors with 5–50 μM PSB36 (50 μM for 15 min incubation time, or 5 μM for 60 min; $n = 5$ airways from 3 rats for 50 μM PSB36 treatment) or 4 μM SLV320 ($n = 9$ airways from 4 rats), or deletion of adenosine with 5 units/ml adenosine deaminase (ADA; $n = 4$ airways from 3 rats) significantly blocked the airway contraction induced by local epithelial injury, whereas inhibition of adenosine A_{2A} receptors with 5 μM ZM241385 ($n = 4$ airways from 2 rats), A_{2B} receptors with 10 μM PSB603 ($n = 6$ airways from 5 rats), or A₃ receptors with 5 μM MRS1334 ($n = 8$ airways from 6 rats) did not. (D) Statistical tests demonstrate that CGS15943 significantly blocked airway contraction induced by 10 μM adenosine (ADO, a nonselective adenosine receptor agonist; $n = 4$ airways from 4 rats for N6-cyclopentyladenosine [CPA] and $n = 4$ airways from 4 rats for CPA + CGS treatment) or 10 μM ATP ($n = 5$ airways from 5 rats for ATP and $n = 4$ airways from 4 rats for ATP + CGS), but not 10 μM ATP-γ-S ($n = 6$ airways from 4 rats for ATP-γ-S and $n = 5$ airways from 4 rats for ATP-γ-S + CGS). As a positive control, 25 mM KCl or 1 μM acetylcholine was used to verify the viability and contractility of airway smooth muscle cells after laser ablation experiments for each treatment. The ratio of lumen area was defined as the minimum cross-sectional area of airways after treatment divided by initial cross-sectional area. Statistical tests of significance of the ratio of lumen area between control airways and each treatment condition were performed with one-way ANOVA, and there are similar numbers of control airways for each treatment condition. * $P < 0.05$ was considered statistically significant.

epithelial injury, but, again, did not block the Ca²⁺ wave in epithelial cells (Figures 1B and 1C, Figure E1C, and Movie E3). In contrast, selective inhibition of the A_{2A} receptor with 5 μM ZM241385 (45, 46), A_{2B} receptor with 10 μM PSB603 (47, 48), or A₃ receptor with 5 μM MRS1334 (49) did not block the local epithelial injury-induced Ca²⁺ oscillations in smooth muscle cells and airway contraction (Figure 1C).

To test the role of adenosine receptors in mediating ATP-induced airway contraction, we investigated the effects of ATP, its nonhydrolyzable analog, ATP-γ-S, and adenosine on airway caliber in the presence and absence of adenosine receptor antagonist CGS15943. We found that 10 μM adenosine, ATP, or ATP-γ-S induced airway contraction (Figure 1D). Inhibition of adenosine receptors with CGS15943 significantly blocked airway contraction induced by adenosine and ATP, but not by ATP-γ-S (Figure 1D).

EP₃ Receptors Participate in the Local Epithelial Injury-Induced Airway Constriction

PGE₂ has been shown to be released by mechanical scratch of the guinea pig tracheal mucosa (19), indicating that epithelial cell damage could possibly increase PGE₂ levels in lung tissue. Thus, we investigated the role of PGE₂ in local epithelial injury-induced airway contraction by inhibiting COX-2, an enzyme that mediates PGE₂ production. We found that inhibition of COX-2 with 20 μM indomethacin, a nonselective COX inhibitor (50–52), or with 10 μM NS-398, a selective COX-2 inhibitor (24, 53, 54), completely blocked Ca²⁺ oscillations in smooth muscle cells and airway contraction induced by local epithelial injury, but did not block the Ca²⁺ wave in epithelial cells (Figure 2A, Figure E1D, and Movie E4).

Because there are four EP receptors presented on airway smooth muscle cells (55), we next attempted to determine whether a selective EP receptor(s) mediates the local epithelial injury-induced airway contraction. We found that 10–25 μM L-798106, a selective EP₃ antagonist (56–59), and 100 μM AH6809, a nonselective rat EP₁, EP₂, and EP₃ receptor inhibitor (59–61), completely blocked Ca²⁺ oscillations in smooth muscle cells and airway contraction induced by local epithelial injury, but did not block the Ca²⁺ wave in epithelial cells (Figure 2B, Figure E1E, and Movie E5).

To investigate the order in which A₁ and EP₃ receptors were activated, we assessed the effects of CPA, a selective A₁R agonist (45, 62), and 11-PGE₂, a stable synthetic analog of PGE₂ that selectively activates EP₃ receptor (61, 63), on airway caliber in the presence and absence of EP₃ receptor antagonist L-798106. We found that both 10 μM CPA and 11-PGE₂ induced airway contraction (Figure 2C). Inhibition of EP₃ receptors with L-798106 significantly blocked airway contraction induced by CPA or 11-PGE₂, but not by 25 mM KCl, a VGCC agonist (Figure 2C). However, inhibition of A₁R with 50 μM PSB36 did not block airway contraction with 11-PGE₂ (Figure 2C). Thus, the role of A₁R is upstream of EP₃ and VGCC in the activation of smooth muscle contraction after airway epithelial injury.

Adenosine A₁R Activates EP₃ Receptors via EGFR/PDGFR Pathway

It has been demonstrated that adenosine A₁ receptor mediates the transactivation of the EGFR in rat cortical neurons (17), whereas activation of receptor tyrosine kinases, such as EGFR and PDGFR, could increase the production of PGE₂ in human or guinea pig tracheal smooth muscle cells (23, 24). Thus, we hypothesized that receptor tyrosine kinases are necessary for adenosine A₁R-mediated EP₃ receptor activation. To assess the role of receptor tyrosine kinases in local epithelial injury-induced airway contraction, we blocked receptor tyrosine kinases with 100 μM AG18, an inhibitor of EGFR and PDGFR (64–66), and 10 μM AG1478 (24, 67, 68) or 10 μM AG1296 (68–70), which are selective inhibitors of EGFR and PDGFR, respectively. We observed that inhibition of EGFR or PDGFR significantly blocked Ca²⁺ oscillations in smooth muscle cells and airway contraction induced by local epithelial injury, but did not block the Ca²⁺ wave in epithelial cells (Figure 3A, Figure E1F, and Movie E6).




Figure 2. Prostaglandin E receptor 3 (EP₃) receptors mediate adenosine-induced airway contraction. (A) Statistical tests demonstrate that inhibition of cyclooxygenase (COX)-2 with 20 μ M indomethacin ($n = 5$ airways from 5 rats) or 10 μ M NS-398 ($n = 4$ airways from 2 rats) completely blocked local epithelial injury-induced airway contraction ($n = 10$ control airways from nine rats). (B) Statistical tests demonstrate that inhibition of EP₃ receptor with 100 μ M AH6809 ($n = 8$ airways from 8 rats) or 10–25 μ M L-798106 (25 μ M for 20 min incubation time or 10 μ M for 45 min; $n = 5$ airways from 2 rats for 25 μ M L-798106 treatment) completely blocked local epithelial injury-induced airway contraction ($n = 14$ control airways from 10 rats). (C) Statistical tests demonstrate that L-798106 significantly blocked airway contraction induced by 10 μ M CPA, an adenosine A₁ receptor agonist ($n = 5$ airways from 3 rats for CPA and $n = 5$ airways from 4 rats for CPA + L-798106), or 10 μ M 11-prostaglandins E₂ (PGE₂), an EP₃ receptor agonist ($n = 9$ airways from 5 rats for 11-PGE₂ and $n = 5$ airways from 4 rats for 11-PGE₂ + L-798106), but not 25 mM KCl, a L-type voltage-dependent Ca²⁺ channel (VGCC) agonist ($n = 6$ airways from 4 rats for KCl and $n = 5$ airways from 4 rats for KCl + L-798106). Inhibition of A₁R with 50 μ M PSB36 did not block airway contraction with 10 μ M 11-PGE₂ ($n = 4$ airways from 2 rats). * $P < 0.05$ was considered statistically significant.

To demonstrate whether the receptor tyrosine kinase pathway is involved in adenosine A₁R-mediated activation of EP₃ receptors, we stimulated A₁ and EP₃ receptors with CPA and 11-PGE₂, respectively, in the presence and absence of EGFR inhibitor, AG1478. We found that inhibition of EGFR with 10 μ M AG1478 significantly blocked airway contraction induced by 10 μ M CPA, but not by 10 μ M 11-PGE₂ or 25 mM KCl (Figure 3C). Thus, the role of EP₃ and VGCC in the activation of smooth muscle contraction is downstream of EGFR.

DISCUSSION

Local epithelial injury induces airway hyperresponsiveness (1); however, the underlying mechanism has not been identified. In this study, we investigated the underlying mechanisms of the rapid (<10 s) communication between local epithelial injury and airway constriction by combining a lung tissue slice model with a femtosecond laser ablation technique. We first identified adenosine as the soluble mediator initiating local epithelial injury-induced airway contraction via A₁R activation. We then revealed a novel signaling pathway that includes the sequence A₁R→EGFR/PDGFR→COX-2→EP₃→VGCCs→CICR, leading to increase in Ca²⁺ oscillations in airway smooth muscle cells and initiation of airway constriction. For the first time, we show a sequential link between adenosine A₁R, receptor tyrosine kinases, including EGFR and PDGFR, and prostaglandin receptor EP₃ in airway smooth muscle cells.

ATP regulates multiple biological responses, such as airway hyperresponsiveness in the lungs (6). As an energy source, ATP is maintained at a very high level in the cytoplasm of airway epithelial cells. Upon epithelial injury, the local concentration of ATP can rapidly increase to 125 μ M (38) and initiate a Ca²⁺ wave in airway epithelium by activating P2Y receptors on epithelial cells (4). However, the effects of local epithelial injury-induced ATP release on airway caliber have not been studied. In this study, we found that local epithelial injury-induced ATP activated both airway epithelial and smooth muscle cells. Locally, ATP activates P2 purinergic receptors on neighboring epithelial cells; however, over the time (~5–10 s [1]) it takes to diffuse to the underlying smooth muscle, ATP is rapidly (<1 s [71]) degraded to adenosine, leading to the stimulation of adenosine A₁ receptors on smooth muscle cells (Figure 4). The hypothesis that ATP is rapidly degraded into adenosine is supported by the results that ATP-induced airway contraction is blocked by inhibition of adenosine receptors (Figure 1D), but not by inhibition of P2 purinergic receptors (1). In the present study, we confirmed the role of adenosine in the local epithelial injury-induced airway contraction by both inhibiting A₁ receptors and neutralization of

adenosine with ADA; however, we cannot rule out the involvement of AMP, because AMP, which has been shown to activate A₁ receptors (9), might also be decreased by ADA.

ATP has been demonstrated to induce airway smooth muscle contraction through either directly activating P2X or P2Y on mouse airway smooth muscle cells (7) or indirectly stimulating P2Y receptors on epithelial cells, which release prostaglandins in guinea pig trachea (20). We can eliminate the direct effect of ATP on airway smooth muscle cells, because inhibition of P2X or P2Y on smooth muscle cells did not block the local epithelial injury-induced smooth muscle contraction (1). Furthermore, we can rule out the indirect effect of ATP on airway epithelial cells, because inhibition of P2 purinergic receptors on epithelial cells significantly decreased the Ca²⁺ wave in the epithelium, but did not block the local epithelial injury-induced airway contraction (1). Thus, we have identified a novel pathway in which adenosine derived from local epithelial injury-released ATP stimulates A₁ receptors on smooth muscle cells to initiate airway contraction in rat lung tissue slices.

The levels of adenosine in bronchoalveolar lavage fluid are increased in asthma, and hyperresponsiveness to adenosine is a hallmark of asthma (72). Adenosine is traditionally thought to induce airway contraction indirectly by activation of adenosine receptors,




Figure 3. Epidermal growth factor receptor (EGFR)/platelet-derived growth factor receptor (PDGFR) mediates A₁R-induced activation of EP₃ receptors. (A) Statistical tests demonstrate that inhibition of EGFR and PDGFR with 100 μ M AG18 ($n = 4$ airways from 3 rats), or inhibition of EGFR or PDGFR with 10 μ M AG1478 ($n = 5$ airways from 3 rats) and 10 μ M AG1296 ($n = 4$ airways from 2 rats), respectively, significantly blocked local epithelial injury-induced airway contraction ($n = 9$ control airways from 5 rats). (B) AG1478 significantly blocked airway contraction induced by 10 μ M CPA ($n = 5$ airways from 2 rats for CPA and $n = 6$ airways from 3 rats for CPA + AG1478), but not 10 μ M 11-PGE₂ ($n = 6$ airways from 4 rats for 11-PGE₂ and $n = 6$ airways from 4 rats for 11-PGE₂ + AG1478) or 25 mM KCl ($n = 6$ airways from 5 rats for KCl and $n = 6$ airways from 3 rats for KCl + AG1478). * $P < 0.05$ was considered statistically significant.




Figure 4. Schematic diagram illustrating the sequential pathway involved in local epithelial injury–induced airway contraction. A single wounded epithelial cell releases ATP that activates P2 purinergic receptors on epithelial cells, and rapidly degrades into adenosine. Adenosine diffuses to the underlying smooth muscle and stimulates adenosine A₁ receptors. The stimulated A₁ receptors activate EGFRs/PDGFRs, which increase the activity of COX-2. COX-2 increases PGE₂ or prostacyclin (PGI₂) production, which, in turn, activates EP₃ receptors. Activated EP₃ receptors stimulate VGCCs, which increase intracellular Ca²⁺ by the Ca²⁺-induced Ca²⁺ release (CICR) mechanism. SMC, smooth muscle cell.

either on neural nerves or on mast cells, which release acetylcholine and prostaglandins, respectively, to induce smooth muscle cell contraction (11). However, we ruled out the involvement of sensory nerves in adenosine-induced airway contraction (11–14), because inhibition of acetylcholine M2 receptors with atropine did not block the local epithelial injury–induced airway contraction (1). We also ruled out the involvement of mast cell degranulation via A_{2B} receptor, because neither inhibition of A_{2B} receptor (Figure 1C) nor inhibition of mast cell degranulation with 300 μM cromolyn sodium (data not shown) blocked the local epithelial injury–induced airway contraction. Thus, we conclude that adenosine directly induces airway smooth muscle contraction by activation of A₁ receptors on the airway smooth muscle cells. Although adenosine has been shown to activate A₁ and A_{2B} receptors on human airway smooth muscle cells and regulate adenylyl cyclase and Ca²⁺ signaling (15, 16), the underlying mechanism leading to mobilization of Ca²⁺ is not clear.

In this study, we show that inhibition of EGFR/PDGFR completely blocks the local epithelial injury–induced airway contraction. We further demonstrate that inhibition of EGFR significantly blocks the A₁R agonist–induced airway contraction, but not EP₃ agonist–induced contraction (Figure 3C). The results indicate that receptor tyrosine kinases, including EGFR and PDGFR, are necessary for A₁R-mediated EP₃ activation. Our results are consistent with those of previous studies in which adenosine A₁ receptors mediated the transactivation of EGFRs in neural cells (17). Because receptor tyrosine kinases regulate many cellular functions, such as cell migration, differentiation, proliferation, apoptosis, and inflammation (73, 74), it is possible that local epithelial injury–induced receptor tyrosine kinase activation may have an even broader impact on pulmonary pathology. Multiple signaling pathways could be involved in EGFR/PDGFR–induced COX-2 production, such as phosphoinositide 3-kinase/Akt/NF-κB (24), mitogen-activated protein kinase kinase/mitogen-activated protein kinase (23, 75), or c-Src (76), and further study is needed to reveal the downstream signaling pathways.

Inhibition of COX-2 with indomethacin and NS-398 completely blocked the local epithelial injury–induced airway contraction, demonstrating the involvement of prostanoids in this process. To confirm this, we found that inhibition of EP₃ receptors also blocked the airway contraction induced by local epithelial injury. Although PGE₂ is traditionally considered an agonist of EP₃ receptors, PGI₂ has recently been shown to activate EP₃ receptors (59), indicating that either PGE₂ or PGI₂ could be released upon local epithelial injury. Our results are consistent with the studies in which activation of receptor tyrosine kinases leads to COX-2 expression (75–77) and PGE₂ generation (24). Our results are also consistent with the studies showing that smooth muscle cell contraction can be stimulated by self-generated prostanoids (23).

VGCC-mediated Ca²⁺ influx induces a large amount of intracellular Ca²⁺ release, which regulates numerous cellular functions, including smooth muscle contraction. In our previous study, we demonstrated that inhibition of VGCCs with nifedipine completely blocked local epithelial injury–induced airway contraction. We further showed that inhibition of A₁ and EP₃ receptors did not block KCl-induced airway contraction (Figures 2C and 3C), whereas inhibition of VGCCs blocked A₁ and EP₃ receptor agonist-induced airway contraction (data not shown). These results suggest that A₁R-, EGFR-, and EP₃ receptor-mediated pathways are upstream and dependent on VGCCs.

We acknowledge the use of relatively high concentrations of some chemical inhibitors in comparison to reported negative log dissociation constants or half maximal inhibitory concentration (IC₅₀) (Table E1), and thus off-target effects for one or more of the compounds are possible. However, in our present study, we report a series of compelling observations (multiple agonists and antagonists for each receptor) that are all consistent with the revealed molecular pathway underlying local epithelial injury–induced airway contraction. Thus, it is essentially impossible that all potential off-target effects would lead to the same set of conclusions.

In conclusion, we have identified a novel sequence of events that provides the underlying mechanism by which local airway epithelial injury can induce global airway smooth muscle contraction. Airway epithelial cell injury releases ATP, which is rapidly degraded to adenosine. Adenosine can diffuse to the underlying smooth muscle and initiates the activation of A₁R, EGFR and PDGFR, COX-2, EP₃, and VGCCs. Our studies provide direct evidence that local epithelial injury could contribute to airway hyperresponsiveness in subjects with asthma, and provides potentially new pharmacological targets for asthma treatment.

Author disclosures are available with the text of this article at www.atsjournals.org.

Acknowledgments: The authors thank members of the laboratory of Professor John Weiss at the University of California, Irvine, especially Dr. Hong Yin for assistance with the preparation of lung tissue slices. They also thank Dr. Tatiana Krasieva for assistance in multiphoton microscopy and femtosecond laser ablation, and the Laser Microbeam and Medical Program at the University of California, Irvine.

References

- Zhou J, Alvarez-Elizondo MB, Botvinick E, George SC. Local small airway epithelial injury induces global smooth muscle contraction and airway constriction. *J Appl Physiol* 2012;112:627–637.
- Frame MK, de Feijter AW. Propagation of mechanically induced intercellular calcium waves via gap junctions and ATP receptors in rat liver epithelial cells. *Exp Cell Res* 1997;230:197–207.
- Woodruff ML, Chaban VV, Worley CM, Dirksen ER. PKC role in mechanically induced Ca²⁺ waves and ATP-induced Ca²⁺ oscillations in airway epithelial cells. *Am J Physiol* 1999;276:L669–L678.
- Isakson BE, Evans WH, Boitano S. Intercellular Ca²⁺ signaling in alveolar epithelial cells through gap junctions and by extracellular ATP. *Am J Physiol Lung Cell Mol Physiol* 2001;280:L221–L228.

5. Eltzschig HK. Adenosine: an old drug newly discovered. *Anesthesiology* 2009;111:904–915.
6. Vliet A, Bove PF. Purinergic signaling in wound healing and airway remodeling. In: Picher M, Boucher RC, editors. Purinergic regulation of respiratory diseases. Heidelberg: Springer Netherlands; 2011. pp. 139–157.
7. Bergner A, Sanderson MJ. Atp stimulates Ca^{2+} oscillations and contraction in airway smooth muscle cells of mouse lung slices. *Am J Physiol Lung Cell Mol Physiol* 2002;283:L1271–L1279.
8. Dubyak GR, el-Moatassim C. Signal transduction via P2-purinergic receptors for extracellular ATP and other nucleotides. *Am J Physiol Cell Physiol* 1993;265:C577–C606.
9. Rittiner JE, Korboukh I, Hull-Ryde EA, Jin J, Janzen WP, Frye SV, Zylka MJ. Amp is an adenosine a1 receptor agonist. *J Biol Chem* 2012;287:5301–5309.
10. Gündüz D, Aslam M, Krieger U, Becker L, Grebe M, Arshad M, Sedding DG, Härtel FV, Abdallah Y, Piper HM, et al. Opposing effects of atp and adenosine on barrier function of rat coronary microvasculature. *J Mol Cell Cardiol* 2012;52:962–970.
11. Polosa R. Adenosine-receptor subtypes: their relevance to adenosine-mediated responses in asthma and chronic obstructive pulmonary disease. *Eur Respir J* 2002;20:488–496.
12. Hua X, Erikson CJ, Chason KD, Rosebrock CN, Deshpande DA, Penn RB, Tilley SL. Involvement of A1 adenosine receptors and neural pathways in adenosine-induced bronchoconstriction in mice. *Am J Physiol Lung Cell Mol Physiol* 2007;293:L25–L32.
13. Reynolds SM, Docherty R, Robbins J, Spina D, Page CP. Adenosine induces a cholinergic tracheal reflex contraction in guinea pigs *in vivo* via an adenosine A1 receptor-dependent mechanism. *J Appl Physiol* 2008;105:187–196.
14. Calzetta L, Spina D, Cazzola M, Page CP, Facciolo F, Rendina EA, Matera MG. Pharmacological characterization of adenosine receptors on isolated human bronchi. *Am J Respir Cell Mol Biol* 2011;45:1222–1231.
15. Mundell SJ, Olah ME, Panettieri RA Jr, Benovic JL, Penn RB. Regulation of G protein-coupled receptor-adenylyl cyclase responsiveness in human airway smooth muscle by exogenous and autocrine adenosine. *Am J Respir Cell Mol Biol* 2001;24:155–163.
16. Ethier MF, Madison JM. Adenosine A1 receptors mediate mobilization of calcium in human bronchial smooth muscle cells. *Am J Respir Cell Mol Biol* 2006;35:496–502.
17. Xie K-q, Zhang L-m, Cao Y, Zhu J, Feng L-y. Adenosine A1 receptor-mediated transactivation of the EGF receptor produces a neuroprotective effect on cortical neurons *in vitro*. *Acta Pharmacol Sin* 2009;30:889–898.
18. Carey MA, Germolec DR, Langenbach R, Zeldin DC. Cyclooxygenase enzymes in allergic inflammation and asthma. *Prostaglandins Leukot Essent Fatty Acids* 2003;69:157–162.
19. Orehek J, Douglas JS, Bouhuys A. Contractile responses of the guinea-pig trachea *in vitro*: modification by prostaglandin synthesis-inhibiting drugs. *J Pharmacol Exp Ther* 1975;194:554–564.
20. Flores-Soto E, Carbajal V, Reyes-García J, García-Hernández L, Figueroa A, Checa M, Barajas-López C, Montaña L. In airways ATP refills sarcoplasmic reticulum via P2X smooth muscle receptors and induces contraction through P2Y epithelial receptors. *Pflügers Arch* 2011;461:261–275.
21. Ruan YC, Zhou W, Chan HC. Regulation of smooth muscle contraction by the epithelium: role of prostaglandins. *Physiology (Bethesda)* 2011; 26:156–170.
22. Delamere F, Holland E, Patel S, Bennett J, Pavord I, Knox A. Production of PGE2 by bovine cultured airway smooth muscle cells and its inhibition by cyclo-oxygenase inhibitors. *Br J Pharmacol* 1994;111: 983–988.
23. Schaafsma D, Gosens R, Bos IS, Meurs H, Zaagsma J, Nelemans SA. Role of contractile prostaglandins and Rho-kinase in growth factor-induced airway smooth muscle contraction. *Respir Res* 2005;6:85.
24. Yang C-M, Lee I-T, Lin C-C, Yang Y-L, Luo S-F, Kou YR, Hsiao L-D. Cigarette smoke extract induces COX-2 expression via a PKC α /c-Src/EGFR, PDGFR/PI3K/Akt/NF- κ B pathway and P300 in tracheal smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 2009;297:L892–L902.
25. Montaña L, Cruz-Valderrama J, Figueroa A, Flores-Soto E, García-Hernández L, Carbajal V, Segura P, Méndez C, Díaz V, Barajas-López C. Characterization of P2Y receptors mediating ATP induced relaxation in guinea pig airway smooth muscle: involvement of prostaglandins and K^{+} channels. *Pflügers Arch* 2011;462:573–585.
26. Cuthbert MF. Effect on airways resistance of prostaglandin E1 given by aerosol to healthy and asthmatic volunteers. *BMJ* 1969;4:723–726.
27. Mathe AA, Hedqvist P. Effect of prostaglandins F2 alpha and E2 on airway conductance in healthy subjects and asthmatic patients. *Am Rev Respir Dis* 1975;111:313–320.
28. Tilley SL, Hartney JM, Erikson CJ, Jania C, Nguyen M, Stock J, McNeisch J, Valancius C, Panettieri RA, Penn RB, et al. Receptors and pathways mediating the effects of prostaglandin E2 on airway tone. *Am J Physiol Lung Cell Mol Physiol* 2003;284:L599–L606.
29. Fortner CN, Breyer RM, Paul RJ. EP2 receptors mediate airway relaxation to substance P, ATP, and PGE2. *Am J Physiol Lung Cell Mol Physiol* 2001;281:L469–L474.
30. Sugimoto Y, Narumiya S. Prostaglandin E receptors. *J Biol Chem* 2007; 282:11613–11617.
31. Chen W, Andom T, Bhattacherjee P, Paterson C. Intracellular calcium mobilization following prostaglandin receptor activation in human ciliary muscle cells. *Curr Eye Res* 1997;16:847–853.
32. Ndukwu IM, White SR, Leff AR, Mitchell RW. EP1 receptor blockade attenuates both spontaneous tone and PGE2-elicited contraction in guinea pig trachealis. *Am J Physiol Lung Cell Mol Physiol* 1997;273: L626–L633.
33. Qian YM, Jones RL, Chan KM, Stock AI, Ho JK. Potent contractile actions of prostanoid EP3-receptor agonists on human isolated pulmonary artery. *Br J Pharmacol* 1994;113:369–374.
34. Bergner A, Sanderson MJ. Acetylcholine-induced calcium signaling and contraction of airway smooth muscle cells in lung slices. *J Gen Physiol* 2002;119:187–198.
35. Bai Y, Sanderson MJ. The contribution of Ca^{2+} signaling and Ca^{2+} sensitivity to the regulation of airway smooth muscle contraction is different in rats and mice. *Am J Physiol Lung Cell Mol Physiol* 2009; 296:L947–L958.
36. Fozard JR, Tigani B, Wolber C, Williams I, Mazzoni L, Hannon JP. Modeling the response of the asthmatic airways to adenosine: mechanisms and receptors. *Drug Dev Res* 2003;59:23–29.
37. Williams M, Francis J, Ghai G, Braunwalder A, Psychoyos S, Stone GA, Cash WD. Biochemical characterization of the triazolquinazoline, CGS 15943, a novel, non-xanthine adenosine antagonist. *J Pharmacol Exp Ther* 1987;241:415–420.
38. Yin J, Xu K, Zhang J, Kumar A, Yu F-SX. Wound-induced ATP release and EGF receptor activation in epithelial cells. *J Cell Sci* 2007;120: 815–825.
39. Thorne JR, Danahay H, Broadley KJ. Analysis of the bronchoconstrictor responses to adenosine receptor agonists in sensitized guinea-pig lungs and trachea. *Eur J Pharmacol* 1996;316:263–271.
40. Pauwels RA, van der Straeten ME. An animal model for adenosine-induced bronchoconstriction. *Am Rev Respir Dis* 1987;136:374–378.
41. Hannon JP, Tigani B, Schuurman H-J, Fozard JR. Suppression of adenosine A3 receptor-mediated hypotension and mast cell degranulation in the rat by dexamethasone. *J Pharmacol Exp Ther* 2002;302:725–730.
42. Abo-Salem OM, Hayallah AM, Bilkei-Gorzo A, Filipek B, Zimmer A, Müller CE. Antinociceptive effects of novel A2B adenosine receptor antagonists. *J Pharmacol Exp Ther* 2004;308:358–366.
43. Töpfer M, Burbiel CE, Müller CE, Knittel J, Verspohl EJ. Modulation of insulin release by adenosine a1 receptor agonists and antagonists in ins-1 cells: The possible contribution of 86rb+ efflux and 45ca2+ uptake. *Cell Biochem Funct* 2008;26:833–843.
44. Kalk P, Eggert B, Relle K, Godes M, Heiden S, Sharkovska Y, Fischer Y, Ziegler D, Bielenberg GW, Hocher B. The adenosine a1 receptor antagonist slv320 reduces myocardial fibrosis in rats with 5/6 nephrectomy without affecting blood pressure. *Br J Pharmacol* 2007; 151:1025–1032.
45. Feoktistov I, Biaggioni I. Adenosine a2b receptors. *Pharmacol Rev* 1997; 49:381–402.
46. Poucher SM, Keddie JR, Singh P, Stogdall SM, Caulkett PW, Jones G, Coll MG. The *in vitro* pharmacology of ZM 241385, a potent, non-xanthine A2A selective adenosine receptor antagonist. *Br J Pharmacol* 1995;115:1096–1102.
47. Borrmann T, Hinz S, Bertarelli DCG, Li W, Florin NC, Scheiff AB, Müller CE. 1-alkyl-8-(piperazine-1-sulfonyl)phenylxanthines: Development

- and characterization of adenosine a2b receptor antagonists and a new radioligand with subnanomolar affinity and subtype specificity. *J Med Chem* 2009;52:3994–4006.
48. van der Hoeven D, Wan TC, Gizewski ET, Kreckler LM, Maas JE, Van Orman J, Ravid K, Auchampach JA. A role for the low-affinity A2B adenosine receptor in regulating superoxide generation by murine neutrophils. *J Pharmacol Exp Ther* 2011;338:1004–1012.
 49. Jiang J-I, van Rhee AM, Chang L, Patchornik A, Ji X-d, Evans P, Melman N, Jacobson KA. Structure–activity relationships of 4-(phenylethynyl)-6-phenyl-1,4-dihydropyridines as highly selective A3 adenosine receptor antagonists. *J Med Chem* 1997;40:2596–2608.
 50. Mounkaila B, Marthan R, Roux E. Biphasic effect of extracellular ATP on human and rat airways is due to multiple P2 purinoceptor activation. *Respir Res* 2005;6:143.
 51. Palomer A, Cabré F, Pascual J, Campos J, Trujillo MA, Entrena A, Gallo MA, García L, Mauleón D, Espinosa A. Identification of novel cyclooxygenase-2 selective inhibitors using pharmacophore models. *J Med Chem* 2002;45:1402–1411.
 52. Yamazaki J, Kitamura K. Cell-to-cell communication via nitric oxide modulation of oscillatory Cl^- currents in rat intact cerebral arterioles. *J Physiol* 2001;536:67–78.
 53. Ogino K, Hatanaka K, Kawamura M, Katori M, Harada Y. Evaluation of pharmacological profile of meloxicam as an anti-inflammatory agent, with particular reference to its relative selectivity for cyclooxygenase-2 over cyclooxygenase-1. *Pharmacology* 1997;55:44–53.
 54. Yamakawa T, Ohnaka K, Tanaka S, Utsunomiya H, Kamei J, Kadonosono K. Cyclooxygenase-2 induction by lysophosphatidylcholine in cultured rat vascular smooth muscle cells: involvement of the p38MAPK pathway. *Biomed Res* 2008;29:1–8.
 55. Chung KF. Evaluation of selective prostaglandin e2 (PGE2) receptor agonists as therapeutic agents for the treatment of asthma. *Sci STKE* 2005;2005:pe47.
 56. Bassil AK, Borman RA, Jarvie EM, McArthur-Wilson RJ, Thangiah R, Sung EZH, Lee K, Sanger GJ. Activation of prostaglandin EP receptors by lubiprostone in rat and human stomach and colon. *Br J Pharmacol* 2008;154:126–135.
 57. Fairbrother SE, Smith JE, Borman RA, Cox HM. Characterization of the EP receptor types that mediate longitudinal smooth muscle contraction of human colon, mouse colon and mouse ileum. *Neurogastroenterol Motil* 2011;23:782–e336.
 58. Juteau H, Gareau Y, Labelle M, Sturino CF, Sawyer N, Tremblay N, Lamontagne S, Carrière M-C, Denis D, Metters KM. Structure–activity relationship of cinnamic acylsulfonamide analogues on the human EP3 prostanoid receptor. *Bioorg Med Chem* 2001;9:1977–1984.
 59. Orie NN, Clapp LH. Role of prostanoid IP and EP receptors in mediating vasorelaxant responses to PGI2 analogues in rat tail artery: evidence for GI/O modulation via EP3 receptors. *Eur J Pharmacol* 2011;654:258–265.
 60. Boie Y, Stocco R, Sawyer N, Slipetz DM, Ungrin MD, Neuschäfer-Rube F, Püschel GP, Metters KM, Abramovitz M. Molecular cloning and characterization of the four rat prostaglandin E2 prostanoid receptor subtypes. *Eur J Pharmacol* 1997;340:227–241.
 61. Jadhav V, Jabre A, Lin S-Z, Lee TJ-F. EP1- and EP3-receptors mediate prostaglandin E2-induced constriction of porcine large cerebral arteries. *J Cereb Blood Flow Metab* 2004;24:1305–1316.
 62. Wolber C, Fozard JR. The receptor mechanism mediating the contractile response to adenosine on lung parenchymal strips from actively sensitised, allergen-challenged brown Norway rats. *Naunyn Schmiedebergs Arch Pharmacol* 2005;371:158–168.
 63. Jia Z, Person MD, Dong J, Shen J, Hensley SC, Stevens JL, Monks TJ, Lau SS. Grp78 is essential for 11-deoxy-16,16-dimethyl PGE2-mediated cytoprotection in renal epithelial cells. *Am J Physiol Renal Physiol* 2004;287:F1113–F1122.
 64. Carmines PK, Fallet RW, Che Q, Fujiwara K. Tyrosine kinase involvement in renal arteriolar constrictor responses to angiotensin II. *Hypertension* 2001;37:569–573.
 65. Gazit A, Yaish P, Gilon C, Levitzki A. Tyrohostins I: synthesis and biological activity of protein tyrosine kinase inhibitors. *J Med Chem* 1989;32:2344–2352.
 66. Soltoff SP. Evidence that tyrohostins AG10 and AG18 are mitochondrial uncouplers that alter phosphorylation-dependent cell signaling. *J Biol Chem* 2004;279:10910–10918.
 67. Kim J, Lee C-K, Park H-J, Kim HJ, So HH, Lee KS, Lee HM, Roh HY, Choi WS, Park TK, *et al.* Epidermal growth factor induces vasoconstriction through the phosphatidylinositol 3-kinase-mediated mitogen-activated protein kinase pathway in hypertensive rats. *J Pharmacol Sci* 2006;101:135–143.
 68. Levitzki A, Gazit A. Tyrosine kinase inhibition: an approach to drug development. *Science* 1995;267:1782–1788.
 69. Shimizu H, Nakagawa Y, Murakami C, Aoki N, Kim-Mitsuyama S, Miyazaki H. Protein tyrosine phosphatase PTPeM negatively regulates PDGF β -receptor signaling induced by high glucose and PDGF in vascular smooth muscle cells. *Am J Physiol Cell Physiol* 2010;299:C1144–C1152.
 70. Yang C-M, Lin C-C, Lee I-T, Lin Y-H, Yang C, Chen W-J, Jou M-J, Hsiao L-D. Japanese encephalitis virus induces matrix metalloproteinase-9 expression via a ROS/c-Src/PDGFR/PI3K/Akt/MAPKs-dependent AP-1 pathway in rat brain astrocytes. *J Neuroinflammation* 2012;9:12.
 71. Dunwiddie TV, Diao L, Proctor WR. Adenine nucleotides undergo rapid, quantitative conversion to adenosine in the extracellular space in rat hippocampus. *J Neurosci* 1997;17:7673–7682.
 72. Polosa R, Rorke S, Holgate ST. Evolving concepts on the value of adenosine hyperresponsiveness in asthma and chronic obstructive pulmonary disease. *Thorax* 2002;57:649–654.
 73. Ricchi P, Zarrilli R, di Palma A, Acquaviva AM. Nonsteroidal anti-inflammatory drugs in colorectal cancer: from prevention to therapy. *Br J Cancer* 2003;88:803–807.
 74. Wong WS. Inhibitors of the tyrosine kinase signaling cascade for asthma. *Curr Opin Pharmacol* 2005;5:264–271.
 75. Xu K, Shu H-KG. EGFR activation results in enhanced cyclooxygenase-2 expression through p38 mitogen-activated protein kinase-dependent activation of the SP1/SP3 transcription factors in human gliomas. *Cancer Res* 2007;67:6121–6129.
 76. Xu K, Kitchen CM, Shu H-KG, Murphy TJ. Platelet-derived growth factor-induced stabilization of cyclooxygenase 2 mRNA in rat smooth muscle cells requires the c-Src family of protein-tyrosine kinases. *J Biol Chem* 2007;282:32699–32709.
 77. Goppelt-Struebe M, Rehm M, Schaefer HJ. Induction of cyclooxygenase-2 by platelet-derived growth factor (PDGF) and its inhibition by dexamethasone are independent of NF- κ B/I κ B transcription factors. *Naunyn Schmiedebergs Arch Pharmacol* 2000;361:636–645.