UC Davis

UC Davis Previously Published Works

Title

Phase-Variance Optical Coherence Tomographic Angiography Imaging of Choroidal Perfusion Changes Associated With Acute Posterior Multifocal Placoid Pigment Epitheliopathy.

Permalink

<https://escholarship.org/uc/item/0c31b6pw>

Journal

JAMA ophthalmology, 134(8)

ISSN

2168-6165

Authors

Park, Susanna S
Thinda, Sumeer
Kim, Dae Yu
et al.

Publication Date

2016-08-01

DOI

10.1001/jamaophthalmol.2016.1645

Peer reviewed

Letters

OBSERVATION

Phase-Variance Optical Coherence Tomographic Angiography Imaging of Choroidal Perfusion Changes Associated With Acute Posterior Multifocal Placoid Pigment Epitheliopathy

Acute posterior multifocal placoid pigment epitheliopathy (APMPPE) is characterized by bilateral multiple placoid white-gray lesions of the posterior pole at the level of the retinal pigment epithelium (RPE) and/or choriocapillaris, which self-resolve with recovery of vision.^{1,2} The etiology is unknown, but APMPPE has been associated with viral prodromes and vasculitis. The placoid lesions are thought to represent focal RPE inflammation or choriocapillaris ischemia based on fluorescein angiogram findings of early hypofluorescence and late hyperfluorescence.

Phase-variance optical coherence tomographic angiography (pv-OCTA) allows noninvasive visualization of the choroidal blood flow in 3 dimensions in vivo.^{3,4} A custom-built OCT system (125-kHz line rate, 855-nm central wavelength, and 4.5- μ m axial resolution) was used to image choroidal perfusion over a 1.5 $\times$ 1.5-mm² area (equally spaced 350 A-scans in both x- and y-axes) in acute APMPPE.

Report of a Case | A woman in her mid to late 20s with viral prodrome presented with blurry vision in the left eye for 1 week. Visual acuity was 20/20 OD and 20/400 OS. The right eye was normal, and the left eye had normal anterior segment and vitreous, but fundus showed multiple white-gray placoid lesions posteriorly with early hypofluorescence and late hyperfluorescence on fluorescein angiogram (**Figure 1**). Patchy hypoperfusion of the choriocapillaris and Sattler layer was noted on pv-OCTA corresponding to focal areas of photoreceptor and RPE irregularity on OCT B-scans (**Figure 2A-C**). Serology test results were negative for syphilis, sarcoidosis, and tuberculosis.

After 2 weeks, similar placoid lesions were noted in the right eye. After 6 weeks, placoid lesions resolved bilaterally to develop pigment clumping (**Figure 1D**) and choroidal perfusion mostly normalized on pv-OCTA. Visual acuity and photoreceptor irregularity on OCT minimally improved (**Figure 2D-F**). By the 6-month follow-up, visual acuity was 20/20 OD and 20/30 OS, with almost normalized photoreceptor layer on OCT (**Figure 2G**).

This study conforms to the World Medical Association Declaration of Helsinki⁵ and subsequent amendments. The research protocol was approved by the University of California, Davis Office of Human Research.

Figure 1. Fundus Photograph and Fluorescein Angiogram of the Left Eye

At presentation, white-gray placoid lesions are seen (A), with early hypofluorescence (B) and late hyperfluorescence (C) on angiography; pigmentary changes developed by 6 weeks' follow-up (D and E).

Figure 2. Left Macula Phase-Variance Optical Coherence Tomographic Angiography (pv-OCTA)

Discussion | Acute posterior multifocal placoid pigment epitheliopathy was first described by Gass in 1968,¹ yet the pathophysiology remains unclear because there is no histopathology of the placoid lesions.² During the acute phase, indocyanine green angiography shows hypofluorescence of the lesions without late hyperfluorescence and large choroidal vessels have been seen coursing through hypofluorescent areas on fluorescein angiogram and indocyanine green angiography, findings suggestive of choriocapillaris ischemia.⁶

In this case of APMPPE, serial pv-OCTA visualized the transient patchy hypoperfusion of the choriocapillaris and Sattler layer associated with acute APMPPE. These changes can be compared with published normal pv-OCTA choroidal perfusion maps.^{3,4} The resolution of the choroidal hypoperfusion on pv-OCT preceded the resolution of photoreceptor abnormality on OCT and visual recovery. Thus, the pathophysiology of APMPPE likely involves a primary hypoperfusion of the

inner choroid with resultant RPE and photoreceptor abnormality and vision loss. Concurrent edema or infiltration of the RPE or choroid is possible.

Perfusion images were produced by en face projection of thin slices from pv-OCTA volumetric data at 8 μm below the Bruch membrane for choriocapillaris visualization and 8 to 16 μm below the Bruch membrane for Sattler layer visualization. By using pv-OCTA signals overlaid on OCT intensity B-scan images, the corresponding retinal structural changes can be studied relative to the perfusion changes (Figure 2A). These focal changes in choroidal flow signal on pv-OCTA do not correlate with shadowing artifacts from overlying RPE changes. The focal lobular areas of choroidal hypoperfusion at the level of the choriocapillaris extend into the deeper Sattler layer and appear to correspond to the placoid lesions.

In conclusion, this case report used pv-OCTA to demonstrate transient inner choroidal flow changes associated with APMPPE.

This and similar OCTA systems can provide new insights into the pathogenesis of this condition. Future larger studies may determine whether these choroidal perfusion changes are universally observed in APMPPE and affect visual prognosis.

Susanna S. Park, MD, PhD
Sumeer Thinda, MD
Dae Yu Kim, PhD
Robert J. Zawadzki, PhD
John S. Werner, PhD

Author Affiliations: Department of Ophthalmology and Vision Science, University of California, Davis Eye Center, Sacramento (Park, Thinda, Zawadzki, Werner); Beckman Laser Institute Korea and Biomedical Engineering, Dankook University, Cheonan, Chungnam, South Korea (Kim).

Corresponding Author: Susanna S. Park, MD, PhD, University of California, Davis Eye Center, 4860 Y St, Ste 2400, Sacramento, CA 95817 (sscpark@ucdavis.edu).

Published Online: June 9, 2016. doi:10.1001/jamaophthalmol.2016.1645.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

Funding/Support: This study was supported in part by grant EYO24239 from the National Eye Institute (Dr Werner) and grant NRF-2015R1D1A1A02061699 from the Basic Science Research Program through the National Research Foundation of Korea.

Role of the Funder/Sponsor: The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Previous Presentation: This article was presented in part as a poster at the Annual Meeting of the Association for Research in Vision and Ophthalmology; May 7, 2015; Denver, Colorado.

Additional Contributions: We thank the patient for granting permission to publish this information.

1. Gass JD. Acute posterior multifocal placoid pigment epitheliopathy. *Arch Ophthalmol*. 1968;80(2):177-185.
2. Agarwal A. *Gass' Atlas of Macular Diseases*. 5th ed. Philadelphia, PA: Saunders; 2011.
3. Kim DY, Fingler J, Zawadzki RJ, et al. Optical imaging of the chorioretinal vasculature in the living human eye. *Proc Natl Acad Sci U S A*. 2013;110(35):14354-14359.
4. Schwartz DM, Fingler J, Kim DY, et al. Phase-variance optical coherence tomography: a technique for noninvasive angiography. *Ophthalmology*. 2014;121(1):180-187.
5. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-2194. doi:10.1001/jama.2013.281053.
6. Howe LJ, Woon H, Graham EM, Fitzke F, Bhandari A, Marshall J. Choroidal hypoperfusion in acute posterior multifocal placoid pigment epitheliopathy: an indocyanine green angiography study. *Ophthalmology*. 1995;102(5):790-798.

COMMENT & RESPONSE

Zika Virus, Microcephaly, and Ocular Findings

To the Editor We read with great interest the article "Ocular Findings in Infants With Microcephaly Associated With Presumed Zika Virus Congenital Infection in Salvador, Brazil" by de Paula Freitas et al¹ as well as the accompanying Invited Commentary "Zika Virus Infection and the Eye" by Jampol and Goldstein.² The authors are to be complimented for addressing the ophthalmologic need and building on their previous report of ophthalmic findings associated with Zika virus (ZIKV) in 3 infants.³

The authors in all 3 publications postulate direct ZIKV-induced ocular findings. However, it remains unclear whether the ocular lesions described in these reports are a direct result of ZIKV or a result of microcephaly itself. For 50 years, there has been well-documented history (beginning in this very journal)⁴ of both sporadic and heritable (autosomal recessive and autosomal dominant) forms of microcephaly associated with ocular findings similar to those recently attributed to ZIKV infection¹⁻³: pigmentary maculopathy, circumscribed chorioretinal atrophy, and optic nerve abnormalities.^{4,5} In addition, microcephaly has been associated with microcornea, microphthalmia, falciform folds, cataracts, retinal dysplasia, persistent fetal vasculature, vascular attenuation, nystagmus, and hyperopia,^{4,5} and we might expect other findings as the ZIKV epidemic unfolds.

Based on the consistency of the findings here with those previously reported, it is possible that ocular findings described as a direct result of the ZIKV infection may be secondary to the microcephaly.¹⁻³ It seems plausible that there could be a dual-mechanism process at work in these patients: microcephaly (secondary to ZIKV)-associated ocular abnormalities as well as directly induced ZIKV ocular abnormalities. We look forward to learning more about this potential dual mechanism as we gain additional experience with patients with ZIKV.

Darius M. Moshfeghi, MD
Homero Augusto de Miranda II, MD
Marcelo Cavalcante Costa, MD

Author Affiliations: Byers Eye Institute, Horngren Family Vitreoretinal Center, Department of Ophthalmology, Stanford University School of Medicine, Palo Alto, California (Moshfeghi); Ophthalmic Telemedicine Santa Casa of São Paulo, São Paulo, Brazil (de Miranda, Costa).

Corresponding Author: Darius M. Moshfeghi, MD, Byers Eye Institute, Horngren Family Vitreoretinal Center, Department of Ophthalmology, Stanford University School of Medicine, 2452 Watson Ct, Room 2277, Palo Alto, CA 94303 (dariusm@stanford.edu).

Published Online: June 2, 2016. doi:10.1001/jamaophthalmol.2016.1303.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Dr Moshfeghi reported serving as a consultant and on a data monitoring board for Alcon Surgical; receiving travel support from Clarity Medical Systems; serving on a steering committee for Iconic Therapeutics; having equity/intellectual property in InSitu Therapeutics, Inc; serving as a consultant to Genentech; having equity in Krypton Vision; being a founder of and having equity in Promisight; serving on scientific advisory boards for Synergetics and Thrombogenics; being a founder and director of and having equity in Versi, Inc; serving on a scientific advisory board, serving as a consultant, and having equity in Visunex Medical Systems Co Ltd; serving on a scientific advisory board for and having equity in Oraya Therapeutics, Inc; and receiving fees as a legal expert on retinopathy of prematurity. No other disclosures were reported.

1. de Paula Freitas B, de Oliveira Dias JR, Prazeres J, et al. Ocular findings in infants with microcephaly associated with presumed Zika virus congenital infection in Salvador, Brazil [published online February 9, 2016]. *JAMA Ophthalmol*. doi:10.1001/jamaophthalmol.2016.0267.
2. Jampol LM, Goldstein DA. Zika virus infection and the eye [published online February 9, 2016]. *JAMA Ophthalmol*. doi:10.1001/jamaophthalmol.2016.0284.
3. Ventura CV, Maia M, Bravo-Filho V, Góis AL, Belfort R Jr. Zika virus in Brazil and macular atrophy in a child with microcephaly. *Lancet*. 2016;387(10015):228.
4. McKusick VA, Stauffer M, Knox DL, Clark DB. Chorioretinopathy with hereditary microcephaly. *Arch Ophthalmol*. 1966;75(5):597-600.
5. Atchaneeyasakul LO, Linck L, Weleber RG. Microcephaly with chorioretinal degeneration. *Ophthalmic Genet*. 1998;19(1):39-48.