# Lawrence Berkeley National Laboratory

**Recent Work** 

## Title

Image simulation and experimental resolution of Lir{sup +} in LiCoO{sub 2}

## Permalink

https://escholarship.org/uc/item/0cn9r3pr

## Authors

Shao-Horn, Yang Croguennec, Laurence Delmas, Claude <u>et al.</u>

# **Publication Date**

2003-01-24

#### Image Simulation and Experimental Resolution of Li<sup>+</sup> in LiCoO<sub>2</sub>

Y. Shao-Horn,\* L. Croguennec,\*\* C. Delmas,\*\* E.C. Nelson,\*\*\* and M.A. O'Keefe\*\*\*

\*Department of Mechanical Engineering, MIT, Cambridge, MA 02139, USA \*\*Institut de Chimie de la Matière Condensée de Bordeaux-CNRS and Ecole Nationale Supérieure de Chimie et Physique de Bordeaux, Université Bordeaux I, France \*\*\*Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

#### DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof, or The Regents of the University of California.

*Ernest Orlando Lawrence Berkeley National Laboratory is an equal opportunity employer.* 

Ernest Orlando Lawrence Berkeley National Laboratory – LBNL-52100 Abs

#### Image Simulation and Experimental Resolution of Li<sup>+</sup> in LiCoO<sub>2</sub>

Y. Shao-Horn,\* L. Croguennec,\*\* C. Delmas,\*\* E.C. Nelson,\*\*\* and M.A. O'Keefe\*\*\*

\*Department of Mechanical Engineering Massachusetts Institute of Technology 77 Massachusetts Ave, Cambridge, MA 02139, USA

\*\*Institut de Chimie de la Matière Condensée de Bordeaux-CNRS and Ecole Nationale Supérieure de Chimie et Physique de Bordeaux Université Bordeaux I, 33608 Pessac cedex France

> \*\*\*Lawrence Berkeley National Laboratory Berkeley, CA 94720, USA

Advances in high-resolution transmission electron microscopes (TEMs) and associated software, particularly exit-surface wave (ESW) reconstruction of the scattered electron wave, push the resolution limit to better than 0.1nm, which provides opportunities to examine whether  $Li^+$  can be imaged and resolved in a transition metal oxide structure. From simulations of the exit-surface wave (ESW) generated by structural models of  $LiCoO_2$ , it was found that a microscope resolution of 0.1nm was needed to experimentally resolve individual lithium ions in the layered structure. The phase of the ESW is a function of the specimen potential (projected atom density). Our simulations show that the ESW phases of the lithium, cobalt and oxygen atom columns change almost linearly with specimen thickness. The simulated ESWs were then compared with experimental ESWs reconstructed from experimental focus-series along the [110] zone axis. Reconstructions were carried out, using the Philips/Brite-Euram software, from several focus-series of 20 TEM images obtained on a modified Philips CM300FEG/UT microscope with a native resolution of 0.17nm and a demonstrated reconstructed ESW resolution of 0.078nm. It was found consistent with the simulation results that the intensity associated with the Li<sup>+</sup> position in the experimental ESWs was the weakest in comparison to those of oxygen and cobalt. Both simulation and experimental results support the fact that Li<sup>+</sup> was resolved atomically in LiCoO<sub>2</sub>.

#### ACKNOWLEDGMENTS

National Science Foundation International Research Fellow Award INT-0000429 is acknowledged. This work was partially supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Materials Science Division, U.S. Department of Energy under Contract No. DE AC-03-76SF00098.