UCSF

UC San Francisco Previously Published Works

Title

Race, Research, and Women's Health: Best Practice Guidelines for Investigators.

Permalink

<https://escholarship.org/uc/item/0cq084nk>

Journal

Obstetrics and gynecology, 134(2)

ISSN

0029-7844

Authors

McLemore, Monica R
Asiodu, Ifeyinwa
Crear-Perry, Joia
et al.

Publication Date

2019-08-01

DOI

10.1097/aog.0000000000003393

Peer reviewed

S, editors. Cochrane handbook for systematic reviews of interventions. Version 5.1.0 (updated March 2011). London (UK): The Cochrane Collaboration; 2011.

5. Viele K, McGlothlin A, Broglio K. Interpretation of clinical trials that stopped early. *JAMA* 2016;315:1646–7.
6. Qin JB, Wang H, Sheng X, Xie Q, Gao S. Assisted reproductive technology and risk of adverse obstetric outcomes in dichorionic twin pregnancies: a systematic review and meta-analysis. *Fertil Steril* 2016;105:1180–92.

In Reply:

We thank Dr. Conde-Agudelo for his interest in our article in the March 2019 issue.¹ He raised some methodologic issues, which, according to his view, could compromise the study's internal and external validity. We indeed used block randomization to achieve balance in participants' allocation. The potential risk for selection bias can be reduced by using random block sizes and keeping the investigators blinded to each block size,² which was exactly what we did. In fact, our trial is graded as low risk of bias.³ Therefore, any imbalance in patients' characteristics is solely by chance.

Out of the 1,113 screened women in our trial, 1% had cervical length less than 20 mm and none had cervical length less than 18 mm. This is completely comparable with other studies.^{4,5} The most likely explanation for the difference with the Nicolaides et al⁶ trial is that they measured cervical length at 20 0/7–24 6/7 weeks of gestation compared with 16 0/7–22 0/7 weeks in our study and others.^{1,4,5}

Indeed, twin pregnancies conceived from in vitro fertilization are at slightly higher risk for complications. However, we do not think that the treatment response would be affected by conception method.

We agree that the reduced sample size, updated with a detailed protocol on ClinicalTrials.gov on August 24, 2017, affected our study power. However, this amendment was approved by the ethics committee and documented in the adjusted protocol in the early phase of the study, as well as being clearly addressed in the Discussion section of our article. We did plan an exploratory subgroup analysis in the original protocol, dated on September 25, 2015. The only post hoc analysis

performed was the composite of poor perinatal outcomes.

We appreciate all the above-mentioned comments from Dr. Conde-Agudelo. It is of interest to note that the study published by El-Refaie et al,⁷ with only non-in vitro fertilization twins and incorporated in an individual patient data meta-analysis⁸ of which Dr. Conde-Agudelo was a co-author, was registered retrospectively.

We believe that our trial has provided new evidence for readers on the prevention of preterm birth in women with twin pregnancies and short cervix.

Financial Disclosure: The authors did not report any potential conflicts of interest.

Vinh Q. Dang, MD

My Duc Hospital and
HOPE Research Center,
Ho Chi Minh City,
Viet Nam

Toan D. Pham, BSc

HOPE Research Center,
Ho Chi Minh City,
Viet Nam

Linh K. Nguyen, MD

My Duc Hospital,
Ho Chi Minh City,
Viet Nam

Ben W. Mol, MD, PhD

Department of
Obstetrics and Gynecology,
Monash University,
Melbourne, Australia

REFERENCES

1. Dang VQ, Nguyen LK, Pham TD, He YTN, Vu KN, Phan MTN, et al. Pessary compared with vaginal progesterone for the prevention of preterm birth in women with twin pregnancies and cervical length less than 38 mm: a randomized controlled trial. *Obstet Gynecol* 2019;133:459–67.
2. Efrid J. Blocked randomization with randomly selected block sizes. *Int J Environ Res Public Health* 2011;8:15–20.
3. Higgins JPT, Altman DG, Sterne JAC. Chapter 8: assessing risk of bias in included studies. In: Higgins JPT, Green S, editors. Cochrane handbook for systematic reviews of interventions. Version 5.1.0 (updated March 2011). London (UK): The Cochrane Collaboration; 2011.
4. Lim AC, Schuit E, Bloemenkamp K, Bernardus RE, Duvekot JJ, Erwich JJ, et al. 17 α -hydroxyprogesterone caproate for the prevention of adverse neonatal out-

come in multiple pregnancies: a randomized controlled trial. *Obstet Gynecol* 2011;118:513–20.

5. Liem S, Schuit E, Hegeman M, Bais J, de Boer K, Bloemenkamp K, et al. Cervical pessaries for prevention of preterm birth in women with a multiple pregnancy (ProTWIN): a multicentre, open-label randomised controlled trial. *Lancet* 2013;382:1341–9.
6. Nicolaides KH, Syngelaki A, Poon LC, de Paco Matallana C, Plasencia W, Molina FS, et al. Cervical pessary placement for prevention of preterm birth in unselected twin pregnancies: a randomized controlled trial. *Am J Obstet Gynecol* 2016;214:3.e1–9.
7. El-Refaie W, Abdelhafez MS, Badawy A. Vaginal progesterone for prevention of preterm labor in asymptomatic twin pregnancies with sonographic short cervix: a randomized clinical trial of efficacy and safety. *Arch Gynecol Obstet* 2016;293:61–7.
8. Romero R, Conde-Agudelo A, El-Refaie W, Rode L, Brizot ML, Cetingoz E, et al. Vaginal progesterone decreases preterm birth and neonatal morbidity and mortality in women with a twin gestation and a short cervix: an updated meta-analysis of individual patient data. *Ultra Obstet Gynecol* 2017;49:303–14.

Race, Research, and Women's Health: Best Practice Guidelines for Investigators

We, black women scholars, clinicians, and public health practitioners, read the article by Ghidei et al published in the April 2019 issue,¹ which outlines their considerations for the appropriate use of race in research on women's health for clinical investigators. We offer four suggestions for the authors and future investigators to consider when deciding whether, when, and how to include race in their research studies. First, we emphasize that black race is not a risk factor; racism is the risk factor for maternal morbidity, mortality, and poor birth outcomes.² Next, we suggest that investigators engage affected communities in exploring the construction of race and how it shows up in their lives.³ There are methods⁴ to facilitate this that can deeply influence how research is carried out and its effects on black people's lives. Third, partnering with critical race scholars, public health practitioners, and transdisciplinary teams is critical.⁵

Finally, we believe research should champion and reflect a diversified health care workforce. Each point is essential to women's health research. We look forward to more comprehensive examinations of racism in research and have cited additional sources that include black thought leaders and resources.

Financial Disclosure: *Dara D. Mendez disclosed receiving money paid to her institution from NIH/NHLBI and the University of Pittsburgh Center for Health Equity-pilot. Karen A. Scott disclosed receiving money paid to her from the California Maternal Quality Care Collaborative/Stanford University. The other authors did not report any potential conflicts of interest.*

Monica R. McLemore, PhD, RN
Ifeyinwa Asiodu, PhD, RN
 Family Health Care Nursing
 Department,
 University of California,
 San Francisco,
 San Francisco, California

Joia Crear-Perry, MD, FACOG
 National Birth Equity Collaborative,
 New Orleans, Louisiana

Dana Ain Davis, MPH, PhD
 Graduate Center,
 City University of New York,
 New York, New York

Michelle Drew, DNP, MPH
 Christiana Care Health Services
 OBGYN Residency Program,
 Wilmington, Delaware

Rachel R. Hardeman, PhD, MPH
 University of Minnesota,
 School of Public Health,
 Minneapolis, Minnesota

Dara D. Mendez, PhD, MPH
 University of Pittsburgh,
 Pittsburgh, Pennsylvania

Lynn Roberts, PhD
 City University of New York
 School of Public Health,
 New York, New York

Karen A. Scott, MD, MPH
 Family Health Care
 Nursing Department,
 University of California,
 San Francisco,
 San Francisco, California

REFERENCES

1. Ghidei L, Murray A, Singer J. Race, research, and women's health best prac-

tice guidelines for investigators. *Obstet Gynecol* 2019;133:815–18.

2. Black Mamas Matter Alliance. Setting the standard for holistic care of and for Black women. Available at: http://blackmamas-matter.org/wp-content/uploads/2018/04/BMMA_BlackPaper_April-2018.pdf. Retrieved May 6, 2019.
3. Ford CL, Airhihenbuwa CO. The public health critical race methodology: praxis for antiracism research. *Soc Sci Med* 2010;71:1390–8.
4. Franck LS, McLemore MR, Cooper N, De Castro B, Gordon AY, Williams S, et al. A novel method for involving women of color at high risk for preterm birth in research priority setting. *J Vis Exp* 2018.
5. Taylor J, Novoa C, Hamm K, Phadke S. Eliminating racial disparities in maternal and infant mortality: a comprehensive policy blueprint. Available at: <https://www.americanprogress.org/issues/women/reports/2019/05/02/469186/eliminating-racial-disparities-maternal-infant-mortality/>. Retrieved May 6, 2019.

In Reply:

Thank you for your letter. We wrote our article¹ in the hopes of generating more discussion surrounding race, research, and women's health. We absolutely agree with the important suggestions you make. We hope the readership of *Obstetrics & Gynecology* continues to contribute to this topic as you have, and we hope that *Obstetrics & Gynecology* and other journals will convene a group to create formal guidelines regarding the appropriate use of race in research.

Financial Disclosure: *The authors did not report any potential conflicts of interest.*

Luwam Ghidei, MD
 Warren Alpert Medical School of
 Brown University,
 Women & Infants Hospital of Rhode
 Island, Providence, Rhode Island

Anne Murray, MD
 Warren Alpert Medical School of
 Brown University,
 Women & Infants Hospital of Rhode
 Island and Providence Community
 Health Centers,
 Providence, Rhode Island

Janet Singer, MSN, CNM
 Warren Alpert Medical School of
 Brown University,
 Women & Infants Hospital of Rhode
 Island, Providence, Rhode Island

REFERENCE

1. Ghidei L, Murray A, Singer J. Race, research, and women's health: best practice guidelines for investigators. *Obstet Gynecol* 2019;133:815–8.

Targeted Treatment of a Rare Vaginal Sarcoma With an Anaplastic Lymphoma Kinase Inhibitor

We wish to update the case report by Forde and Tewari¹ from the February 2016 issue. This was a case of a 34-year-old woman with a history of an epithelioid inflammatory myofibroblastic vaginal sarcoma that recurred 2 months after initial surgical excision. The patient had declined repeat excision and had progression of disease with radiation therapy. Tumor testing identified anaplastic lymphoma kinase gene rearrangements, and the patient was placed on crizotinib 250 mg orally twice daily for 20 weeks. The patient had rapid resolution of her vaginal tumor. A few years after completing treatment, she became pregnant and delivered a healthy neonate. At last follow-up, she remains without evidence of disease for 164 weeks.

This case report is just one of many great examples of how precision medicine has changed management of difficult and rare tumors. Nagumo et al and Butrynski et al have reported cases that show drastic improvement in tumor size with use of crizotinib in patients with anaplastic lymphoma kinase rearrangement.^{2,3} Studies such as these have led the National Comprehensive Cancer Network to include crizotinib as standard treatment for patients with inflammatory myofibroblastic sarcomas and anaplastic lymphoma kinase rearrangements.⁴

Given the longevity of our patient and the success of others, we strongly encourage health care providers to perform fluorescence in situ hybridization testing of rare inflammatory myofibroblastic sarcomas to assess for anaplastic lymphoma kinase rearrangements. We hope to encourage other health care providers who treat rare tumor types to continue to advance the field and provide personalized medicine for our patients.

Financial Disclosure: *Nolan Kinne disclosed that White Memorial Medical Center–Adventist Health is his employer for residency. The other*