Lawrence Berkeley National Laboratory

Recent Work

Title

Contradictory effects of silver nanoparticles on activated sludge wastewater treatment.

Permalink

<https://escholarship.org/uc/item/0cw9w0r5>

Authors

Sheng, Zhiya
Van Nostrand, Joy D
Zhou, Jizhong
et al.

Publication Date

2018

DOI

10.1016/j.jhazmat.2017.07.051

Peer reviewed

Contradictory effects of silver nanoparticles on activated sludge wastewater treatment

Author links open overlay panel [ZhiyaSheng^a](#) [Joy D. Van Nostrand^b](#) [JizhongZhou^b](#) [YangLiu^a](#)
Show more

<https://doi.org/10.1016/j.jhazmat.2017.07.051> Get rights and content

Highlights

- Ag-NPs, especially freshly prepared, can have positive effects.
- Ag-NPs can help to maintain [microbial community](#) diversity in [activated sludge](#).
- Improved [sludge](#) settleability can be important in the positive effects observed.
- The hormesis model may need to be considered for the [toxicology](#) of Ag-NPs.

Abstract

Increased amount of nano-silver will be released into domestic and industrial waste streams due to its extensive application. However, great controversy still exists on the effects of silver nanoparticle (Ag-NP) on biological wastewater treatment processes and a toxicology model has not been built yet. Four sequencing batch reactors with activated sludge has been run for over three months with different silver species at a concentration of 1 mg Ag/L in influent. Both freshly prepared Ag-NPs and aged Ag-NPs were tested with released silver ion as control. Results in this study showed that Ag-NPs, especially freshly prepared Ag-NPs, can help to maintain or even increase the diversity of microbial community in activated sludge and the biomass concentration even under long-term treatment. It indicates that the hormesis model need to be considered for the toxicology of Ag-NPs.

- [Previous article](#)
- [Next article](#)

Keywords

Silver nanoparticles (Ag-NPs)

Activated sludge

Microbial diversity

Functional diversity

Geochip

Pyrosequencing

1. Introduction

Nano-silver is inevitably released into domestic and [industrial waste](#) streams as it is one of the most commonly used [nano materials](#) in consumer products [1]. Considerable attention has been paid to the potential adverse effects on [biological wastewater treatment](#) system (BWTS) due to the antimicrobial properties of [silver nanoparticles](#) (Ag-NPs). A general conclusion can be made from previous research that the effects of Ag-NPs depend on the dose and time period applied as well as the property of Ag-NPs and the system Ag-NPs are applied to. However, great controversy still exists on how each of these parameters affects the impacts of Ag-NPs, and a sophisticated [toxicology](#) model has not been built at all. Previous research covers only a tip of the iceberg of all possible combination of these parameters. Not to mention that the mechanisms behind the phenomena are poorly understood.

Higher concentration of Ag-NPs often results in more significant adverse effects [2], [3], [4], [5]. Hormetic effects under sublethal concentration have been reported occasionally but stayed as a marginalized concept [6], [7], [8], [9], [10]. Properties of Ag-NPs that affect its toxicity include [nanoparticle](#) size, shape and coating. Smaller Ag-NPs tend to be more toxic [11], [12], [13]. Spherical Ag-NPs and polyvinylpyrrolidone (PVP) coating tend to have weaker bactericidal action [14], [15]. However, the effects of shape and coating have not been well-studied yet. Great controversy also exists on if the toxicity of Ag-NPs come from the released Ag⁺ ion or the nanoparticle form itself [3], [7], [14], [16], [17], [18].

Properties of the BWTS are even more complicated. The effects of Ag-NPs on [activated sludge](#) and [biofilm](#) in BWTS have been studied previously [19]. Ordered from the most resistant to Ag-NPs to the least, the [microbial communities](#) in BWTS include biofilm/activated [sludge](#), planktonic mixed culture and pure culture of single [strains](#) [2], [9], [20], [21]. Potential ligands in BWTS can bind with Ag-NPs or released Ag⁺ ions and lower the dissolution of Ag-NPs and their bactericidal effects [22], [23], [24], although some [anions](#) may accelerate Ag-NP dissolution [25]. These ligands range from organic matter such as [dissolved organic carbon](#) to inorganic ions such as chloride and [sulfide](#). It has been reported that [ammonia](#) oxidizing bacteria (AOB) are more vulnerable towards Ag-NPs treatment, as compared to [nitrite](#) oxidizing bacteria (NOB) and organic oxidation [heterotrophs](#) [20], [26], [27], [28]. More recent

studies tend to focus on [long-term effects](#) of Ag-NPs under conditions mimicking the real-world conditions in BWTs [\[21\]](#), [\[29\]](#), [\[30\]](#). Acute inhibition is often observed at the beginning of Ag-NP addition, but the system usually recovers in the long term [\[4\]](#), [\[5\]](#). For instance, in a membrane [bioreactor](#) activated sludge system with 0.1 mg/L Ag-NPs in the reactor influent, the silver resistance gene (*silE*) increased at the beginning of the Ag-NPs addition and then decreased to the initial level; and the reactor performance was not significantly affected by Ag-NPs [\[31\]](#). In a separate study, with the addition of 1 and 5 mg/L Ag-NPs in bioreactors, phosphorus removal decreased and the microbial community changed at the beginning of the study but then stabilized with persistent exposure [\[30\]](#). The adverse effects of Ag-NPs are minimal especially when sulfidation plays an important role in most of the BWTs [\[32\]](#), [\[33\]](#), [\[34\]](#), [\[35\]](#), [\[36\]](#), [\[37\]](#). A toxicological model to estimate the effects of Ag-NPs is beginning to take shape. This raises the question: should the hormesis model be considered here?

This study examines the response of the microbial community to a potentially “least-toxic” combination, which is the case in most of our current BWTs in practical operation: low dose, long-term, spherical Ag-NPs with PVP coating in activated sludge bioreactors fed with synthetic [municipal wastewater](#). No significant effects were seen on [pollutants removal](#). However, interestingly, Ag-NPs helped to maintain the microbial community diversity in the activated sludge. 16S rRNA gene based pyrosequencing was used to monitor the bacterial community and GeoChip was used to directly examine the functional diversity of the microbial community. Properties of the sludge, accumulation of silver species inside the sludge and characteristics of the Ag-NPs were examined to explain this phenomenon.

2. Material and methods

2.1. Reactor setup

Four [sequencing batch reactors](#) were operated for over three months. The total volume of the reactors was 1 L and the effective volume was 700 mL. The reactors were run on a 12 hour cycle (5 min of influent filling, 11 h of [aeration](#), 30 min settling, 5 min [effluent](#) withdraw and 20 min idle). Hydraulic retention time was 24 h. [Sludge](#) was wasted through effluent withdraw by gravity. Solids retention time was monitored but not controlled. Solids retention time (SRT) was 17 days at the steady state for all four reactors. However, after the addition of silver species started, almost no sludge was wasted from the reactor with fresh Ag-NPs added, indicating that SRT was dramatically increased after the addition of fresh Ag-NPs. The reactor feed was prepared according

to Alito and Gunsch [5] and contains an average COD of 450 mg/L and ammonia of 40 mg/L with pH adjusted to 7.3 ± 0.5 .

2.2. Silver species addition

Self-dispersing silver nanopowder was purchased from SkySpring Nanomaterials, Inc. (Houston, USA). According to the Ag-NP product description, the particle size is less than 15 nm, and the particle composition is 25% silver (99.99% purity) and 75% polyvinylpyrrolidone (PVP), similar to Ag-NPs commonly used in commercial products. Fresh and aged Ag-NP suspensions were examined by transmission electron microscopy (TEM) according to the method described in previous studies [38]. Spherical aggregates of Ag-NPs were observed. The particle size and zeta potential of Ag-NPs were characterized using a Malvern Zetasizer Nano-ZS (Model: ZEN3600, Malvern Instruments Ltd, Worcestershire, UK). Ag⁺ ion dissolution was characterized as described in Section 2.5. PVP and silver species addition started after the reactor reached steady state for over two weeks (27 days after start-up). PVP, aged Ag-NPs, fresh Ag-NPs and Ag⁺ ion released from fresh Ag-NPs were added to each of the four reactors respectively. Aged and fresh Ag-NPs were added at a concentration of 1 mg Ag/L in influent. This concentration resulted in 0.5 mg Ag/L in the reactor, which falls within a representative range [39] while it was high enough to see significant effects based on previous tests (data not shown). PVP was added to the control reactor at the concentration of 3 mg/L which is the same as in reactors with Ag-NP addition. Aged Ag-NPs stock suspension was prepared when reactors were started up and kept at 4 °C in dark and was added into the influent tank and kept under room temperature in dark for one week. Fresh Ag-NPs suspension (3.5 g Ag L⁻¹) was prepared everyday and 0.1 mL suspension was spiked into the reactor during influent filling in each cycle, producing a concentration equalled to 1 mg Ag/L in influent. For the fourth reactor, to test Ag⁺ ion released from fresh Ag-NPs, 0.1 mL of the freshly prepared Ag-NP suspension was added into a dialysis unit (Slide-A-Lyzer™ MINI Dialysis Device, 2 K MWCO, 0.1 mL, Thermo Scientific, USA) and the dialysis unit was put into the reactor during influent filling in each cycle and float in the activated sludge for 12 h before changing to a new one. Equal amount of PVP (3 mg PVP L⁻¹) was added into the control reactor. The tests of Ag-NP and PVP addition have been performed for over two months. Similar operation of reactors and Ag-NP addition were repeated for several times.

2.3. Reactor performance monitoring

Effluent quality was monitored in terms of COD and [ammonium](#) removal using Hach methods 8000 and 10205 [40]. SVI and MLSS were measured according to the standard methods [41]. [Reaction kinetics](#) of COD and ammonium removal and nitrate production was also performed using the substrate depletion method [42]. Mixed liquor samples were collected at 15, 30, 35, 60, 90, 120, 150, 180, 210, 240, 300, 480 and 660 min, centrifuged at 3000 g for 10 min at 4 °C, filtered (0.45 µm) and analyzed for COD, ammonium and nitrate. Nitrate was measured using [ion chromatography](#) (IC).

2.4. Microbial community analysis

Activated sludge samples were collected in duplicates and [genomic DNA](#) was extracted using a Powersoil® DNA Isolation Kit from MO BIO Laboratories, Inc. (Carlsbad, USA). DNA was analyzed with pyrosequencing and GeoChip.

Paired-end [sequencing](#) based on the 16S rRNA gene was performed at the Research and Testing Laboratory (Lubbock, TX, USA), using the Illumina MiSeq platform [43]. Primers 28F (5'-GAGTTTGATCNTGGCTCAG-3') and 519R (5'-GTNTTACNGCGGCKGCTG-3') were used, which covered V1–V3 hypervariable regions [44]. [Chimeras](#) and poor quality sequences were removed from the denoised sequence reads. The remaining sequences were clustered into operational taxonomic units (OTUs) with 0% divergence using USEARCH. Taxonomic information was assigned to OTUs based on a database of high quality sequences derived from the NCBI using a distributed. NET algorithm that utilizes BLASTN+ (Kraken BLAST, www.krakenblast.com). A principal coordinates analysis (PCoA) of [microbial community](#) diversity was performed using the QIIME pipeline (<http://qiime.org>) with the beta diversity metrics of weighted unifracs [45]. The canonical [correspondence analysis](#) (CCA) was carried out using the XLSTAT software version 2016 to evaluate the correlation between microbial communities and silver species addition. Ag-NP treatment was converted into four quantitative variables: silver accumulation, silver added as Ag⁺ ion, silver added in the [nanoparticle](#) form, and PVP added.

DNA (1 µg) was labeled with Cy3 and hybridized to the GeoChip 5 microarray synthesized by NimbleGen (Madison, WI, USA) and processed as previously described by Lu et al. [46]. The [signal-to-noise ratio](#) threshold for a spot to be considered positive was ≥2 as described previously [47]. Detrended correspondence analysis (DCA) was performed based on sample scores. Hierarchical clustering analysis based on the Bray-Curtis dissimilarity indices was performed and a corresponding heatmap was built with the top 30 abundant functional genes in each sample.

2.5. Silver accumulation and release and nanoparticle characterization

Silver species accumulation and release was measured with [inductively coupled plasma mass spectrometry](#) (ICP-MS) using the ELAN 9000 ICP [mass spectrometer](#) (PerKinElmer, Canada). Samples were analyzed directly to measure dissolved Ag⁺ ions and digested to measure total silver. Microwave digestion was performed as described by Wu *et al.* [48] and briefly summarized here. 10 mL concentrated [nitric acid](#) and 2 mL ultra-pure water were added to 1 g [biofilm](#) (wet weight) or 1 mL suspension and kept at room temperature overnight for pre-digestion. Microwave digestion was then carried out using ETHOS EZ Microwave [Solvent Extraction](#) Labstation (Milestone Inc., USA) with the following heating program: heat to 190 °C within 15 min and then hold at 190 °C for 10 min. The particle size and zeta potential of Ag-NPs was characterized using a Malvern Zetasizer Nano-ZS (Model: ZEN3600, Malvern Instruments Ltd, Worcestershire, UK). Since PVP dissolves in water completely, parameters of silver were adopted for the analysis: the [refractive index](#) was 2.0 and the absorption coefficient was 0.320 [49].

3. Results

3.1. Effects of Ag-NPs on reactor performance

No significant change in [pollutant removal](#) was observed after Ag-NP addition in any of the four reactors ([Fig. 1A](#) and B). COD removal rate was maintained at above 90% and [ammonium](#) removal was above 99% in each reactor all the time. [Fig. 1C, D](#) and E show the COD, ammonium and nitrate concentration change against time in four reactors on Day 63. Only slight difference in [reaction kinetics](#) has been observed. The majority of COD was removed from each reactor within the first 30 min and ammonium removal was completed within 4 h. The reaction rate constant for COD removal remained almost the same in all reactors (p -value > 0.05) after two months of Ag-NP addition. Nevertheless, minor difference can be seen in ammonium removal kinetics. The rate constant for ammonium removal decreased to 0.35 h⁻¹ in the PVP control and 0.34 h⁻¹ in the reactor with aged Ag-NP addition (p -value = 0.03), compared to the initial value of 0.55 h⁻¹ before Ag-NP addition. However, this rate constant increased to 0.66 h⁻¹ in the reactor fed with fresh Ag-NPs (p -value = 0.11). The rate constant decreased slightly to 0.48 h⁻¹ in reactor fed with Ag⁺ ion released from fresh Ag-NPs (p -value = 0.18). Fresh Ag-NP helped to maintain the ammonium oxidizing reaction rate in the reactors. However, it should be noted that if the biomass concentration is taken into account, the ammonium uptake rate decreased in all reactors, compared with the initial

values (p -value < 0.05). The initial ammonium uptake rate was $3.0 \text{ mg N g}^{-1}\text{VSS}^{-1}$; the ammonium uptake rate decreased to 2.2, 2.0, 1.3, $2.3 \text{ mg N g}^{-1}\text{VSS h}^{-1}$ in reactor fed with PVP, Aged Ag-NPs, fresh Ag-NPs and Ag^+ ion, respectively. Similarly, the reactor fed with fresh Ag-NPs also had the lowest COD uptake rate. This indicates that the metabolic activity of microbes in the reactor fed with fresh Ag-NPs didn't increase proportionally to the biomass concentration. As a matter of fact, the ammonium oxidizing reaction rate constant in the reactor fed with fresh Ag-NPs stayed the highest (above 0.6 h^{-1}) among all the reactors during the two-month treatment period. The ammonium uptake rate in the reactor fed with fresh Ag-NPs was also the highest ($2.5 \text{ mg N g}^{-1} \text{ VSS h}^{-1}$) in all four reactors on Day 22, before the significant biomass increase occurred. It appeared that [nutrient uptake](#) rates in the reactor fed with fresh Ag-NPs hit a bottleneck when the biomass and silver accumulation increased significantly.

1. [Download high-res image \(392KB\)](#)
2. [Download full-size image](#)

Fig. 1. Performance of each reactor. (A) Effluent COD concentration; (B) [Effluent ammonium](#) concentration; (C) COD removal kinetics; (D) Ammonium removal kinetics; (E) Nitrate production kinetics.

3.2. Ag-NPs improved sludge settleability and increased biomass concentration

The biomass concentration in the reactor with fresh Ag-NPs increased significantly after two months of operation as indicated by mixed liquor suspended solids (MLSS) concentration in [Fig. 2A](#). MLSS concentration reached the peak of 4866 mg/L on Day 56 in the reactor with fresh Ag-NPs while stabilized at about 2400 mg/L in the others. Meanwhile, the [sludge](#) volume index (SVI) in this reactor is much lower than the others as well ([Fig. 2B](#)). SVI stabilized at about 100 mL/g in the reactor with fresh Ag-NPs and stayed below 220 mL/g in reactors fed with aged Ag-NPs and Ag⁺ released from fresh Ag-NPs. However, SVI increased to 295.6 mL/g in PVP control and remained above 240 mL/g until the end of the test. Big chunks of sludge can be seen by naked eye in the PVP control (irregular shape with the longest side over 2 cm). Sludge flocs may be physically bridged by of the PVP [polymer](#) and therefore they form chunks that are very light and fluffy, which makes them difficult to settle down. This is not hard to explain since PVP can help Ag-NPs to disperse in water using the same mechanism. Sludge flocs are bigger in the reactor with fresh Ag-NPs. All the reactors have one peak with chord length below 5 μm , while the reactor with fresh Ag-NPs has another major peak at about 20 μm ([Fig. 2C](#)). The density of sludge flocs in this reactor is also slightly larger than those in the other reactors (Fig. S1). More extracellular polymeric substances (EPS) were produced by the PVP control, especially protein ([Fig. 2D](#)). This also contributes to the poor settleability of the sludge in the PVP control, since excess EPS will result in sludge bulking [\[50\]](#).

1. [Download high-res image \(487KB\)](#)
2. [Download full-size image](#)

Fig. 2. [Sludge](#) property in each reactor. (A) MLSS concentration; (B) SVI; (C) Floc size on day 64; (D) Sludge EPS concentration on day 64.

3.3. Ag-NPs helped to maintain compositional diversity of the microbial community

Bacteria families vary in each sample as shown in [Fig. 3A](#). After the thirty-day startup stage (sample S1), the [microbial community](#) composition changed slightly from the initial [inoculum](#) (sample S0). More significant shifts in community structure can be seen in the samples at the end of the test (S4-S7). Based on the principal coordinate analysis (PCoA) with weighted unifrac ([Fig. 3B](#)), along the axis accounts for 72% of variance samples S0 and S1 are close to each other while samples S4-S7 are clustered together. This divergence from the initial inoculum is very likely caused by operation time period in the lab since laboratory condition is different from that in the full-scale [wastewater treatment plant](#), which is confirmed by the canonical [correspondence analysis](#)(CCA) as shown in [Fig. 3C](#). If looking closer into axes PC2 and PC3 ([Fig. 3C](#)), it can be seen that sample S6 (reactor with fresh Ag-NPs, day 64) locates the closest to the initial inoculum S0. S6 is also close to sample S1, which is the initial state before Ag-NP addition for all reactors. This indicates that fresh Ag-NP may help to maintain the microbial community.

1. [Download high-res image \(1MB\)](#)
2. [Download full-size image](#)

Fig. 3. Microbial diversity analysis. (A) [Relative abundance](#) of bacteria families; (B) principal coordinate analysis (PCoA) of [microbial community](#) diversity; (C) Canonical [correspondence analysis](#) (CCA) triplot of dominant bacteria families and test conditions; (D) The number of genes detected in each functional category; (E) Detrended correspondence analysis of functional gene diversity; (F) Heatmap of the top 30 abundant functional genes; Sample S0: The initial [activated sludge inoculum](#); S1: [Sludge](#) after the thirty-day startup stage; S2: Sludge in reactor with aged Ag-NPs, day 8; S3: Sludge in reactor with fresh Ag-NPs, day 14; S4: Sludge in reactor with only PVP as control, day 64; S5: Sludge in reactor with aged Ag-NPs, day 64; S6: Sludge in reactor with fresh Ag-NPs, day 64; S7: Sludge in reactor with Ag⁺ released from fresh Ag-NPs, day 64.

CCA triplot in [Fig. 3C](#) illustrated the significant correlations ($p = 0.021$) between dominant bacteria families (accounting for over 70% sequence abundance in each sample) and the test conditions: operation time, silver accumulation and silver added as the [nanoparticle](#) form (NP added). Correlation of dominant bacteria families with silver added as Ag⁺ ion was less significant and no significant correlation with PVP addition was seen in CCA. The 'Time' vector almost overlap with the F1 axes and the 'NP added' vector almost overlap with the F2 axes. Based on the CCA silver accumulation was positively associated with time and NP added, while time and NP added were almost orthogonal. The two axes F1 and F2 combined explained 92% of the variances in microbial community, indicating that time, silver added as the NP form and silver accumulation are the major factors shaping the microbial community. The family *Xanthomonadaceae* (F14) located close to the origin of the plot, indicating that this family has little response to the test conditions. Dominant families *Comamonadaceae* and *Rhodocyclaceae* (F9 and F10) fell near the F1 axes, indicating that time is the major factor that resulted in the changes of these two families. F2 (a family in the order of *Cytophagales*) and F8 (*Sphingomonadaceae*) increased with silver accumulation. *Cytophagales* has been reported to dominant in certain conditions with high metal concentration and may potentially be used in [phytoremediation](#) [51]. *Sphingomonadaceae* is a family of bacteria with sphingolipids in the outer membrane. Sphingolipids may potentially work as signaling molecules, although the endogenous functions of sphingolipids in bacteria are still unknown [52]. This signaling pathway may likely be associated with the synergetic effects of maintaining community diversity found in this study.

In terms of [diversity indices](#) (richness and evenness, [Table 1](#)), both richness and evenness were lost at the beginning of the Ag-NP addition (S2, reactor with aged Ag-NPs, day 8; S3, reactor with fresh Ag-NPs, day 14). However, the loss of evenness recovered with time. After two months of Ag-NP addition, fresh Ag-NP (S6) maintained the highest evenness without relatively less loss in richness compared with the initial inoculum (S0).

Table 1. [Diversity index](#) of [microbial community](#) and functional genes.

Sample	Microbial community (bacteria families)		Functional genes	
	Richness	Evenness	Richness	Evenness
S0	126	0.554	–	–
S1	122	0.407	904	0.796
S2	104	0.381	813	0.802
S3	93	0.296	906	0.797
S4	88	0.538	888	0.796
S5	86	0.558	891	0.796
S6	97	0.626	918	0.796
S7	110	0.552	915	0.795

3.4. Ag-NPs helped to maintain functional diversity of the microbial community

In terms of functional diversity, reactor with fresh Ag-NPs (S6) has the highest number of genes detected ([Fig. 3D](#)) and richness ([Table 1](#)). Evenness is almost identical for all samples for functional genes, which is consistent with the fact that there is no significant change in gene distribution in each function category. Detrended correspondence analysis (DCA) showed that ([Fig. 3E](#)) samples are clustered by time period along DCA1 and samples with Ag-NP addition (S5-7) are closer to the initial state (S1) compared to the PVP control (S4). This indicates that Ag-NP addition helped to maintain the functional diversity of the microbial community in the reactors. Fresh Ag-NP even increased functional diversity in terms of richness. Heatmap of the top 30 abundant functional genes also indicated that S6 (reactor with fresh Ag-NPs, day 64) and S7 (reactor with Ag⁺ released from fresh Ag-NPs, day 64) are clustered with the initial state S1 ([Fig. 3F](#)). Some [antioxidant enzyme](#) and metal and stress resistant genes has been elevated by fresh Ag-NPs (S6). More significant increase in these genes was seen in sample S5 (reactor with aged Ag-NPs, day 64), which makes sense since aged Ag-NPs release more Ag⁺ ions and Ag⁺ ion plays an important role in the toxicity of Ag-NPs [\[3\]](#), [\[53\]](#), [\[54\]](#), [\[55\]](#). Particle size and [zeta potential](#) of aged and fresh Ag-NPs are at

the same scale as shown in [Table 2](#). The data also confirmed that much more Ag⁺ ion was released from Aged Ag-NPs.

Table 2. Particle characterization and silver dissolution.

	Particle size (nm)	Zeta potential in synthetic wastewater (mV)	Silver ion concentration in influent (mg/L)
Aged Ag-NP	74.2 ± 5.1	-9.37 ± 1.50	0.70 ± 0.10
Fresh Ag-NP	67.9 ± 1.0	-4.17 ± 1.03	0.10 ± 0.02

3.5. Silver species accumulation and release

Silver concentration in [effluent](#) and sludge was monitored. Dissolved Ag⁺ [ion concentration](#) in the effluent in all reactors with Ag-NP addition remained below 0.01 mg/L all the time ([Fig. 4A](#)). Total silver release with effluent was the highest (0.017 mg/L) on the first day of Ag-NP addition in the reactor with Ag⁺ ion released from fresh Ag-NPs. The reactor adapted to the silver addition within two weeks and total silver release never exceeded 0.01 mg/L thereafter. Total silver release increased with time in the reactors fed with fresh and aged Ag-NP and reached 0.76 and 0.89 mg/L respectively at the end of the test. The only difference is the release increased in steps in the reactor with fresh Ag-NPs, indicating that this reactor has more capacity to maintain Ag-NPs in the sludge, which is consistent with the total silver accumulation in sludge ([Fig. 4B](#)). Silver accumulation in sludge appeared to be periodical to some extent which is coupled with biomass concentration change. When the accumulation reached a threshold and the cells started to die, more silver is released through effluent and after the concentration in sludge decreased after the release, cell growth recovered and a new round of accumulation starts. More biomass was maintained in the reactor with fresh Ag-NPs and more silver was accumulated in the sludge in that reactor.

1. [Download high-res image \(260KB\)](#)
2. [Download full-size image](#)

Fig. 4. Silver species accumulation and release. (A) Silver release via [effluent](#); (B) Silver accumulation in [activated sludge](#). Silver concentrations in effluent and [sludge](#) from the PVP control reactor were both below the detection limit.

4. Discussion

There is ample evidence that bacteria growth can be stimulated when sublethal concentration of antimicrobial agents are applied [56], [57]. An inverted U-shaped [dose-response relationship](#), so called hormesis model, have been reported for many [antibiotics](#) [58]. Most of these research were done with pure-cultured bacteria. Similar dose response has also been seen in studies with Ag-NPs [6], [7], [8], [9]. However, controversy exists and these findings did not attract a fair amount of attention. Majority of literature reports non-significant to adverse effects of Ag-NP when tested at the mg/L scale [3], [18], [59], [60], [61]. In this study, increase in biomass concentration was clearly observed in reactors with fresh Ag-NPs and it was stably repeated in several batch of tests under similar Ag-NP concentrations. Results in this study and literature indicate that the effect of Ag-NP may conform to the hormesis model as well. When tested with pure-cultured bacteria, the hormetic effects were less significant and difficult to replicate. However, when tested with sophisticated bacteria community such

as [activated sludge](#) used in this study, the positive effects appeared to be evident and stable. The maintenance of community diversity played a very important role in this hormetic effect observed here. The increased [microbial community](#) diversity by fresh Ag-NP is the most significant phenomenon observed besides the increase in biomass concentration. Acute effects at the beginning of addition come from loss of both richness and evenness. This is also consistent with the hormesis model where initial decrease in growth was followed by the adaptive rebound response [\[58\]](#). More experiments with systematically designed dosages and treatment time need to be carried out to confirm the hormesis model in Ag-NP [toxicology](#).

In this study, better maintenance of [sludge](#) in the reactor may play a more important role other than stimulation of growth. In the reactor with fresh Ag-NPs, increased floc size and density led to better settleability and therefore less sludge was lost through [effluent](#) withdraw and more biomass was accumulated in the reactor. In addition, higher diversity of the microbial community makes the microbes more resistant to stress [\[62\]](#), [\[63\]](#). As a result, the increased survival contributes to the higher biomass concentration. It has been reported that nano zero-valent iron can be used to control sludge bulking [\[64\]](#), and Ag-NPs can be another option. This is also supported by the fact that [nutrient uptake](#) rates didn't increase with biomass concentration in the reactor fed with fresh Ag-NPs. More cells were maintained in the reactor but not all of them are metabolically active or they may not be in their most active state. However, it should be noted that the lack of sufficient nutrients in the reactor might also contribute to the bottleneck in nutrient uptake rate increase.

Without considering time, the microbial community structure in samples with fresh and aged Ag-NPs closer resembled the initial [inoculum](#), compared to the sample with only Ag⁺ ion released from fresh Ag-NPs. This indicated that the [nanoparticle](#) form may play a more important role in the maintenance of microbial community diversity. In addition, Ag⁺ ion doesn't improve sludge settling and increase biomass concentration in the reactor. However, in terms of reactor performance, especially [ammonium](#) removal kinetics, the reactor with Ag⁺ ion performed more similar to the reactor with fresh Ag-NPs instead of aged Ag-NPs. This indicates, to some extent, that the released Ag⁺ ion played an important role in maintaining the [pollutant removal](#) capacity. To better understand the mechanism behind the positive effects of Ag-NPs, tests with equal Ag⁺ [ion concentration](#) but different nanoparticle concentration can be run in the future, along with additional control without treatment to verify the effects of operation time period. Longer startup stage can also be considered. High influent ammonium concentration may help to test if Ag-NP can improve high-ammonium concentration removal.

Hormetic effects can be triggered by many kinds of toxic substances, such as radiation and chemical reagents including antibiotics. The term “hormesis” was initially used to describe effects caused by low [dose radiation](#) and is now generally used to describe the inverted U-shape biological dose-response to stress [\[56\]](#), [\[65\]](#). Reports on the hormetic dose-response to antibiotics date back to the 1950s and various antibiotics were found to be able to cause hormetic effects [\[56\]](#), [\[57\]](#). Low-intensity pulsed [ultrasound](#) (LIPUS) was also found to stimulate cell growth and [antibody](#) production, although high dose of ultrasound can kill cells [\[66\]](#). Agents that can cause hormetic effects can be classified into two types: 1) chemical toxins that can leave residuals and accumulate in cells, such as Ag-NP and other antibiotics; 2) agents that can physically change cells and leave no residuals, such as LIPUS. Radiation is more complicated and can cause both physical and [chemical effects](#). Ag-NPs are chemically toxic to cells and can be accumulated in or near cells, therefore, the positive effects caused by Ag-NPs can be limited. This is consistent with the result that biomass concentration started to decrease in the reactor fed with fresh Ag-NPs at the end of the test, which may result from the high concentration of silver species accumulated in the sludge. Longer term of operation will help to verify the effects of silver accumulation. This is also consistent with the result that nutrient uptake didn't increase with biomass concentration in the reactor fed with fresh Ag-NPs. The toxicity of Ag-NPs may induce a dormancy state of the cells therefore no enhanced metabolic activity was observed although more cells were maintained in the reactor. LIPUS work with a different mechanism as it increase cell permeability which lead to better circulation, faster cell metabolism and enhanced antibody secretion. Because no residual is left by LIPUS, repeated stimulation with LIPUS can cause more significant beneficial effects than Ag-NP. Substances that can cause hormetic effects can also be classified as non-selective and selective. LIPUS can work on many kinds of cells and is non-selective. Ag-NP, which can be taken as a broad-spectrum antibiotics, is also relatively non-selective. Therefore, agents such as Ag-NP and LIPUS tend to work equally on various kinds of cells. This explains why Ag-NP increased the diversity of microbial community in activated sludge without significant changes in the distribution of genes in each functional group. On the contrary, selective agents such as narrow-spectrum antibiotics, are more inclined to cause selective effects such as changes in dominant species in microbial community. As reported in literature, changes in [gene expression](#) caused by subinhibitory concentration of broad-spectrum antibiotics are often termed with “enhance”, “increase” or “stimulation”; effects caused by subinhibitory concentration of narrow-spectrum antibiotics are often described by the word “inhibited” or “reduced” [\[56\]](#). In regard to a complicated ecosystem such as activated sludge

in [wastewater treatment](#), it is easier for non-selective agents with no residuals (such as LIPUS) to cause significant beneficial effects; non-selective agents with residuals (such as Ag-NP) ranks the second, but there are limitations caused by the toxicity and accumulation; it is relatively difficult for selective agents with residuals (such as narrow-spectrum antibiotics) to cause significant beneficial effects on such a complicated microbial community.

Acknowledgments

This work was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), and the Alberta Innovates Graduate Student Scholarship (PhD in Nanotechnology).

References

[1]

Consumer products inventory

Project-on-Emerging-Nanotechnologies (2014)

<http://www.nanotechproject.org/cpi/about/analysis/> (Accessed 03.07. 2017)

[2]

J.S. Kim, E. Kuk, K.N. Yu, J.H. Kim, S.J. Park, H.J. Lee, S.H. Kim, Y.K. Park, Y.H. Park, C.Y. Hwang, Y.K. Kim, Y.S. Lee, D.H. Jeong, M.H. Cho **Antimicrobial effects of silver nanoparticles** Nanomedicine, 3 (2007), pp. 95-101

[ArticleDownload PDFView Record in Scopus](#)

[3]

J.H. Priester, A. Singhal, B. Wu, G.D. Stucky, P.A. Holden **Integrated approach to evaluating the toxicity of novel cysteine-capped silver nanoparticles to *Escherichia coli* and *Pseudomonas aeruginosa***

Analyst, 139 (2014), pp. 954-963

[CrossRefView Record in Scopus](#)

[4]

H. Chen, X. Zheng, Y. Chen, H. Mu **Long-term performance of enhanced biological phosphorus removal with increasing concentrations of silver nanoparticles and ions**

RSC Adv., 3 (2013), pp. 9835-9842

[CrossRefView Record in Scopus](#)

[5]

C.L. Alito, C.K. Gunsch **Assessing the effects of silver nanoparticles on biological nutrient removal in bench-scale activated sludge sequencing batch reactors**

Environ. Sci. Technol., 48 (2014), pp. 970-976

[CrossRefView Record in Scopus](#)

[6]

Y. Yang, J. Wang, Z. Xiu, P.J.J. Alvarez **Impacts of silver nanoparticles on cellular and transcriptional activity of nitrogen-cycling bacteria**

Environ. Toxicol. Chem., 32 (2013), pp. 1488-1494

[View Record in Scopus](#)

[7]

Z. Xiu, Q. Zhang, H.L. Puppala, V.L. Colvin, P.J.J. Alvarez **Negligible particle-specific antibacterial activity of silver nanoparticles**

Nano Lett., 12 (2012), pp. 4271-4275

[CrossRefView Record in Scopus](#)

[8]

J. Fabrega, J.C. Renshaw, J.R. Lead **Interactions of silver nanoparticles with *Pseudomonas putida* biofilms**

Environ. Sci. Technol., 43 (2009), pp. 9004-9009

[CrossRefView Record in Scopus](#)

[9]

Z. Sheng, Y. Liu **Effects of silver nanoparticles on wastewater biofilms**

Water Res., 45 (2011), pp. 6039-6050

[ArticleDownload PDFView Record in Scopus](#)

[10]

Y. Yang, P.J.J. Alvarez **Sublethal concentrations of silver nanoparticles stimulate biofilm development**

Environ. Sci. Technol. Lett., 2 (2015), pp. 221-226

[CrossRefView Record in Scopus](#)

[11]

J.R. Morones, J.L. Elechiguerra, A. Camacho, K. Holt, J.B. Kouri, J.T. Ramirez, M.J. Yacaman **The bactericidal effect of silver nanoparticles**

Nanotechnology, 16 (2005), pp. 2346-2353

[CrossRefView Record in Scopus](#)

[12]

O. Choi, Z. Hu **Size dependent and reactive oxygen species related nanosilver toxicity to nitrifying bacteria**

Environ. Sci. Technol., 42 (2008), pp. 4583-4588

[CrossRefView Record in Scopus](#)

[13]

Y. Zhang, H. Peng, W. Huang, Y. Zhou, D. Yan **Facile preparation and characterization of highly antimicrobial colloid Ag or Au nanoparticles**

J. Colloid. Interf. Sci., 325 (2008), pp. 371-376

[ArticleDownload PDFView Record in Scopus](#)

S. Pal, Y.K. Tak, J.M. Song **Does the antibacterial activity of silver nanoparticles depend on the shape of the nanoparticle? a study of the gram-negative bacterium *Escherichia coli***
Appl. Environ. Microb., 73 (2007), pp. 1712-1720
[CrossRefView Record in Scopus](#)

[15]

C.L. Arnaout, C.K. Gunsch **Impacts of silver nanoparticle coating on the nitrification potential of *Nitrosomonas europaea***
Environ. Sci. Technol., 46 (2012), pp. 5387-5395
[CrossRefView Record in Scopus](#)

[16]

Y. Yang, J. Quensen, J. Mathieu, Q. Wang, J. Wang, M. Li, J.M. Tiedje, P.J. Alvarez **Pyrosequencing reveals higher impact of silver nanoparticles than Ag⁺ on the microbial community structure of activated sludge**
Water Res., 48 (2014), pp. 317-325
[ArticleDownload PDFCrossRefView Record in Scopus](#)

[17]

C.N. Lok, C.M. Ho, R. Chen, Q.Y. He, W.Y. Yu, H. Sun, P.K. Tam, J.F. Chiu, C.M. Che **Silver nanoparticles: partial oxidation and antibacterial activities**
J. Biol. Inorg. Chem., 12 (2007), pp. 527-534
[CrossRefView Record in Scopus](#)

[18]

O. Choi, K.K. Deng, N.J. Kim, L. Ross Jr., R.Y. Surampalli, Z. Hu **The inhibitory effects of silver nanoparticles, silver ions, and silver chloride colloids on microbial growth**
Water Res., 42 (2008), pp. 3066-3074
[ArticleDownload PDFView Record in Scopus](#)

[19]

Z. Sheng, Y. Liu **Potential impacts of silver nanoparticles on bacteria in the aquatic environment**
J. Environ. Manage., 191 (2017), pp. 290-296
[ArticleDownload PDFView Record in Scopus](#)

[20]

X. Sun, Z. Sheng, Y. Liu **Effects of silver nanoparticles on microbial community structure in activated sludge**
Sci. Total Environ., 443 (2013), pp. 828-835
[ArticleDownload PDFView Record in Scopus](#)

[21]

D. Kumar, J. Kumari, S. Pakrashi, S. Dalai, A.M. Raichur, T.P. Sastry, A.B. Mandal, N.Chandrasekaran, A. Mukherjee **Qualitative toxicity assessment of silver nanoparticles on the fresh water bacterial isolates and consortium at low level of exposure concentration**

Ecotoxicol. Environ. Saf., 108 (2014), pp. 152-160

[ArticleDownload](#) [PDFView](#) [Record in Scopus](#)

[22]

J. Fabrega, S.R. Fawcett, J.C. Renshaw, J.R. Lead **Silver nanoparticle impact on bacterial growth: effect of pH, concentration, and organic matter**

Environ. Sci. Technol., 43 (2009), pp. 7285-7290

[CrossRefView](#) [Record in Scopus](#)

[23]

J.W. Anderson, L. Semprini, T.S. Radniecki **Influence of water hardness on silver ion and silver nanoparticle fate and toxicity toward *Nitrosomonas europaea***

Environ. Eng. Sci., 31 (2014), pp. 403-409

[CrossRefView](#) [Record in Scopus](#)

[24]

C.N. Lok, C.M. Ho, R. Chen, Q.Y. He, W.Y. Yu, H. Sun, P.K. Tam, J.F. Chiu, C.M. Che **Proteomic analysis of the mode of antibacterial action of silver nanoparticles**

J. Proteome Res., 5 (2006), pp. 916-924

[CrossRefView](#) [Record in Scopus](#)

[25]

C. Zhang, Z. Hu, B. Deng **Silver nanoparticles in aquatic environments: physiochemical behavior and antimicrobial mechanisms**

Water Res., 88 (2016), pp. 403-427

[ArticleDownload](#) [PDFView](#) [Record in Scopus](#)

[26]

Z. Liang, A. Das, Z. Hu **Bacterial response to a shock load of nanosilver in an activated sludge treatment system**

Water Res., 44 (2010), pp. 5432-5438

[ArticleDownload](#) [PDFView](#) [Record in Scopus](#)

[27]

Y. Yang, M. Li, C. Michels, H. Moreira-Soares, P.J.J. Alvarez **Differential sensitivity of nitrifying bacteria to silver nanoparticles in activated sludge**

Environ. Toxicol. Chem., 33 (2014), pp. 2234-2239

[CrossRefView](#) [Record in Scopus](#)

[28]

E. Jeong, W. Im, D. Kim, M. Kim, S. Kang, H. Shin, S. Chae **Different susceptibilities of bacterial community to silver nanoparticles in wastewater treatment systems**

J. Environ. Sci. Heal. A, 49 (2014), pp. 685-693

[CrossRefView Record in Scopus](#)

[29]

A.B. Smetana, K.J. Klabunde, G.R. Marchin, C.M. Sorensen **Biocidal activity of nanocrystalline silver powders and particles**

Langmuir, 24 (2008), pp. 7457-7464

[CrossRefView Record in Scopus](#)

[30]

Z. Yuan, X. Yang, A. Hu, C. Yu **Long-term impacts of silver nanoparticles in an anaerobic-anoxic-oxic membrane bioreactor system**

Chem. Eng. J., 276 (2015), pp. 83-90

[ArticleDownload PDFView Record in Scopus](#)

[31]

C. Zhang, Z. Liang, Z. Hu **Bacterial response to a continuous long-term exposure of silver nanoparticles at sub-ppm silver concentrations in a membrane bioreactor activated sludge system**

Water Res., 50 (2014), pp. 350-358

[ArticleDownload PDFView Record in Scopus](#)

[32]

B. Kim, C. Park, M. Murayama, M.F. Hochella **Discovery and characterization of silver sulfide nanoparticles in final sewage sludge products**

Environ. Sci. Technol., 44 (2010), pp. 7509-7514

[CrossRefView Record in Scopus](#)

[33]

C. Levard, E.M. Hotze, G.V. Lowry Jr., G.E. Brown **Environmental transformations of silver nanoparticles: impact on stability and toxicity**

Environ. Sci Technol., 46 (2012), pp. 6900-6914

[CrossRefView Record in Scopus](#)

[34]

C.L. Doolette, M.J. McLaughlin, J.K. Kirby, D.J. Batstone, H.H. Harris, H. Ge, G. Cornelis **Transformation of PVP coated silver nanoparticles in a simulated wastewater treatment process and the effect on microbial communities**

Chem. Cent. J., 7 (2013)

[35]

R. Kaegi, A. Voegelin, C. Ort, B. Sinnet, B. Thalmann, J. Krismer, H. Hagendorfer, M. Elumelu, E. Mueller **Fate and transformation of silver nanoparticles in urban wastewater systems**

Water Res., 47 (2013), pp. 3866-3877

[ArticleDownload PDFView Record in Scopus](#)

[36]

J. Hedberg, C. Baresel, I.O. Wallinder **Transport and fate of silver as polymer-stabilised nanoparticles and ions in a pilot wastewater treatment plant, followed by sludge digestion and disposal of sludge/soil mixtures: a case study**

J. Environ. Sci. Health. A, 49 (2014), pp. 1416-1424

[CrossRefView Record in Scopus](#)

[37]

R. Ma, C. Levard, J.D. Judy, J.M. Unrine, M. Durenkamp, B. Martin, B. Jefferson, G.V. Lowry **Fate of zinc oxide and silver nanoparticles in a pilot wastewater treatment plant and in processed biosolids**

Environ. Sci. Technol., 48 (2014), pp. 104-112

[CrossRefView Record in Scopus](#)

[38]

Z. Sheng, J.D. Van Nostrand, J. Zhou, Y. Liu **The effects of silver nanoparticles on intact wastewater biofilms**

Front. Microbiol., 6 (2015), p. 680

[39]

T.S. Radniecki, D.P. Stankus, A. Neigh, J.A. Nason, L. Semprini **Influence of liberated silver from silver nanoparticles on nitrification inhibition of *Nitrosomonas europaea***

Chemosphere, 85 (2011), pp. 43-49

[ArticleDownload PDFView Record in Scopus](#)

[40]

L. Racz, T. Datta, R. Goel **Effect of organic carbon on ammonia oxidizing bacteria in a mixed culture**

Bioresour. Technol., 101 (2010), pp. 6454-6460

[ArticleDownload PDFView Record in Scopus](#)

[41]

APHA, standard methods for the examination of water and wastewater

Water Environment Federation (WEF) (20th ed.), American Public Health Association (APHA), American Water Works Association (AWWA), Washington, DC (1998)

[42]

J.P. Bassin, R. Kleerebezem, A.S. Rosado, M.C. van Loosdrecht, M. Dezotti **Effect of different operational conditions on biofilm development nitrification, and nitrifying microbial population in moving-bed biofilm reactors**

Environ. Sci. Technol., 46 (2012), pp. 1546-1555

[CrossRefView Record in Scopus](#)

[43]

K.R. Zodrow, E. Bar-Zeev, M.J. Giannetto, M. Elimelech **Biofouling and microbial communities in membrane distillation and reverse osmosis**

Environ. Sci. Technol., 48 (2014), pp. 13155-13164

[CrossRefView Record in Scopus](#)

[44]

B.J. Campbell, D.L. Kirchman **Bacterial diversity, community structure and potential growth rates along an estuarine salinity gradient**

ISME J., 7 (2013), pp. 210-220

[CrossRefView Record in Scopus](#)

[45]

P.A. Crawford, J.R. Crowley, N. Sambandam, B.D. Muegge, E.K. Costello, M. Hamady, R. Knight, J.I. Gordon **Regulation of myocardial ketone body metabolism by the gut microbiota during nutrient deprivation**

Proc. Natl. Acad. Sci. U.S.A, 106 (2009), pp. 11276-11281

[CrossRefView Record in Scopus](#)

[46]

Z. Lu, Y. Deng, J.D. Van

Nostrand, Z. He, J. Voordeckers, A. Zhou, Y.J. Lee, O.U. Mason, E.A. Dubinsky, K.L. Chavarria, L. M. Tom, J.L. Fortney, R. Lamendella, J.K. Jansson, P.D'Haeseleer, T.C. Hazen, J. Zhou **Microbial gene functions enriched in the Deepwater Horizon deep-sea oil plume**

ISME J., 6 (2012), pp. 451-460

[CrossRefView Record in Scopus](#)

[47]

Z. He, Y. Deng, J.D. Van

Nostrand, Q. Tu, M. Xu, C.L. Hemme, X. Li, L. Wu, T.J. Gentry, Y. Yin, J. Liebich, T.C. Hazen, J. Zhou **GeoChip 3.0 as a high-throughput tool for analyzing microbial community composition, structure and functional activity**

ISME J., 4 (2010), pp. 1167-1179

[CrossRefView Record in Scopus](#)

[48]

S. Wu, X. Feng, A. Wittmeier **Microwave digestion of plant and grain reference materials in nitric acid or a mixture of nitric acid or a mixture of nitric acid and hydrogen peroxide for the determination of multi-elements by inductively coupled plasma mass spectrometry**

J. Anal. Atom. Spectrom., 12 (1997), pp. 797-806

[CrossRefView Record in Scopus](#)

[49]

I. Sur, D. Cam, M. Kahraman, A. Baysal, M. Culha **Interaction of multi-functional silver nanoparticles with living cells**

Nanotechnology, 21 (2010)

[50]

B. van den Akker, H. Beard, U. Kaeding, S. Giglio, M.D. Short **Exploring the relationship between viscous bulking and ammonia-oxidiser abundance in activated sludge: a comparison of conventional and IFAS systems**

Water Res., 44 (2010), pp. 2919-2929

[ArticleDownload PDFView Record in Scopus](#)

[51]

N. Terry, S.V. Sambukumar, D.L. LeDuc **Biotechnological approaches for enhancing phytoremediation of heavy metals and metalloids**

Acta. Biotechnol., 23 (2003), pp. 281-288

[CrossRefView Record in Scopus](#)

[52]

C. Goodman **Sphingolipids: microbes talk to T cells**

PLoS Biol., 9 (2013), p. 529

[CrossRef](#)

[53]

A. Bradford, R.D. Handy, J.W. Readman, A. Atfield, M. Mühling **Impact of silver nanoparticle contamination on the genetic diversity of natural bacterial assemblages in estuarine sediments**

Environ. Sci. Technol., 43 (2009), pp. 4530-4536

[CrossRefView Record in Scopus](#)

[54]

B.P. Colman, S.Y. Wang, M. Auffan, M.R. Wiesner, E.S. Bernhardt **Antimicrobial effects of commercial silver nanoparticles are attenuated in natural streamwater and sediment**

Ecotoxicology, 21 (2012), pp. 1867-1877

[CrossRefView Record in Scopus](#)

[55]

S. Kittler, C. Greulich, J. Diendorf, M. Köller, M. Eppe **Toxicity of silver nanoparticles increases during storage because of slow dissolution under release of silver ions**

Chem. Mater., 22 (2010), pp. 4548-4554

[CrossRefView Record in Scopus](#)

[56]

J. Davies, G.B. Spiegelman, G. Yim **The world of subinhibitory antibiotic concentrations**

Curr. Opin. Microbiol., 9 (2006), pp. 445-453

[ArticleDownload PDFView Record in Scopus](#)

[57]

E.J. Calabrese **Hormesis: a fundamental concept in biology**

Microb. Cell, 1 (2014), pp. 145-149

[CrossRefView Record in Scopus](#)

[58]

E.J. Calabrese, L.A. Baldwin **Hormesis U-shaped dose responses and their centrality in toxicology**

Trends Pharmacol. Sci., 22 (2001), pp. 285-291

[ArticleDownload PDFView Record in Scopus](#)

[59]

V. Lavakumar, K. Masilamani, V. Ravichandiran, N. Venkateshan, D.V.R. Saigopal, C.K.Ashok Kumar, C. Sowmya **Promising upshot of silver nanoparticles primed from *Gracilaria crassa* against bacterial pathogens**

Chem. Cent. J., 9 (2015), p. 42

[60]

M. Raffi, F. Hussain, T.M. Bhatti, H.A. Akhter **Antibacterial characterization of silver nanoparticles against E. Coli ATCC-15224**

J. Mater. Sci. Technol., 24 (2008), p. 192

[View Record in Scopus](#)

[61]

S. Rajesh, V. Dharanishanthi, A.V. Kanna **Antibacterial mechanism of biogenic silver nanoparticles of *Lactobacillus acidophilus***

J. Exp. Nanosci., 10 (2015), pp. 1143-1152

[CrossRefView Record in Scopus](#)

[62]

B.E. Rittmann, P.L. McCarty **Environmental Biotechnology: Principles and Applications**
McGraw-Hill, Boston (2001)

[63]

S. Siripong, B.E. Rittmann **Diversity study of nitrifying bacteria in full-scale municipal wastewater treatment plants**

Water Res., 41 (2007), pp. 1110-1120

[ArticleDownload PDFView Record in Scopus](#)

[64]

S. Xu, M. Sun, C. Zhang, R. Surampalli, Z. Hu **Filamentous sludge bulking control by nano zero-valent iron in activated sludge treatment systems**

Environ. Sci. Process. Impacts, 16 (2014), pp. 2721-2728

[CrossRefView Record in Scopus](#)

[65]

E.J. Calabrese, L.A. Baldwin **Defining hormesis**

Hum. Exp. Toxicol., 21 (2002), pp. 91-97

[CrossRefView Record in Scopus](#)

[66]

Y. Zhao, J. Xing, J.Z. Xing, W.T. Ang, J. Chen **Applications of low-intensity pulsed ultrasound to increase monoclonal antibody production in CHO cells using shake flasks or wavebags**

Ultrasonics, 54 (2014), pp. 1439-1447

[ArticleDownload PDFView Record in Scopus](#)