

UC Berkeley

UC Berkeley Previously Published Works

Title

Scale setting the Möbius domain wall fermion on gradient-flowed HISQ action using the omega baryon mass and the gradient-flow scales t_0 and w_0

Permalink

<https://escholarship.org/uc/item/0dp3d0dg>

Journal

Physical Review D, 103(5)

ISSN

2470-0010

Authors

Miller, Nolan
Carpenter, Logan
Berkowitz, Evan
[et al.](#)

Publication Date

2021-03-01

DOI

10.1103/physrevd.103.054511

Peer reviewed

Scale setting the Möbius domain wall fermion on gradient-flowed HISQ action using the omega baryon mass and the gradient-flow scales t_0 and w_0

Nolan Miller,¹ Logan Carpenter,² Evan Berkowitz,^{3,4} Chia Cheng Chang (張家丞),^{5,6,7} Ben Hörz,⁶ Dean Howarth,^{8,6} Henry Monge-Camacho,^{9,1} Enrico Rinaldi,^{10,5} David A. Brantley,⁸ Christopher Körber,^{7,6} Chris Bouchard,¹¹ M.A. Clark,¹² Arjun Singh Gambhir,^{13,6} Christopher J. Monahan,^{14,15} Amy Nicholson,^{1,6} Pavlos Vranas,^{8,6} and André Walker-Loud^{6,8,7}

¹*Department of Physics and Astronomy, University of North Carolina, Chapel Hill, NC 27516-3255, USA*

²*Department of Physics, Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

³*Department of Physics, University of Maryland, College Park, MD 20742, USA*

⁴*Institut für Kernphysik and Institute for Advanced Simulation, Forschungszentrum Jülich, 52425 Jülich Germany*

⁵*Interdisciplinary Theoretical and Mathematical Sciences Program*

(iTHEMS), RIKEN, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

⁶*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA*

⁷*Department of Physics, University of California, Berkeley, CA 94720, USA*

⁸*Physics Division, Lawrence Livermore National Laboratory, Livermore, CA 94550, USA*

⁹*Escuela de Física, Universidad de Costa Rica, 11501 San José, Costa Rica*

¹⁰*Arithmer Inc., R&D Headquarters, Minato, Tokyo 106-6040, Japan*

¹¹*School of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, UK*

¹²*NVIDIA Corporation, 2701 San Tomas Expressway, Santa Clara, CA 95050, USA*

¹³*Design Physics Division, Lawrence Livermore National Laboratory, Livermore, CA 94550, USA*

¹⁴*Department of Physics, The College of William & Mary, Williamsburg, VA 23187, USA*

¹⁵*Theory Center, Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA*

(Dated: April 16, 2021 - 1:31)

We report on a subpercent scale determination using the omega baryon mass and gradient-flow methods. The calculations are performed on 22 ensembles of $N_f = 2 + 1 + 1$ highly improved, rooted staggered sea-quark configurations generated by the MILC and CalLat Collaborations. The valence quark action used is Möbius domain wall fermions solved on these configurations after a gradient-flow smearing is applied with a flowtime of $t_{\text{gf}} = 1$ in lattice units. The ensembles span four lattice spacings in the range $0.06 \lesssim a \lesssim 0.15$ fm, six pion masses in the range $130 \lesssim m_\pi \lesssim 400$ MeV and multiple lattice volumes. On each ensemble, the gradient-flow scales t_0/a^2 and w_0/a and the omega baryon mass am_Ω are computed. The dimensionless product of these quantities is then extrapolated to the continuum and infinite volume limits and interpolated to the physical light, strange and charm quark mass point in the isospin limit, resulting in the determination of $\sqrt{t_0} = 0.1422(14)$ fm and $w_0 = 0.1709(11)$ fm with all sources of statistical and systematic uncertainty accounted for. The dominant uncertainty in both results is the stochastic uncertainty, though for $\sqrt{t_0}$ there are comparable continuum extrapolation uncertainties. For w_0 , there is a clear path for a few-per-mille uncertainty just through improved stochastic precision, as recently obtained by the Budapest-Marseille-Wuppertal Collaboration.

I. INTRODUCTION

Lattice QCD (LQCD) has become a prominent theoretical tool for calculations of hadronic quantities, and many calculations have reached a level of precision to be able to supplement and/or complement experimental determinations [1]. Precision calculations of Standard Model processes, for example, are crucial input for experimental tests of fundamental symmetries in searches for new physics.

Lattice calculations receive only dimensionless bare parameters as input, so the output is inherently dimensionless. In some cases, dimensionless quantities or ratios of quantities may be directly computed without the need to determine any dimensionful scale. Calculations of g_A and F_K/F_π are examples for which a precise scale setting is not necessary to make a precise, final prediction. However, there are many quantities for which a precise scale setting is desirable, such as the hadron spectrum,

the nucleon axial radius, the hadronic contribution to the muon $g - 2$ [2] and many others.

In these cases, a quantity which is dimensionful (after multiplying or dividing by an appropriate power of the lattice spacing) is calculated and compared to experiment, following extrapolations to the physical point in lattice spacing, volume, and pion mass. Because the precision of any calculations of further dimensionful quantities is limited by the statistical and systematic uncertainties of this scale setting, quantities which have low stochastic noise and mild light quark mass dependence, such as the omega baryon mass m_Ω , are preferred. The lattice spacing on each ensemble may then be determined by comparing the quantity calculated on a given ensemble to the continuum value.

However, the most precise quantities one may calculate are not necessarily accessible experimentally. For example, the Sommer scale r_0 [3] has been one of the most commonly used scales. This scale requires a determina-

tion of the heavy-quark potential which is susceptible to fitting systematic uncertainties. More recently, the gradient flow scales t_0 [4] and w_0 [5] have been used for a more precise determination of the lattice spacing [6–14]. In this case, a well-controlled extrapolation of these quantities to the physical point is also necessary.

In this paper we present a precision scale setting for our mixed lattice action [15] which uses $N_f = 2 + 1 + 1$ highly improved, rooted staggered sea-quark (HISQ) configurations generated by the MILC [16] and CalLat Collaborations and Möbius domain wall fermions for the valence sector. We compute the dimensionless products $\sqrt{t_0}m_\Omega$ and $m_\Omega w_0$ on each ensemble and extrapolate them to the physical point resulting in the determinations

$$\begin{aligned}\sqrt{t_0}m_\Omega &= 1.2051(82)^s(15)^\chi(46)^a(00)^V(21)^{\text{phys}}(61)^M \\ &= 1.205(12), \\ \frac{\sqrt{t_0}}{\text{fm}} &= 0.1422(09)^s(02)^\chi(05)^a(00)^V(02)^{\text{phys}}(07)^M \\ &= 0.1422(14),\end{aligned}\tag{1.1}$$

$$\begin{aligned}w_0m_\Omega &= 1.4483(82)^s(15)^\chi(45)^a(00)^V(26)^{\text{phys}}(18)^M \\ &= 1.4483(97) \\ \frac{w_0}{\text{fm}} &= 0.1709(10)^s(02)^\chi(05)^a(00)^V(03)^{\text{phys}}(02)^M \\ &= 0.1709(11),\end{aligned}\tag{1.2}$$

with the statistical (s), chiral (χ), continuum-limit (a), infinite volume (V), physical-point (phys), and model selection uncertainties (M).

We then perform an interpolation of the values of t_0/a^2 and w_0/a to the physical quark-mass limit and extrapolation to infinite volume which allows us to provide a precise, quark mass independent scale setting for each lattice spacing, with our final results in Table V. In the following sections we provide details of our lattice setup, our methods for extrapolation, and our results with uncertainty breakdown. We conclude with a discussion in the final section.

II. DETAILS OF THE LATTICE CALCULATION

A. MDWF on gradient-flowed HISQ

The lattice action we use is the mixed-action [20, 21] with Möbius [22] domain wall fermions [23–25] solved on $N_f = 2 + 1 + 1$ highly improved staggered quarks [26] after they are gradient-flow smeared [27–29] (corresponding to an infinitesimal stout-smearing procedure [30]) to a flow-time of $t_{\text{gf}}/a^2 = 1$ [15]. The choice to hold the flow-time fixed in lattice units is important to ensure that as the continuum limit is taken, effects arising from finite flow-time also extrapolate to zero.

This action has been used to compute the nucleon axial coupling, g_A , with a 1% total uncertainty [31–34], the

$\pi^- \rightarrow \pi^+$ matrix elements relevant to neutrinoless double beta-decay [35] and most recently, F_K/F_π [36]. Our calculation of F_K/F_π was obtained with a total uncertainty of 0.4% which provides an important benchmark for our action, as the result is consistent with other determinations in the literature [8, 11, 37–44] (and the FLAG average [1]), and also contributes to the test of the universality of lattice QCD results in the continuum limit.

Our plan to compute the axial and other elastic form factors of the nucleon with this mixed-action, as well as other quantities, leads to a desire to have a scale setting with sufficiently small uncertainty that it does not increase the final uncertainty of such quantities. It has been previously observed that both w_0 [5, 12] and the omega baryon mass [14, 45–51] have mild quark mass dependence and that they can be determined with high statistical precision with relatively low computational cost. The input parameters of our action on all ensembles are provided in Table I.

B. Correlation function construction and analysis

For the scale setting computation, we have to determine four or five quantities on each ensemble, the pion, kaon and omega masses, the gradient-flow scale w_0 and the pion decay constant F_π . For m_π , m_K and F_π , we take the values from our F_K/F_π computation for the 18 ensembles in common. For the four new ensembles in this work (a15m310L, a12m310XL, a12m220ms, a12m180L), we follow the same analysis strategy described in Ref. [36].

The a12m220ms ensemble is identical to a12m220 except that the strange quark mass is roughly 60% of the physical value rather than being near the physical value. The a15m310L ensemble has identical input parameters as the a15m310 ensemble but $L = 24$ (3.6 fm) instead of $L = 16$ (2.4 fm), while the a12m310XL ensemble is identical to the a12m310 ensemble but with $L = 48$ (5.8 fm) instead of $L = 24$ (2.9 fm). The a12m180L and a12m310XL ensembles have a lattice volume that is the same size as a12m130 but pion masses of roughly $m_\pi \simeq 180$ and 310 MeV. These new ensembles provide important lever arms for the various extrapolations. The a12m220ms provides a unique lever arm for varying the strange quark mass significantly from its physical value, the a15m310L and a12m310XL provide other pion masses where we can perform a volume study and the a12m180L ensemble provides an additional light pion mass ensemble to help with the physical pion mass extrapolation. The first of these is important for this scale setting while the latter three will be more important for future work.

The omega baryon correlation functions are constructed similarly to the pion and kaon. A source for the propagator is constructed with the gauge invariant Gaussian smearing routine in Chroma [52] (GAUGE_INV_GAUSSIAN). Then, correlation functions are constructed using both a point sink as well as the same

TABLE I. Input parameters for our lattice action. The abbreviated ensemble name [17] indicates the approximate lattice spacing in fm and pion mass in MeV. The S, L, XL which come after an ensemble name denote a relatively small, large and extra-large volume with respect to $m_\pi L = 4$.

Ensemble	β	N_{cfg}	Volume	am_l	am_s	am_c	L_5/a	aM_5	b_5, c_5	am_l^{val}	$am_l^{\text{res}} \times 10^4$	am_s^{val}	$am_s^{\text{res}} \times 10^4$	σ	N	N_{src}
a15m400 ^a	5.80	1000	$16^3 \times 48$	0.0217	0.065	0.838	12	1.3	1.50, 0.50	0.0278	9.365(87)	0.0902	6.937(63)	3.0	30	8
a15m350 ^a	5.80	1000	$16^3 \times 48$	0.0166	0.065	0.838	12	1.3	1.50, 0.50	0.0206	9.416(90)	0.0902	6.688(62)	3.0	30	16
a15m310	5.80	1000	$16^3 \times 48$	0.013	0.065	0.838	12	1.3	1.50, 0.50	0.0158	9.563(67)	0.0902	6.640(44)	4.2	45	24
a15m310L ^a	5.80	1000	$24^3 \times 48$	0.013	0.065	0.838	12	1.3	1.50, 0.50	0.0158	9.581(50)	0.0902	6.581(37)	4.2	45	4
a15m220	5.80	1000	$24^3 \times 48$	0.0064	0.064	0.828	16	1.3	1.75, 0.75	0.00712	5.736(38)	0.0902	3.890(25)	4.5	60	16
a15m135XL ^a	5.80	1000	$48^3 \times 64$	0.002426	0.06730	0.8447	24	1.3	2.25, 1.25	0.00237	2.706(08)	0.0945	1.860(09)	3.0	30	32
a12m400 ^a	6.00	1000	$24^3 \times 64$	0.0170	0.0509	0.635	8	1.2	1.25, 0.25	0.0219	7.337(50)	0.0693	5.129(35)	3.0	30	8
a12m350 ^a	6.00	1000	$24^3 \times 64$	0.0130	0.0509	0.635	8	1.2	1.25, 0.25	0.0166	7.579(52)	0.0693	5.062(34)	3.0	30	8
a12m310	6.00	1053	$24^3 \times 64$	0.0102	0.0509	0.635	8	1.2	1.25, 0.25	0.0126	7.702(52)	0.0693	4.950(35)	3.0	30	8
a12m310XL ^a	6.00	1000	$48^3 \times 64$	0.0102	0.0509	0.635	8	1.2	1.25, 0.25	0.0126	7.728(22)	0.0693	4.927(21)	3.0	30	8
a12m220S	6.00	1000	$24^4 \times 64$	0.00507	0.0507	0.628	12	1.2	1.50, 0.50	0.00600	3.990(42)	0.0693	2.390(24)	6.0	90	4
a12m220	6.00	1000	$32^3 \times 64$	0.00507	0.0507	0.628	12	1.2	1.50, 0.50	0.00600	4.050(20)	0.0693	2.364(15)	6.0	90	4
a12m220ms	6.00	1000	$32^3 \times 64$	0.00507	0.0304	0.628	12	1.2	1.50, 0.50	0.00600	3.819(26)	0.0415	2.705(20)	6.0	90	8
a12m220L	6.00	1000	$40^3 \times 64$	0.00507	0.0507	0.628	12	1.2	1.50, 0.50	0.00600	4.040(26)	0.0693	2.361(19)	6.0	90	4
a12m180L ^a	6.00	1000	$48^3 \times 64$	0.00339	0.0507	0.628	14	1.2	1.75, 0.75	0.00380	3.038(13)	0.0693	1.888(11)	3.0	30	16
a12m130	6.00	1000	$48^3 \times 64$	0.00184	0.0507	0.628	20	1.2	2.00, 1.00	0.00195	1.642(09)	0.0693	0.945(08)	3.0	30	32
a09m400 ^a	6.30	1201	$32^3 \times 64$	0.0124	0.037	0.44	6	1.1	1.25, 0.25	0.0160	2.532(23)	0.0491	1.957(17)	3.5	45	8
a09m350 ^a	6.30	1201	$32^3 \times 64$	0.00945	0.037	0.44	6	1.1	1.25, 0.25	0.0121	2.560(24)	0.0491	1.899(16)	3.5	45	8
a09m310	6.30	780	$32^3 \times 96$	0.0074	0.037	0.44	6	1.1	1.25, 0.25	0.00951	2.694(26)	0.0491	1.912(15)	6.7	167	8
a09m220	6.30	1001	$48^3 \times 96$	0.00363	0.0363	0.43	8	1.1	1.25, 0.25	0.00449	1.659(13)	0.0491	0.834(07)	8.0	150	6
a09m135 ^a	6.30	1010	$64^3 \times 96$	0.001326	0.03636	0.4313	12	1.1	1.50, 0.50	0.00152	0.938(06)	0.04735	0.418(04)	3.5	45	16
a06m310L ^a	6.72	1000	$72^3 \times 96$	0.0048	0.024	0.286	6	1.0	1.25, 0.25	0.00617	0.225(03)	0.0309	0.165(02)	3.5	45	8

^a Additional ensembles generated by CalLat using the MILC code. The m350 and m400 ensembles were made on the Vulcan supercomputer at LLNL while the a12m310XL, a12m180L, a15m135XL, a09m135, and a06m310L ensembles were made on the Sierra and Lassen supercomputers at LLNL and the Summit supercomputer at OLCF using QUDA [18, 19]. These configurations are available to any interested party upon request, and will be available for easy anonymous downloading—hopefully soon.

gauge invariant Gaussian smearing routine with the same parameters as the source. The values of the “smearing width” (σ) and the number of iterations (N) used to approximate the exponential smearing profile are provided in Table I. The correlation functions constructed with the point sink are referred to as PS and those with the smeared sink as SS.

Local spin wave functions are constructed following Refs. [53, 54]. Both positive- and negative-parity omega-baryon correlation functions are constructed with the upper and lower spin components of the quark propagators in the Dirac basis. The negative-parity correlation functions are time-reversed with an appropriate sign flip of the correlation function, effectively doubling the statistics with no extra inversions. The four different spin projections of the omega are averaged as well to produce the final spin and parity averaged two-point correlation functions.

The reader will notice that the values of σ and N do not follow an obvious pattern. This is because in our first computations of g_A [31, 32], we applied an “aggressive” smearing with a larger value of σ and correspondingly larger number of iterations, which led to a large suppression of excited states, but also showed evidence of “over smearing” such that the non-positive-definite PS correlation functions displayed symptoms of having a relatively large negative overlap factor for excited states (there were wiggles in the PS effective masses). In a subsequent paper studying the two-nucleon system on the a12m350 ensemble [55], where we utilized matrix Prony [56] to form linear combinations of PS and SS nucleons to construct a

“calm” nucleon which is ground-state dominated earlier in time, we observed that using a milder smearing with smaller width and fewer iterations provided a much more stable extraction of the ground state and did not show signs of large negative overlap factors. Hence, many but not all of the ensembles have been rerun with our improved choices of σ and N . We have observed the choice $\sigma = 3.0$ and $N = 30$ works well for the a15 and a12 ensembles and that $\sigma = 3.5$ with $N = 45$ works well for the a09 and a06 ensembles.

In order to determine the omega baryon mass on each ensemble, we perform a stability analysis of the extracted ground state mass as a function of t_{min} used in the fit as well as the number of states used in the analysis. The correlation functions are analyzed in a Bayesian framework with constraints [57]. We choose normally distributed priors for the ground-state energy and all overlap factors, and log-normal distributions for excited-state energy priors. The ground-state energy and overlap factors are motivated by the plateau values of the effective masses with the priors taken to be roughly 10 times larger than the stochastic uncertainty of the respective effective mass data in the plateau region. The excited-state energy splittings are set to the value of two pion masses with a width allowing for fluctuations down to one pion mass within one standard deviation.

In Fig. 1 we show sample extractions of the ground state mass on our three physical pion mass ensembles. In the left plot, we show the effective mass data from the two correlation functions. The weights are normalized on a given time slice by the largest Bayes factor at that




FIG. 1. Stability plots of the ground state omega baryon mass on the three physical pion mass ensembles. The left plots show the effective mass data (in lattice units) and reconstructed effective mass from the chosen fit for both the SS and PS correlation functions. The dark gray and colored band are displayed for the region of time used in the analysis and an extrapolation beyond t_{\max} is shown after a short break in the fit band. The horizontal gray band is the prior used for the ground state mass. The right plots show the corresponding value of E_0 as a function of t_{\min} and the number of states n_s used in the analysis, as well as the corresponding Q value and relative weight as a function of n_s for a given t_{\min} , where the weight is set by the Bayes Factor. See Appendix A for more detail on the selection of the final fit. The chosen fit is denoted with a filled black symbol and the horizontal band is the value of E_0 from the chosen fit. The y -range of the upper panel of the stability plots is equal to the prior of the ground state energy (the horizontal gray band in the left plot).

t_{\min} value. We have not implemented a more thorough algorithm to weight fits against each other that utilize different amounts of data, as described for example in Ref. [58]. Rather, we have chosen a fit for a given ensemble (the filled black symbol in the right panels highlighted by the horizontal colored band) that has a good fit qual-

ity, the maximum or near maximum relative weight, and consistency with the late-time data. We tried to optimize this choice over all ensembles simultaneously, with t_{\min} held nearly fixed for a given lattice spacing, rather than hand-picking the optimal fit on each ensemble separately, in order to minimize the possible bias introduced

by analysis choices. Good fits are obtained on all ensembles with $n_s = 2$, simplifying the model function and reducing the chance of overfitting the correlation functions, which is most relevant on ensembles with the more aggressive choices of smearing parameters. In Appendix C, we show the corresponding stability plots for all remaining ensembles. In Table II we show the resulting values of am_Ω on all ensembles used in this work.

C. Calculation of t_0 and w_0

In order to efficiently compute the value of t_0/a^2 and w_0/a on each ensemble, we have implemented the Symanzik flow in the QUDA library [18, 19, 59]. We used the tree-level improved action and the symmetric, cloverleaf definition of the field-strength tensor, following the MILC implementation [60]. We used a step size of $\epsilon = 0.01$ in the Runge-Kutta algorithm proposed by Lüscher [4], which leads to negligibly small integration errors. The scales t_0 and w_0 are defined by the equations

$$\begin{aligned} t^2 \langle E(t) \rangle \Big|_{t=t_0, \text{orig}} &= 0.3, \\ W_{\text{orig}} \equiv t \frac{d}{dt} (t^2 \langle E(t) \rangle) \Big|_{t=w_0^2, \text{orig}} &= 0.3, \end{aligned} \quad (2.1)$$

where $\langle E(t) \rangle$ is the gluonic action density at flow time t . In Fig. 2, we show the determination of the $w_{0, \text{orig}}/a$ on the two physical pion mass ensembles that we have generated. The uncertainties are determined by observing a saturation of the uncertainty as the bin-size is increased when binning the results from configurations close in Monte Carlo time. These uncertainties were cross-checked with an autocorrelation study using the Γ -method [61] implemented in the `unew` Python package [62].

Reference [64] determined the tree-level in lattice perturbation theory improvement coefficients for the determination of these gradient flow scales through $\mathcal{O}(a^8/t^4)$ for various choices of the gauge action, the gradient flow action and the definition of the field-strength tensor. As defined in Ref. [64], we have implemented the Symanzik-Symanzik-Clover (SSC) scheme with the relevant improvement coefficients (see Table 1 of Ref. [64])

$$\begin{aligned} C_2 &= -\frac{19}{72}, & C_4 &= \frac{145}{1536}, \\ C_6 &= -\frac{12871}{276480}, & C_8 &= \frac{52967}{1769472}. \end{aligned} \quad (2.2)$$

One can then determine improved scales, $t_{0, \text{imp}}$ and $w_{0, \text{imp}}$ in which one has perturbatively removed the leading discretization effects in these flow observables,

$$\begin{aligned} \frac{t^2 \langle E(t) \rangle}{1 + \sum_n C_{2n} \frac{a^{2n}}{t^n}} \Big|_{t=t_{0, \text{imp}}} &= 0.3, \\ t \frac{d}{dt} \left(\frac{t^2 \langle E(t) \rangle}{1 + \sum_n C_{2n} \frac{a^{2n}}{t^n}} \right) \Big|_{t=w_{0, \text{imp}}^2} &= 0.3, \end{aligned} \quad (2.3)$$




FIG. 2. Determination of $w_{0, \text{orig}}$ on the two new physical pion mass ensembles.

In the present work, we explore using both the original and improved versions of t_0 and w_0 when performing our scale setting analysis. For the improved versions, we have implemented the fourth order improvement [up to and including the $C_8(a^2/t)^4$ correction]. This is the same implementation performed by MILC in Ref. [12].

In Table II, we list the values of t_0/a^2 and w_0/a for the original and improved definitions as well as the dimensionless products of $\sqrt{t_0}m_\Omega$ and w_0m_Ω .

III. EXTRAPOLATION FUNCTIONS

This work utilizes 22 different ensembles, each with $\mathcal{O}(1000)$ configurations (Table I), to control the systematic uncertainties in the LQCD calculation of the scale. This allows us to address:

1. The physical light and strange quark mass limit;
2. The physical charm quark mass limit;
3. The continuum limit;
4. The infinite volume limit.

TABLE II. The omega baryon mass (m_Ω) and gradient flow scales ($t_{0,\text{orig}}$, $t_{0,\text{imp}}$, $w_{0,\text{orig}}$ and $w_{0,\text{imp}}$) determined on each ensemble are listed as well as their dimensionless products. Additionally, in the bottom panel, we list the parameters used to control the physical point extrapolation: $l_\Omega^2 = m_\pi^2/m_\Omega^2$, $s_\Omega^2 = (2m_K^2 - m_\pi^2)/m_\Omega^2$, $l_F^2 = m_\pi^2/(4\pi F_\pi)^2$, $s_F^2 = (2m_K^2 - m_\pi^2)/(4\pi F_\pi)^2$, $m_\pi L$, $\epsilon_a = a/(2w_{0,\text{orig}})$ and α_S . The values of α_S are taken from Table III of Ref. [12] which were determined with a heavy-quark potential method [63]. An HDF5 file is provided with this publication which includes the resulting bootstrap samples of all these quantities which can be used to construct the correlated uncertainties.

Ensemble	am_Ω	$t_{0,\text{orig}}/a^2$	$t_{0,\text{imp}}/a^2$	$\sqrt{t_{0,\text{orig}}}m_\Omega$	$\sqrt{t_{0,\text{imp}}}m_\Omega$	$w_{0,\text{orig}}/a$	$w_{0,\text{imp}}/a$	$w_{0,\text{orig}}m_\Omega$	$w_{0,\text{imp}}m_\Omega$
a15m400	1.2437(43)	1.1905(15)	0.9563(11)	1.3570(47)	1.2162(42)	1.1053(12)	1.0868(14)	1.3747(49)	1.3516(50)
a15m350	1.2331(31)	1.2032(14)	0.9660(11)	1.3526(35)	1.2120(32)	1.1154(11)	1.0988(13)	1.3754(37)	1.3550(38)
a15m310L	1.2287(31)	1.2111(07)	0.9720(05)	1.3521(34)	1.2113(30)	1.1219(06)	1.1066(07)	1.3784(35)	1.3596(35)
a15m310	1.2312(36)	1.2121(09)	0.9727(07)	1.3555(40)	1.2143(36)	1.1230(07)	1.1079(09)	1.3826(41)	1.3640(41)
a15m220	1.2068(26)	1.2298(08)	0.9861(07)	1.3383(30)	1.1984(27)	1.1378(07)	1.1255(08)	1.3731(31)	1.3582(31)
a15m135XL	1.2081(19)	1.2350(04)	0.9897(03)	1.3425(21)	1.2018(19)	1.1432(03)	1.1319(04)	1.3811(22)	1.3674(22)
a12m400	1.0279(25)	1.6942(11)	1.4172(10)	1.3380(33)	1.2237(30)	1.3616(07)	1.3636(08)	1.3997(35)	1.4017(35)
a12m350	1.0139(26)	1.7091(14)	1.4298(12)	1.3255(35)	1.2124(32)	1.3728(10)	1.3755(11)	1.3918(38)	1.3946(38)
a12m310XL	1.0072(41)	1.7221(07)	1.4407(06)	1.3217(54)	1.2089(50)	1.3831(05)	1.3865(06)	1.3930(58)	1.3964(58)
a12m310	1.0112(32)	1.7213(19)	1.4398(17)	1.3267(42)	1.2134(39)	1.3830(13)	1.3863(12)	1.3985(46)	1.4019(46)
a12m220ms	0.8896(92)	1.7891(15)	1.4977(13)	1.190(12)	1.089(11)	1.4339(12)	1.4406(12)	1.276(13)	1.282(13)
a12m220S	0.9970(26)	1.7466(20)	1.4614(17)	1.3177(35)	1.2053(32)	1.4021(15)	1.4069(16)	1.3980(39)	1.4027(39)
a12m220L	0.9944(30)	1.7489(09)	1.4633(08)	1.3150(40)	1.2028(37)	1.4041(06)	1.4090(07)	1.3962(43)	1.4010(43)
a12m220	0.9924(60)	1.7498(14)	1.4641(12)	1.3127(80)	1.2007(73)	1.4047(10)	1.4096(11)	1.3940(85)	1.3988(86)
a12m180L	0.9924(26)	1.7553(05)	1.4686(05)	1.3148(35)	1.2026(32)	1.4093(05)	1.4145(05)	1.3985(38)	1.4037(38)
a12m130	0.9801(26)	1.7628(07)	1.4749(06)	1.3013(34)	1.1903(31)	1.4155(05)	1.4211(06)	1.3873(37)	1.3928(37)
a09m400	0.7716(23)	2.9158(42)	2.6040(33)	1.3176(41)	1.2451(38)	1.8602(26)	1.8686(27)	1.4353(48)	1.4418(48)
a09m350	0.7561(35)	2.9455(37)	2.6301(34)	1.2977(61)	1.2262(58)	1.8810(25)	1.8900(26)	1.4222(69)	1.4291(70)
a09m310	0.7543(36)	2.9698(32)	2.6521(29)	1.2998(63)	1.2283(59)	1.8970(22)	1.9066(22)	1.4308(71)	1.4381(71)
a09m220	0.7377(30)	3.0172(16)	2.6952(15)	1.2814(53)	1.2111(50)	1.9282(12)	1.9388(12)	1.4224(59)	1.4302(60)
a09m135	0.7244(25)	3.0390(12)	2.7147(11)	1.2629(44)	1.1936(42)	1.9450(10)	1.9563(11)	1.4091(50)	1.4172(50)
a06m310L	0.5069(21)	6.4079(45)	6.0606(44)	1.2830(54)	1.2478(52)	2.8958(20)	2.9053(20)	1.4678(62)	1.4726(62)

Ensemble	l_F^2	s_F^2	l_Ω^2	s_Ω^2	$m_\pi L$	ϵ_a^2	α_S
a15m400	0.09216(33)	0.2747(10)	0.05928(42)	0.1767(12)	4.85	0.20462(44)	0.58801
a15m350	0.07505(28)	0.2915(09)	0.04609(25)	0.1790(09)	4.24	0.20096(39)	0.58801
a15m310L	0.06018(23)	0.2984(11)	0.03630(18)	0.1800(09)	5.62	0.19864(20)	0.58801
a15m310	0.06223(17)	0.3035(09)	0.03675(23)	0.1792(11)	3.78	0.19825(26)	0.58801
a15m220	0.03269(11)	0.3253(09)	0.01877(09)	0.1868(08)	3.97	0.19311(23)	0.58801
a15m135XL	0.01319(05)	0.3609(11)	0.00726(03)	0.1986(06)	4.94	0.19129(10)	0.58801
a12m400	0.08889(30)	0.2648(10)	0.05610(28)	0.1671(08)	5.84	0.13484(15)	0.53796
a12m350	0.07307(37)	0.2810(13)	0.04454(24)	0.1713(09)	5.14	0.13266(19)	0.53796
a12m310XL	0.05904(23)	0.2909(12)	0.03506(29)	0.1727(14)	9.05	0.13069(10)	0.53796
a12m310	0.05984(25)	0.2933(12)	0.03482(23)	0.1707(11)	4.53	0.13071(24)	0.53796
a12m220ms	0.03400(16)	0.2000(09)	0.02229(46)	0.1311(27)	4.25	0.12159(20)	0.53796
a12m220S	0.03384(19)	0.3210(19)	0.01849(13)	0.1754(09)	3.25	0.12716(27)	0.53796
a12m220L	0.03289(15)	0.3195(14)	0.01816(12)	0.1765(11)	5.36	0.12681(11)	0.53796
a12m220	0.03314(15)	0.3202(15)	0.01831(23)	0.1769(22)	4.30	0.12670(18)	0.53796
a12m180L	0.02277(09)	0.3319(13)	0.01220(07)	0.1779(10)	5.26	0.12588(08)	0.53796
a12m130	0.01287(08)	0.3429(14)	0.00687(05)	0.1832(10)	3.90	0.12477(09)	0.53796
a09m400	0.08883(32)	0.2638(09)	0.05512(34)	0.1637(10)	5.80	0.07225(20)	0.43356
a09m350	0.07256(32)	0.2827(11)	0.04358(42)	0.1698(16)	5.05	0.07066(19)	0.43356
a09m310	0.06051(22)	0.2946(10)	0.03481(34)	0.1695(16)	4.50	0.06947(16)	0.43356
a09m220	0.03307(14)	0.3278(13)	0.01761(15)	0.1746(14)	4.70	0.06724(08)	0.43356
a09m135	0.01346(08)	0.3500(17)	0.00674(05)	0.1752(12)	3.81	0.06608(07)	0.43356
a06m310L	0.06141(35)	0.2993(17)	0.03481(29)	0.1696(14)	6.81	0.02981(04)	0.29985

1. Physical light and strange quark mass limit

The ensembles have a range of light quark masses which correspond roughly to $130 \lesssim m_\pi \lesssim 400$ MeV. We have three lattice spacings at $m_\pi \simeq m_\pi^{\text{phys}}$ such that the light quark mass extrapolation is really an interpolation.

On all but one of the 22 ensembles, the strange quark mass is close to its physical value, allowing us to perform a simple interpolation to the physical strange quark mass point. One ensemble has a strange quark mass of roughly 2/3 its physical value (a12m220ms), allowing us to explore systematics in this strange quark mass inter-

polation.

To parametrize the light and strange quark mass dependence, we utilize two sets of small parameters

$$\Lambda = F : \quad l_F^2 = \frac{m_\pi^2}{\Lambda_\chi^2}, \quad s_F^2 = \frac{2m_K^2 - m_\pi^2}{\Lambda_\chi^2}, \quad (3.1)$$

$$\Lambda = \Omega : \quad l_\Omega^2 = \frac{m_\pi^2}{m_\Omega^2}, \quad s_\Omega^2 = \frac{2m_K^2 - m_\pi^2}{m_\Omega^2}, \quad (3.2)$$

where we have defined

$$\Lambda_\chi \equiv 4\pi F_\pi. \quad (3.3)$$

Using the Gell-Mann–Oakes–Renner relation [65], the numerators in these parameters correspond roughly to the light and strange quark mass, $m_\pi^2 = 2B\hat{m}$ and $2m_K^2 - m_\pi^2 = 2Bm_s$, where $\hat{m} = \frac{1}{2}(m_u + m_d)$. The first set of parameters, Eq. (3.1), is inspired by χ PT and commonly used as a set of small expansion parameters in extrapolating LQCD results. The second set of small parameters, Eq. (3.2), is inspired by Ref. [47]. In Fig. 3, we plot the values of these parameters in comparison with the physical point. Since we are working in the isospin limit in this work, we define the physical point as

$$\begin{aligned} m_\pi^{\text{phys}} &= \bar{M}_\pi = 134.8(3) \text{ MeV}, \\ m_K^{\text{phys}} &= \bar{M}_K = 494.2(3) \text{ MeV}, \\ F_\pi^{\text{phys}} &= F_{\pi^+}^{\text{phys}} = 92.07(57) \text{ MeV}, \\ m_\Omega^{\text{phys}} &= 1672.43(32) \text{ MeV}, \end{aligned} \quad (3.4)$$

with the first three values from the FLAG report [1] and the omega baryon mass from the PDG [66]. The values of $l_{F,\Omega}$ and $s_{F,\Omega}$ are given in Table II for all ensembles.

2. Physical charm quark mass limit

The FNAL and MILC Collaborations have provided a determination of the input value of the charm quark mass that reproduces the ‘‘physical’’ charm quark mass for each of the four lattice spacings used in this work. The mass of the D_s meson was used to tune the input charm quark mass until the physical D_s mass was reproduced (with the already tuned values of the input strange quark mass), defining the ‘‘physical’’ charm quark masses [41],

$$\begin{aligned} a_{15}m_c^{\text{phys}} &= 0.8447(15), & a_{12}m_c^{\text{phys}} &= 0.6328(8), \\ a_{09}m_c^{\text{phys}} &= 0.4313(6), & a_{06}m_c^{\text{phys}} &= 0.2579(4). \end{aligned} \quad (3.5)$$

Comparing to Table I, the simulated charm quark mass is mistuned by less than 2% of the physical charm quark mass for all ensembles used in this work except the a06m310L ensemble, whose simulated charm quark is almost 10% heavier than its physical value.

In order to test the sensitivity of our results to this mistuning of the charm quark mass, we perform a reweighting [67] study of the a06m310L correlation functions and

extracted pion, kaon and omega baryon masses. While the relative shift of the charm quark mass is small, this shift is approximately equal to the value of the physical strange quark mass,

$$\begin{aligned} \delta a_{06}m_c &= a_{06}m_c^{\text{phys}} - a_{06}m_c^{\text{HMC}} \\ &= -0.0281(4). \end{aligned} \quad (3.6)$$

As the reweighting factor is provided by a ratio of the charm quark fermion determinant, it is an extensive quantity, and the relatively large volume we have used to generate the a06m310L ensemble causes some challenges in accurately determining the reweighting factor.

The summary of our study is that our scale setting is not sensitive to this mistuning of the charm quark mass, in line with prior expectation. For example, we find

$$\begin{aligned} a_{06}m_\Omega &= \begin{cases} 0.5069(21), & \text{unweighted} \\ 0.5065(29), & \text{reweighted} \end{cases}, \\ a_{06}(m_\Omega^{\text{reweighted}} - m_\Omega^{\text{unweighted}}) &= -0.0004(34), \end{aligned} \quad (3.7)$$

where the splitting is determined under bootstrap. We provide extensive details in Appendix A.

3. Continuum limit

In order to control the continuum extrapolation, we utilize four lattice spacings ranging from $0.057 \lesssim a \lesssim 0.15$ fm. For most of the pion masses, we have three values of a with four values at $m_\pi \sim 310$ MeV and one value at $m_\pi \sim 180$ MeV. The parameter space is depicted in Fig. 4. The small dimensionless parameter we utilize to extrapolate to the continuum limit is

$$\epsilon_a^2 = \frac{a^2}{(2w_{0,\text{orig}})^2}. \quad (3.8)$$

As noted in Ref. [36], this choice is convenient as the values of ϵ_a^2 span a similar range as l_F^2 . This allows us to test the ansatz of our assumed power counting that treats corrections of $O(l_F^2) = O(\epsilon_a^2)$ which we found to be the case for F_K/F_π [36].

An equally valid way to define the small parameter characterizing the discretization corrections is to utilize the gradient flow scale that is also used to define the observable y being extrapolated. The following normalizations are comparable to our standard choice, Eq. (3.8)

$$\epsilon_a^2 = \begin{cases} \frac{a^2}{(2w_{0,\text{orig}})^2}, & y = w_{0,\text{orig}}m_\Omega \\ \frac{a^2}{(2w_{0,\text{imp}})^2}, & y = w_{0,\text{imp}}m_\Omega \\ \frac{a^2}{4t_{0,\text{orig}}}, & y = \sqrt{t_{0,\text{orig}}}m_\Omega \\ \frac{a^2}{4t_{0,\text{imp}}}, & y = \sqrt{t_{0,\text{imp}}}m_\Omega \end{cases}. \quad (3.9)$$

While these choices do not exhaust the possibilities, they are used to explore possible systematic uncertainties arising from this choice. Unless otherwise noted, the fixed choice in Eq. (3.8) is used in subsequent results and plots.




FIG. 3. The parameter space of s_F^2 versus l_F^2 (left) and s_Ω^2 versus l_Ω^2 used in this calculation. The vertical and horizontal gray lines represent the physical point defined by Eq. (3.4).




FIG. 4. Parameter space of pion mass and lattice spacing utilized in this work expressed in terms of l_F^2 and ϵ_a^2 .

4. Infinite volume limit

The leading sensitivity of m_Ω , t_0 and w_0 to the size of the volume is exponentially suppressed for sufficiently large $m_\pi L$ [68]. We have ensembles with multiple volumes at a15m310, a12m310 and a12m220 to test the predicted finite volume corrections against the observed ones. We derive the predicted volume dependence of $w_0 m_\Omega$ to the first two nontrivial orders in Sec. III B.

A. Light and strange quark mass dependence

The light and strange quark mass dependence of the omega baryon has been derived in SU(3) heavy baryon χ PT (HB χ PT) [69, 70] to next-to-next-to-leading order (N²LO) which is $O(m_{\pi,K,\eta}^4)$ [71–73]. It has been shown that SU(3) HB χ PT does not produce a converging expansion at the physical quark masses [46, 74–77], and so using these formulas to obtain a precise, let alone sub-percent, determination, at the physical pion mass is not

possible when incorporating systematic uncertainties associated with the truncation of SU(3) HB χ PT.

However, many LQCD calculations, including this one, keep the strange quark mass fixed near its physical value. Therefore, a simple interpolation in the strange quark mass is possible. Further, as the omega is an isosinglet, it will have a simpler, and likely more rapidly converging chiral expansion of the light-quark mass dependence than baryons with one or more light valence quarks. This has motivated the construction of an SU(2) HB χ PT for hyperons which considers only the pion as a light degree of freedom [78–82]. In particular, the chiral expansion for the omega baryon mass was determined to $O(m_\pi^6)$ [79]

$$m_\Omega = m_0 + \alpha_2 \frac{m_\pi^2}{\Lambda_\chi} + \frac{m_\pi^4}{\Lambda_\chi^3} [\alpha_4 \lambda_\pi + \beta_4] + \frac{m_\pi^6}{\Lambda_\chi^5} [\alpha_6 \lambda_\pi^2 + \beta_6 \lambda_\pi + \gamma_6], \quad (3.10)$$

where

$$\lambda_\pi = \lambda_\pi(\mu) = \ln \left(\frac{m_\pi^2}{\mu^2} \right), \quad (3.11)$$

α_n , β_n and γ_6 are linear combinations of μ -dependent dimensionless low energy constants (LECs) of the theory, and m_0 is the mass of the omega baryon in the SU(2) chiral limit at the physical strange quark mass. The renormalization group [83] restricts the coefficient of the \ln^2 term to be linearly dependent on α_2 and α_4 , with the relation provided in Ref. [79]. In standard HB χ PT power counting, in which the expansion includes odd powers of the pion mass, this order would be called next-to-next-to-next-to-next-to-leading order (N⁴LO), where leading order (LO) is the $O(m_\pi^2/\Lambda_\chi)$ contribution, next-to-leading order (NLO) would be an $O(m_\pi^3/\Lambda_\chi^2)$ contribution, which vanishes for m_Ω , etc.

The light quark mass dependence for t_0 and w_0 has also been determined in χ PT through $O(m_\pi^4)$ [84] which

is N²LO in the meson chiral power counting. For example

$$w_0 = w_{0,\text{ch}} \left\{ 1 + k_1 \frac{m_\pi^2}{\Lambda_\chi^2} + k_3 \frac{m_\pi^4}{\Lambda_\chi^4} + k_2 \frac{m_\pi^4}{\Lambda_\chi^4} \lambda_\pi \right\}, \quad (3.12)$$

where the LO term, $w_{0,\text{ch}}$, is the value in the chiral limit and the k_i are linear combinations of dimensionless LECs. The expression for t_0 is identical in form and will have different numerical values of the LECs.

From these expressions, we can see both m_Ω and t_0 and w_0 depend only upon even powers of the pion mass through the order we are working: m_Ω receives a chiral correction that scales as $\mathcal{O}(m_\pi^2)$ from a double-sunset two-loop diagram [79] and the next correction to t_0 and w_0 will appear at $\mathcal{O}(m_\pi^6)$. We can multiply these expressions together, Eqs. (3.10) and (3.12), in order to form an expression describing the light-quark mass dependence of $w_0 m_\Omega$. As the characterization of the order of the expansion with respect to the order of m_π^2 is not the same for w_0 and m_Ω , we define the contributions to $w_0 m_\Omega$ as

$$w_0 m_\Omega = c_0 + \delta_{ls,\Lambda}^{\text{NLO}} + \delta_{ls,\Lambda}^{\text{N}^2\text{LO}} + \delta_{ls,\Lambda}^{\text{N}^3\text{LO}}, \quad (3.13)$$

with a similar expression for $\sqrt{t_0} m_\Omega$. We add polynomial terms in $s_{\Lambda,\Omega}^2$ such that

$$\begin{aligned} \delta_{ls,\Lambda}^{\text{NLO}} &= l_\Lambda^2 c_l + s_\Lambda^2 c_s, \\ \delta_{ls,\Lambda}^{\text{N}^2\text{LO}} &= l_\Lambda^4 (c_{ll} + c_{ll}^n \lambda_\pi) + l_\Lambda^2 s_\Lambda^2 c_{ls} + s_\Lambda^4 c_{ss}, \\ \delta_{ls,\Lambda}^{\text{N}^3\text{LO}} &= l_\Lambda^6 (c_{lll} + c_{lll}^n \lambda_\pi + c_{lll}^{n^2} \lambda_\pi^2) + l_\Lambda^4 s_\Lambda^2 \lambda_\pi c_{lls}^n \\ &\quad + l_\Lambda^4 s_\Lambda^2 c_{lls} + l_\Lambda^2 s_\Lambda^4 c_{lss} + s_\Lambda^6 c_{sss}. \end{aligned} \quad (3.14)$$

We will consider both $\Lambda = F$ and $\Lambda = \Omega$ for the two choices of small parameters. For convenience, we set $\mu = \Lambda_\chi$ and $\mu = m_\Omega$ respectively for these choices. For a detailed discussion how one can track the consequence of such a quark mass dependent choice for the dim-reg scale, see Ref. [36].

B. Finite volume corrections

The finite-volume (FV) corrections for m_Ω are determined at one loop through the modification to the tadpole integral [85, 86]

$$\frac{(4\pi)^2}{m^2} \mathcal{I}^{\text{FV}} = \ln \left(\frac{m^2}{\mu^2} \right) + 4k_1(m_\pi L) \quad (3.15)$$

where $k_1(x)$ is given by

$$k_1(x) = \sum_{|\mathbf{n}| \neq 0} c_n \frac{K_1(x|\mathbf{n}|)}{x|\mathbf{n}|}. \quad (3.16)$$

$K_1(x)$ is a modified Bessel function of the second kind and c_n are multiplicity factors for the number of ways the integers (n_x, n_y, n_z) can form a vector of length $|\mathbf{n}|$; see Table III for the first few.

TABLE III. Multiplicity factors for the finite volume corrections of the first ten vector lengths, $|\mathbf{n}|$.

$ \mathbf{n} $	1	$\sqrt{2}$	$\sqrt{3}$	$\sqrt{4}$	$\sqrt{5}$	$\sqrt{6}$	$\sqrt{7}$	$\sqrt{8}$	$\sqrt{9}$	$\sqrt{10}$
c_n	6	12	8	6	24	24	0	12	30	24

At N³LO, the finite volume corrections for m_Ω are also trivially determined, as the only two-loop integral that contributes is a double-tadpole with un-nested momentum integrals; see Fig. 2 of Ref. [79]. The N³LO correction to w_0 is not known. However, the isoscalar nature of w_0 means that at the two-loop order, just like the correction to m_Ω , it will only receive contributions from trivial two-loop integrals with factorizable momentum integrals. Therefore, the N³LO FV correction can also be determined from the square of the tadpole integral

$$\begin{aligned} \frac{(4\pi)^4}{m^4} \left[(\mathcal{I}^{\text{FV}})^2 - (\mathcal{I}^\infty)^2 \right] \\ = 8\lambda_\pi k_1(m_\pi L) + 16k_1^2(m_\pi L), \end{aligned} \quad (3.17)$$

resulting in

$$\begin{aligned} \delta_{L,F}^{\text{N}^2\text{LO}}(l_F, m_\pi L) &= c_{ll}^n l_F^4 4k_1(m_\pi L) \\ \delta_{L,F}^{\text{N}^3\text{LO}}(l_F, m_\pi L) &= c_{lll}^{n^2} l_F^6 \ln(l_F^2) 8k_1(m_\pi L) \\ &\quad + c_{lll}^n l_F^6 16k_1(m_\pi L). \end{aligned} \quad (3.18)$$

The FV correction through N³LO arising from loop corrections to w_0 and m_Ω is given by

$$\delta_{L,F} = \delta_{L,F}^{\text{N}^2\text{LO}} + \delta_{L,F}^{\text{N}^3\text{LO}}, \quad (3.19)$$

with a similar expression for $\delta_{L,\Omega}$.

We are neglecting a few FV corrections at $\delta_{L,F}^{\text{N}^3\text{LO}}$. The NLO $l_{F,\Omega}^2$ correction to the omega mass is from a quark mass operator, which has been converted to an m_π^2 correction, $2B\hat{m}_l = m_\pi^2 + \mathcal{O}(m_\pi^4/\Lambda_\chi^2)$. This choice for organizing the perturbative expansion induces corrections in what we have called N²LO and N³LO. At N²LO, the corrections arise from single tadpole diagrams and so the FV corrections are accounted for through Eq. (3.18). At the next order, the corrections to the pion self-energy involve more complicated two-loop diagrams [87] and so the FV corrections arising from these are not captured in our parametrization. Similar corrections arise from expressing $4\pi F_0 = 4\pi F_\pi + \mathcal{O}(m_\pi^2/\Lambda_\chi^2)$ when using l_F^2 to track the light-quark mass corrections [F_0 is F_π in the SU(2) chiral limit].

While we have neglected these contributions, the FV corrections to m_Ω are suppressed by an extra power in the chiral power counting compared with many observables, beginning with an m_π^4/Λ_χ^4 prefactor, Eq. (3.18). In Fig. 5, we show the predicted FV correction along with the results at three volumes on the a12m220 ensembles. As can be observed, the predicted FV corrections are very small and consistent with the numerical results.




FIG. 5. Predicted finite volume corrections (band) compared with the results on the a12m220 ensembles. The $N^3\text{LO}$ result is from Eq. (3.19) using l_F^2 as the small parameter.

C. Discretization corrections

A standard method of incorporating discretization effects into the extrapolation formula used for hadronic observables is to follow the strategy of Sharpe and Singleton [88]:

1. For a given lattice action, one first constructs the Symanzik effective theory (SET) by expanding the discretized action about the continuum limit. This results in a local effective action in terms of quark and gluon fields [89, 90];
2. With this continuum effective theory, one builds a chiral effective theory by using spurion analysis to construct not only operators with explicit quark mass dependence, but also operators with explicit lattice spacing dependence.

Such an approach captures the leading discretization effects in a local hadronic effective theory. At the level of the SET, radiative corrections generate logarithmic dependence upon the lattice spacing. The leading corrections can be resummed such that, for an $O(a)$ improved action, the leading discretization effects then scale as $a^2 \alpha_S^{n+\hat{\gamma}_1}$ [91, 92] where $n = 0$ for an otherwise unimproved action, $n = 1$ for a tree-level improved action and $n = 2$ for a one-loop improved action. The coefficient $\hat{\gamma}_1$ is an anomalous dimension which has been determined for Yang-Mills and Wilson actions [93].¹

For mixed-action setups [20, 21] such as the one used in this work, a low-energy mixed-action effective field theory (MAEFT) [21, 94–102] can be constructed to capture the manifestation of infrared radiative corrections from the

discretization². Corrections come predominantly from a modification of the pseudoscalar meson spectrum as well as from “hairpin” interactions [104] that are proportional to the lattice spacing in rooted-staggered [105] and mixed-action theories [96]; in partially quenched theories, these hairpins are proportional to the difference in the valence and sea quark masses [106–108].

In our analysis of F_K/F_π , we observed that the use of continuum chiral perturbation theory with corrections polynomial in ϵ_a^2 was highly favored over the use of the MAEFT expression, as measured by the Bayes-Factor, though the results from both were consistent within a fraction of 1 standard deviation [36]. Similar findings have been observed by other groups for various quantities; see for example Refs. [109–111]. Therefore, in this work, we restrict our analysis to a continuum-like expression enhanced by polynomial discretization terms.

The dynamical HISQ ensembles have a perturbatively improved action such that the leading discretization effects (before resumming the radiative corrections [91–93]) scale as $O(\alpha_S a^2)$ [26]. The MDWF action, in the limit of infinite extent in the fifth dimension, has no chiral symmetry breaking other than that from the quark mass. Consequently, the leading discretization corrections begin at $O(a^2)$ [112, 113]. For finite L_5 , the $O(a)$ corrections are proportional to am_{res} which is sufficiently small that these terms are numerically negligible. Therefore, we parametrize our discretization corrections with the following terms where we count $\epsilon_a^2 \sim l_\Lambda^2 \sim s_\Lambda^2$:

$$\begin{aligned} \delta_{a,\Lambda} &= \delta_{a,\Lambda}^{\text{NLO}} + \delta_{a,\Lambda}^{\text{N}^2\text{LO}} + \delta_{a,\Lambda}^{\text{N}^3\text{LO}}, \\ \delta_{a,\Lambda}^{\text{NLO}} &= d_a \epsilon_a^2 + d'_a \alpha_S \epsilon_a^2, \\ \delta_{a,\Lambda}^{\text{N}^2\text{LO}} &= d_{aa} \epsilon_a^4 + \epsilon_a^2 (d_{al} l_\Lambda^2 + d_{as} s_\Lambda^2), \\ \delta_{a,\Lambda}^{\text{N}^3\text{LO}} &= d_{aaa} \epsilon_a^6 + \epsilon_a^4 (d_{aal} l_\Lambda^2 + d_{aas} s_\Lambda^2) \\ &\quad + \epsilon_a^2 (d_{all} l_\Lambda^4 + d_{als} l_\Lambda^2 s_\Lambda^2 + d_{ass} s_\Lambda^4). \end{aligned} \quad (3.20)$$

IV. EXTRAPOLATION DETAILS AND UNCERTAINTY ANALYSIS

We perform our extrapolation analysis under a Bayesian model-averaging framework as described in detail in Refs. [33, 36, 114], which is more extensively discussed for lattice QFT analysis in Ref. [58]. We consider a variety of extrapolation functions by working to different orders in the power counting, using the l_F, s_F or l_Ω, s_Ω small parameters, by including or excluding the chiral logarithms associated with pion loops, and by including

¹ See also the presentation by N. Husung at the MIT Virtual Lattice Field Theory Colloquium Series, <http://ctp.lns.mit.edu/latticecolloq/>.

² While this might seem counterintuitive, it is analogous to the infrared sensitivity of hadronic quantities to the Higgs vacuum expectation value (vev): hadronic quantities have infrared (logarithmic) sensitivity to the pion mass from radiative pion loops, and the squared pion mass is proportional to the light quark mass which is proportional to the Higgs vev [103].

or excluding discretization corrections scaling as $\alpha_S a^2$. The resulting Bayes factors are then used to weight the fits with respect to each other and perform a model averaging. In this section, we discuss the selection of the priors for the various LECs and then present an uncertainty analysis of the results.

A. Prior widths of LECs

In our F_K/F_π analysis [36], we observed that using $\epsilon_\pi^2 = l_F^2$, ϵ_K^2 and ϵ_a^2 as the small parameters in the expansion,³ the LECs were naturally of $O(1)$. We therefore have a prior expectation that this may hold for $\sqrt{t_0}m_\Omega$ and w_0m_Ω as well.

Let us use $w_{0,\text{orig}}m_\Omega$ to guide the discussion. The $a \sim 0.12$ fm ensembles were simulated with a fixed strange quark mass. Therefore, the entire change in $w_{0,\text{orig}}m_\Omega$ between the a12m130 and a12m180L ensembles can be attributed to the change in l_F . This allows us to “eyeball” the c_l prior to be $c_l \simeq 1$ if we assume the dominant contribution comes from the NLO $c_l l_F^2$ term.⁴ Motivated by SU(3) flavor symmetry considerations, we can roughly expect $c_s \sim c_l$. In order to be conservative, we set the prior for these LECs as

$$\tilde{c}_l = \tilde{c}_s = N(\mu = 1, \sigma = 1), \quad (4.1)$$

where $N(\mu, \sigma)$ denotes a normal distribution with mean μ and width σ . A similar observation is made for $w_{0,\text{imp}}m_\Omega$ and the original and improved values of $\sqrt{t_0}m_\Omega$. We observe (with a full analysis) that the log-Bayes-Factor (logGBF) prefers even tighter priors, with logGBF continuing to increase as the width is taken down to 0.1 on these NLO LECs.⁵

The observation that m_Ω increases with increasing values of l_F and s_F (normalized by any and all gradient flow scales) allows us to conservatively estimate the LO prior,

$$\tilde{c}_0 = N(1, 1). \quad (4.2)$$

We then conservatively estimate the priors for all of the higher order l_F and s_F LECs to be

$$\tilde{c}_i = N(0, 1). \quad (4.3)$$

We observe, with a full analysis, that this choice is near the optimal value as measured by the logGBF weighting.

For the discretization corrections, see Fig. 7 in Sec. IV B, as we change the gradient flow scale from

$w_{0,\text{orig}}$ to the improved version to using the original and improved versions of $\sqrt{t_0}$, the approach to the continuum limit can change sign. We also observe, the convexity of the approach to the continuum limit (the ϵ_a^4 contributions) can change sign. Therefore, we perform a prior-optimization study for the discretization LECs in which we change the prior width of the NLO and N²LO LECs in concert, such that the priors are given by

$$\tilde{d}_i = N(0, \sigma_a^{\text{opt}}), \quad (4.4)$$

with the optimized values

$$\sigma_a^{\text{opt}} = \begin{cases} 1.2, & w_{0,\text{orig}}m_\Omega \\ 1.4, & w_{0,\text{imp}}m_\Omega \\ 1.8, & \sqrt{t_{0,\text{orig}}}m_\Omega \\ 1.4, & \sqrt{t_{0,\text{imp}}}m_\Omega \end{cases}, \quad (4.5)$$

maximizing the respective logGBF factors. We set the N³LO priors for discretization LECs to

$$\tilde{d}_i = N(0, 1). \quad (4.6)$$

Although we find using tighter priors increases the logGBF, the final results are unchanged.

When we add the $\alpha_S \epsilon_a^2$ term in the analysis, this introduces a fourth class of discretization corrections. As we only have four lattice spacings in this work, we perform an independent prior-width optimization for this LEC. For all four choices of gradient-flow scales, we find the choice,

$$\tilde{d}'_a = N(0, 0.5), \quad (4.7)$$

to be near-optimal, with three of the analyses preferring an even tighter prior width. An empirical Bayes study [57] in which the widths of all the chiral and all the discretization priors are varied together at a given order leads to similar choices of all the priors.

In Table IV we list the values of all the priors used in the final analysis. The full analysis demonstrates that these choices also result in no tension between the priors and the final posterior values of the LECs, further indication that our choices are reasonable.

When we use l_Ω^2 and s_Ω^2 as the small parameters instead of l_F^2 and s_F^2 , we note that since $(m_\Omega/\Lambda_\chi)^2 \sim 2$, we can use the same prescription, except to double the mean and width of all the NLO priors (which scale linearly in l_Ω^2 and s_Ω^2), set the widths to be 4 times larger for the N²LO priors and 8 times larger for the N³LO LECs. The mixed contributions which scale with some power of ϵ_a^2 and l_Ω^2 and s_Ω^2 are scaled accordingly; see also Table IV.

B. Extrapolation analysis

For each of the four quantities, $w_{0,\text{orig}}m_\Omega$, $w_{0,\text{imp}}m_\Omega$, $\sqrt{t_{0,\text{orig}}}m_\Omega$ and $\sqrt{t_{0,\text{imp}}}m_\Omega$, we consider several reasonable choices of extrapolation functions to perform the

³ The small parameter $\epsilon_K^2 = \frac{m_K^2}{\Lambda_\chi^2} = \frac{1}{2}(s_F^2 + l_F^2)$.

⁴ This is analogous to using the effective mass and effective overlap factors to choose conservative priors for the ground state parameters in the correlation function analysis.

⁵ For fixed data, $\exp\{\log\text{GBF}\}$ provides a relative weight of the likelihood of one model versus another.

TABLE IV. The values of the priors used in the analysis.

LEC	$\Lambda = F$	$\Lambda = \Omega$
c_0	$N(1, 1)$	$N(1, 1)$
c_l, c_s	$N(1, 1)$	$N(2, 2)$
$c_{ll}, c_{ll}^{\text{ln}}, c_{ls}, c_{ss}$	$N(0, 1)$	$N(0, 4)$
$c_{lll}, c_{lll}^{\text{ln}}, c_{lll}^{\text{ln}^2}, c_{lls}, c_{lls}^{\text{ln}}, c_{lss}, c_{sss}$	$N(0, 1)$	$N(0, 8)$
d_a, d_{aa}	$N(0, \sigma_a^{\text{opt}})$	$N(0, \sigma_a^{\text{opt}})$
d'_a	$N(0, 0.5)$	$N(0, 0.5)$
d_{al}, d_{as}	$N(0, \sigma_a^{\text{opt}})$	$N(0, 2\sigma_a^{\text{opt}})$
d_{aaa}	$N(0, 1)$	$N(0, 1)$
d_{aal}, d_{aas}	$N(0, 1)$	$N(0, 2)$
$d_{all}, d_{als}, d_{ass}$	$N(0, 1)$	$N(0, 4)$

continuum, infinite volume and physical quark-mass limits. The final result for each extrapolation is then determined through a model average in which the relative weight of each model is given by the exponential of the

corresponding logGBF value. The various choices we consider in the extrapolations consists of

Include the $\ln(m_\pi)$ terms or counterterm only :	$\times 2$
Expand to N ² LO, or N ³ LO :	$\times 2$
Include/exclude finite volume corrections :	$\times 2$
Include/exclude the $\alpha_S a^2$ term :	$\times 2$
Use the $\Lambda = F$ or $\Lambda = \Omega$ expansion :	$\times 2$
total choices :	32

We find that there is very little dependence upon the particular model chosen. In Fig. 6, we show the stability of the final result of $\sqrt{t_{0,\text{imp}}}m_\Omega$ and $w_{0,\text{imp}}m_\Omega$ as various options from the above list are turned on and off. In addition, we show the impact of including or excluding the a12m220ms ensemble, whose strange quark mass is $m_s \sim 0.6 \times m_s^{\text{phys}}$, as well as the impact of including the a06m310L ensemble. We observe a small variation of the result when either of these ensembles is dropped, but the results are still consistent with our final result (top of the figure). Using the fixed definition of ϵ_a^2 , Eq. (3.8), we find

$$\begin{aligned}
\sqrt{t_{0,\text{orig}}}m_\Omega &= 1.2107(78)^s(14)^\chi(40)^a(00)^V(22)^{\text{phys}}(07)^M \rightarrow \frac{\sqrt{t_0}}{\text{fm}} = 0.1429(09)^s(02)^\chi(05)^a(00)^V(03)^{\text{phys}}(01)^M, \\
\sqrt{t_{0,\text{imp}}}m_\Omega &= 1.2052(73)^s(13)^\chi(40)^a(00)^V(22)^{\text{phys}}(08)^M \rightarrow \frac{\sqrt{t_0}}{\text{fm}} = 0.1422(09)^s(02)^\chi(05)^a(00)^V(03)^{\text{phys}}(01)^M, \\
w_{0,\text{orig}}m_\Omega &= 1.4481(81)^s(14)^\chi(46)^a(00)^V(26)^{\text{phys}}(11)^M \rightarrow \frac{w_0}{\text{fm}} = 0.1709(10)^s(02)^\chi(05)^a(00)^V(03)^{\text{phys}}(01)^M, \\
w_{0,\text{imp}}m_\Omega &= 1.4467(83)^s(15)^\chi(44)^a(00)^V(24)^{\text{phys}}(10)^M \rightarrow \frac{w_0}{\text{fm}} = 0.1707(10)^s(02)^\chi(05)^a(00)^V(03)^{\text{phys}}(01)^M, \quad (4.8)
\end{aligned}$$

with the statistical (s), chiral interpolation (χ), continuum-limit (a), infinite-volume (V), physical-point (phys), and model selection uncertainties (M). The conversion to physical units is performed with Eq. (3.4).

As discussed in Sec. III 3, we explore potential systematics in the use of one definition of the small param-

eter used to characterize the discretization corrections, Eq. (3.8), versus another equally valid choice, Eq. (3.9). For each choice of w_0m_Ω and $\sqrt{t_0}m_\Omega$ that is extrapolated to the physical point, we repeat the above model averaging procedure, but we also include the variation of using these two definitions of ϵ_a^2 for which we find

$$\begin{aligned}
\sqrt{t_{0,\text{orig}}}m_\Omega &= 1.2050(87)^s(16)^\chi(46)^a(00)^V(21)^{\text{phys}}(61)^M \rightarrow \frac{\sqrt{t_0}}{\text{fm}} = 0.1422(10)^s(02)^\chi(05)^a(00)^V(02)^{\text{phys}}(07)^M, \\
\sqrt{t_{0,\text{imp}}}m_\Omega &= 1.2052(77)^s(14)^\chi(46)^a(00)^V(21)^{\text{phys}}(10)^M \rightarrow \frac{\sqrt{t_0}}{\text{fm}} = 0.1422(09)^s(02)^\chi(05)^a(00)^V(02)^{\text{phys}}(01)^M, \\
w_{0,\text{orig}}m_\Omega &= 1.4481(81)^s(14)^\chi(46)^a(00)^V(26)^{\text{phys}}(11)^M \rightarrow \frac{w_0}{\text{fm}} = 0.1709(10)^s(02)^\chi(05)^a(00)^V(03)^{\text{phys}}(01)^M, \\
w_{0,\text{imp}}m_\Omega &= 1.4485(82)^s(15)^\chi(44)^a(00)^V(25)^{\text{phys}}(18)^M \rightarrow \frac{w_0}{\text{fm}} = 0.1709(10)^s(02)^\chi(05)^a(00)^V(03)^{\text{phys}}(02)^M. \quad (4.9)
\end{aligned}$$

For all choices of the gradient-flow scale besides $\sqrt{t_{0,\text{orig}}}$,

the average over the choice of how to define ϵ_a^2 has mini-




FIG. 6. Model breakdown and comparison using the scale-improved, gradient-flow derived quantities (i.e., $\sqrt{t_{0,\text{imp}}}m_\Omega$, $w_{0,\text{imp}}m_\Omega$). The vertical band is our model average result to guide the eye. **$\chi\text{pt-full}$** : χPT model average, including $\ln(m_\pi^2/\mu^2)$ corrections. **$\chi\text{pt-ct}$** : χPT model average with counterterms only, excluding $\ln(m_\pi^2/\mu^2)$ corrections. **$\text{N}^3\text{LO}/\text{N}^2\text{LO}$** : model average restricted to specified order. **Λ_χ** : model average with specified chiral cutoff. **incl./excl. α_S** : model average with/without α_S corrections. **fixed/variable ϵ_a^2** : model average with Eqs. (3.8) or (3.9), respectively. **excl. a06m310L**: model average excluding $a = 0.06$ fm ensemble (a06m310L). **excl. a12m220ms**: model average excluding small strange quark mass ensemble (a12m220ms). **Below solid line**: results from other collaborations for various numbers of dynamical fermions: BMWc [2020] [14], MILC [2015] [12], HPQCD [2013] [8], CLS [2017] [13], QCDSF-UKQCD [2015] [9], RBC [2014] [11], HotQCD [2014] [10], BMWc [2012] [6] and ALPHA [2013] [7].

mal impact on the final result, as can be seen comparing Eq. (4.8) and (4.9). In the case of $\sqrt{t_{0,\text{orig}}}$, the two choices for the cutoff-effect expansion parameter lead to a slight difference in the continuum-extrapolated value, which is reflected in the model-averaging uncertainty.⁶ In all

⁶ Rather than performing a model average over the two definitions of ϵ_a^2 as defined in Eqs. (3.8) and (3.9), one might instead consider a model average over the choices for fixed ϵ_a^2 , i.e.

$$\epsilon_a^2 = \left\{ \left(\frac{a}{2w_{0,\text{orig}}} \right)^2, \left(\frac{a}{2w_{0,\text{impr}}} \right)^2, \frac{a^2}{4t_{0,\text{orig}}}, \frac{a^2}{4t_{0,\text{impr}}} \right\}.$$

Performing the model average in this manner instead yields

cases, the dominant uncertainty is statistical, suggesting a straightforward path to reducing the uncertainty to a few per-mille.

To arrive at our final determination of $\sqrt{t_0}$, Eq. (1.1) and w_0 , Eq. (1.2), we perform an average of the results in Eq. (4.9). As the data between the two choices differ slightly, we can not perform this final averaging step under the Bayes model-averaging procedure; instead we treat each result with equal weight. We would add half

results for $\sqrt{t_0}, w_0$ consistent within a fraction of a sigma of Eqs. (1.1) and (1.2).




FIG. 7. Example extrapolations of $\sqrt{t_{0,\text{orig}}}m_\Omega$ and $w_{0,\text{orig}}m_\Omega$ versus l_F^2 (top) and ϵ_a^2 (bottom) using the N³LO χ PT analysis. For the continuum extrapolation plots (bottom) we also show the results using the “improved” determinations of the scales, Eq. (2.3). The numerical results have been shifted from the original data points as described in the text.

the difference between the central values as an additional discretization uncertainty, but as is evident from Eq. (4.9), the central values are essentially the same.

In Fig. 6, we also compare our result with other values in the literature. All the results, except the most recent one from BMWc [14], have been determined in the isospin symmetric limit. Our results are in good agreement with the more recent and precise results, though one notes, there is some tension in the values of $\sqrt{t_0}$ and w_0 reported.

In Fig. 7, we show the resulting extrapolation of $\sqrt{t_{0,\text{orig}}}m_\Omega$ and $w_{0,\text{orig}}m_\Omega$ projected into the l_F^2 plane using the N³LO analysis including the $\ln(m_\pi)$ type corrections. The finite lattice spacing bands are plotted with a value of ϵ_a^2 taken from the near-physical pion mass ensembles from Table II, a15m135XL, a12m130 and a09m135. For the $a \sim 0.06$ fm band, we use the value,

$$\frac{w_{0,\text{orig}}}{a_{06}} = 3.0119(19), \quad (4.10)$$

from Table IV of Ref. [12] with $m'_l/m'_s = 1/27$ and use this to construct ϵ_a^2 . The data points are plotted after being shifted to the extrapolated values of all the parameters using the posterior values of the LECs from the N³LO fit.

The lower panel of Fig. 7 is similarly constructed by

shifting all the data points to the infinite volume limit, l_F^{phys} , s_F^{phys} and the value of ϵ_a^2 from the particular ensemble with the corresponding band in this same limit and only varying ϵ_a^2 . We plot the continuum extrapolation of both the original and improved values to demonstrate the impact of the improvement at finite lattice spacing, noting the agreement in the continuum limit.

For w_0 , there is very little difference between the original and improved values with very similar continuum extrapolations. In contrast, there is a striking difference between the original and improved values using $\sqrt{t_0}$, though they agree in the continuum limit. We also observe that the use of $\sqrt{t_{0,\text{orig}}}$ is susceptible to larger model-extrapolation uncertainties arising from different choices of parametrizing the continuum extrapolation; see Eq. (4.9). Additional results at $a \lesssim 0.06$ fm will be required to control the continuum extrapolation using $\sqrt{t_0}$ in order to obtain a few-per-mille level of precision.

C. Interpolation of t_0 and w_0

With our determination of t_0 and w_0 , Eqs. (1.1) and (1.2), we can determine the lattice spacing for each bare coupling. We could use the near-physical pion mass

ensemble values of the gradient-flow scales, or alternatively, we could interpolate the results to the physical quark mass point using the predicted quark mass dependence [84]. The interpolation can be performed for each lattice spacing separately, or in a combined analysis of all lattice spacings simultaneously. The latter is preferable in order for us to determine the lattice spacing a_{06} as we only have results at a single pion mass at this lattice spacing. To perform the global analysis, we use an N²LO extrapolation function (which has the same form for t_0),

$$\frac{w_0}{a} = \frac{w_{0,\text{ch}}}{a} \left\{ 1 + k_l l_F^2 + k_s s_F^2 + k_a \epsilon_{a,\text{ch}}^2 + k_{ll} l_F^4 + k_{lln} l_F^4 \ln(l_F^2) + k_{l_s} l_F^2 s_F^2 + k_{ss} s_F^4 + k_{aa} \epsilon_{a,\text{ch}}^4 + k_{al} l_F^2 \epsilon_{a,\text{ch}}^2 + k_{as} s_F^2 \epsilon_{a,\text{ch}}^2 \right\}, \quad (4.11)$$

$$\epsilon_{a,\text{ch}} \equiv \frac{1}{(2w_{0,\text{ch}}/a)}. \quad (4.12)$$

This global analysis treats the value of LO parameter $\frac{w_{0,\text{ch}}}{a}$ for each lattice spacing as a separate unknown parameter, and then assumes that the remaining dimensionless LECs are shared between all lattice spacings. We use this LO parameter to also construct $\epsilon_{a,\text{ch}}$ which controls the discretization corrections rather than using ϵ_a , as ϵ_a is half the inverse of the left-hand side of Eq. (4.11).

It is tempting to think of this as a combined chiral and continuum extrapolation analysis of w_0 , but it is not as one normally thinks of them. Because we do not know the lattice spacings already, there remains an ambiguity in the interpretation of $w_{0,\text{ch}}/a$ and the LECs k_a , k_{aa} , k_{la} and k_{sa} : we are not able to interpret $w_{0,\text{ch}}/a$ as the chiral limit value of w_0 divided by the lattice spacing. We perform this analysis for all four gradient-flow scales with independent LECs for each scale as well as a similar parametrization of the $\epsilon_{a,\text{ch}}$ parameter.

When we perform the interpolation for each lattice spacing separately, we utilize this same expression except that we set all parameters proportional to any power of $\epsilon_{a,\text{ch}}$ to zero. When the individual interpolations are used, the resulting values of a_{15} , a_{12} and a_{09} are compatible with those from the global analysis well within 1 standard deviation. In Fig. 8, we show an example interpolation of $w_{0,\text{orig}}/a$ using the global analysis of all ensembles to the l_F^{phys} and s_F^{phys} point for each lattice spacing. The open circles show the raw values of w_0/a while the open squares show the values shifted to $\frac{w_0}{a}(l_F^{\text{ens}}, s_F^{\text{phys}}, \epsilon_a^{\text{ens}})$ using the resultant parameters determined in the global analysis.

A quark-mass independent determination of the lattice spacing can be made by using the determination of $\sqrt{t_0}$, Eq. (1.1) or w_0 , Eq. (1.2) at the physical point, combined with the physical-quark mass interpolated values of t_0/a^2 or w_0/a from either the original or improved values of these gradient flow scales. Each choice represents a different scheme for setting the lattice spacing. The continuum extrapolated value of some observable quantity,




FIG. 8. Interpolation of w_0/a values to the physical l_F and s_F point. The open circles are the raw data from Table II and the open squares have been shifted to the infinite volume and s_F^{phys} point. The vertical gray line is at $l_F^{2,\text{phys}}$.

using any of these schemes, should agree in the continuum limit, while at finite lattice spacing, the results can be substantially different, as is evident in Fig. 7.

In Table V, we provide the determination of the lattice spacing for each bare coupling, expressed in terms of the approximate lattice spacing. It is interesting to note that the determination of the lattice spacing with $\sqrt{t_{0,\text{orig}}}$ is substantially different than with the other gradient-flow scales, while the scale determined with from the three remaining auxiliary scales are very similar.

V. SUMMARY AND DISCUSSION

We have performed a precise scale setting with our MDWF on gradient-flowed HISQ action [15] achieving a total uncertainty of $\sim 0.6\%$ – 0.8% for each lattice spacing, Table V. The scale setting was performed by extrapolating the quantities $\sqrt{t_0} m_\Omega(l_F, s_F, \epsilon_a, m_\pi L)$ and $w_0 m_\Omega(l_F, s_F, \epsilon_a, m_\pi L)$ to the continuum ($\epsilon_a \rightarrow 0$), infinite volume ($m_\pi L \rightarrow \infty$) and physical quark mass limits ($l_F \rightarrow l_F^{\text{phys}}$ and $s_F \rightarrow s_F^{\text{phys}}$), and using the experimental determination of m_Ω to determine the scales $\sqrt{t_0}$ and w_0 in fm. The values of $\sqrt{t_{0,\text{orig}}}/a$, $\sqrt{t_{0,\text{imp}}}/a$, $w_{0,\text{orig}}/a$

TABLE V. The value of the lattice spacing in fm using the interpolation of the four different determinations of the gradient-flow scales, combined with the determination of $\sqrt{t_0}$ and w_0 at the physical point, Eq. (1.1) and Eq. (1.2) respectively. Any one scheme for determining the lattice spacing can be picked as a quark-mass independent scale setting that can be used to convert results from dimensionless lattice units to MeV. The different schemes will result in different approaches to the continuum; however all choices should agree in the continuum limit.

Scheme	a_{15}/fm	a_{12}/fm	a_{09}/fm	a_{06}/fm
$t_{0,\text{orig}}/a^2$	0.1284(10)	0.10788(83)	0.08196(64)	0.05564(44)
$t_{0,\text{imp}}/a^2$	0.1428(10)	0.11735(87)	0.08632(65)	0.05693(44)
$w_{0,\text{orig}}/a$	0.1492(10)	0.12126(87)	0.08789(71)	0.05717(51)
$w_{0,\text{imp}}/a$	0.1505(10)	0.12066(88)	0.08730(70)	0.05691(51)

and $w_{0,\text{imp}}/a$ were interpolated to the infinite volume and physical quark mass limits for each lattice spacing, allowing for the quark-mass independent determination of a for each bare coupling β , expressed in terms of the approximate lattice spacing; see Table V.

Of note, the approach to the continuum limit of $\sqrt{t_{0,\text{orig}}}m_\Omega$ and $\sqrt{t_{0,\text{imp}}}m_\Omega$ are quite different, Fig. 7, with the use of $\sqrt{t_{0,\text{imp}}}$ leading to an almost flat continuum extrapolation. The two different extrapolations agree quite nicely in the continuum limit, as they must if all systematic uncertainties are under control. In contrast, the use of the original and improved values of w_0 leads to very similar continuum extrapolations of $w_{0,\text{orig}}m_\Omega$ and $w_{0,\text{imp}}m_\Omega$, which also agree very nicely in the continuum limit.

We also observe that the use of l_Ω and s_Ω as small parameters to control the quark-mass interpolation are relatively heavily penalized as compared to the use of l_F and s_F ; see Table VII for an example. We observe the same qualitative weighting with all choices of the gradient-flow scale. Perhaps this is an indication that this parametrization is suboptimal.

Our final uncertainty using w_0 is dominated by the stochastic uncertainty, Eq. (1.2), providing a clear path to reducing the uncertainty by almost a factor of 3 before an improved understanding of the various systematic uncertainties becomes relevant. At such a level of precision, a systematic study of the effect of isospin breaking on the scale setting, as has been performed by BMWc [14], is likely required to retain full control of the uncertainty. For $\sqrt{t_0}$, we observe the model-selection uncertainty is comparable to the stochastic uncertainty, Eq. (1.1), which arises from the different ways to parameterize the continuum extrapolation; see Eq. (4.9). Therefore, additional results at $a \lesssim 0.06$ fm will be required to obtain a fer-per-mille precision with $\sqrt{t_0}$.

The pursuit of our physics program of determining the nucleon elastic structure functions and improving the precision of our g_A result [33, 34] will naturally lead to an improved scale setting precision. The current precision is already expected to be subdominant for most of the

results we will obtain, but a further improved precision is welcome.

ACKNOWLEDGMENTS

We thank M. Goltermann and O. Bär for useful discussion and correspondence about the chiral corrections to w_0 . We thank Andrea Shindler for helpful discussions regarding reweighting. We thank Peter Lepage for helpful correspondence regarding `lsqfit` and the log Gaussian Bayes factor. We thank K. Orginos for use of `wm.chroma` that was used to compute some of the correlation functions used in this work. We thank the MILC Collaboration for providing some of the HISQ configurations used in this work, and A. Bazavov, C. Detar and D. Toussaint for guidance on using their code to generate the new HISQ ensembles also used in this work. We thank R. Sommer for helpful correspondence and encouragement to determine t_0 as well as w_0 .

Computing time for this work was provided through the Innovative and Novel Computational Impact on Theory and Experiment (INCITE) program and the LLNL Multiprogrammatic and Institutional Computing program for Grand Challenge allocations on the LLNL supercomputers. This research utilized the NVIDIA GPU-accelerated Titan and Summit supercomputers at Oak Ridge Leadership Computing Facility at the Oak Ridge National Laboratory, which is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC05-00OR22725 as well as the Surface, RZHasGPU, Pascal, Lassen, and Sierra supercomputers at Lawrence Livermore National Laboratory.

The computations were performed utilizing LALIBE [115] which utilizes the Chroma software suite [52] with QUDA solvers [18, 19] and HDF5 [116] for I/O [117]. They were efficiently managed with METAQ [118, 119] and status of tasks logged with EspressoDB [120]. The HMC was performed with the MILC Code [60], and for the ensembles new in this work, running on GPUs using QUDA. The final extrapolation analysis utilized `gvar` v11.2 [121] and `lsqfit` v11.5.1 [122]. The analysis and data for this work can be found at this git repo: https://github.com/callat-qcd/project_scale_setting_mdwf_hisq.

This work was supported by the NVIDIA Corporation (MAC), the Alexander von Humboldt Foundation through a Feodor Lynen Research Fellowship (CK), the DFG and the NSFC Sino-German CRC110 (EB), the RIKEN Special Postdoctoral Researcher Program (ER), the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under Awards No. DE-AC02-05CH11231 (CCC, CK, BH, AWL), No. DE-AC52-07NA27344 (DAB, DH, ASG, PV), No. DE-FG02-93ER-40762 (EB), No. DE-AC05-06OR23177 (CM); the Nuclear Physics Double Beta Decay Topical Collaboration (DAB, HMC, AN, AWL); the U.K. Science and Technology Facilities Council Grants No. ST/S005781/1 and

No. ST/T000945/1 (CB); and the DOE Early Career Award Program (CCC, AWL).

Appendix A: CHARM QUARK MASS REWEIGHTING

The use of reweighting [67] to estimate a correlation function with a slightly different sea-quark mass than the one simulated with is very common in LQCD; see for example Refs. [11, 123, 124]. A nice discussion of mass reweighting, including single flavor reweighting is found in Refs. [125, 126].

In our case, we are interested in reweighting the computation from the hybrid Monte Carlo (HMC) simulated charm quark mass, m_c^{HMC} , to the physical quark mass, m_c^{phys} which requires an estimate of the ratio of the fermion determinant with the physical mass to the determinant with the HMC mass. If the mass shift is a $\delta m_c = m_c^{\text{phys}} - m_c^{\text{HMC}}$ then up to $\mathcal{O}(\delta m^2)$ this ratio can be written (including the quarter-root arising from rooted-staggered fermions)

$$w_i^4 = \det [\mathbb{1} + \delta m (D[U_i] + m_c^{\text{phys}})^{-1}] \quad (\text{A1})$$

for each configuration U_i and observables may be computed using the weight w ,

$$\langle O \rangle_{\text{phys}} = \frac{\langle Ow \rangle_{\text{HMC}}}{\langle w \rangle_{\text{HMC}}}. \quad (\text{A2})$$

We can use two methods to stochastically estimate w for each configuration. First, by rewriting the determinant as the exponential of a trace-log, one finds

$$w_i^{\text{tr}} = \exp \frac{\delta m}{4} \text{tr} [(D[U_i] + m_c^{\text{phys}})^{-1}] + \mathcal{O}(\delta m^2) \quad (\text{A3})$$

and we can use vectors of complex Gaussian noise η to estimate the trace,

$$\text{tr} [(D[U_i] + m_c^{\text{phys}})^{-1}] \approx \frac{1}{N_\eta} \sum_i \eta_i^\dagger (D[U_i] + m_c^{\text{phys}})^{-1} \eta_i, \quad (\text{A4})$$

$$\eta \sim \frac{1}{\pi^V} \exp -\eta^\dagger \eta, \quad (\text{A5})$$

where V is the size of each η vector.

Alternatively, we may estimate the determinant in the reweighting factor (A1) using the identity,

$$\det A = \pi^{-V} \int \mathcal{D}\eta \mathcal{D}\eta^\dagger e^{-\eta^\dagger A^{-1} \eta}, \quad (\text{A6})$$

which is often used to implement pseudofermions. Up to $\mathcal{O}(\delta m^2)$, this becomes

$$w_i^4 = \pi^{-V} \int \mathcal{D}\eta \mathcal{D}\eta^\dagger e^{-\eta^\dagger \eta} e^{\delta m \eta^\dagger (D[U_i] + m_c^{\text{phys}})^{-1} \eta}, \quad (\text{A7})$$

which tells us to draw η according to the same Gaussian (A5) and estimate

$$w_i^{\text{ps}} = \left(\frac{1}{N_\eta} \sum_j e^{\delta m \eta_j^\dagger (D[U_i] + m_c^{\text{phys}})^{-1} \eta_j} \right)^{1/4}. \quad (\text{A8})$$

Both the trace method (A3) and the pseudofermion method (A8) are only valid to $\mathcal{O}(\delta m^2)$; when they agree we assume those corrections are under control. In order to stabilize the numerical estimate of the reweighting factors, it is also common to split the reweighting factor into a product of reweighting factors where each is computed with a fraction of the full mass shift [127–129]. For example, with a simulated mass of m_1 and target mass of $m_1 + \Delta m$, one could use two steps of $\Delta m/2$ and estimate the reweighting factor with the trace method,

$$w_i^{\text{tr}} = w_{i,1}^{\text{tr}} w_{i,2}^{\text{tr}}, \quad (\text{A9})$$

$$w_{i,1}^{\text{tr}} = \exp \frac{\Delta m/2}{4} \frac{1}{N_\eta} \sum_j \eta_j^\dagger \left(D[U_i] + m_1 + \frac{\Delta m}{2} \right)^{-1} \eta_j,$$

$$w_{i,2}^{\text{tr}} = \exp \frac{\Delta m/2}{4} \frac{1}{N_\theta} \sum_j \theta_j^\dagger (D[U_i] + m_1 + \Delta m)^{-1} \theta_j,$$

using independently sampled complex Gaussian noise η and θ . Of course, one may split the shift Δm into finer steps if needed, for increased computational cost.

The reweighting factor accounts for a change in the action and is exponential in the spacetime volume. This can lead to numerical under- or overflow. As a cure, we factor out the average reweighting factor. Recognizing the trace of the inverse Dirac operator on a configuration U_i as the scalar quark density times the lattice volume,

$$(V\bar{q}q)_i = \text{tr} [(D[U_i] + m_q)^{-1}], \quad (\text{A10})$$

we can rescale w , shifting by the ensemble average $V\langle \bar{c}c \rangle$ computed via (A4). For example, rescaling the trace method (A3) gives

$$w_i^{\text{tr}} = \exp \frac{\delta m}{4} (\text{tr} [(D[U_i] + m_c^{\text{phys}})^{-1}] - V\langle \bar{c}c \rangle); \quad (\text{A11})$$

such a rescaling cancels exactly in the reweighting procedure (A2). A similar rescaling cures the pseudofermion method (A8). If we split the mass shift as in (A9), each factor of the weight may be independently so stabilized.

On the a06m310L ensemble the lattice volume and shift in the mass are $V = 72^3 \times 96$ and

$$\begin{aligned} \delta a_{06} m_c &= a_{06} m_c^{\text{phys}} - a_{06} m_c^{\text{HMC}} \\ &= -0.0281(4). \end{aligned} \quad (\text{A12})$$

While the shift in the mass is only about 10% of the physical charm quark mass, it is of the order of the physical strange quark mass. In order to stabilize the numerical estimate of the reweighting factors, we split this mass shift into ten equal steps, and for each step, we used




FIG. 9. Distribution of reweighting factors, r , for the a06m310L ensemble. The reweighting was performed with ten equal steps of the mass difference from $a_0 m_c^{\text{HMC}} = 0.286$ to $a_0 m_c^{\text{phys}} = 0.2579$. The reweighting factors from each step are shown in the bottom ten subpanels with the product reweighting factor shown in the top panel. The a06m310L ensemble was generated with two different streams of equal length. For more information on the ensembles, see Ref. [36].

$N_\eta = 128$ independent Gaussian random noise vectors. For each step in the reweighting, we used the same Naik value of $\epsilon_N = -0.0533$ as was used in the original HMC. This ensures that the Dirac operator only differs from one mass to the next by the quark mass itself. As the Naik term is used to improve the approach to the continuum limit, this is a valid choice to make as it results in a slightly different approach to the continuum than if one simulated at the physical charm quark mass with the optimized Naik value for that mass.

In Fig. 9, we show the reweighting factors for each of the mass steps, with the bottom panel having a mass $a_0 m_c = 0.28319$ closest to the HMC mass and the second from top panel having $a_0 m_c^{\text{phys}} = 0.2579$, scaled for numerical stability. The top panel is the resulting product reweighting factor normalized by the average reweighting factor. One observes that there are a few large reweighting factors of $O(100)$. We have verified that the trace estimation method (A3) and the pseudofermion method (A8) produce comparable normalized reweighting factors. The large reweighting factors are likely due to the parent HMC distribution of configurations having a suboptimal overlap with the physical, target distribution.

1. Reweighted spectrum

The next task is to understand how this reweighting impacts the extracted spectrum. To aid in this discussion, we reiterate our strategy for fitting the correlation functions. Because the noise of the omega baryon correlation function grows in Euclidean time, it is the most challenging to fit and so we focus our discussion on the omega. Our strategy is to find a good quality fit to the correlation function for which the extracted ground state energy is stable against the number of states and the time-range used in the analysis. For a given t_{\min} , we opt to choose the simplest model which satisfies this criteria, which amounts to picking the minimum number of excited states possible.

The SS correlation functions are positive definite, therefore implying that the excited state contamination of the effective mass must come from above. When examining the SS omega-baryon effective mass on the a06m310L ensemble, one observes that around $t = 25$, the effective mass stops decreasing, and even increases a little. Because this is not allowed for a positive definite correlation function, we can conclude this must be due to a correlated stochastic fluctuation; see Fig. 10. In the reweighted effective mass, one observes more dramatic behavior of the effective mass beginning around the same time. To be conservative, we set $t_{\max} = 30$ in our analysis as this allows the analysis to be sensitive to these stochastic fluctuations, which fluctuate in the opposite direction between the unweighted and reweighted configurations.

As a comparison, we also show the reweighting factors and reweighted omega baryon effective mass on the a12m130 ensemble (Fig. 11) where the charm quark was 2% different from its physical value. In this case, the reweighting factors are much easier to estimate, and we do not observe any large values.

The lower panels in Fig. 10 show the extracted ground state mass as a function of t_{\min} and n_{state} . We observe that an $n_{\text{state}} = 2$ fit beginning at $t_{\min} = 19$ and 15 for the unweighted and reweighted correlation functions satisfies our optimization criteria. These lead to the estimate value of m_Ω given in Eq. (3.7). In Table VI, we also show the reweighted value of m_π and m_K on this a06m310L ensemble. While the pion and kaon have a statistically significant shift from the reweighting, when we use the reweighted values of m_π , m_K and m_Ω from this ensemble, the final extrapolated value of $w_0 m_\Omega$ is within 1 standard deviation of the completely unweighted analysis. As the a06m310L ensemble has the largest potential change from reweighting, we conclude that at the level of precision we currently have, our results are not sensitive to the slight mistuning of the charm quark mass from its physical value on each of the configurations.




FIG. 10. Same as Fig. 1 for the a06m310L ensemble.




FIG. 11. The reweighting factors for the a12m130 ensemble (left) and the reweighted omega baryon effective mass (right).

TABLE VI. The pion, kaon and omega masses on the a06m310L ensemble with and without reweighting as well as the correlated difference.

State	Unweighted	Reweightd	Difference
am_π	0.09456(06)	0.09419(14)	-0.00037(14)
am_K	0.16204(07)	0.16173(16)	-0.00032(15)
am_Ω	0.5069(21)	0.5065(29)	-0.0004(34)

Appendix B: MODELS INCLUDED IN AVERAGE

We use $w_{0,\text{orig}}m_\Omega$ as an example to demonstrate the model averaging and naming conventions for the various models. The other gradient-flow scale studies have identical in form, extrapolation functions. Table VII provides the various models and their relative weights for $w_{0,\text{orig}}m_\Omega$. Of note, in all four gradient-flow scale extrapolations, the extrapolations which make use of l_Ω and s_Ω versus l_F and s_F , all have relatively smaller weights in the averaging, such that in all cases, these extrapola-

tions could be dropped from the model averaging without effecting the final result. A reproduction of this model averaging as well as for the other three gradient-flow scales can be obtained by downloading and running the

associated analysis software at https://github.com/callat-qcd/project_scale_setting_mdwf_hisq.

A few example extrapolation formula are given below to demonstrate the naming convention:

$$w_0 m_\Omega = \underbrace{c_0}_{\text{LO}} + \underbrace{\delta^{\text{NLO}}(l_F, s_F)}_{\text{chiral NLO}} + \underbrace{\delta_{a,F}^{\text{NLO}}}_{\text{disc NLO}} \quad [\text{xpt_nlo_fv_F}] \quad (\text{B1a})$$

$$w_0 m_\Omega = \underbrace{c_0}_{\text{LO}} + \underbrace{\delta^{\text{NLO}}(l_F, s_F)}_{\text{chiral NLO}} + \underbrace{\delta^{\text{N}^2\text{LO}}(l_F, s_F)}_{\text{chiral N}^2\text{LO}} + \underbrace{\delta_{\text{ln}}^{\text{N}^2\text{LO}}}_{\text{chiral log N}^2\text{LO}} + \underbrace{\delta_{a,F}^{\text{NLO}}}_{\text{disc NLO}} + \underbrace{\delta_{a,F}^{\text{N}^2\text{LO}}}_{\text{disc N}^2\text{LO}} + \underbrace{\delta_{L,F}^{\text{N}^2\text{LO}}}_{\text{FV N}^2\text{LO}} \quad [\text{xpt_n2lo_fv_F}] \quad (\text{B1b})$$

$$w_0 m_\Omega = \underbrace{c_0}_{\text{LO}} + \underbrace{\delta^{\text{NLO}}(l_F, s_F)}_{\text{chiral NLO}} + \underbrace{\delta^{\text{N}^2\text{LO}}(l_F, s_F)}_{\text{chiral N}^2\text{LO}} + \underbrace{\delta^{\text{N}^3\text{LO}}(l_F, s_F)}_{\text{chiral N}^3\text{LO}} + \underbrace{\delta_{\text{ln}}^{\text{N}^2\text{LO}}}_{\text{chiral log N}^2\text{LO}} + \underbrace{\delta_{\text{ln}}^{\text{N}^3\text{LO}}}_{\text{chiral logs N}^3\text{LO}} + \underbrace{\delta_{a,F}^{\text{NLO}}}_{\text{disc NLO}} + \underbrace{\delta_{a,F}^{\text{N}^2\text{LO}}}_{\text{disc N}^2\text{LO}} + \underbrace{\delta_{a,F}^{\text{N}^3\text{LO}}}_{\text{disc N}^3\text{LO}} + \underbrace{\delta_{L,F}^{\text{N}^2\text{LO}}}_{\text{FV N}^2\text{LO}} + \underbrace{\delta_{L,F}^{\text{N}^3\text{LO}}}_{\text{FV N}^3\text{LO}} \quad [\text{xpt_n3lo_fv_F}] \quad (\text{B1c})$$

where the **chiral**, **discretization** and **finite volume** corrections are defined in Eqs. (3.14), (3.20) and (3.18) respectively.

-
- [1] S. Aoki *et al.* (Flavour Lattice Averaging Group), “FLAG Review 2019: Flavour Lattice Averaging Group (FLAG),” *Eur. Phys. J. C* **80**, 113 (2020), [arXiv:1902.08191 \[hep-lat\]](https://arxiv.org/abs/1902.08191).
- [2] T. Aoyama *et al.*, “The anomalous magnetic moment of the muon in the Standard Model,” *Phys. Rept.* **887**, 1–166 (2020), [arXiv:2006.04822 \[hep-ph\]](https://arxiv.org/abs/2006.04822).
- [3] R. Sommer, “A New way to set the energy scale in lattice gauge theories and its applications to the static force and alpha-s in SU(2) Yang-Mills theory,” *Nucl. Phys. B* **411**, 839–854 (1994), [arXiv:hep-lat/9310022](https://arxiv.org/abs/hep-lat/9310022).
- [4] Martin Lüscher, “Properties and uses of the Wilson flow in lattice QCD,” *JHEP* **1008**, 071 (2010), [arXiv:1006.4518 \[hep-lat\]](https://arxiv.org/abs/1006.4518).
- [5] Szabolcs Borsanyi *et al.*, “High-precision scale setting in lattice QCD,” *JHEP* **09**, 010 (2012), [arXiv:1203.4469 \[hep-lat\]](https://arxiv.org/abs/1203.4469).
- [6] Albert Deuzeman and Urs Wenger, “Gradient flow and scale setting for twisted mass fermions,” *PoS LATTICE2012*, 162 (2012).
- [7] Mattia Bruno and Rainer Sommer (ALPHA), “On the N_f -dependence of gluonic observables,” *PoS LATTICE2013*, 321 (2014), [arXiv:1311.5585 \[hep-lat\]](https://arxiv.org/abs/1311.5585).
- [8] R. J. Dowdall, C. T. H. Davies, G. P. Lepage, and C. McNeile, “Vus from pi and K decay constants in full lattice QCD with physical u, d, s and c quarks,” *Phys. Rev. D* **88**, 074504 (2013), [arXiv:1303.1670 \[hep-lat\]](https://arxiv.org/abs/1303.1670).
- [9] V.G. Bornyakov *et al.*, “Wilson flow and scale setting from lattice QCD,” (2015), [arXiv:1508.05916 \[hep-lat\]](https://arxiv.org/abs/1508.05916).
- [10] A. Bazavov *et al.* (HotQCD), “Equation of state in (2+1)-flavor QCD,” *Phys. Rev. D* **90**, 094503 (2014), [arXiv:1407.6387 \[hep-lat\]](https://arxiv.org/abs/1407.6387).
- [11] T. Blum *et al.* (RBC, UKQCD), “Domain wall QCD with physical quark masses,” *Phys. Rev. D* **93**, 074505 (2016), [arXiv:1411.7017 \[hep-lat\]](https://arxiv.org/abs/1411.7017).
- [12] A. Bazavov *et al.* (MILC), “Gradient flow and scale setting on MILC HISQ ensembles,” *Phys. Rev. D* **93**, 094510 (2016), [arXiv:1503.02769 \[hep-lat\]](https://arxiv.org/abs/1503.02769).
- [13] Mattia Bruno, Tomasz Korzec, and Stefan Schaefer, “Setting the scale for the CLS 2 + 1 flavor ensembles,” *Phys. Rev. D* **95**, 074504 (2017), [arXiv:1608.08900 \[hep-lat\]](https://arxiv.org/abs/1608.08900).
- [14] Sz. Borsanyi *et al.*, “Leading hadronic contribution to the muon 2 magnetic moment from lattice QCD,” *Nature* (2021), [10.1038/s41586-021-03418-1](https://doi.org/10.1038/s41586-021-03418-1), [arXiv:2002.12347 \[hep-lat\]](https://arxiv.org/abs/2002.12347).
- [15] E. Berkowitz, C. Bouchard, C.C Chang, M.A. Clark, B. Joo, T. Kurth, C. Monahan, A. Nicholson, K. Orginos, E. Rinaldi, P. Vranas, and A. Walker-Loud, “Möbius Domain-Wall fermions on gradient-flowed dynamical HISQ ensembles,” *Phys. Rev. D* **96**, 054513 (2017), [arXiv:1701.07559 \[hep-lat\]](https://arxiv.org/abs/1701.07559).
- [16] A. Bazavov *et al.* (MILC), “Lattice QCD ensembles with four flavors of highly improved staggered quarks,” *Phys. Rev. D* **87**, 054505 (2013), [arXiv:1212.4768 \[hep-lat\]](https://arxiv.org/abs/1212.4768).
- [17] Tanmoy Bhattacharya, Vincenzo Cirigliano, Saul Cohen, Rajan Gupta, Anosh Joseph, Huey-Wen Lin, and Boram Yoon (PNDME), “Iso-vector and Iso-scalar Tensor Charges of the Nucleon from Lattice QCD,” *Phys. Rev. D* **92**, 094511 (2015), [arXiv:1506.06411 \[hep-lat\]](https://arxiv.org/abs/1506.06411).

TABLE VII. Models included in final model average of $w_{0,\text{orig}}m_\Omega$.

Moddel	chi2/dof	Q	logGBF	weight	w_0m_Ω	$w_0[\text{fm}]$
xpt_n3lo_alphaS_F	1.065	0.378	64.919	0.077	1.448(10)	0.1709(12)
xpt_n3lo_F	1.070	0.372	64.916	0.077	1.4467(91)	0.1707(11)
taylor_n3lo_alphaS_F	1.089	0.350	64.863	0.073	1.449(10)	0.1709(12)
xpt_n3lo_FV_alphaS_F	1.068	0.374	64.861	0.073	1.448(10)	0.1709(12)
xpt_n3lo_FV_F	1.074	0.367	64.858	0.073	1.4467(91)	0.1707(11)
taylor_n3lo_FV_alphaS_F	1.088	0.350	64.857	0.073	1.449(10)	0.1709(12)
taylor_n3lo_F	1.094	0.343	64.857	0.073	1.4470(90)	0.1707(11)
taylor_n3lo_FV_F	1.094	0.343	64.851	0.072	1.4470(90)	0.1707(11)
xpt_n2lo_alphaS_F	1.124	0.310	64.610	0.057	1.449(10)	0.1710(12)
xpt_n2lo_F	1.130	0.304	64.600	0.056	1.4475(91)	0.1708(11)
xpt_n2lo_FV_alphaS_F	1.128	0.306	64.551	0.054	1.449(10)	0.1710(12)
xpt_n2lo_FV_F	1.134	0.299	64.540	0.053	1.4475(91)	0.1708(11)
taylor_n2lo_alphaS_F	1.161	0.272	64.410	0.047	1.450(10)	0.1711(12)
taylor_n2lo_FV_alphaS_F	1.161	0.272	64.403	0.046	1.450(10)	0.1711(12)
taylor_n2lo_F	1.168	0.265	64.392	0.046	1.4480(90)	0.1708(11)
taylor_n2lo_FV_F	1.167	0.266	64.386	0.045	1.4480(90)	0.1708(11)
xpt_n2lo_FV_alphaS_O	0.971	0.498	59.864	0.000	1.441(14)	0.1700(16)
xpt_n3lo_FV_alphaS_O	0.937	0.545	59.813	0.000	1.440(13)	0.1699(16)
taylor_n3lo_alphaS_O	1.089	0.349	59.792	0.000	1.444(13)	0.1704(15)
xpt_n2lo_alphaS_O	0.985	0.480	59.789	0.000	1.441(13)	0.1701(16)
xpt_n3lo_FV_O	0.937	0.545	59.515	0.000	1.440(12)	0.1699(15)
taylor_n3lo_FV_alphaS_O	1.089	0.349	59.394	0.000	1.444(13)	0.1704(15)
xpt_n3lo_alphaS_O	0.949	0.529	59.380	0.000	1.441(13)	0.1700(16)
xpt_n2lo_FV_O	0.971	0.498	59.234	0.000	1.441(12)	0.1700(14)
xpt_n2lo_O	0.985	0.480	59.172	0.000	1.441(13)	0.1700(15)
taylor_n3lo_O	1.090	0.348	58.874	0.000	1.444(12)	0.1703(14)
taylor_n3lo_FV_O	1.090	0.349	58.798	0.000	1.444(12)	0.1703(14)
taylor_n2lo_alphaS_O	1.131	0.303	58.742	0.000	1.445(13)	0.1705(15)
taylor_n2lo_FV_O	1.131	0.302	58.607	0.000	1.444(12)	0.1704(14)
xpt_n3lo_O	0.949	0.528	58.329	0.000	1.440(12)	0.1699(14)
taylor_n2lo_FV_alphaS_O	1.131	0.303	58.292	0.000	1.445(13)	0.1705(15)
taylor_n2lo_O	1.132	0.302	58.070	0.000	1.444(12)	0.1704(14)
Model average					1.4481(98)(11)	0.1709(12)(01)

- [18] M.A. Clark, R. Babich, K. Barros, R.C. Brower, and C. Rebbi, “Solving Lattice QCD systems of equations using mixed precision solvers on GPUs,” *Comput.Phys.Commun.* **181**, 1517–1528 (2010), <https://github.com/lattice/quda>, arXiv:0911.3191 [hep-lat].
- [19] R. Babich, M.A. Clark, B. Joo, G. Shi, R.C. Brower, *et al.*, “Scaling Lattice QCD beyond 100 GPUs,” (2011), arXiv:1109.2935 [hep-lat].
- [20] Dru Bryant Renner, W. Schroers, R. Edwards, George Tamminga Fleming, Ph. Hagler, John W. Negele, K. Orginos, A. V. Pochinski, and D. Richards (LHP), “Hadronic physics with domain-wall valence and improved staggered sea quarks,” *Lattice field theory. Proceedings, 22nd International Symposium, Lattice 2004, Batavia, USA, June 21-26, 2004*, *Nucl. Phys. Proc. Suppl.* **140**, 255–260 (2005), [,255(2004)], arXiv:hep-lat/0409130 [hep-lat].
- [21] Oliver Bar, Gautam Rupak, and Noam Shoresh, “Simulations with different lattice Dirac operators for valence and sea quarks,” *Phys. Rev.* **D67**, 114505 (2003), arXiv:hep-lat/0210050 [hep-lat].
- [22] Richard C. Brower, Harmut Neff, and Kostas Orginos, “The möbius domain wall fermion algorithm,” *Comput. Phys. Commun.* **220**, 1–19 (2017), arXiv:1206.5214 [hep-lat].
- [23] David B. Kaplan, “A Method for simulating chiral fermions on the lattice,” *Phys. Lett.* **B288**, 342–347 (1992), arXiv:hep-lat/9206013 [hep-lat].
- [24] Yigal Shamir, “Chiral fermions from lattice boundaries,” *Nucl. Phys.* **B406**, 90–106 (1993), arXiv:hep-lat/9303005 [hep-lat].
- [25] Vadim Furman and Yigal Shamir, “Axial symmetries in lattice QCD with Kaplan fermions,” *Nucl. Phys.* **B439**, 54–78 (1995), arXiv:hep-lat/9405004 [hep-lat].
- [26] E. Follana, Q. Mason, C. Davies, K. Hornbostel, G. P. Lepage, J. Shigemitsu, H. Trottier, and K. Wong (HPQCD, UKQCD), “Highly improved staggered quarks on the lattice, with applications to charm physics,” *Phys. Rev.* **D75**, 054502 (2007), arXiv:hep-lat/0610092 [hep-lat].
- [27] R. Narayanan and H. Neuberger, “Infinite N phase transitions in continuum Wilson loop operators,” *JHEP* **0603**, 064 (2006), arXiv:hep-th/0601210 [hep-th].
- [28] M. Lüscher and P. Weisz, “Perturbative analysis of the gradient flow in non-abelian gauge theories,” *JHEP* **1102**, 051 (2011), arXiv:1101.0963 [hep-th].
- [29] Martin Lüscher, “Chiral symmetry and the Yang–Mills gradient flow,” *JHEP* **1304**, 123 (2013), arXiv:1302.5246 [hep-lat].
- [30] Colin Morningstar and Mike J. Peardon, “Analytic

- smearing of SU(3) link variables in lattice QCD,” *Phys.Rev.* **D69**, 054501 (2004), [arXiv:hep-lat/0311018 \[hep-lat\]](#).
- [31] Chris Bouchard, Chia Cheng Chang, Thorsten Kurth, Kostas Orginos, and Andre Walker-Loud, “On the Feynman-Hellmann Theorem in Quantum Field Theory and the Calculation of Matrix Elements,” *Phys. Rev.* **D96**, 014504 (2017), [arXiv:1612.06963 \[hep-lat\]](#).
- [32] E. Berkowitz, C. Bouchard, D. B. Brantley, C.C Chang, M.A. Clark, N. Garron, B. Joó, T. Kurth, C. Monahan, H. Monge-Camacho, A. Nicholson, K. Orginos, E. Rinaldi, P. Vranas, and A. Walker-Loud, “An accurate calculation of the nucleon axial charge with lattice QCD,” (2017), [arXiv:1704.01114 \[hep-lat\]](#).
- [33] C. C. Chang, A. Nicholson, E. Rinaldi, E. Berkowitz, N. Garron, D. A. Brantley, H. Monge-Camacho, C. J. Monahan, C. Bouchard, M. A. Clark, B. Joó, T. Kurth, K. Orginos, P. Vranas, and A. Walker-Loud, “A percent-level determination of the nucleon axial coupling from quantum chromodynamics,” *Nature* **558**, 91–94 (2018), [arXiv:1805.12130 \[hep-lat\]](#).
- [34] Evan Berkowitz, M. A. Clark, A. Gambhir, K. McElvain, A. Nicholson, E. Rinaldi, P. Vranas, A. Walker-Loud, C. C. Chang, B. Joó, T. Kurth, and K. Orginos, “Simulating the weak death of the neutron in a femtoscale universe with near-Exascale computing,” (2018), [10.1109/SC.2018.00058](#), [arXiv:1810.01609 \[hep-lat\]](#).
- [35] A. Nicholson, E. Berkowitz, H. Monge-Camacho, D. Brantley, N. Garron, C. C. Chang, E. Rinaldi, M. A. Clark, B. Joó, T. Kurth, B. Tiburzi, P. Vranas, and A. Walker-Loud, “Heavy physics contributions to neutrinoless double beta decay from QCD,” *Phys. Rev. Lett.* **121**, 172501 (2018), [arXiv:1805.02634 \[nucl-th\]](#).
- [36] Nolan Miller *et al.*, “ F_K/F_π from M⁰bius Domain-Wall fermions solved on gradient-flowed HISQ ensembles,” *Phys. Rev. D* **102**, 034507 (2020), [arXiv:2005.04795 \[hep-lat\]](#).
- [37] E. Follana, C.T.H. Davies, G.P. Lepage, and J. Shigemitsu (HPQCD, UKQCD), “High Precision determination of the pi, K, D and D(s) decay constants from lattice QCD,” *Phys. Rev. Lett.* **100**, 062002 (2008), [arXiv:0706.1726 \[hep-lat\]](#).
- [38] B. Blossier *et al.* (ETM), “Pseudoscalar decay constants of kaon and D-mesons from $N_f = 2$ twisted mass Lattice QCD,” *JHEP* **07**, 043 (2009), [arXiv:0904.0954 \[hep-lat\]](#).
- [39] S. Durr, Z. Fodor, C. Hoelbling, S.D. Katz, S. Krieg, T. Kurth, L. Lellouch, T. Lippert, A. Ramos, and K.K. Szabo, “The ratio F_K/F_π in QCD,” *Phys. Rev. D* **81**, 054507 (2010), [arXiv:1001.4692 \[hep-lat\]](#).
- [40] N. Carrasco *et al.*, “Leptonic decay constants f_K, f_D , and f_{D_s} with $N_f = 2+1+1$ twisted-mass lattice QCD,” *Phys. Rev.* **D91**, 054507 (2015), [arXiv:1411.7908 \[hep-lat\]](#).
- [41] A. Bazavov *et al.* (Fermilab Lattice, MILC), “Charmed and light pseudoscalar meson decay constants from four-flavor lattice QCD with physical light quarks,” *Phys. Rev.* **D90**, 074509 (2014), [arXiv:1407.3772 \[hep-lat\]](#).
- [42] Stephan Dürr *et al.*, “Leptonic decay-constant ratio f_K/f_π from lattice QCD using 2+1 clover-improved fermion flavors with 2-HEX smearing,” *Phys. Rev. D* **95**, 054513 (2017), [arXiv:1601.05998 \[hep-lat\]](#).
- [43] V.G. Bornyakov, R. Horsley, Y. Nakamura, H. Perlt, D. Pleiter, P.E.L. Rakow, G. Schierholz, A. Schiller, H. Stüben, and J.M. Zanotti (QCDSF–UKQCD), “Flavour breaking effects in the pseudoscalar meson decay constants,” *Phys. Lett. B* **767**, 366–373 (2017), [arXiv:1612.04798 \[hep-lat\]](#).
- [44] A. Bazavov *et al.*, “ B - and D -meson leptonic decay constants from four-flavor lattice QCD,” *Phys. Rev.* **D98**, 074512 (2018), [arXiv:1712.09262 \[hep-lat\]](#).
- [45] C. Allton *et al.* (RBC-UKQCD), “Physical Results from 2+1 Flavor Domain Wall QCD and SU(2) Chiral Perturbation Theory,” *Phys. Rev.* **D78**, 114509 (2008), [arXiv:0804.0473 \[hep-lat\]](#).
- [46] A. Walker-Loud *et al.*, “Light hadron spectroscopy using domain wall valence quarks on an Asqtad sea,” *Phys. Rev.* **D79**, 054502 (2009), [arXiv:0806.4549 \[hep-lat\]](#).
- [47] Huey-Wen Lin *et al.* (Hadron Spectrum), “First results from 2+1 dynamical quark flavors on an anisotropic lattice: Light-hadron spectroscopy and setting the strange-quark mass,” *Phys. Rev. D* **79**, 034502 (2009), [arXiv:0810.3588 \[hep-lat\]](#).
- [48] S. Durr *et al.*, “Ab-Initio Determination of Light Hadron Masses,” *Science* **322**, 1224–1227 (2008), [arXiv:0906.3599 \[hep-lat\]](#).
- [49] C. Alexandrou, R. Baron, J. Carbonell, V. Drach, P. Guichon, K. Jansen, T. Korzec, and O. Pene (ETM), “Low-lying baryon spectrum with two dynamical twisted mass fermions,” *Phys. Rev. D* **80**, 114503 (2009), [arXiv:0910.2419 \[hep-lat\]](#).
- [50] C. Alexandrou, V. Drach, K. Jansen, C. Kallidonis, and G. Koutsou, “Baryon spectrum with $N_f = 2 + 1 + 1$ twisted mass fermions,” *Phys. Rev. D* **90**, 074501 (2014), [arXiv:1406.4310 \[hep-lat\]](#).
- [51] David A. Brantley, Balint Joo, Ekaterina V. Mastropas, Emanuele Mereghetti, Henry Monge-Camacho, Brian C. Tiburzi, and Andre Walker-Loud, “Strong isospin violation and chiral logarithms in the baryon spectrum,” (2016), [arXiv:1612.07733 \[hep-lat\]](#).
- [52] Robert G. Edwards and Balint Joo (SciDAC Collaboration, LHPC Collaboration, UKQCD Collaboration), “The Chroma software system for lattice QCD,” *Nucl.Phys.Proc.Suppl.* **140**, 832 (2005), [arXiv:hep-lat/0409003 \[hep-lat\]](#).
- [53] S. Basak, R. G. Edwards, G. T. Fleming, U. M. Heller, C. Morningstar, D. Richards, I. Sato, and S. Wallace, “Group-theoretical construction of extended baryon operators in lattice QCD,” *Phys. Rev.* **D72**, 094506 (2005), [arXiv:hep-lat/0506029 \[hep-lat\]](#).
- [54] Subhasish Basak, Robert Edwards, George T. Fleming, Urs M. Heller, Colin Morningstar, David Richards, Ikuro Sato, and Stephen J. Wallace (Lattice Hadron Physics (LHPC)), “Clebsch-Gordan construction of lattice interpolating fields for excited baryons,” *Phys. Rev.* **D72**, 074501 (2005), [arXiv:hep-lat/0508018 \[hep-lat\]](#).
- [55] Evan Berkowitz, Amy Nicholson, Chia Cheng Chang, Enrico Rinaldi, M. A. Clark, Bálint Joó, Thorsten Kurth, Pavlos Vranas, and André Walker-Loud, “Calm Multi-Baryon Operators,” in *35th International Symposium on Lattice Field Theory (Lattice 2017) Granada, Spain, June 18-24, 2017* (2017) [arXiv:1710.05642 \[hep-lat\]](#).
- [56] Silas R. Beane, William Detmold, Thomas C. Luu, Kostas Orginos, Assumpta Parreno, Martin Savage, Aaron Torok, and Andre Walker-Loud, “High Statistics Analysis using Anisotropic Clover Lattices: (I) Single Hadron Correlation Functions,” *Phys.Rev.* **D79**, 114502

- (2009), [arXiv:0903.2990 \[hep-lat\]](#).
- [57] G.P. Lepage, B. Clark, C.T.H. Davies, K. Hornbostel, P.B. Mackenzie, C. Morningstar, and H. Trotter, “Constrained curve fitting,” *Nucl. Phys. B Proc. Suppl.* **106**, 12–20 (2002), [arXiv:hep-lat/0110175](#).
- [58] William I. Jay and Ethan T. Neil, “Bayesian model averaging for analysis of lattice field theory results,” (2020), [arXiv:2008.01069 \[stat.ME\]](#).
- [59] The MILC Collaboration, “QUDA: a library for performing calculations in lattice qcd on graphics processing units (gpus), leveraging nvidia’s cuda platform.” <https://github.com/lattice/quda>.
- [60] “MILC Collaboration code for lattice qcd calculations,” https://github.com/milc-qcd/milc_qcd.
- [61] Ulli Wolff (ALPHA), “Monte Carlo errors with less errors,” *Comput. Phys. Commun.* **156**, 143–153 (2004), [Erratum: *Comput. Phys. Commun.* 176, 383 (2007)], [arXiv:hep-lat/0306017](#).
- [62] Barbara De Palma, Marco Erba, Luca Mantovani, and Nicola Mosco, “A Python program for the implementation of the Γ -method for Monte Carlo simulations,” *Comput. Phys. Commun.* **234**, 294–301 (2019), [arXiv:1703.02766 \[hep-lat\]](#).
- [63] C. Davies *et al.*, “The Determination of $\alpha(s)$ from lattice QCD with 2+1 flavors of dynamical quarks,” *Nucl. Phys. B Proc. Suppl.* **119**, 595–597 (2003), [arXiv:hep-lat/0209122](#).
- [64] Zoltan Fodor, Kieran Holland, Julius Kuti, Santanu Mondal, Daniel Nogradi, and Chik Him Wong, “The lattice gradient flow at tree-level and its improvement,” *JHEP* **09**, 018 (2014), [arXiv:1406.0827 \[hep-lat\]](#).
- [65] Murray Gell-Mann, R. J. Oakes, and B. Renner, “Behavior of current divergences under $SU(3) \times SU(3)$,” *Phys. Rev.* **175**, 2195–2199 (1968).
- [66] M. Tanabashi *et al.* (Particle Data Group), “Review of Particle Physics,” *Phys. Rev. D* **98**, 030001 (2018).
- [67] A.M. Ferrenberg and R.H. Swendsen, “New Monte Carlo Technique for Studying Phase Transitions,” *Phys. Rev. Lett.* **61**, 2635–2638 (1988).
- [68] M. Luscher, “Volume Dependence of the Energy Spectrum in Massive Quantum Field Theories. 1. Stable Particle States,” *Commun. Math. Phys.* **104**, 177 (1986).
- [69] Elizabeth Ellen Jenkins and Aneesh V. Manohar, “Baryon chiral perturbation theory using a heavy fermion Lagrangian,” *Phys. Lett.* **B255**, 558–562 (1991).
- [70] Elizabeth Ellen Jenkins and Aneesh V. Manohar, “Chiral corrections to the baryon axial currents,” *Phys. Lett.* **B259**, 353–358 (1991).
- [71] Richard F. Lebed, “Baryon decuplet mass relations in chiral perturbation theory,” *Nucl. Phys. B* **430**, 295–318 (1994), [arXiv:hep-ph/9311234](#).
- [72] Richard F. Lebed and Markus A. Luty, “Baryon masses at second order in chiral perturbation theory,” *Phys. Lett. B* **329**, 479–485 (1994), [arXiv:hep-ph/9401232](#).
- [73] Brian C. Tiburzi and Andre Walker-Loud, “Decuplet baryon masses in partially quenched chiral perturbation theory,” *Nucl. Phys.* **A748**, 513–536 (2005), [arXiv:hep-lat/0407030 \[hep-lat\]](#).
- [74] Aaron Torok *et al.* (NPLQCD), “Meson-Baryon Scattering Lengths from Mixed-Action Lattice QCD,” *Phys. Rev.* **D81**, 074506 (2010), [arXiv:0907.1913 \[hep-lat\]](#).
- [75] K. I. Ishikawa *et al.* (PACS-CS), “ $SU(2)$ and $SU(3)$ chiral perturbation theory analyses on baryon masses in 2+1 flavor lattice QCD,” *Phys. Rev.* **D80**, 054502 (2009), [arXiv:0905.0962 \[hep-lat\]](#).
- [76] Elizabeth Ellen Jenkins, Aneesh V. Manohar, John W. Negele, and Andre Walker-Loud, “A Lattice Test of $1/N(c)$ Baryon Mass Relations,” *Phys. Rev.* **D81**, 014502 (2010), [arXiv:0907.0529 \[hep-lat\]](#).
- [77] Andre Walker-Loud, “Evidence for non-analytic light quark mass dependence in the baryon spectrum,” *Phys. Rev.* **D86**, 074509 (2012), [arXiv:1112.2658 \[hep-lat\]](#).
- [78] Silas R. Beane *et al.* (NPLQCD), “Hyperon nucleon scattering from fully-dynamical lattice QCD,” *Nucl. Phys.* **A794**, 62–72 (2007), [arXiv:hep-lat/0612026](#).
- [79] Brian C. Tiburzi and Andre Walker-Loud, “Hyperons in Two Flavor Chiral Perturbation Theory,” *Phys. Lett.* **B669**, 246–253 (2008), [arXiv:0808.0482 \[nucl-th\]](#).
- [80] Fu-Jiun Jiang and Brian C. Tiburzi, “Hyperon Axial Charges in Two-Flavor Chiral Perturbation Theory,” *Phys. Rev. D* **80**, 077501 (2009), [arXiv:0905.0857 \[nucl-th\]](#).
- [81] Fu-Jiun Jiang, Brian C. Tiburzi, and Andre Walker-Loud, “Kaon Thresholds and Two-Flavor Chiral Expansions for Hyperons,” *Phys. Lett. B* **695**, 329–336 (2011), [arXiv:0911.4721 \[nucl-th\]](#).
- [82] Fu-Jiun Jiang and Brian C. Tiburzi, “Hyperon Electromagnetic Properties in Two-Flavor Chiral Perturbation Theory,” *Phys. Rev. D* **81**, 034017 (2010), [arXiv:0912.2077 \[nucl-th\]](#).
- [83] Steven Weinberg, “Phenomenological Lagrangians,” *Physica* **A96**, 327–340 (1979).
- [84] Oliver Bar and Maarten Golterman, “Chiral perturbation theory for gradient flow observables,” *Phys. Rev. D* **89**, 034505 (2014), [Erratum: *Phys. Rev. D* 89, 099905 (2014)], [arXiv:1312.4999 \[hep-lat\]](#).
- [85] J. Gasser and H. Leutwyler, “Light Quarks at Low Temperatures,” *Phys. Lett.* **B184**, 83–88 (1987).
- [86] Gilberto Colangelo and Christoph Haefeli, “An Asymptotic formula for the pion decay constant in a large volume,” *Phys. Lett. B* **590**, 258–264 (2004), [arXiv:hep-lat/0403025](#).
- [87] Johan Bijnens and Thomas Rössler, “Finite Volume at Two-loops in Chiral Perturbation Theory,” *JHEP* **01**, 034 (2015), [arXiv:1411.6384 \[hep-lat\]](#).
- [88] Stephen R. Sharpe and Robert L. Singleton, Jr, “Spontaneous flavor and parity breaking with Wilson fermions,” *Phys. Rev.* **D58**, 074501 (1998), [arXiv:hep-lat/9804028 \[hep-lat\]](#).
- [89] K. Symanzik, “Continuum Limit and Improved Action in Lattice Theories. 1. Principles and ϕ^4 Theory,” *Nucl. Phys.* **B226**, 187–204 (1983).
- [90] K. Symanzik, “Continuum Limit and Improved Action in Lattice Theories. 2. $O(N)$ Nonlinear Sigma Model in Perturbation Theory,” *Nucl. Phys.* **B226**, 205–227 (1983).
- [91] Janos Balog, Ferenc Niedermayer, and Peter Weisz, “Logarithmic corrections to $O(a^2)$ lattice artifacts,” *Phys. Lett. B* **676**, 188–192 (2009), [arXiv:0901.4033 \[hep-lat\]](#).
- [92] Janos Balog, Ferenc Niedermayer, and Peter Weisz, “The Puzzle of apparent linear lattice artifacts in the 2d non-linear sigma-model and Symanzik’s solution,” *Nucl. Phys. B* **824**, 563–615 (2010), [arXiv:0905.1730 \[hep-lat\]](#).
- [93] Nikolai Husung, Peter Marquard, and Rainer Sommer, “Asymptotic behavior of cutoff effects in Yang–Mills theory and in Wilson’s lattice QCD,” *Eur. Phys. J. C*

- 80**, 200 (2020), [arXiv:1912.08498 \[hep-lat\]](#).
- [94] Oliver Bar, Gautam Rupak, and Noam Shores, “Chiral perturbation theory at $\mathcal{O}(a^2)$ for lattice QCD,” *Phys. Rev. D* **70**, 034508 (2004), [arXiv:hep-lat/0306021 \[hep-lat\]](#).
- [95] Oliver Bar, Claude Bernard, Gautam Rupak, and Noam Shores, “Chiral perturbation theory for staggered sea quarks and Ginsparg-Wilson valence quarks,” *Phys. Rev. D* **72**, 054502 (2005), [arXiv:hep-lat/0503009 \[hep-lat\]](#).
- [96] Maarten Golterman, Taku Izubuchi, and Yigal Shamir, “The Role of the double pole in lattice QCD with mixed actions,” *Phys. Rev. D* **71**, 114508 (2005), [arXiv:hep-lat/0504013](#).
- [97] Brian C. Tiburzi, “Baryons with Ginsparg-Wilson quarks in a staggered sea,” *Phys. Rev. D* **72**, 094501 (2005), [Erratum: *Phys. Rev. D* **79**, 039904 (2009)], [arXiv:hep-lat/0508019 \[hep-lat\]](#).
- [98] Jiunn-Wei Chen, Donal O’Connell, Ruth S. Van de Water, and Andre Walker-Loud, “Ginsparg-Wilson pions scattering on a staggered sea,” *Phys. Rev. D* **73**, 074510 (2006), [arXiv:hep-lat/0510024 \[hep-lat\]](#).
- [99] Jiunn-Wei Chen, Donal O’Connell, and Andre Walker-Loud, “Two Meson Systems with Ginsparg-Wilson Valence Quarks,” *Phys. Rev. D* **75**, 054501 (2007), [arXiv:hep-lat/0611003 \[hep-lat\]](#).
- [100] Kostas Orginos and Andre Walker-Loud, “Mixed meson masses with domain-wall valence and staggered sea fermions,” *Phys. Rev. D* **77**, 094505 (2008), [arXiv:0705.0572 \[hep-lat\]](#).
- [101] Jiunn-Wei Chen, Donal O’Connell, and Andre Walker-Loud, “Universality of mixed action extrapolation formulae,” *JHEP* **04**, 090 (2009), [arXiv:0706.0035 \[hep-lat\]](#).
- [102] Jiunn-Wei Chen, Maarten Golterman, Donal O’Connell, and Andre Walker-Loud, “Mixed Action Effective Field Theory: An Addendum,” *Phys. Rev. D* **79**, 117502 (2009), [arXiv:0905.2566 \[hep-lat\]](#).
- [103] (2005), Example given by M. Savage in a discussion on the inclusion of discretization effects in an effective field theory.
- [104] Claude W. Bernard and Maarten F.L. Golterman, “Chiral perturbation theory for the quenched approximation of QCD,” *Phys. Rev. D* **46**, 853–857 (1992), [arXiv:hep-lat/9204007](#).
- [105] Claude W. Bernard and Maarten F. L. Golterman, “Partially quenched gauge theories and an application to staggered fermions,” *Phys. Rev. D* **49**, 486–494 (1994), [arXiv:hep-lat/9306005 \[hep-lat\]](#).
- [106] Stephen R. Sharpe, “Enhanced chiral logarithms in partially quenched QCD,” *Phys. Rev. D* **56**, 7052–7058 (1997), [Erratum: *Phys. Rev. D* **62**, 099901 (2000)], [arXiv:hep-lat/9707018](#).
- [107] Stephen R. Sharpe and Noam Shores, “Physical results from unphysical simulations,” *Phys. Rev. D* **62**, 094503 (2000), [arXiv:hep-lat/0006017 \[hep-lat\]](#).
- [108] Stephen R. Sharpe and Noam Shores, “Partially quenched chiral perturbation theory without Φ_0 ,” *Phys. Rev. D* **64**, 114510 (2001), [arXiv:hep-lat/0108003 \[hep-lat\]](#).
- [109] Szabolcs Borsanyi, Stephan Durr, Zoltan Fodor, Stefan Krieg, Andreas Schafer, Enno E Scholz, and Kalman K Szabo, “SU(2) chiral perturbation theory low-energy constants from 2+1 flavor staggered lattice simulations,” *Phys. Rev. D* **88**, 014513 (2013), [arXiv:1205.0788 \[hep-lat\]](#).
- [110] B. Colquhoun, R.J. Dowdall, J. Koponen, C.T.H. Davies, and G.P. Lepage, “ $B \rightarrow \pi \ell \nu$ at zero recoil from lattice QCD with physical u/d quarks,” *Phys. Rev. D* **93**, 034502 (2016), [arXiv:1510.07446 \[hep-lat\]](#).
- [111] R.J. Dowdall, C.T.H. Davies, R.R. Horgan, G.P. Lepage, C.J. Monahan, J. Shigemitsu, and M. Wingate, “Neutral B-meson mixing from full lattice QCD at the physical point,” *Phys. Rev. D* **100**, 094508 (2019), [arXiv:1907.01025 \[hep-lat\]](#).
- [112] Stephen R. Sharpe, “Future of Chiral Extrapolations with Domain Wall Fermions,” in *Workshop on Domain Wall Fermions at Ten Years* (2007) [arXiv:0706.0218 \[hep-lat\]](#).
- [113] Y. Aoki *et al.* (RBC, UKQCD), “Continuum Limit Physics from 2+1 Flavor Domain Wall QCD,” *Phys. Rev. D* **83**, 074508 (2011), [arXiv:1011.0892 \[hep-lat\]](#).
- [114] Robert E. Kass and Adrian E. Raftery, “Bayes factors,” *Journal of the American Statistical Association* **90**, 773–795 (1995).
- [115] “LALIBE: Callat Collaboration code for lattice qcd calculations,” <https://github.com/callat-qcd/lalibe>.
- [116] The HDF Group, “Hierarchical Data Format, version 5,” (1997-NNNN), <http://www.hdfgroup.org/HDF5/>.
- [117] Thorsten Kurth, Andrew Pochinsky, Abhinav Sarje, Sergey Syritsyn, and Andre Walker-Loud, “High-Performance I/O: HDF5 for Lattice QCD,” *Proceedings, 32nd International Symposium on Lattice Field Theory (Lattice 2014): Brookhaven, NY, USA, June 23-28, 2014*, PoS **LATTICE2014**, 045 (2015), [arXiv:1501.06992 \[hep-lat\]](#).
- [118] Evan Berkowitz, “METAQ: Bundle Supercomputing Tasks,” (2017), <https://github.com/evanberkowitz/metaq>, [arXiv:1702.06122 \[physics.comp-ph\]](#).
- [119] Evan Berkowitz, Gustav R. Jansen, Kenneth McElvain, and André Walker-Loud, “Job Management and Task Bundling,” *EPJ Web Conf.* **175**, 09007 (2018), [arXiv:1710.01986 \[hep-lat\]](#).
- [120] Chia Cheng Chang, Christopher Körber, and André Walker-Loud, “EspressoDB: A Scientific Database for Managing High-Performance Computing Workflow,” *J. Open Source Softw.* **5**, 2007 (2020), [arXiv:1912.03580 \[hep-lat\]](#).
- [121] Peter Lepage, “gplepage/gvar: gvar version 11.2,” (2020), <https://github.com/gplepage/gvar>.
- [122] Peter Lepage, “gplepage/lqfit: lqfit version 11.5.1,” (2020), <https://github.com/gplepage/lqfit>.
- [123] S. Aoki *et al.* (PACS-CS), “Physical Point Simulation in 2+1 Flavor Lattice QCD,” *Phys. Rev. D* **81**, 074503 (2010), [arXiv:0911.2561 \[hep-lat\]](#).
- [124] Martin Luscher and Stefan Schaefer, “Lattice QCD with open boundary conditions and twisted-mass reweighting,” *Comput. Phys. Commun.* **184**, 519–528 (2013), [arXiv:1206.2809 \[hep-lat\]](#).
- [125] Jacob Finkenrath, Francesco Knechtli, and Björn Leder, “One flavor mass reweighting in lattice QCD,” *Nucl. Phys. B* **877**, 441–456 (2013), [Erratum: *Nucl. Phys. B* **880**, 574–575 (2014)], [arXiv:1306.3962 \[hep-lat\]](#).
- [126] Björn Leder, Jacob Finkenrath, and Francesco Knechtli, “One flavor mass reweighting: foundations,” *PoS LATTICE2013*, 035 (2014), [arXiv:1401.1079 \[hep-lat\]](#).

- [127] Martin Hasenbusch, “Speeding up the hybrid Monte Carlo algorithm for dynamical fermions,” *Phys. Lett. B* **519**, 177–182 (2001), [arXiv:hep-lat/0107019](#).
- [128] Anna Hasenfratz, Roland Hoffmann, and Stefan Schaefer, “Reweighting towards the chiral limit,” *Phys. Rev. D* **78**, 014515 (2008), [arXiv:0805.2369 \[hep-lat\]](#).
- [129] Qi Liu, Norman H. Christ, and Chulwoo Jung, “Light Quark Mass Reweighting,” *Phys. Rev. D* **87**, 054503

(2013), [arXiv:1206.0080 \[hep-lat\]](#).

Appendix C: STABILITY PLOTS OF THE OMEGA GROUND STATE MASS

Here we present the stability plots for the remaining Omega correlator fits used in our analysis, which are presented in Fig. 12–25.




FIG. 12. Same as Fig. 1 for the a15m400 ensemble.




FIG. 13. Same as Fig. 1 for the a15m350 ensemble.




FIG. 14. Same as Fig. 1 for the a15m310 and a15m310L ensembles.




FIG. 15. Same as Fig. 1 for the a15m220 ensemble.




FIG. 16. Same as Fig. 1 for the a12m400 ensemble.




FIG. 17. Same as Fig. 1 for the a12m350 ensemble.




FIG. 18. Same as Fig. 1 for the a12m310 and a12m310XL ensembles.




FIG. 19. Same as Fig. 1 for the a12m220 ensembles.




FIG. 20. Same as Fig. 1 for the a12m220ms ensembles.




FIG. 21. Same as Fig. 1 for the a12m180L ensembles.




FIG. 22. Same as Fig. 1 for the a09m400 ensemble.




FIG. 23. Same as Fig. 1 for the a09m350 ensemble.




FIG. 24. Same as Fig. 1 for the a09m310 ensemble.




FIG. 25. Same as Fig. 1 for the a09m220 ensemble.