

UCSF

UC San Francisco Previously Published Works

Title

Vascular endothelial function is impaired by aerosol from a single IQOS HeatStick to the same extent as by cigarette smoke

Permalink

<https://escholarship.org/uc/item/0fq6p93h>

Journal

Tobacco Control, 27(Suppl 1)

ISSN

0964-4563

Authors

Nabavizadeh, Pooneh

Liu, Jiangtao

Havel, Christopher M

et al.

Publication Date

2018-11-01

DOI

10.1136/tobaccocontrol-2018-054325

Peer reviewed

Vascular endothelial function is impaired by aerosol from a single IQOS HeatStick to the same extent as by cigarette smoke

Pooneh Nabavizadeh¹, Jiangtao Liu¹, Christopher M. Havel², Sharina Ibrahim^{3*}, Ronak Derakhshandeh¹, Peyton Jacob, III^{2,4}, Matthew L. Springer^{1,3,4}

1 Cardiovascular Research Institute, University of California, San Francisco, San Francisco, CA 94143, USA.

2 Division of Clinical Pharmacology and Experimental Therapeutics, University of California, San Francisco, CA 94110, USA.

3 Division of Cardiology, University of California, San Francisco, San Francisco, CA 94143, USA.

4 Center for Tobacco Control Research and Education, University of California, San Francisco, San Francisco, CA 94143, USA.

*Current address: Cardiovascular Research, Vrije Universiteit Medical Center, School of Medical Sciences, Amsterdam, Netherlands

Corresponding author: Matthew L. Springer
Division of Cardiology, Box 0124, 513 Parnassus Ave.
University of California, San Francisco
San Francisco, CA 94143-0124
415-502-8404 (Telephone)
415-476-9802 (Fax)
matt.springer@ucsf.edu

ABSTRACT:

Background Heated tobacco products (also called “heat-not-burn” products) heat tobacco at temperatures below that of combustion, causing nicotine and other compounds to aerosolize. One such product, IQOS from Philip Morris International, is being marketed internationally with claims of harm reduction. We sought to determine whether exposure to IQOS aerosol impairs arterial flow-mediated dilation (FMD), a measure of vascular endothelial function that is impaired by tobacco smoke.

Methods We exposed anesthetized rats (n=8/group) via nose cone to IQOS aerosol from single HeatSticks, mainstream smoke from single Marlboro Red cigarettes, or clean air, for a series of consecutive 30-second cycles over 1.5 to 5 minutes. Each cycle consisted of 15 or 5 seconds of exposure followed by removal from the nose cone. We measured pre- and post-exposure FMD, and post-exposure serum nicotine and cotinine.

Results FMD was impaired comparably by ten 15-second exposures and ten 5-second exposures to IQOS aerosol and to cigarette smoke, but not by clean air. Serum nicotine levels were similar to plasma levels after humans have smoked one cigarette, confirming that exposure conditions had real-world relevance. Post-exposure nicotine levels were ~4.5-fold higher in rats exposed to IQOS than to cigarettes, despite nicotine being measured in the IQOS aerosol at ~63% the amount measured in smoke. When IQOS exposure was briefer, leading to comparable serum nicotine levels to the cigarette group, FMD was still comparably impaired.

Conclusions Acute exposures to IQOS aerosol impairs FMD in rats. IQOS use does not necessarily avoid the adverse cardiovascular effects of smoking cigarettes.

KEY WORDS: Electronic Nicotine Delivery Systems; heated tobacco products, heat-not-burn tobacco products; modified risk tobacco products; IQOS; endothelium; flow-mediated dilation

WHAT THIS PAPER ADDS:

- IQOS is a new heated tobacco product from Philip Morris International that has been marketed in 30 countries and has been portrayed to consumers and regulatory agencies as modified risk tobacco product.
- Brief exposures to IQOS aerosol cause endothelial dysfunction, a well-established physiological measure of cardiovascular risk that is also caused by exposure to cigarette smoke to the same extent as cigarette smoke in rats.
- Acute exposure to IQOS aerosol impairs FMD in rats.
- Use of IQOS does not necessarily avoid the adverse cardiovascular effects of cigarette smoking.

INTRODUCTION:

Heated tobacco products (also called “heat-not-burn” tobacco products) heat tobacco at temperatures that avoid combustion but cause the nicotine to aerosolize. Philip Morris International’s (PMI’s) heated tobacco product IQOS is now marketed in at least 30 non-US countries and has been considerably more successful than similar products introduced at various times over the last three decades.¹ In 2017, Philip Morris Products S.A. submitted modified risk tobacco product (MRTP) applications to the FDA to market IQOS in the US with reduced risk claims.²

IQOS is composed of three main parts: HeatStick, holder, and pocket charger (Figure 1).

HeatSticks, which contain long strips of processed and reformed tobacco packed together (Figure 1), are inserted in the holder for use. The holder contains an electronic heating blade that is activated by pressing a side button on the holder and heats the mixture of tobacco and other compounds in the HeatSticks to 350°C. Aerosol is generated as the user inhales through the HeatStick/holder combination.

Despite harm reduction claims by the tobacco industry and the clinical and basic research reported by PMI researchers,³⁻²³ the cardiovascular health effects of IQOS and similar products are incompletely understood. Notably, industry-supported studies of potential cardiovascular consequences of IQOS aerosol exposure published to date have not included some common measures of adverse effects of smoke exposure, such as vascular endothelial function tested in vivo. As IQOS has been marketed widely, it is important to study the health effects of this product so that regulatory agencies such as the FDA as well as consumers can make informed decisions.

Endothelial function assessed by arterial flow-mediated dilation (FMD) is a validated measure of cardiovascular health effects and is defined as the percent by which the arteries vasodilate in response to an increase in blood flow.²⁴ Endothelial cells lining the vessel sense increases in flow and respond by producing nitric oxide, which triggers the artery to grow in diameter to accommodate the increased demand. In humans, FMD is determined by temporarily interrupting and restoring blood flow to the forearm, and using ultrasound to measure the increase in brachial artery diameter caused by the resulting sudden increase in flow. FMD is a clinical prognostic indicator of endothelial function and cardiovascular health.²⁵⁻²⁷ Acute and chronic exposures to secondhand smoke and active smoking impair FMD in humans.²⁸⁻³¹ While such studies have typically focused on exposure to smoke, one group has reported that exposure to e-cigarette

aerosol can impair FMD³², raising the possibility that this effect is not limited to smoke from combustion and suggesting that inhalation of other non-combustible tobacco products may also have similar consequences.

We previously developed and validated an in vivo rat model for FMD measurements using high resolution ultrasound and microsurgical techniques.³³ We showed that exposure of rats to sidestream smoke (from the smoldering tip; an accepted approximation of secondhand smoke) of cigarettes, little cigars, and marijuana impairs FMD in an endothelium-dependent manner (endothelium-independent vasodilation induced by nitroglycerin is not impaired).³⁴⁻³⁶ The consistent responses of humans and rats to smoke and non-combustion aerosols from multiple sources led us to hypothesize that IQOS aerosol can similarly cause endothelial dysfunction. We report here that exposure to IQOS aerosol impairs FMD to the same extent as mainstream cigarette smoke.

MATERIALS AND METHODS:

Animals:

We used male Sprague-Dawley rats (Charles River, MI) at 8-10 weeks of age with body weights of 200 to 250 g as has been the standard condition for our previous studies on smoke exposure. Rats were anesthetized with intraperitoneal injection of ketamine (100 mg/kg)/xylazine (5 mg/kg) for these experiments. During all procedures, rats were kept on a heating pad to prevent hypothermia. Frequency and depth of respiration were closely monitored to ensure full anesthesia and supplemental IP anesthetic was given if necessary. All animal procedures were approved by the University of California, San Francisco Institutional Animal Care and Use Committee.

Mainstream smoke/aerosol generation and exposure:

Russian IQOS Parliament branded HeatSticks and an IQOS device from Ukraine were used for the first part of the project. Due to heating blade breaking incidences during cleaning, IQOS devices obtained from Japan and subsequently from South Korea were used for later exposures.

We exposed rats (n=8/group) via nose cone to IQOS aerosol, Marlboro cigarette mainstream smoke, or clean air as control. The exposure regimen consisted of a series of consecutive 30-second cycles, each consisting of 15 or 5 seconds of exposure followed by removal of the nose cone for the rest of the 30 second interval. Each rat was exposed to either 10 cycles over 5 minutes or 3 cycles over 1.5 minutes, depending on the experiment, to approximate the consumption of a single IQOS HeatStick or less. To generate the aerosol and mainstream smoke, we used a manual syringe pump system for the initial experiment and then purchased a Gram Universal Vaping Machine version 5.0 (Gram Research, Oakland, CA) for the subsequent experiments (Figure 2). The vaping machine contains an automatic syringe pump and has been shown to reproducibly generate aerosols from e-cigarette under controlled conditions.³⁷ Both systems generated 35 ml of aerosol over 2 seconds and the vents on the side of the cigarette filters remained unoccluded. Separate sets of syringe pump, nose cone, and connecting tubes and valves were used to avoid cross contamination. The IQOS holder was fully cleaned and recharged after each use.

Endothelial function measurements:

We used our previously established living rat model³³ to measure FMD before and after each exposure in individual animals.

We made a 1 cm incision on the rat's groin and surgically dissected around the common iliac artery. We then placed an arterial loop occluder consisting of a 5-0 Prolene filament under the artery and passed it through a 15-cm PE-90 tubing to enable transient occlusion of blood flow after suturing the skin. A series of diameter images of the femoral artery and accompanying Doppler blood flow images were recorded with a 35-MHz ultrasound transducer (Vevo660, VisualSonics, Toronto, Canada) before transient surgical occlusion at baseline. We induced a transient limb ischemia for 5 minutes and obtained ultrasound imaging immediately after reperfusion, and then every 30 seconds for 5 minutes

We used an automated program (Brachial Analyzer 5; Medical Imaging Applications, Coralville, IA) to measure baseline artery diameter and peak post ischemia diameter during diastole. FMD was calculated as the percent increase in the diameter of the artery after the transient ischemia. FMD was measured before and after exposures to smoke or IQOS aerosol, as summarized schematically in Figure 3A,B. The investigator was blinded to exposure conditions during FMD procedure, analysis of ultrasound images, and subsequent calculations.

Serum nicotine and cotinine measurements:

For measurement of nicotine and its metabolite cotinine, we collected blood immediately after exposure (from IQOS- and cigarette-exposed rats not being used for FMD measurement; n=5/group) and 20 minutes later (from IQOS- and cigarette-exposed rats after the final FMD measurement; n=8/group) via thoracotomy and cardiac puncture using a 23G blood collection set (BD Vacutainer) and untreated tubes. We centrifuged the blood samples at 800 rcf for 15 minutes and immediately stored the serum samples at -80°C. Concentrations of nicotine and cotinine were determined by gas chromatography - tandem mass spectrometry (GC-MS/MS).³⁸

The limits of quantitation were 1 ng/ml for nicotine and 10 ng/ml for cotinine.

Aerosol and mainstream smoke nicotine measurements:

We measured nicotine in both gas and particle phases of the IQOS aerosol and mainstream smoke from Marlboro Red cigarettes and 1R6F research cigarettes using the method described above. Particle phase nicotine was captured in mainstream aerosol and smoke on Cambridge filter pads placed downstream of the cigarette/IQOS filter, using modified ISO conditions of one 35 ml puff every 30 seconds for a total of 10 puffs (n=3/group). The filtered aerosol went through an impinger filled with 0.036N HCl to trap the gas phase nicotine. We also analyzed the post-use residual material in the HeatSticks as well as unused HeatSticks for nicotine levels. The filters and the contents of the HeatStick were extracted with 20 mL of an extraction buffer containing 25% by volume of t-butanol, 0.5 M citric acid, and 20 mM ascorbic acid by sonicating for 60 minutes at 60°C. The tubes were centrifuged at 3000 x g for 5 minutes and aliquots of the supernatant were analyzed. Liquid chromatography-tandem mass spectrometry (LC-MS/MS)³⁷ was used to measure the nicotine levels in the samples.

Statistics:

We used paired t-tests to compare the FMD values in each group before and after exposures. Errors are presented in this report as SD. A two-way repeated measures ANOVA was run to examine the effect of type and duration of exposure on percent FMD reduction. P value of <0.05 was considered statistically significant. Calculations were done with Stata 13.1.

RESULTS:

Comparable impairment of FMD by exposure to IQOS aerosol and cigarette mainstream smoke

Exposure in the first experiment (Figure 3C) consisted of 10 cycles of 15 seconds exposure + 15 seconds break (out of the nose cone). A subsequent experiment (Figure 3D) used the more realistic regimen of 10 cycles of 5 seconds exposure + 25 seconds break.

FMD was reduced comparably by ten 15-second exposures to IQOS aerosol ($10.6 \pm 2.9\%$ pre-exposure vs. $4.5 \pm 1.9\%$ post-exposure, $p = .0009$) and cigarette smoke ($10.6 \pm 2.0\%$ pre-exposure vs. $4.6 \pm 1.3\%$ post-exposure, $p = .0004$). FMD was not affected in the clean air control group ($8.3 \pm 1.9\%$ vs. $8.8 \pm 4.5\%$, $p = .82$)

FMD was also impaired comparably by ten 5-second exposures to IQOS aerosol and cigarette smoke ($10.8 \pm 1.0\%$ pre-exposure vs. $3.8 \pm 2.6\%$ post-exposure, $p = .0001$; and $11.2 \pm 2.6\%$ pre-exposure vs. $4.2 \pm 2.3\%$ post-exposure, $p = .0006$, respectively). FMD was not affected in the air control group ($9.5 \pm 3.0\%$ vs. $8.1 \pm 1.8\%$, $p = .85$).

A two-way repeated measures ANOVA revealed that there was not a significant interaction between type and duration of exposure on percent FMD reduction, $F(2, 41) = 0.30$, $p = .73$. The percent FMD impairment was not significantly different in groups exposed for 5 seconds compared with 15 seconds ($p = .27$), which suggests that the endothelial response was saturated with a single HeatStick or cigarette.

Serum nicotine was comparable to that resulting from smoking by humans:

To evaluate the accuracy of the exposure conditions in our rats and its relevance to real world levels, we compared the post cigarette exposure serum nicotine levels in rats to that reported for humans (Figure 4). In rats subjected to the 10 x (5 + 25 seconds) regimen, mean nicotine levels

in samples immediately after exposure were 70.3 ± 26.3 and 15.0 ± 7.7 ng/ml in the IQOS and cigarette groups respectively. Average nicotine levels immediately after exposure in the cigarette group was similar to the levels in humans after smoking one cigarette (~ 10 - 50 ng/ml),^{39, 40} confirming that the exposure conditions were relevant to real-world smoking. In the serum samples taken 20 minutes after the exposure, mean nicotine levels were 39.7 ± 18.9 and 5.6 ± 2.5 ng/ml in IQOS and cigarette groups. Serum cotinine levels were 4.6 ± 1.9 and 6.5 ± 2.8 ng/ml in IQOS-exposed group immediately after exposure and 20 minutes later. In the cigarette-exposed group, cotinine was undetectable after exposure and 0.8 ± 1.4 ng/ml after 20 minutes.

Serum nicotine and cotinine levels were significantly higher in the IQOS-exposed group compared with the cigarette-exposed group at all times. Nicotine and cotinine was not detected in the air groups.

Comparable FMD impairment from lower level IQOS exposure:

Because the nicotine levels in IQOS group were substantially higher than in the cigarette group, impairment of FMD in the IQOS group could potentially be attributed to excessive nicotine exposure. Therefore, we titrated the IQOS exposure timing down to conditions that led to serum nicotine comparable to the cigarette group (three 5+25 second cycles rather than ten cycles led to 14.8 ± 11.6 ng/ml serum nicotine initially). Average FMD in a group of rats exposed to three cycles of IQOS aerosol decreased significantly from 11.0 ± 4.2 to 4.5 ± 1.5 ($p = .0019$). A group of four air-exposed rats that was included for blinding purposes showed no significant change in FMD (Figure 5).

Higher nicotine in Marlboro Red smoke than IQOS aerosol:

To determine if the higher serum nicotine levels resulting from IQOS exposure than smoke exposure reflected higher nicotine concentrations in IQOS aerosol, we measured nicotine in both kinds of emissions, using both Marlboro Red cigarettes and 1R6F research cigarettes for comparison. Total nicotine levels in smoke from the Marlboro cigarettes was significantly higher than that in IQOS aerosol (1.07 ± 0.05 vs. 0.67 ± 0.02 , $p=.0006$). Nicotine levels in the 1R6F smoke were similar to IQOS aerosol levels (Table 1).

Nicotine in the IQOS aerosol gas phase was below the detection limits. Analysis of HeatSticks before and after use also revealed that the nicotine in the aerosol is mainly in the particle phase.

DISCUSSION:

Historically, the tobacco industry has made a major effort to introduce their so-called heat-not-burn tobacco products to consumers as reduced-risk products. Unlike earlier generation products such as R.J. Reynolds' Premier (1988), Eclipse (1996, 2003), and Revo (2015), and Philip Morris' Accord (1998) and Heatbar (2006), IQOS has been successfully marketed in at least 30 countries.^{1, 41, 42} Although IQOS is currently not available in the US, industry analysts are making optimistic predictions regarding successful future US marketing of IQOS.^{41, 43} Our aim was to study the vascular effects of exposure to IQOS aerosol before this product achieves more widespread use. To our knowledge, this is the first study of the effects of IQOS aerosol on vascular endothelial function in a physiologic model.

PMI strongly implies in publications^{21, 23, 44} and in their MRTTP applications⁴⁵ that IQOS causes less endothelial dysfunction than conventional cigarettes. Their conclusions were based

only on cultured cell behavior²¹ and circulating sICAM-1 protein²³ that indirectly reflect processes relevant to vascular function, but do not directly assess endothelial function physiologically. PMI's physiological cardiovascular studies were limited to the demonstration that long-term IQOS aerosol exposure led to less aortic plaque than cigarette smoke in transgenic mice predisposed to atherosclerosis.⁴⁴ In contrast, our direct evaluation of IQOS aerosol's acute effects showed that brief exposures to IQOS aerosol from a single HeatStick cause rapid impairment of vascular endothelial function in rats comparable to that caused by cigarette smoke. We assessed endothelial function with FMD, a validated measure of cardiovascular health effects²⁴⁻²⁷ that has been used by numerous groups to evaluate cardiovascular effects of smoke exposure in humans and animals.^{28-31, 34-36} Therefore, our approach provides a robust readout of potential adverse effects of IQOS aerosol on endothelial function, and indicates that IQOS use is likely to have rapid adverse vascular effects comparable to those from cigarette smoking.

We chose to use a nose cone smoke/aerosol delivery system, rather than a whole body exposure system, to prevent exposure via the ocular and oral routes and also to prevent subsequent oral exposure from licking fur and paws. Establishment of an exact relationship between our rat exposure conditions and human exposure is limited by differences in route of administration (rats are obligate nose breathers with complex nasal topography⁴⁶ while humans inhale cigarette smoke orally), differences between human intentional aerosol inhalation volume and anesthetized rat tidal (inhalation) volumes,^{47, 48} and differences in airway clearance rates.⁴⁹ Sophisticated models have been developed to convert inhalation dosimetry between rats and humans,⁴⁹ but smoke does not present a straightforward situation due to the combination of gas and highly heterogeneous particles that behave differently in such models. Therefore, to assess whether our exposure conditions were relevant to real-world smoking, we measured serum

nicotine levels and confirmed that nicotine in the rats that were exposed to cigarette smoke for ten 5+25 second cycles were similar to the levels found in humans immediately after smoking one cigarette, validating our exposure model for cigarettes.

However, the serum nicotine level in the IQOS-exposed group under identical conditions was more than four times higher than in the cigarette group. This was surprising; our results revealed that the nicotine content in the IQOS aerosol in our generation system was much lower than that in the Marlboro mainstream smoke (Table 1). Moreover, nicotine content in the filler and aerosol is reported in the literature to be roughly comparable for HeatSticks and cigarettes.^{50, 51} The underlying reason for this remains unclear and deserves further investigation. One potential explanation for this difference could be particle size difference, which determines to what extent the particles reach the respiratory zone. Nonetheless, reducing the number of IQOS exposure cycles from ten to three to result in comparable serum nicotine concentrations to the cigarette group still led to impairment of FMD. Of note, the extents of FMD impairment in the three IQOS and cigarette groups were quite similar, indicating an extremely rapid saturation of the response to mainstream levels of smoke and IQOS aerosol.

A limitation of this study is that all functional measurements were obtained in anesthetized rats, which was necessary for the FMD procedure to be carried out. However, as in our previous reports using ketamine/xylazine,³⁴⁻³⁶ as well as our prior study unrelated to smoke in which isoflurane was used,⁵² negative control groups showed no significant alteration of FMD from pre- to post-intervention, and the relative responses of experimental and control groups have been consistent and reproducible. Moreover, prospective human exposure experiments have shown similar impairment of FMD by smoke inhalation in conscious individuals.^{30, 31, 53}

Regarding potential variability between experiments, we used a manual syringe-pump-driven

system to generate smoke and IQOS aerosol for our initial experiment, and a commercially available automatic vaping system for our subsequent experiments. The manual system was modelled after the automatic system; that is, the relevant volumes and aerosol paths were the same in all experiments. The volume of air drawn through the cigarettes and HeatSticks, and the time over which the syringe pump plunger was moved, were the same in all experiments; albeit slightly more accurate with the automatic system. Similarly, due to broken heating blades as described in the Methods, we used IQOS holders from three different countries over the course of the project. The characteristics of these electronic devices are presumably highly controlled and would not be expected to vary greatly by country. We did not perform comparative analyses of aerosols generated from the three devices. When it was necessary to switch from a broken device to a new one, rats in both experimental and control groups were exposed to aerosol from the new device.

We conclude that mainstream IQOS aerosol from a single HeatStick can rapidly and substantially impair endothelial function in rats comparably to smoke from a cigarette. While these findings do not prove that inhalation of IQOS aerosol causes endothelial dysfunction in humans, the results underscore that integrative physiological assays of function can reveal adverse health effects not noted in PMI's biomarker and cell culture studies,² and suggest that at least some of the adverse health effects of cigarettes may not be avoided by using IQOS.

ACKNOWLEDGEMENTS:

Gas chromatography-tandem mass spectrometry analyses were performed by Kristina Bello and Lawrence Chan in the Clinical Pharmacology Research Laboratory at UCSF, supervised by Peyton Jacob, III. We thank Dorie Apollonio and Minji Kim for helping us obtain some of the

IQOS devices used in this study, and Stanton Glantz and Xiaoyin Wang for helpful discussions during the design of the project and critical comments on the manuscript.

FUNDING:

This work was supported by grant R01HL120062 from the National Heart, Lung, and Blood Institute at the National Institutes of Health (NIH) and the US Food and Drug Administration Center for Tobacco Products (FDA CTP), grant 25IR-0030 from the California Tobacco-Related Disease Research Program, and grant P50CA180890 from the National Cancer Institute at the NIH and FDA CTP. Laboratory resources for analytical chemistry were supported by grant P30 DA012393 from the National Institute on Drug Abuse at the NIH. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or the FDA.

DISCLOSURE:

None

REFERENCES:

- [1] Philip Morris International. Designing a Smoke-Free Future (Accessed January 24, 2018). <https://www.pmi.com>.
- [2] U.S. Food and Drug Administration. Philip Morris Products S.A. Modified Risk Tobacco Product (MRTP) Applications (5/24/17; accessed 6/9/17). <https://www.fda.gov/TobaccoProducts/Labeling/MarketingandAdvertising/ucm546281.htm>.
- [3] Schorp MK, Tricker AR, Dempsey R. Reduced exposure evaluation of an Electrically Heated Cigarette Smoking System. Part 1: Non-clinical and clinical insights. *Regul Toxicol Pharmacol* 2012;**64**(2 Suppl): S1-10.

- [4] Zenzen V, Diekmann J, Gerstenberg B, *et al.* Reduced exposure evaluation of an Electrically Heated Cigarette Smoking System. Part 2: Smoke chemistry and in vitro toxicological evaluation using smoking regimens reflecting human puffing behavior. *Regul Toxicol Pharmacol* 2012;**64**(2 Suppl): S11-34.
- [5] Tricker AR, Stewart AJ, Leroy CM, *et al.* Reduced exposure evaluation of an Electrically Heated Cigarette Smoking System. Part 3: Eight-day randomized clinical trial in the UK. *Regul Toxicol Pharmacol* 2012;**64**(2 Suppl): S35-44.
- [6] Tricker AR, Jang IJ, Martin Leroy C, *et al.* Reduced exposure evaluation of an Electrically Heated Cigarette Smoking System. Part 4: Eight-day randomized clinical trial in Korea. *Regul Toxicol Pharmacol* 2012;**64**(2 Suppl): S45-53.
- [7] Tricker AR, Kanada S, Takada K, *et al.* Reduced exposure evaluation of an Electrically Heated Cigarette Smoking System. Part 5: 8-Day randomized clinical trial in Japan. *Regul Toxicol Pharmacol* 2012;**64**(2 Suppl): S54-63.
- [8] Tricker AR, Kanada S, Takada K, *et al.* Reduced exposure evaluation of an Electrically Heated Cigarette Smoking System. Part 6: 6-Day randomized clinical trial of a menthol cigarette in Japan. *Regul Toxicol Pharmacol* 2012;**64**(2 Suppl): S64-73.
- [9] Martin Leroy C, Jarus-Dziedzic K, Ancerewicz J, *et al.* Reduced exposure evaluation of an Electrically Heated Cigarette Smoking System. Part 7: A one-month, randomized, ambulatory, controlled clinical study in Poland. *Regul Toxicol Pharmacol* 2012;**64**(2 Suppl): S74-84.
- [10] Urban HJ, Tricker AR, Leyden DE, *et al.* Reduced exposure evaluation of an Electrically Heated Cigarette Smoking System. Part 8: Nicotine bridging--estimating smoke constituent exposure by their relationships to both nicotine levels in mainstream cigarette smoke and in smokers. *Regul Toxicol Pharmacol* 2012;**64**(2 Suppl): S85-97.
- [11] Smith MR, Clark B, Ludicke F, *et al.* Evaluation of the Tobacco Heating System 2.2. Part 1: Description of the system and the scientific assessment program. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S17-S26.
- [12] Schaller JP, Keller D, Poget L, *et al.* Evaluation of the Tobacco Heating System 2.2. Part 2: Chemical composition, genotoxicity, cytotoxicity, and physical properties of the aerosol. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S27-S47.
- [13] Schaller JP, Pijnenburg JP, Ajithkumar A, *et al.* Evaluation of the Tobacco Heating System 2.2. Part 3: Influence of the tobacco blend on the formation of harmful and potentially harmful constituents of the Tobacco Heating System 2.2 aerosol. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S48-S58.
- [14] Wong ET, Kogel U, Veljkovic E, *et al.* Evaluation of the Tobacco Heating System 2.2. Part 4: 90-day OECD 413 rat inhalation study with systems toxicology endpoints demonstrates reduced exposure effects compared with cigarette smoke. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S59-S81.
- [15] Sewer A, Kogel U, Talikka M, *et al.* Evaluation of the Tobacco Heating System 2.2 (THS2.2). Part 5: microRNA expression from a 90-day rat inhalation study indicates that exposure to THS2.2 aerosol causes reduced effects on lung tissue compared with cigarette smoke. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S82-S92.
- [16] Oviedo A, Lebrun S, Kogel U, *et al.* Evaluation of the Tobacco Heating System 2.2. Part 6: 90-day OECD 413 rat inhalation study with systems toxicology endpoints demonstrates reduced exposure effects of a mentholated version compared with mentholated and non-mentholated cigarette smoke. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S93-S122.

- [17] Kogel U, Titz B, Schlage WK, *et al.* Evaluation of the Tobacco Heating System 2.2. Part 7: Systems toxicological assessment of a mentholated version revealed reduced cellular and molecular exposure effects compared with mentholated and non-mentholated cigarette smoke. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S123-S138.
- [18] Haziza C, de La Bourdonnaye G, Skiada D, *et al.* Evaluation of the Tobacco Heating System 2.2. Part 8: 5-Day randomized reduced exposure clinical study in Poland. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S139-S150.
- [19] Martin F, Talikka M, Ivanov NV, *et al.* Evaluation of the tobacco heating system 2.2. Part 9: Application of systems pharmacology to identify exposure response markers in peripheral blood of smokers switching to THS2.2. *Regul Toxicol Pharmacol* 2016;**81 Suppl 2**: S151-S157.
- [20] van der Toorn M, Frentzel S, De Leon H, *et al.* Aerosol from a candidate modified risk tobacco product has reduced effects on chemotaxis and transendothelial migration compared to combustion of conventional cigarettes. *Food Chem Toxicol* 2015;**86**: 81-87.
- [21] Poussin C, Laurent A, Peitsch MC, *et al.* Systems toxicology-based assessment of the candidate modified risk tobacco product THS2.2 for the adhesion of monocytic cells to human coronary arterial endothelial cells. *Toxicology* 2016;**339**: 73-86.
- [22] Ludicke F, Picavet P, Baker G, *et al.* Effects of Switching to the Tobacco Heating System 2.2 Menthol, Smoking Abstinence, or Continued Cigarette Smoking on Biomarkers of Exposure: A Randomized, Controlled, Open-Label, Multicenter Study in Sequential Confinement and Ambulatory Settings (Part 1). *Nicotine Tob Res* 2018;**20**(2): 161-172.
- [23] Ludicke F, Picavet P, Baker G, *et al.* Effects of Switching to the Menthol Tobacco Heating System 2.2, Smoking Abstinence, or Continued Cigarette Smoking on Clinically Relevant Risk Markers: A Randomized, Controlled, Open-Label, Multicenter Study in Sequential Confinement and Ambulatory Settings (Part 2). *Nicotine Tob Res* 2018;**20**(2): 173-182.
- [24] Pyke KE, Tschakovsky ME. The relationship between shear stress and flow-mediated dilatation: implications for the assessment of endothelial function. *J Physiol* 2005;**568**(Pt 2): 357-369.
- [25] Flammer AJ, Anderson T, Celermajer DS, *et al.* The assessment of endothelial function: from research into clinical practice. *Circulation* 2012;**126**(6): 753-767.
- [26] Celermajer DS, Sorensen KE, Gooch VM, *et al.* Non-invasive detection of endothelial dysfunction in children and adults at risk of atherosclerosis. *Lancet* 1992;**340**(8828): 1111-1115.
- [27] Nabel EG, Selwyn AP, Ganz P. Large coronary arteries in humans are responsive to changing blood flow: an endothelium-dependent mechanism that fails in patients with atherosclerosis. *J Am Coll Cardiol* 1990;**16**(2): 349-356.
- [28] Celermajer DS, Sorensen KE, Georgakopoulos D, *et al.* Cigarette smoking is associated with dose-related and potentially reversible impairment of endothelium-dependent dilation in healthy young adults. *Circulation* 1993;**88**(5 Pt 1): 2149-2155.
- [29] Celermajer DS, Adams MR, Clarkson P, *et al.* Passive smoking and impaired endothelium-dependent arterial dilatation in healthy young adults. *N Engl J Med* 1996;**334**(3): 150-154.
- [30] Kato T, Inoue T, Morooka T, *et al.* Short-term passive smoking causes endothelial dysfunction via oxidative stress in nonsmokers. *Can J Physiol Pharmacol* 2006;**84**(5): 523-529.
- [31] Heiss C, Amabile N, Lee AC, *et al.* Brief secondhand smoke exposure depresses endothelial progenitor cells activity and endothelial function: sustained vascular injury and blunted nitric oxide production. *J Am Coll Cardiol* 2008;**51**(18): 1760-1771.

- [32] Carnevale R, Sciarretta S, Violi F, *et al.* Acute Impact of Tobacco vs Electronic Cigarette Smoking on Oxidative Stress and Vascular Function. *Chest* 2016;**150**(3): 606-612.
- [33] Heiss C, Sievers RE, Amabile N, *et al.* In vivo measurement of flow-mediated vasodilation in living rats using high resolution ultrasound. *Am J Physiol Heart Circ Physiol* 2008;**294**(2): H1086-1093.
- [34] Pinnamaneni K, Sievers RE, Sharma R, *et al.* Brief exposure to secondhand smoke reversibly impairs endothelial vasodilatory function. *Nicotine Tob Res* 2014;**16**(5): 584-590.
- [35] Liu J, Wang X, Narayan S, *et al.* Impairment of endothelial function by little cigar secondhand smoke. *Tob Regul Sci* 2016;**2**(1): 56-63.
- [36] Wang X, Derakhshandeh R, Liu J, *et al.* One minute of marijuana secondhand smoke exposure substantially impairs vascular endothelial function. *J Am Heart Assoc* 2016;**5**: e003858.
- [37] Havel CM, Benowitz NL, Jacob P, 3rd, *et al.* An Electronic Cigarette Vaping Machine for the Characterization of Aerosol Delivery and Composition. *Nicotine Tob Res* 2017;**19**(10): 1224-1231.
- [38] Jacob P, 3rd, Wu S, Yu L, *et al.* Simultaneous determination of mecamylamine, nicotine, and cotinine in plasma by gas chromatography-mass spectrometry. *J Pharm Biomed Anal* 2000;**23**(4): 653-661.
- [39] Russell MA, Feyerabend C, Cole PV. Plasma nicotine levels after cigarette smoking and chewing nicotine gum. *British Medical Journal* 1976;**1**(6017): 1043-1046.
- [40] Benowitz NL, Jacob P, 3rd. Daily intake of nicotine during cigarette smoking. *Clin Pharmacol Ther* 1984;**35**(4): 499-504.
- [41] Caputi TL. Industry watch: heat-not-burn tobacco products are about to reach their boiling point. *Tob Control* 2017;**26**(5): 609-610.
- [42] Dutra LM, Grana R, Glantz SA. Philip Morris research on precursors to the modern e-cigarette since 1990. *Tob Control* 2017;**26**(e2): e97-e105.
- [43] Herzog B, Kanada P, Scott A. "Tobacco Talk " -- Q1 U.S. Retailer Survey, 4/26/17. 2017.
- [44] Phillips B, Veljkovic E, Boué S, *et al.* An 8-month systems toxicology inhalation/cessation study in Apoe^{-/-} mice to investigate cardiovascular and respiratory exposure effects of a candidate modified risk tobacco product, THS 2.2, compared with conventional cigarettes. *Toxicological Sciences* 2015;**149**(2): 411-432.
- [45] Administration USFaD. Philip Morris Products S.A. Modified Risk Tobacco Product (MRTP) Applications Appendix A6.1.4.3
<https://www.fda.gov/TobaccoProducts/Labeling/MarketingandAdvertising/ucm546281.htm>.
- [46] Reznik GK. Comparative anatomy, physiology, and function of the upper respiratory tract. *Environ Health Perspect* 1990;**85**: 171-176.
- [47] Lai YL, Hildebrandt J. Respiratory mechanics in the anesthetized rat. *J Appl Physiol Respir Environ Exerc Physiol* 1978;**45**(2): 255-260.
- [48] Whitehead GS, Kimmel EC, Reboulet JE, *et al.* Pulmonary function in normal rats. *Naval Health Research Center Detachment Report No Toxdet 99-5*. Bethesda, MD: Naval Medical Research and Development Command 1999.
- [49] Kuempel ED, Sweeney LM, Morris JB, *et al.* Advances in Inhalation Dosimetry Models and Methods for Occupational Risk Assessment and Exposure Limit Derivation. *J Occup Environ Hyg* 2015;**12 Suppl 1**: S18-40.

- [50] Bekki K, Inaba Y, Uchiyama S, *et al.* Comparison of Chemicals in Mainstream Smoke in Heat-not-burn Tobacco and Combustion Cigarettes. *Journal of UOEH* 2017;**39**(3): 201-207.
- [51] Auer R, Concha-Lozano N, Jacot-Sadowski I, *et al.* Heat-not-burn tobacco cigarettes: Smoke by any other name. *JAMA Internal Medicine* 2017;**177**(7): 1050-1052.
- [52] Chen Q, Sievers RE, Varga M, *et al.* Pharmacological inhibition of S-nitrosoglutathione reductase improves endothelial vasodilatory function in rats in vivo. *J Appl Physiol* 2013;**114**(6): 752-760.
- [53] Frey PF, Ganz P, Hsue PY, *et al.* The exposure-dependent effects of aged secondhand smoke on endothelial function. *J Am Coll Cardiol* 2012;**59**(21): 1908-1913.

Table 1. Nicotine concentration in IQOS aerosol, cigarette mainstream smoke, and IQOS tobacco

Material	Total Nicotine (mg) Mean±SD
IQOS Aerosol (n=3)	
<i>Particle phase</i>	0.67±0.02*
<i>Gas phase</i>	BLQ
Marlboro Red MS (n=3)	
<i>Particle phase</i>	1.07±0.05*
1R6F MS (n=3)	
<i>Particle phase</i>	0.65±0.04
Used IQOS HeatStick (residual; n=3)	3.16±0.67
Unused IQOS HeatStick (n=4)	3.92±0.11

**p=.0006 between starred values; BLQ=Below Level of Quantification; MS=Mainstream Smoke.*

FIGURE LEGENDS

Figure 1. IQOS. IQOS is composed of three main parts: HeatStick, holder, and pocket charger. HeatSticks are inserted in the holder, which contains an electronic heating blade to heat tobacco and release aerosol. HeatSticks contain strips of processed and reformed tobacco. Photos by M.L.S and P.N.

Figure 2. Aerosol generation and exposure systems. A. Manual exposure system; B. Analytical vaping machine made by Gram Research; C. IQOS aerosol coming out of nose cone; D. Rat's nose placed in the nose cone. Photos in A, C, and D by P.N.; photo in B from Gram Research with permission.

Figure 3. Arterial flow-mediated dilation was impaired by mainstream cigarette smoke and IQOS aerosol. A. Ultrasound imaging of rat femoral artery. B. FMD experimental design: FMD was measured pre- and post-exposure in each rat. C. FMD after ten cycles of 15-second exposure + 15-second break. D. FMD after ten cycles of 5-second exposure + 25-second break. Colored lines denote individual rats pre- and post-exposure; bars denote group means; p values are derived from paired 2-tailed t-tests.

Figure 4. Serum nicotine and cotinine levels immediately and 20 min post-exposure. Samples were taken after ten cycles of 5 seconds exposure + 25 seconds break. p values are derived from 2-tailed student t-tests. IQOS x10: 10 IQOS exposure cycles, Cigarette x10: 10 cigarette exposure cycles, Air x10: 10 air exposure cycles, IQOS x3: 3 IQOS exposure cycles, BLQ: Below Level of Quantification.

Figure 5. Reduced IQOS exposure to match nicotine absorption level of the cigarette group

still impairs FMD to a comparable extent. Colored lines denote individual rats pre- and post-exposure; bars denote group means; p values are derived from paired 2-tailed t-tests. Exposure conditions were 3 cycles of 5-second exposure + 25-second break.

FIGURE LEGENDS



Figure 1. IQOS. IQOS is composed of three main parts: HeatStick, holder, and pocket charger. HeatSticks are inserted in the holder, which contains an electronic heating blade to heat tobacco and release aerosol. HeatSticks contain strips of processed and reformed tobacco. Photos by M.L.S and P.N.




Figure 2. Aerosol generation and exposure systems. A. Manual exposure system; B. Analytical vaping machine made by Gram Research; C. IQOS aerosol coming out of nose cone; D. Rat's nose placed in the nose cone. Photos in A, C, and D by P.N.; photo in B from Gram Research with permission.




Figure 3. Arterial flow-mediated dilation was impaired by mainstream cigarette smoke and IQOS aerosol. A. Ultrasound imaging of rat femoral artery. B. FMD experimental design: FMD was measured pre- and post-exposure in each rat. C. FMD after ten cycles of 15-second exposure + 15-second break. D. FMD after ten cycles of 5-second exposure + 25-second break. Colored lines denote individual rats pre- and post-exposure; bars denote group means; p values are derived from paired 2-tailed t-tests.




Figure 4. Serum nicotine and cotinine levels immediately and 20 min post-exposure.

Samples were taken after ten cycles of 5 seconds exposure + 25 seconds break. p values are derived from 2-tailed student t-tests. IQOS x10: 10 IQOS exposure cycles, Cigarette x10: 10 cigarette exposure cycles, Air x10: 10 air exposure cycles, IQOS x3: 3 IQOS exposure cycles, BLQ: Below Level of Quantification.




Figure 5. Reduced IQOS exposure to match nicotine absorption level of the cigarette group still impairs FMD to a comparable extent. Colored lines denote individual rats pre- and post-exposure; bars denote group means; p values are derived from paired 2-tailed t-tests. Exposure conditions were 3 cycles of 5-second exposure + 25-second break.