

UC Riverside

UC Riverside Previously Published Works

Title

Peripheral blood non-canonical small non-coding RNAs as novel biomarkers in lung cancer

Permalink

<https://escholarship.org/uc/item/0fz218m0>

Journal

Molecular Cancer, 19(1)

ISSN

1476-4598

Authors

Gu, Wanjun

Shi, Junchao

Liu, Hui

et al.

Publication Date

2020-12-01

DOI

10.1186/s12943-020-01280-9

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>


Peer reviewed

LETTER TO THE EDITOR

Open Access



Peripheral blood non-canonical small non-coding RNAs as novel biomarkers in lung cancer

Wanjun Gu^{1*†}, Junchao Shi^{2†}, Hui Liu^{3†}, Xudong Zhang^{2†}, Jin J. Zhou⁴, Musheng Li⁵, Dandan Zhou⁵, Rui Li³, Jingzhu Lv³, Guoxia Wen¹, Shanshan Zhu¹, Ting Qi¹, Wei Li⁶, Xiaojing Wang⁶, Zhaohua Wang^{7*}, Hua Zhu⁸, Changcheng Zhou², Kenneth S. Knox⁹, Ting Wang⁹, Qi Chen^{2*}, Zhongqing Qian^{3*} and Tong Zhou^{5*} 

Abstract

One unmet challenge in lung cancer diagnosis is to accurately differentiate lung cancer from other lung diseases with similar clinical symptoms and radiological features, such as pulmonary tuberculosis (TB). To identify reliable biomarkers for lung cancer screening, we leverage the recently discovered non-canonical small non-coding RNAs (i.e., tRNA-derived small RNAs [tsRNAs], rRNA-derived small RNAs [rsRNAs], and YRNA-derived small RNAs [ysRNAs]) in human peripheral blood mononuclear cells and develop a molecular signature composed of distinct ts/rs/ysRNAs (TRY-RNA). Our TRY-RNA signature precisely discriminates between control, lung cancer, and pulmonary TB subjects in both the discovery and validation cohorts and outperforms microRNA-based biomarkers, which bears the diagnostic potential for lung cancer screening.

Keywords: Lung cancer, Tuberculosis, tsRNA, rsRNA, ysRNA

One unmet challenge in current lung cancer diagnosis is to accurately differentiate lung cancer from other lung diseases with similar clinical symptoms and radiological features. Imaging-based screening methods, such as low-dose computed topography (LDCT), could sometimes be false positives, as indeterminate pulmonary nodules may also be caused by other lung diseases such as pulmonary

tuberculosis (TB) [1], which is especially concerning for clinical practice in TB-endemic countries/regions. Therefore, additional noninvasive diagnostic procedures are much-needed to avoid a misdiagnosis in patients with lung cancer mimicking pulmonary TB, or vice versa. Here, we aim to develop a peripheral blood mononuclear cell (PBMC)-based molecular signature to differentiate lung cancer patients from healthy controls and pulmonary TB patients by harnessing the novel small non-coding RNAs (sncRNAs).

Recent sncRNA sequencing (sncRNA-seq) attempts have ubiquitously detected several non-canonical sncRNA types, which are fragments derived from canonically transcribed parent large RNAs, including tRNA-derived small RNAs (tsRNAs), rRNA-derived small RNAs (rsRNAs), and YRNA-derived small RNAs (ysRNAs) [2]. ts/rs/ysRNAs have been discovered in a wide range of species [2]. The biological functions of tsRNAs have become a recent highlight and been linked with various human diseases [3], including cancers [4–6],

* Correspondence: wanjungu@seu.edu.cn; qi.chen@medsch.ucr.edu; qzq7778@bbmc.edu.cn; tongz@med.unr.edu

[†]Zhaohua Wang is deceased.

¹State Key Laboratory of Bioelectronics, School of Biological Sciences and Medical Engineering, Southeast University, 2 Sipailou, Nanjing 210096, Jiangsu, China

²Division of Biomedical Sciences, School of Medicine, University of California, Riverside, 900 University Avenue, Riverside, CA 92521, USA

³Anhui Province Key Laboratory of Immunology in Chronic Diseases, Anhui Key Laboratory of Infection and Immunity, Department of Laboratory Medicine, Bengbu Medical College, 2600 Donghaidadao, Bengbu 233003, Anhui, China

⁵Department of Physiology and Cell Biology, University of Nevada, Reno School of Medicine, 1664 North Virginia Street, Reno, Nevada 89557, USA
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

while rsRNAs and ysRNAs show sensitive response to pathophysiological conditions [6, 7]. In this study, we develop a diagnostic signature composed of distinct ts/rs/ysRNAs (TRY-RNA) in human PBMCs. Our TRY-RNA signature accurately discriminates between control, lung cancer, and pulmonary TB subjects in both the discovery and validation cohorts and outperforms microRNA (miRNA)-based biomarkers. Fig. S1 (Additional file 1) provides an overview of the experimental design.

Dysregulated non-canonical sncRNAs in lung cancer

We performed sncRNA-seq for the PBMC samples collected from 59 human subjects in the discovery cohort, including 13 healthy controls, 10 pulmonary TB patients, and 36 lung cancer patients (Additional file 2: Table S1). The raw sequencing data were processed by our newly developed computational framework, *SPORTS1.0*, which was designed to optimize the annotation and quantification of non-canonical sncRNAs (i.e., tsRNAs, rsRNAs, and ysRNAs) in addition to miRNAs [2]. In total, 6673 tsRNA species, 20,172 rsRNA species, 1238 ysRNA species, and 973 miRNA species were identified in human PBMCs (Additional file 1: Fig. S2). We investigated the co-expression pattern of tsRNAs across the PBMC samples in the discovery cohort by grouping tsRNA species into subcategories according to their parent tRNA types. We found that the expression of the tsRNAs derived from the tRNAs of alanine (tsRNA-Ala), asparagine (tsRNA-Asn), leucine (tsRNA-Leu), lysine (tsRNA-Lys), and tyrosine (tsRNA-Tyr) was strongly and positively correlated with that of each other (*Spearman's* rank correlation test: $\rho > 0.700$ and $P < 10^{-9}$) (Fig. 1a), suggesting shared biogenesis pathways among these tsRNAs. Interestingly, tsRNA-Ala, tsRNA-Asn, tsRNA-Leu, tsRNA-Lys, and tsRNA-Tyr were the only five tsRNA groups that were upregulated in the lung cancer patients relative to the controls (adjusted $P < 0.05$) (Fig. 1b). We further found that the expression of these five tsRNA groups was also significantly higher in the lung cancer patients than in the pulmonary TB subjects ($P < 0.05$) (Fig. 1b). We next grouped rsRNA and ysRNA species into subcategories according to their parent rRNA/YRNA types and found that the rsRNAs derived from rRNA-5S (rsRNA-5S) were significantly upregulated in the lung cancer patients relative to the controls, while the ysRNAs originating from YRNA-RNY1 (ysRNA-RNY1) were downregulated in the lung cancer patients compared with the controls (adjusted $P < 0.05$) (Fig. 1b). More interestingly, the expression of rsRNA-5S and ysRNA-RNY1 showed a completely inverse pattern in the pulmonary TB patients: rsRNA-5S was significantly downregulated in the TB patients relative to the controls, while ysRNA-RNY1 was upregulated in the TB patients

compared with the controls ($P < 0.05$) (Fig. 1b). We further mapped the individual tsRNA-Ala, tsRNA-Asn, tsRNA-Leu, tsRNA-Lys, tsRNA-Tyr, rsRNA-5S, and ysRNA-RNY1 species to the corresponding parent RNAs and identified a nonrandom fragmentation pattern (Fig. 1c-d and Additional file 1: Fig. S3), suggesting highly regulated biogenesis of these sncRNAs. In addition, we investigated the association of these non-canonical sncRNA expression with cancer stage, histological type, lymph node status, metastasis status, and smoking history, but no significant difference was observed (Additional file 1: Fig. S4-S8).

The molecular signature composed of non-canonical sncRNAs

We next develop a molecular signature of sncRNAs by harnessing the above prioritized sncRNA subcategories (i.e., tsRNA-Ala, tsRNA-Asn, tsRNA-Leu, tsRNA-Lys, tsRNA-Tyr, rsRNA-5S, and ysRNA-RNY1). In total, nine tsRNA species, eight rsRNA species, and eight ysRNA species (Additional file 1: Fig. S9A-C) were selected, which consisted of a molecular signature with 25 distinct non-canonical sncRNAs (Additional file 1: Fig. S9D and Additional file 2: Table S2), referred to as the TS/RS/YS-RNA (TRY-RNA) signature. Both principal component analysis (Additional file 1: Fig. S9E) and hierarchical clustering on RNA expression (Fig. 1e) confirmed the discriminative power of the TRY-RNA signature between the control, lung cancer, and TB groups in the discovery cohort. To systematically evaluate the classification power of the TRY-RNA signature, a TRY-RNA index was assigned to each subject based on the expression of the ts/rs/ysRNAs within the TRY-RNA signature (Additional file 3: Methods). The TRY-RNA index was a linear combination of the expression values of the sncRNA species within the TRY-RNA signature. A higher TRY-RNA index implies a higher likelihood of lung cancer. We found that the TRY-RNA index was significantly higher in the lung cancer patients than in the healthy controls, while the TRY-RNA index of the pulmonary TB patients was significantly lower than that of the controls (*t*-test: $P < 10^{-5}$) (Additional file 1: Fig. S10A). The area under the receiver operating characteristic (ROC) curve (*AUC*) was 1.000 between the cancer and non-cancer subjects and 0.994 between the TB and non-TB subjects (Additional file 1: Fig. S10B). In addition, we investigated the association of the expression of the individual RNA species within the TRY-RNA signature with cancer stage, histological type, lymph node status, metastasis status, and smoking history, but significant difference was only observed for ysRNA-RNY1-28 and ysRNA-RNY1-29a between adenocarcinoma and squamous cell carcinoma patients (Additional file 1: Fig. S11-S15).





Fig. 1 The dysregulated non-canonical sncRNAs in lung cancer. **a** The co-expression pattern tsRNA subcategories across the PBMC samples in the discovery cohort. The correlation coefficient was calculated by *Spearman's* rank correlation test. tsRNA-Ala, tsRNA-Asn, tsRNA-Leu, tsRNA-Lys, and tsRNA-Tyr were strongly and positively correlated with each other. **b** The expression profile of tsRNA-Ala, tsRNA-Asn, tsRNA-Leu, tsRNA-Lys, tsRNA-Tyr, rsRNA-5S, and ysRNA-RNY1 among the control, lung cancer, and TB subjects. *RPM*: reads per million. **c** and **d** The coverage profile of the PBMC rsRNA-5S and ysRNA-RNY1 sequences along rRNA-5S and YRNA-RNY1, respectively. The solid curves indicate the mean *RPM* values for the control, lung cancer, and TB groups. The colored bands represent the 95% confidence interval. nt: nucleotide. rsRNA-5S sequences were primarily derived from the 3'-end of rRNA-5S, whereas ysRNA-RNY1 sequences were largely derived from the 5'-end of YRNA-RNY1 with a small portion of fragments mapped to the 3'-end of YRNA-RNY1. **e** Expression heatmap of the sncRNA species within the TRY-RNA signature in the discovery cohort

The performance of the TRY-RNA signature in the validation cohort

We further assessed the TRY-RNA signature in the validation cohort with 35 human PBMC samples collected from 12 healthy controls, 15 lung cancer patients, and 8 pulmonary TB patients (Additional file 2: Table S3). Un-supervised hierarchical clustering and principal component analysis demonstrated a totally distinct expression

pattern of the TRY-RNA signature between the lung cancer and TB subjects, with the controls largely falling in between in the validation cohort (Fig. 2a-b). The TRY-RNA index in the validation cohort was significantly higher in the lung cancer patients than in the healthy controls, while the TRY-RNA index of the TB patients was significantly lower than that of the controls (*t*-test: *P* < 0.005) (Fig. 2c). The *AUC* was 0.930 between the cancer and non-cancer



subjects and 1.000 between the TB and non-TB subjects (Fig. 2d), which suggests the strong classification power of the TRY-RNA signature for both lung cancer and pulmonary TB screening.

Comparison between the TRY-RNA and miRNA-based signatures

We also compared the expression profile of miRNAs among the control, lung cancer, and pulmonary TB subjects in the discovery cohort and identified a signature with 43 miRNA species, referred to as the MIR signature (Additional file 1: Fig. S16 and Additional file 2: Table S4). Similar to the TRY-RNA signature, we assigned a MIR index to each subject based on the expression of the miRNAs within the MIR signature. We found that in both the discovery and validation cohorts, while the MIR index can differentiate the lung cancer patients from the control and TB subjects (Additional file 1: Fig. S17 and S18A-B), a resampling test (Additional file 3: Methods) demonstrated a superior classification power of the TRY-RNA signature compared to the MIR signature (Additional file 1: Fig. S18C). We further investigated whether the MIR signature provided additive classification power to the TRY-RNA signature by combining both signatures (referred to as the

TRY-RNA \cup MIR signature). Although the performance of the TRY-RNA \cup MIR signature was fairly good, the TRY-RNA \cup MIR signature didn't outperform the TRY-RNA signature and barely provided additive information for the classification (Additional file 1: Fig. S18D-F).

Conclusions

Our TRY-RNA signature derived from the repertoire of PBMC non-canonical sncRNAs makes it possible for the early diagnosis of lung cancer and pulmonary TB, which may reflect the host responses to different antigens and would represent an improvement over the previous studies focusing solely on tsRNAs in cancer tissues [4, 5, 8]. Interestingly, the performance of the TRY-RNA signature shows superiority over the miRNA-based signature, which could be due to the more complex layer of non-canonical sncRNAs. For example, tsRNAs and rsRNAs exhibit an unexpected complexity in regards to their RNA modifications as well as their sequence diversities [9]. Our previous study suggests that both tsRNAs and rsRNAs are involved in mammalian epigenetic inheritance, which form a "RNA code" to convey environmental clue to the offspring [7, 10]. Also, tsRNAs are thought to regulate translation process and ribosome biogenesis in versatile ways, including the fine-tuning of the ribosome composition that may affect the

translational specificity on a selective pool of mRNAs (also referred to as ribosome heterogeneity). In other words, change in tsRNA (and perhaps rsRNA/ysRNA as well) composition may result in altered ribosome heterogeneity that directs the cell to a specific functional state [9]. The complexity and possible permutations of different tsRNA/rsRNA/ysRNAs may endow the superior information capacity and specificity that are needed to distinguish complex diseases, and as being harnessed here, represent a “disease RNA code” in lung cancer screening.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12943-020-01280-9>.

Additional file 1: Supplementary Figures. **Fig. S1.** The workflow of the study. **Fig. S2.** The landscape of non-canonical sncRNAs in human PBMCs. **Fig. S3.** The mapping profile of tsRNAs. **Fig. S4.** Comparison of the expression of the prioritized sncRNA subcategories between lung cancer stages. **Fig. S5.** Comparison of the expression of the prioritized sncRNA subcategories between lung cancer histological types. **Fig. S6.** Comparison of the expression of the prioritized sncRNA subcategories between the lung cancer patients with and without lymph node involvement. **Fig. S7.** Comparison of the expression of the prioritized sncRNA subcategories between the lung cancer patients with and without distant metastasis. **Fig. S8.** Comparison of the expression of the prioritized sncRNA subcategories between the lung cancer patients with and without smoking history. **Fig. S9.** The TRY-RNA signature. **Fig. S10.** The TRY-RNA index in the discovery cohort. **Fig. S11.** Comparison of the expression of the sncRNA species within the TRY-RNA signature between lung cancer stages. **Fig. S12.** Comparison of the expression of the sncRNA species within the TRY-RNA signature between lung cancer histological types. **Fig. S13.** Comparison of the expression of the sncRNA species within the TRY-RNA signature between the lung cancer patients with and without lymph node involvement. **Fig. S14.** Comparison of the expression of the sncRNA species within the TRY-RNA signature between the lung cancer patients with and without distant metastasis. **Fig. S15.** Comparison of the expression of the sncRNA species within the TRY-RNA signature between the lung cancer patients with and without smoking history. **Fig. S16.** Expression heatmap of the MIR signature in the discovery cohort. **Fig. S17.** The MIR index in the discovery cohort. **Fig. S18.** Comparison between the TRY-RNA and MIR signatures.

Additional file 2: Supplementary Tables. **Table S1.** The human subjects of the discovery cohort. **Table S2.** The TRY-RNA signature. **Table S3.** The human subjects of the validation cohort. **Table S4.** The MIR signature.

Additional file 3. Methods.

Abbreviations

LDCT: Low-dose computed topography; TB: Tuberculosis; miRNA: MicroRNA; PBMC: Peripheral blood mononuclear cell; EV: Extracellular vesicle; sncRNA: Small non-coding RNA; sncRNA-seq: Small non-coding RNA sequencing; tsRNA: tRNA-derived small RNA; rsRNA: rRNA-derived small RNAs; ysRNA: YRNA-derived small RNAs; nt: Nucleotides; RPM: Reads per million; ROC: Receiver operating characteristic; AUC: Area under the receiver operating characteristic curve

Acknowledgements

Not applicable.

Authors' contributions

WG, QC, ZQ, and TZ conceived this study. WL, XW, ZW, and ZQ recruited the human subjects. HL, RL, and JL processed the RNA samples. JJZ, and TZ designed the statistical analyses. WG, JS, XZ, ML, DZ, GW, SZ, TQ, and TZ processed the sequencing data. WG, HZ, CZ, KSK, TW, QC, ZQ, and TZ

interpreted the results. WG, JS, JJZ, HZ, CZ, KSK, TW, QC, ZQ, and TZ wrote the manuscript. The author (s) read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The datasets generated and analyzed during the current study are available in the Gene Expression Omnibus (<https://www.ncbi.nlm.nih.gov/geo/>) under the accession number GSE148861 and GSE148862 for the discovery and validation cohorts, respectively.

Ethics approval and consent to participate

The Ethics Committee of Bengbu Medical College approved this study, with written informed consent obtained from all subjects, which conformed to the standard indicated by the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

We have filed provisional patent application for the TRY-RNA and MIR signatures.

Author details

¹State Key Laboratory of Bioelectronics, School of Biological Sciences and Medical Engineering, Southeast University, 2 Sipailou, Nanjing 210096, Jiangsu, China. ²Division of Biomedical Sciences, School of Medicine, University of California, Riverside, 900 University Avenue, Riverside, CA 92521, USA. ³Anhui Province Key Laboratory of Immunology in Chronic Diseases, Anhui Key Laboratory of Infection and Immunity, Department of Laboratory Medicine, Bengbu Medical College, 2600 Donghaidadao, Bengbu 233003, Anhui, China. ⁴Department of Epidemiology and Biostatistics, University of Arizona, Tucson, AZ 85721, USA. ⁵Department of Physiology and Cell Biology, University of Nevada, Reno School of Medicine, 1664 North Virginia Street, Reno, Nevada 89557, USA. ⁶Anhui Clinical and Preclinical Key Laboratory of Respiratory Disease, Department of Respiration, First Affiliated Hospital, Bengbu Medical College, Bengbu 233000, Anhui, China. ⁷The Infectious Disease Hospital of Bengbu City, Bengbu 233000, Anhui, China. ⁸Department of Surgery, The Ohio State University, Columbus, OH 43210, USA. ⁹Department of Internal Medicine, College of Medicine Phoenix, University of Arizona, Phoenix, AZ 85004, USA.

Received: 20 August 2020 Accepted: 3 November 2020

Published online: 12 November 2020

References

- Paras L, Baskaran S. Deep learning at chest radiography: automated classification of pulmonary tuberculosis by using convolutional neural networks. *Radiology*. 2017;284:574–82.
- Shi J, Ko EA, Sanders KM, Chen Q, Zhou T. SPORTS1.0: a tool for annotating and profiling non-coding RNAs optimized for rRNA- and tRNA-derived small RNAs. *Genomics Proteomics Bioinformatics*. 2018;16:144–51.
- Su Z, Wilson B, Kumar P, Dutta A. Noncanonical roles of tRNAs: tRNA fragments and beyond. *Annu Rev Genet*. 2020. Online ahead of print.
- Balatti V, Nigita G, Veneziano D, Drusco A, Stein GS, Messier TL, Farina NH, Lian JB, Tomasello L, Liu CG, et al. tsRNA signatures in cancer. *Proc Natl Acad Sci U S A*. 2017;114:8071–6.
- Farina NH, Scalia S, Adams CE, Hong D, Fritz AJ, Messier TL, Balatti V, Veneziano D, Lian JB, Croce CM, et al. Identification of tRNA-derived small RNA (tsRNA) responsive to the tumor suppressor, RUNX1, in breast cancer. *J Cell Physiol*. 2020;235:5318–27.
- Dhabhi JM, Spindler SR, Atamna H, Boffelli D, Martin DI. Deep sequencing of serum small RNAs identifies patterns of 5' tRNA half and YRNA fragment expression associated with breast Cancer. *Biomark Cancer*. 2014;6:37–47.
- Zhang Y, Zhang X, Shi J, Tuorto F, Li X, Liu Y, Liebers R, Zhang L, Qu Y, Qian J, et al. Dnm2 mediates intergenerational transmission of paternally acquired metabolic disorders through sperm small non-coding RNAs. *Nat Cell Biol*. 2018;20:535–40.
- Pekarsky Y, Balatti V, Palamarchuk A, Rizzotto L, Veneziano D, Nigita G, Rassenti LZ, Pass HI, Kipps TJ, Liu CG, Croce CM. Dysregulation of a family of

short noncoding RNAs, tsRNAs, in human cancer. *Proc Natl Acad Sci U S A*. 2016;113:5071–6.

9. Shi J, Zhang Y, Zhou T, Chen Q. tsRNAs: the Swiss Army knife for translational regulation. *Trends Biochem Sci*. 2019;44:185–9.
10. Zhang Y, Shi J, Rassoulzadegan M, Tuorto F, Chen Q. Sperm RNA code programmes the metabolic health of offspring. *Nat Rev Endocrinol*. 2019;15:489–98.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

