

UCSF

UC San Francisco Previously Published Works

Title

The Strongest Brønsted Acid: Protonation of Alkanes by H(CHB11F11) at Room Temperature

Permalink

<https://escholarship.org/uc/item/0gc6w7mx>

Journal

Angewandte Chemie International Edition, 53(4)

ISSN

1433-7851

Authors

Nava, Matthew

Stoyanova, Irina V

Cummings, Steven

et al.

Publication Date

2014-01-20

DOI

10.1002/anie.201308586

Peer reviewed



# HHS Public Access

Author manuscript

*Angew Chem Int Ed Engl.* Author manuscript; available in PMC 2016 August 22.

Published in final edited form as:

*Angew Chem Int Ed Engl.* 2014 January 20; 53(4): 1131–1134. doi:10.1002/anie.201308586.

## The Strongest Brønsted Acid: Protonation of Alkanes by H(CHB<sub>11</sub>F<sub>11</sub>) at Room Temperature\*\*

**Matthew Nava<sup>+</sup>,**

Center for s and p Block Chemistry, Department of Chemistry University of California, Riverside, CA 92521 (USA)

**Irina V. Stoyanova,**

Center for s and p Block Chemistry, Department of Chemistry University of California, Riverside, CA 92521 (USA)

**Dr. Steven Cummings,**

Center for s and p Block Chemistry, Department of Chemistry University of California, Riverside, CA 92521 (USA)

**Prof. Dr. Evgenii S. Stoyanov,** and

Center for s and p Block Chemistry, Department of Chemistry University of California, Riverside, CA 92521 (USA)

**Christopher A. Reed**

Center for s and p Block Chemistry, Department of Chemistry University of California, Riverside, CA 92521 (USA), Homepage: <http://reedgrouplab.ucr.edu/>

Christopher A. Reed: [chris.reed@ucr.edu](mailto:chris.reed@ucr.edu)

### Abstract

What is the strongest acid? Can a simple Brønsted acid be prepared that can protonate an alkane at room temperature? Can that acid be free of the complicating effects of added Lewis acids that are typical of common, difficult-to-handle superacid mixtures? The carborane superacid H-(CHB<sub>11</sub>F<sub>11</sub>) is that acid. It is an extremely moisture-sensitive solid, prepared by treatment of anhydrous HCl with [Et<sub>3</sub>Si-H-SiEt<sub>3</sub>][CHB<sub>11</sub>F<sub>11</sub>]. It adds H<sub>2</sub>O to form [H<sub>3</sub>O][CHB<sub>11</sub>F<sub>11</sub>] and benzene to form the benzenium ion salt [C<sub>6</sub>H<sub>7</sub>][CHB<sub>11</sub>F<sub>11</sub>]. It reacts with butane to form a crystalline tBu<sup>+</sup> salt and with n-hexane to form an isolable hexyl carbocation salt. Carbocations, which are thus no longer transient intermediates, react with NaH either by hydride addition to reform an alkane or by deprotonation to form an alkene and H<sub>2</sub>. By protonating alkanes at room temperature, the reactivity of H(CHB<sub>11</sub>F<sub>11</sub>) opens up new opportunities for the easier study of acid-catalyzed hydrocarbon reforming.

\*\*We thank Prof. Dr. Pingyun Feng and Dr. Richard W. Kondrat for kind assistance with H<sub>2</sub> detection and GC/MS, respectively. This work was supported by the National Science Foundation (CHE 1144838 and CHE 0742001).

Correspondence to: Christopher A. Reed, [chris.reed@ucr.edu](mailto:chris.reed@ucr.edu).

<sup>+</sup>Current address: Department of Chemistry, Massachusetts Institute of Technology 77 Massachusetts Ave., Cambridge, MA 02139-4307 (USA)

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201308586>.

## Keywords

Brønsted acids; carbocations; carboranes; hydrocarbons

For the past decade, the strongest acid to be isolated and fully characterized has been the chlorinated carborane acid,  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$ .<sup>[1, 2]</sup> More recently, this has been equaled by the comparably strong, isoelectronic, all-boron diprotic superacid,  $\text{H}_2(\text{B}_{12}\text{Cl}_{12})$ .<sup>[3]</sup> The superior strength of these Brønsted acids has been established in all phases. In solution, the fact that both readily protonate benzene (whereas oxyacids do not) places them ahead of  $\text{HFSO}_3$ , the previous strongest pure acid measured on the  $H_o$  scale.<sup>[4]</sup> In the gas phase, the calculated<sup>[5]</sup> and measured<sup>[6]</sup> deprotonation enthalpy of  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$  is the lowest of any available acid. The  $\nu\text{NH}$  anion basicity scale<sup>[7]</sup> indicates that carborane anions are less basic than the oxyanions of traditional acids. This same scale indicates that fluorinated carborane anions are less basic than chlorinated carborane anions, so the conjugate acid  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  should be a stronger acid than  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$ . Calculated gas-phase deprotonation energies concur.<sup>[5]</sup> There is, of course, no guarantee that weaker anion basicity leads to an isolable stronger conjugate acid. The anion must also be chemically stable to  $\text{H}^+$ .<sup>[8]</sup> However, the stability of fluorinated carborane anions<sup>[9]</sup> towards the potent Lewis acidity of trialkylsilylium ions<sup>[10]</sup> augurs well for their stability towards  $\text{H}^+$ .

Indeed, there was a preliminary report in 2007 of two fluorinated carborane acids,  $\text{H}(\text{HCB}_{11}\text{F}_{11})$  and  $\text{H}(\text{EtCB}_{11}\text{F}_{11})$ .<sup>[10]</sup> As evidence of formulation, an IR spectrum of the latter was given. By analogy to  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$ , which is known from X-ray analysis to have a linear polymeric structure with  $\text{H}^+$  bridges between anion Cl atoms, and from IR spectroscopy to have low-barrier H-bonding,<sup>[11]</sup> the expected IR features for  $\text{H}(\text{EtCB}_{11}\text{F}_{11})$  were present. These include two bands at about  $1705$  and  $1620\text{ cm}^{-1}$  as candidates for  $\nu\text{FHF}$  and a band at about  $920\text{ cm}^{-1}$  for  $\delta\text{FHF}$ , along with expected  $\nu\text{BB}$  and  $\nu\text{BF}$  bands from the anion at circa  $1300$  and  $700\text{ cm}^{-1}$ , respectively.

Given that there has been no follow-up to this preliminary report in the intervening six years, we were curious whether we could reproduce this work and prepare a fluorinated carborane acid in synthetic amounts for reactivity studies with hydrocarbons. We chose the non-alkylated anion  $\text{CHB}_{11}\text{F}_{11}^-$  for our studies, based on concern for the chemical stability of the ethyl group in  $\text{EtCB}_{11}\text{F}_{11}^-$ . From preliminary investigations into the reactivity of the chlorinated carborane acid  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$  with alkanes at somewhat elevated temperatures, we had reason to believe that fluorinated carborane acids might react with alkanes at room temperature, thus giving  $\text{H}(\text{EtCB}_{11}\text{F}_{11})$  the capability of reacting with its own ethyl group. An indication that this may be occurring is suggested by the IR spectrum reported for  $\text{H}(\text{EtCB}_{11}\text{F}_{11})$ <sup>[10]</sup> where two of the three  $\nu\text{CH}$  bands of the ethyl group in the  $2900\text{--}3020\text{ cm}^{-1}$  region are nearly absent in comparison to other  $\text{EtCB}_{11}\text{F}_{11}^-$  salts (see Figure S8 in the Supporting Information of Ref. [10]).

The synthesis of  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  (Scheme 1) was eventually achieved using much the same procedure as that for its chlorinated analogue  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$ ,<sup>[12]</sup> although the preparation of significant amounts of clean product eluded us for a long time. We now understand this difficulty in terms of  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  being a considerably stronger acid than  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$ ,

and thus capable of reacting rapidly with all organic species, and being the ultimate desiccant. So, purity of starting materials and the scrupulous exclusion of water were imperative. In particular, we found that the trityl salt of  $\text{CHB}_{11}\text{F}_{11}^-$  must be carefully purified by successive recrystallizations from *o*-dichlorobenzene to remove occluded organic compounds. Another critical step was to perform the reaction of  $[\text{Et}_3\text{Si}-\text{H}-\text{SiEt}_3][\text{CHB}_{11}\text{F}_{11}]$  with HCl twice (sequentially) to completely remove silane byproducts. Once isolated, small quantities of  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  were obtained in high purity by sublimation under high vacuum at about 160 °C, a temperature similar to that used for the sublimation of  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$ . Otherwise, synthetic amounts were typically made in 100–200 mg batches and, under the best conditions, were estimated, from the virtual absence of  $\nu\text{OH}$  in the IR, to contain less than 2 % hydrated impurity. In our experience, samples can be stored for no more than a few days, even in a good dry box, before becoming significantly degraded by hydration and/or reaction with trace solvent vapors. Once hydrated, the acid cannot be dehydrated by vacuum sublimation.


$\text{H}(\text{CHB}_{11}\text{F}_{11})$  is expected to be isostructural with  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$  and have a linear polymeric structure with two-coordinate, low-barrier  $\text{H}^+$  bridging between F atoms. As shown in Figure 1a and b, the IR spectrum is somewhat related to that of the symmetrical bifluoride ion,  $\text{FHF}^-$ ,<sup>[13]</sup> and is diagnostic of low-barrier H-bonding.<sup>[14]</sup> Instead of  $\nu\text{HF}$  at about  $2500\text{ cm}^{-1}$  as might be expected for a terminal H–F bond,  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  shows a broad band at about  $1605\text{ cm}^{-1}$  assigned to  $\nu\text{FHF}$  and absorptions in the  $1000\text{--}900\text{ cm}^{-1}$  region assigned to  $\delta\text{FHF}$  in a nearly symmetrical proton-bridged structure. The corresponding bands in  $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$  are at about  $1100$  and  $615\text{ cm}^{-1}$ , respectively.<sup>[11]</sup> Deuteration, achieved by stirring the protio acid in liquid DCl, led to the loss of both the  $1605$  and  $1000\text{--}900\text{ cm}^{-1}$  bands. The corresponding  $\nu\text{DFD}$  and  $\delta\text{DFD}$  bands, expected at about  $1140$  and  $710\text{--}640\text{ cm}^{-1}$  by harmonic oscillator calculations, are both largely masked by the very strong  $\nu\text{BB}$  and  $\nu\text{BF}$  bands of the anion. Otherwise, the IR spectrum of  $\text{D}(\text{CHB}_{11}\text{F}_{11})$  shows only bands expected from the anion:  $\nu\text{CH}$  at  $3030\text{ cm}^{-1}$  and multiple absorptions centered near  $1300$  and  $700\text{ cm}^{-1}$  from  $\nu\text{BB}$  and  $\nu\text{BF}$ , respectively.

Upon exposure to traces of water vapor,  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  is instantly hydrated. With minimal hydration, the IR spectrum (Figure 1c; Supporting Information, Figure S11) is consistent with the formation of an  $\text{H}_3\text{O}^+$  salt. Broad  $\nu\text{OH}$  bands at about  $3300$  and  $3175\text{ cm}^{-1}$  are assigned to  $\nu_s$  and  $\nu_{as}$ , respectively. Their frequencies are higher than those in the chlorinated analogue  $[\text{H}_3\text{O}][\text{CHB}_{11}\text{Cl}_{11}]$  at  $3225$ ,  $2910\text{ cm}^{-1}$ ,<sup>[15]</sup> indicating weaker H-bonding of the  $\text{H}_3\text{O}^+$  ion to the less-basic fluorinated carborane anion. The band at about  $1625\text{ cm}^{-1}$  is assigned to  $\delta(\text{H}_3\text{O})$ . The other major bands in the spectrum are  $\nu\text{BB}$  and  $\nu\text{BF}$  at circa  $1300$  and  $700\text{ cm}^{-1}$ , respectively.

As expected for such a strong acid, solid  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  reacts instantly on contact with benzene to give an isolable benzenium ion salt,  $[\text{C}_6\text{H}_7^+][\text{CHB}_{11}\text{F}_{11}^-]$  (for the IR spectrum, see the Supporting Information, Figure S12). The circa  $10\text{ cm}^{-1}$  increase in frequency observed for the  $\nu(\text{CC}) + \delta(\text{CCH})$  band near  $1600\text{ cm}^{-1}$  relative to free benzene is diagnostic of the benzenium ion.<sup>[16]</sup> The  $\nu\text{CH}$  bands of the acidic  $\text{sp}^3\text{CH}_2$  group occur at low frequency ( $2818$  and  $2794\text{ cm}^{-1}$ ) owing to  $\text{C}-\text{H}\cdots\text{anion}$  H-bonding.<sup>[17]</sup> Their frequencies are somewhat higher than that in  $[\text{C}_6\text{H}_7][\text{CHB}_{11}\text{Cl}_{11}]$  ( $2776\text{ cm}^{-1}$ ), consistent

with weaker H-bonding to the less-basic fluorinated carborane anion relative to the chlorinated analogue. The structure and remarkable stability of these so-called Wheland intermediates of electrophilic aromatic substitution have been discussed previously.<sup>[16]</sup>

In a clear demonstration of the extremely high acid strength of this Brønsted-only acid, we find that H(CHB<sub>11</sub>F<sub>11</sub>) reacts with alkanes at room temperature, like mixed Brønsted/Lewis acids such as “magic acid” (HFSO<sub>3</sub>/SbF<sub>5</sub>).<sup>[18]</sup> When solid H(CHB<sub>11</sub>F<sub>11</sub>) is stirred in suspension with *n*-hexane, the appearance of a diagnostic low frequency ν<sub>CH</sub> band at 2758 cm<sup>-1</sup> in the IR spectrum of the product (the low frequency arising from hyperconjugation and C–H···anion hydrogen bonding)<sup>[17, 19]</sup> indicates that about 50% of the solid acid is converted into a microcrystalline carbocation salt within 2 h (Supporting Information, Figure S14). This is the expected outcome of protonation of an alkane and elimination of H<sub>2</sub> [Eq. (1)]:



Evolved H<sub>2</sub> was detected by gas chromatography (Supporting Information, Figure S13). A similar protonation experiment with a suspension of H(CHB<sub>11</sub>F<sub>11</sub>) in liquefied *n*-butane indicates that the acid converts butane into a micro-crystalline salt of *tert*-butyl cation [Eq. (2)]:



(2)

The distinctive low frequency and shape of the ν<sub>CH</sub> absorptions in the IR spectrum of the product (Supporting Information, Figures S15–S17) closely matches that of known [*t*Bu<sup>+</sup>][CHB<sub>11</sub>Cl<sub>11</sub><sup>-</sup>],<sup>[17]</sup> except that the ν<sub>max</sub> frequency at 2823 cm<sup>-1</sup> of the fluorinated anion salt is higher than that of the chlorinated anion salt (2788 cm<sup>-1</sup>). This is readily understood in terms of weaker C–H···anion H-bonding of *t*Bu<sup>+</sup> to the fluorinated anion. The tertiary isomer of butyl cation is the expected result of facile 1,2 shifts from initially formed primary or secondary carbocation-like species.<sup>[20]</sup> The X-ray structures of two *tert*-butyl salts with carborane anions have been reported previously.<sup>[17, 21]</sup>

Additional characterization of these C<sub>4</sub> and C<sub>6</sub> carbocation salts was obtained from reactivity studies with NaH, anticipating two possible outcomes: 1) hydride transfer to reform an alkane; or b) hydride acting as a base, deprotonating the carbocation and forming an alkene.

When a sample of the presumed C<sub>6</sub> carbocation salt, obtained as above from the protonation of *n*-hexane with H(CHB<sub>11</sub>F<sub>11</sub>), was treated with a NaH in liquid SO<sub>2</sub>, a single hydrocarbon product was detected by gas chromatographic analysis (Supporting Information, Figure S18). Its identity as a C<sub>6</sub> alkene was shown by mass spectroscopy (*m/z* 84.086; see the

Supporting Information, Figure S19). Electron impact fragments corresponding to loss of methyl, ethyl, and propyl groups were observed and comparison of this fragmentation pattern to those catalogued for C<sub>6</sub> alkenes shows the closest match to that of 3,3-dimethyl-1-butene, although it shares substantial similarity to those of 2-methyl-2-pentene, 2,3-dimethyl-1-butene, 2,3-dimethyl-2-butene, and 4-methyl-2-pentene (Supporting Information, Figure S20). Thus, a C<sub>6</sub> carbocation was the product of protonation of *n*-hexane, most likely with a tertiary cationic center arising from the well-known rearrangement of initially formed primary or secondary carbocation-like species to a more stable tertiary cation by rapid 1,2 shifts.<sup>[20]</sup>

Under somewhat different experimental conditions, NaH acts as a hydride transfer agent rather than a base. Intimate co-grinding of the presumed C<sub>6</sub> carbocation salt with NaH<sub>(s)</sub> led to evolution of four different C<sub>6</sub> alkanes by GC analysis: 2,2-dimethylbutane, 2,3-dimethylbutane, 2-methylpentane, and 3-methylpentane (Supporting Information, Figure S21). This mixture presumably reflects the rapid rearrangement equilibria possible in all tertiary alkyl cations.<sup>[20]</sup> Similarly, intimate co-grinding with NaH<sub>(s)</sub> of the *t*Bu<sup>+</sup> salt derived from protonation of *n*-butane led to *iso*-butane as the major alkane detectable by GC/MS in the head space (Supporting Information, Figure S22). *tert*-Butyl cations produced in a different manner,<sup>[22]</sup> by protonation of butyl chloride with H(CHB<sub>11</sub>Cl<sub>11</sub>), behaved in identical fashion.

In contrast to these room-temperature reactions of H-(CHB<sub>11</sub>F<sub>11</sub>) with alkanes, the chlorinated carborane acid H(CHB<sub>11</sub>Cl<sub>11</sub>) scarcely reacts with *n*-hexane or butane under comparable conditions. Even after 24 h stirring as a suspension in *n*-hexane, less than 25 % conversion of H(CHB<sub>11</sub>Cl<sub>11</sub>) into carbocation salts was detected by IR spectroscopy. Thus, the fluorinated carborane acid is now the strongest pure acid known.

The existence of a Brønsted-only acid that can protonate alkanes at room temperature offers new opportunities for studying acid-catalyzed hydrocarbon chemistry related to hydrocarbon reforming on zeolites at high temperatures. Potential advantages are that a sublimable solid acid is easier to handle than a liquid acid and a Brønsted-only acid is free of the potentially complicating effects of a Lewis acid. Room-temperature access to alkane protonation chemistry should also allow the easier application of investigative techniques that are difficult to apply to zeolites;<sup>[23]</sup> the acidity of zeolites is much lower<sup>[24]</sup> and their reactions with alkanes must be carried out at high temperatures.<sup>[25]</sup> Such room-temperature techniques have already been applied with unexpected results to chloroalkanes, which are more easily protonated than alkanes. Peculiar stoichiometric relationships in reversible, carbocation-like oligomerization sequences were observed.<sup>[22]</sup>

## Experimental Section

All manipulations were carried out under very dry conditions using Schlenkware or an inert atmosphere glovebox (H<sub>2</sub>O, O<sub>2</sub> < 0.5 ppm). Solvents were dried by standard methods. HF was dried by condensing gaseous HF into a PFA vessel containing potassium hexafluoro-nickelate. HCl was dried by passing it through a 25 cm column of P<sub>2</sub>O<sub>5</sub> plugged at both ends with glass wool. 10% F<sub>2</sub> in N<sub>2</sub> was used without additional purification. <sup>1</sup>H, <sup>11</sup>B,

and  $^{19}\text{F}$  spectra were obtained on a Bruker Avance 300 MHz spectrometer. Attenuated Reflectance spectra (ATR) IR spectra were run on ABB MB3000 spectrometer in the 525–4000  $\text{cm}^{-1}$  frequency range using a diamond crystal. Mass spectra were collected using a Waters GCT GC/MS operating at 15 °C.

$\text{Cs}(\text{CHB}_{11}\text{F}_{11})$ : **Caution!** HF and  $\text{F}_2$  are extremely toxic and are used under pressure. All reactions should be carried out with appropriate apparatus and safety precautions in a well-ventilated hood using full-body protective clothing and the “buddy” system. Adapting the method of Solntsev and Strauss,<sup>[9]</sup> a 300 mL Teflon-lined Monel bomb equipped with a Teflon stir bar was charged with 2 g of  $\text{Cs}(\text{CHB}_{11}\text{H}_{11})$ <sup>[26]</sup> and placed in an oven at 80 °C for 4 h. It was then connected to a PFA vacuum manifold, evacuated, and cooled to dry ice temperature in a dry ice/methanol bath. Approximately 35 mL of high purity hydrogen fluoride was condensed into the reactor and the reaction was stirred at room temperature for 24 h. After cooling in a dry ice/methanol bath, the reactor was evacuated and then placed on a stainless steel vacuum manifold. 280 PSI (20 bar) of 10 %  $\text{F}_2$  in  $\text{N}_2$  was introduced into the bomb, sealed, and allowed to warm to room temperature. After stirring for 8 h, the bomb was cooled in a dry ice/methanol bath and the excess pressure vented. The process of adding fluorine, allowing it to react, and venting the reactor was repeated five times. Then, hydrogen fluoride was condensed out of the bomb and the reactor was carefully opened and allowed to stand in the air. After 1 hour, the contents were dissolved in 80 mL of dry acetonitrile and filtered through fine frit, to remove insoluble  $\text{CsBF}_4$ . The filtrate was evaporated to dryness on a rotary evaporator, yielding a sticky yellow powder. This powder was placed in an inert atmosphere glove box, suspended in dry dichloromethane (100 mL), and stirred overnight. The suspension was filtered off to yield a white powder, which was dried under vacuum for 1 hour (2.74 g; 80%).  $^1\text{H}$  NMR ( $[\text{D}_6]$ acetone):  $\delta = 4.35$  ppm (s, 1H, CH).  $^{11}\text{B}$  NMR ( $[\text{D}_6]$ acetone, unreferenced):  $\delta = -7.75$  (s, 1B), 15.86 (s, 5B), 17.09 ppm (s, 5B) (SI, Figure S3).  $^{19}\text{F}$  NMR ( $[\text{D}_6]$ acetone, unreferenced):  $\delta = -251.59$  (m, 1F),  $-255.63$  ppm (m, 10F) (SI, Figure S2). The completeness of fluorination was initially judged by the absence of  $\nu\text{BH}$  bands near 2550  $\text{cm}^{-1}$  in the IR spectrum of the product (SI, Figure S1).  $^{19}\text{F}$  and  $^{11}\text{B}$  NMR spectra were found to be less reliable indicators of completeness of fluorination. The most sensitive indicator of anion purity was negative ion electrospray mass spectrometry (SI, Figure S4).

$\text{Ag}(\text{CHB}_{11}\text{F}_{11}) \cdot 2 \text{C}_6\text{H}_6$ :  $\text{Cs}(\text{CHB}_{11}\text{F}_{11})$  (2.78 g, 5.87 mmol) and  $\text{AgNO}_3$  (4.98 g, 29.93 mmol) were suspended in dry benzene (300 mL) and stirred magnetically. After 24 h, the benzene was decanted and a fresh 300 mL aliquot added to the remaining solids. This process was repeated three times. The combined ( $3 \times 300$  mL) benzene fractions were filtered through a fine frit and evaporated to dryness to yield an off-white solid that was dried under vacuum for 1 hour (2.18 g, 83 %).  $^1\text{H}$  NMR: (300 MHz,  $\text{CD}_2\text{Cl}_2$ ):  $\delta = 3.946$  (s, 1H, CH), 7.36 ppm (s, 6H, benzene) (SI, Figure S6). ATR-IR (SI, Figure S5).

$[(\text{C}_6\text{H}_5)_3\text{C}][\text{CHB}_{11}\text{F}_{11}]$ : In the glove box,  $\text{Ag}(\text{CHB}_{11}\text{F}_{11}) \cdot 2 \text{C}_6\text{H}_6$  (2.48 g, 4.14 mmol) and  $(\text{C}_6\text{H}_5)_3\text{CBr}$  (1.48 g, 4.55 mmol) were dissolved in 25 mL of dry dichloromethane and stirred overnight. The yellow precipitate of  $\text{AgBr}$  was filtered off with a fine frit, washing with 5 mL of dichloromethane. The filtrate was evaporated to dryness under vacuum, yielding a bright yellow waxy solid. This was recrystallized twice by layering 10 mL 1,2-

dichlorobenzene solutions with 15 mL of hexane. The solid was filtered off and dried under vacuum for 1 h (2.2 g, 89 %).  $^1\text{H NMR}$  (300 MHz,  $\text{CD}_2\text{Cl}_2$ ):  $\delta = 2.776$  (s, 1H, C-H), 7.705, 7.938, 8.325 ppm (phenyl groups) (SI, Figure S8). ATR-IR (SI, Figure S7).

$[(\text{Et}_3\text{Si})_2\text{H}][\text{CHB}_{11}\text{F}_{11}]$ :  $[(\text{C}_6\text{H}_5)_3\text{C}][\text{CHB}_{11}\text{F}_{11}]$  (200 mg, 0.342 mmol) was covered with dry 1,2-dichlorobenzene (0.5 mL). To this suspension 1 mL of  $\text{Et}_3\text{SiH}$  was added and the solution stirred for 0.5 h. Dry hexane (5 mL) was added gradually while stirring. The resulting precipitate was filtered and washed with dry hexanes (5 mL) (0.175 g, 89%). ATR-IR (SI, Figure S9). The broad band at  $1873\text{ cm}^{-1}$  is diagnostic of the hydride-bridged disilyl cation.<sup>[27]</sup>

$\text{H}(\text{CHB}_{11}\text{F}_{11})$ : Extra-dry HCl gas was condensed (ca. 2 mL) onto  $[(\text{Et}_3\text{Si})_2\text{H}][\text{CHB}_{11}\text{F}_{11}]$  (200 mg) using liquid  $\text{N}_2$ , and the mixture was stirred at  $0^\circ\text{C}$  for 0.5 hour in a sealed heavy-walled Schlenk tube having a wide bore Teflon stopcock below the joint. The HCl was removed under vacuum and a new portion of HCl (2 mL) was added, stirring for 0.5 hour at  $0^\circ\text{C}$ . The HCl allowed to escape anaerobically and the remnant solid was pumped under vacuum for 0.5 h (96 mg, 81%). The deuterated acid,  $\text{D}(\text{CHB}_{11}\text{F}_{11})$ , was prepared by stirring the protio acid in liquid DCl followed by evaporation of DCl/HCl and drying under vacuum. The IR spectrum is compared to the protio form in the Supporting Information, Figure S10.

## Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

## References


1. Juhasz M, Hoffmann SP, Stoyanov ES, Kim KC, Reed CA. *Angew Chem.* 2004; 116:5466–5469. *Angew Chem Int Ed.* 2004; 43:5352–5355.
2. Reed CA. *Chem N Z.* 2011 Oct.;174–179.
3. Avelar A, Tham FS, Reed CA. *Angew Chem.* 2009; 121:3543–3545. *Angew Chem Int Ed.* 2009; 48:3491–3493.
4. Gillespie RJ. *Acc Chem Res.* 1968; 1:202–209.
5. Lipping L, Leito I, Koppel I, Koppel IA. *J Phys Chem A.* 2009; 113:12972–12978. [PubMed: 19807147]
6. Meyer MM, Wang XB, Reed CA, Wang LS, Kass SR. *J Am Chem Soc.* 2009; 131:18050–18051. [PubMed: 19950932]
7. Stoyanov ES, Kim KC, Reed CA. *J Am Chem Soc.* 2006; 128:8500–8508. [PubMed: 16802816]
8. Reed CA. *Chem Commun.* 2005:1669–1677.
9. Ivanov SV, Rockwell JJ, Polyakov OG, Gaudinski CM, Anderson OP, Solntsev KA, Strauss SH. *J Am Chem Soc.* 1998; 120:4224–4225.
10. Küppers TK, Bernhardt E, Eujen R, Willner H, Lehmann CW. *Angew Chem.* 2007; 119:6462–6465. *Angew Chem Int Ed.* 2007; 46:6346–6349.
11. Stoyanov ES, Hoffmann SP, Juhasz M, Reed CA. *J Am Chem Soc.* 2006; 128:3160–3161. [PubMed: 16522093]
12. See the Supporting Information in: Reed CA. *Acc Chem Res.* 2010; 43:121–128. [PubMed: 19736934]
13. Kawaguchi K, Hirota E. *J Chem Phys.* 1987; 87:6838–6841.



14. Stasko D, Hoffmann SP, Kim KC, Fackler NLP, Larsen AS, Drovetskaya T, Tham FS, Reed CA, Rickard CEF, Boyd PDW, Stoyanov ES. *J Am Chem Soc.* 2002; 124:13869–13876. [PubMed: 12431118]
15. Stoyanov ES, Kim KC, Reed CA. *J Am Chem Soc.* 2006; 128:1948–1958. [PubMed: 16464096]
16. Reed CA, Kim KC, Stoyanov ES, Stasko D, Tham FS, Mueller LJ, Boyd PDW. *J Am Chem Soc.* 2003; 125:1796–1804. [PubMed: 12580605]
17. Stoyanov ES, Stoyanova IV, Tham FS, Reed CA. *Angew Chem.* 2012; 124:9283–9285. *Angew Chem Int Ed.* 2012; 51:9149–9151.
18. Olah, GA.; Molinár, Á. *Hydrocarbon Chemistry.* Wiley; New York: 2003.
19. Reed CA, Stoyanov ES, Tham FS. *Org Biomol Chem.* 2013; 11:3797–3802. [PubMed: 23632995]
20. Saunders M, Kates MR. *J Am Chem Soc.* 1978; 100:7082–7083.
21. Kato T, Reed CA. *Angew Chem.* 2004; 116:2968–2971. *Angew Chem Int Ed.* 2004; 43:2908–2911.
22. Stoyanov ES, Stoyanova IV, Reed CA. *J Am Chem Soc.* 2011; 133:8452–8454. [PubMed: 21568318]
23. Sani Souna Sido A, Barbiche J, Sommer J. *Chem Commun.* 2010; 46:2913–2914.
24. Haw JF. *Phys Chem Chem Phys.* 2002; 4:5431–5441.
25. Haw JF, Nicholas JB, Xu T, Beck LW, Ferguson DB. *Acc Chem Res.* 1996; 29:259–267.
26. Plešek J, Jelínek T, Drdáková E, He mánek S, Štíbr B. *Collect Czech Chem Commun.* 1984; 49:1559–1562.
27. Hoffmann SP, Kato T, Tham FS, Reed CA. *Chem Commun.* 2006:767–769.



**Figure 1.** IR spectra of a)  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  purified by sublimation, b)  $\text{H}(\text{CHB}_{11}\text{F}_{11})$  from bulk synthesis, and c)  $[\text{H}_3\text{O}][\text{CHB}_{11}\text{F}_{11}]$  from minimal uptake of water. Spectra (a) and (c) were recorded in transmission mode, (b) in ATR mode.

**Scheme 1.**

Synthesis of the carborane acid HY ( $\text{Y}=\text{CHB}_{11}\text{F}_{11}^-$ ).