

UCLA

UCLA Previously Published Works

Title

Membrane estrogen signaling in female reproduction and motivation

Permalink

<https://escholarship.org/uc/item/0h22s98g>

Authors

Johnson, Caroline S
Micevych, Paul E
Mermelstein, Paul G

Publication Date

2022

DOI

10.3389/fendo.2022.1009379

Peer reviewed



OPEN ACCESS

EDITED BY
Allan Herbison,
University of Cambridge,
United Kingdom

REVIEWED BY
Kyle Flippo,
The University of Iowa, United States

*CORRESPONDENCE
Caroline S. Johnson
joh20852@umn.edu

SPECIALTY SECTION
This article was submitted to
Reproduction,
a section of the journal
Frontiers in Endocrinology

RECEIVED 01 August 2022
ACCEPTED 13 September 2022
PUBLISHED 29 September 2022

CITATION
Johnson CS, Micevych PE and
Mermelstein PG (2022) Membrane
estrogen signaling in female
reproduction and motivation.
Front. Endocrinol. 13:1009379.
doi: 10.3389/fendo.2022.1009379

COPYRIGHT
© 2022 Johnson, Micevych and
Mermelstein. This is an open-access
article distributed under the terms of
the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Membrane estrogen signaling in female reproduction and motivation

Caroline S. Johnson^{1*}, Paul E Micevych²
and Paul G. Mermelstein¹

¹Department of Neuroscience, University of Minnesota, Minneapolis, MN, United States, ²Laboratory of Neuroendocrinology, Department of Neurobiology, David Geffen School of Medicine at University of California, Los Angeles, Los Angeles, CA, United States

Estrogen receptors were initially identified in the uterus, and later throughout the brain and body as intracellular, ligand-regulated transcription factors that affect genomic change upon ligand binding. However, rapid estrogen receptor signaling initiated outside of the nucleus was also known to occur *via* mechanisms that were less clear. Recent studies indicate that these traditional receptors, estrogen receptor- α and estrogen receptor- β , can also be trafficked to act at the surface membrane. Signaling cascades from these membrane-bound estrogen receptors (mERs) not only rapidly effect cellular excitability, but can and do ultimately affect gene expression, as seen through the phosphorylation of CREB. A principal mechanism of neuronal mER action is through glutamate-independent transactivation of metabotropic glutamate receptors (mGluRs), which elicits multiple signaling outcomes. The interaction of mERs with mGluRs has been shown to be important in many diverse functions in females, including, but not limited to, reproduction and motivation. Here we review membrane-initiated estrogen receptor signaling in females, with a focus on the interactions between these mERs and mGluRs.

KEYWORDS

estrogen, estrogen receptors, membrane estrogen receptors, metabotropic glutamate (mGlu) receptors, estrogen receptor signaling

Introduction

The estrogen receptors, estrogen receptor- α (ER α) and estrogen receptor- β (ER β) were initially identified as intracellular, ligand-regulated transcription factors (1), members of the larger nuclear receptor superfamily (2, 3). Originally identified in the uterus (4, 5), these estrogen receptors are expressed throughout the body, including in a multitude of brain regions (6, 7). Estradiol binding to these receptors was initially demonstrated to induce transcriptional changes at estrogen response elements (EREs) (8). However, this classical signaling pathway is not the only mechanism through which

estrogen receptors directly elicit genomic change. Many estrogen-regulated genes lack ERE sequences (9, 10), which led to the discovery of additional genomic actions occurring *via* other response elements and transcription factors (11, 12). However, even with multiple pathways leading to direct genomic effects, this was still insufficient to fully explain the plethora of actions estradiol was observed to induce both inside and outside the nervous system.

Membrane-initiated signaling

The first clues that estrogen signaling could be initiated outside the nucleus came from Szego & Davis in the late 1960s. Following ovariectomy (ovx) in rats, acute exogenous estradiol treatment resulted in an increase in uterine cAMP accumulation within seconds, concentrations indistinguishable from intact animals (13). The speed at which these changes occurred eliminated the possibility of nuclear-initiated action and strongly suggested the recruitment of a surface-initiated second messenger signaling pathway. Rapid effects of estradiol were subsequently noted within the nervous system, first in female preoptic-septal neurons in the hypothalamus. Within seconds of application, estradiol modulated firing rates, returning to experimental baseline when the steroid was removed (14). The use of estradiol conjugated to bovine serum albumen (BSA) further implicated membrane-associated estrogen receptors (15). However, skepticism remained, as there was suspicion that estradiol might be cleaved from BSA (16). Thus, large dendrimer macromolecules conjugated to estrogens were produced. These conjugates avoided the potential for cleaving and were unable to cross the cellular membrane, precluding the activation of nuclear ERs, but still resulted in rapid estradiol signaling (17). While in 2000 a novel estrogen receptor potentially located at the membrane was identified, i.e. G protein-coupled estrogen receptor 1 (GPER1) (18), overexpression of both ER α and ER β (19), along with immunohistochemical (20) and co-immunoprecipitation studies (21) also indicated that a subpopulation of these classical receptors are trafficked to the membrane (19). The development of transgenic mice allowed researchers to explore the effects of rapid signaling *in vivo*. In transgenic knockout mice devoid of ER α , and/or ER β , rapid estradiol signaling was eliminated in a brain-region and signaling pathway-dependent manner, suggesting that these receptors are responsible for many of the membrane signaling effects (22).

Membrane-initiated estrogen receptor signaling does not preclude downstream influences on gene expression. Particularly prominent is estradiol activation of PKC-MAPK signaling, ultimately resulting in the phosphorylation of CREB (23–26). Serine-133 phosphorylation of CREB can initiate a diverse array of transcriptional and behavioral changes, including by estradiol-mediated CREB activation *via*

membrane ER (mER) interactions with metabotropic glutamate receptors (mGluRs) (23). Initial findings in hippocampal neurons found estradiol acting through ER α -mGluR1a leads to MAPK-dependent CREB phosphorylation. This study elucidated a separate second pathway whereby activation of ER α or ER β associated with mGluR2 (and Gi/o-mediated inhibition of cAMP) resulted in a decrease in L-type calcium-channel mediated CREB phosphorylation (23). Follow-up studies found mER signaling through mGluR activation throughout the brain, which appears to be a mechanism allowing for diverse signaling outcomes. Not only does mER activation of group I or group II mGluRs activate separate cell signaling pathways, but mER pairing with different group I or II mGluRs (i.e. mGluR1 vs. mGluR5 and mGluR2 vs. mGluR3) can differentially impact neuronal function as well (27).

The interaction of mERs with mGluRs requires caveolin (CAV) (28–30), a family of scaffolding proteins involved in trafficking receptors to the membrane (31). The particular ER-mGluR pairing is mediated through the CAV isoform associated with the ER (28) (Figure 1). A single point mutation in ER α that disrupts receptor localization with CAV1 inhibited estradiol-induced CREB phosphorylation. Reducing CAV1 expression through siRNA knockdown inhibited estradiol-induced CREB phosphorylation while leaving the estradiol-induced L-type calcium channel-dependent decrease in CREB phosphorylation intact. In reciprocal experiments, siRNA knockdown of CAV3 inhibited estradiol-dependent activation of group II mGluRs without affecting estradiol-mediated CREB phosphorylation. In both cases, siRNA knockdown did not grossly impact mGluR signaling, demonstrating the essential nature of caveolin proteins to mER signaling (28). These data contribute to the understanding that CAV1 mediates ER α interactions with group I mGluRs through clustering the receptor at the membrane (28, 29, 33), while CAV3 is involved in the interactions between mERs and group II mGluRs (28). Additionally, an alternatively spliced form of ER α , ER $\alpha\Delta 4$, is highly expressed in membrane fractions derived from cultured cells. This receptor has been shown to associate with both mGluR2/3 and CAV3 in ARH membrane fractions (34).

Following these experiments, the precise mechanism of action linking ERs to mGluRs and to the membrane remained unclear, though palmitoylation was an attractive hypothesis. Palmitoylation is a reversible, post-transcriptional modification involved in the trafficking and function of proteins both within and outside the nervous system (35). Global pharmacological blockade of palmitoylation inhibited the downstream outcomes of membrane estradiol signaling, while introducing single point mutations at palmitoylation sites in both ER α and ER β was sufficient to inhibit membrane signaling (36). Two palmitoyl acetyltransferases (DHHC-7 and DHHC-21) have been shown to be crucial in ER membrane localization (37). Disrupting expression of either was sufficient to inhibit estradiol-dependent CREB phosphorylation (36), and to prevent ER α




FIGURE 1

mER transactivation of group I and group II mGluRs. 17β -Estradiol (17β E) binds to membrane-bound estrogen receptors (ER) to activate distinct signaling pathways via Group I (A) or Group II (B) mGluRs. (A) Membrane-ER interactions with Group I mGluRs, dependent on CAV1, activates G_q -mediated signaling through protein lipase C (PLC) and protein kinase C (PKC), subsequent activation of MEK, MAPK, and RSK, and ultimately the phosphorylation of CREB. PLC also activates IP₃, which binds to the IP₃ receptor (IP₃R) to result in the release of intracellular calcium (Ca^{2+}). (B) Membrane-ER activation of Group II mGluRs, dependent on CAV3, results in the $G_{i/o}$ -mediated inhibition of adenylyl cyclase (AC), decreasing the activity (indicated by dashed lines) of protein kinase A (PKA). This results in reduced L-type calcium channel currents and L-type calcium channel-dependent CREB phosphorylation. CaM, Calmodulin; CaMKIV, calmodulin-dependent protein kinase IV; CAV, caveolin; IP₃, inositol triphosphate; MEK, MAPK/ERK kinase; MAPK, mitogen-activated protein kinase; RSK, ribosomal S6 kinase. Figure adapted from (32).

from associating with CAV1 (38). siRNA knockdown of these enzymes together, but not separately, was sufficient to decrease the palmitoylation of CAV1 itself (38). These data suggest that palmitoylation is a crucial component in the interaction of mERs with CAV proteins, the coupling of mERs with mGluRs, and the subsequent signaling cascades.

While estrogen-mediated signaling plays a crucial role in the female brain, estrogen-mediated signaling is not absent in the male brain. Estrogen plays an important role in masculinizing the brain (39), and rapid estradiol signaling occurs in adult males, including through mGluRs. Estradiol activation of mGluR1a through ER β modulates sexual behavior in male quails (40, 41), and rodent studies have confirmed rapid mER-mGluR signaling in both the male and female adult cerebellum (42). In females, though, rapid signaling of estradiol, including through mGluRs, has been shown to be incredibly important in driving reproduction, including in the development of the luteinizing hormone surge which stimulates ovulation, the central event in female reproduction. In rodents, and certain other species, rapid membrane signaling is also crucial in the physical display of the principal reproductive behavior, lordosis. Finally, rapid membrane signaling has been shown to play an important role in female motivation for reproduction.

Ovulation and the luteinizing hormone surge

Ovulation is the central event in female reproduction, controlled by a network of neurons and astrocytes in the hypothalamus that act as a pattern generator, releasing gonadotropin-releasing hormone (GnRH) onto luteinizing hormone (LH) neurons in the anterior pituitary in small, rhythmic pulses (43, 44). Rising estradiol concentrations *via* ovarian release, trigger a switch from an estrogen-negative to an

estrogen-positive feedback loop. This estrogen-positive feedback loop, which is unique to females, is crucial in the surge release of LH that ultimately triggers ovulation (45). The preovulatory rise in circulating estradiol sharply increases GnRH neuronal activity and the release of LH from the pituitary to elicit ovulation (46, 47). Blocking either progesterone receptors or progesterone synthesis prevents the surge release of both GnRH and LH and halts the estrous cycle (48, 49). While GnRH neurons do not express ER α or nuclear progesterone receptors (45, 50, 51), kisspeptin neurons that are upstream regulators of GnRH signaling do express the necessary steroid receptors (52–54).

Classically it has been understood that both estradiol and progesterone released from the ovaries orchestrate the LH surge, but it has become apparent that progesterone is also synthesized *de novo* in the brain (55–57), and that it is this neuroprogesterone (neuroP) that is vital in the LH surge that ultimately leads to ovulation (58). Neuroprogesterone is synthesized in hypothalamic astrocytes that express mER α and mGluRs, and it has been shown that the LH surge relies upon the mER-mGluR signaling in these astrocytes (55, 59). Estradiol activation of mER α directly leads to the activation of mGluR1 and its downstream signaling cascades. mGluR1 activity increases inositol triphosphate and allows for the release of intracellular calcium ($[Ca^{2+}]_i$) stores (59, 60). The release of $[Ca^{2+}]_i$ activates a Ca^{2+} -sensitive adenylyl cyclase (AC-1), which increases the production of cAMP. This cAMP activates protein kinase A (60), leading to the synthesis of neuroP (56, 57, 59). Blocking neuroP synthesis in rats that had both ovaries and adrenals removed is sufficient to prevent the LH surge (61). Cell culture experiments in astrocytes isolated this signaling pathway. Blocking mGluR1a activity, or any part of the cell signaling cascade initiated by the ER α activation of mGluR1, in astrocytes inhibits neuroP synthesis (49, 55, 59, 60). Further, in the absence of estradiol, activating mGluR1a directly is sufficient to release $[Ca^{2+}]_i$ and induce neuroP synthesis (59, 62).


Lordosis

Another important aspect of reproduction controlled by ER interactions with mGluRs is lordosis. Lordosis is a reflexive behavior that is acutely triggered by mounting from a conspecific male. This behavior consists of an arching of the spine, the raising of both the head and the hindquarters, and the lifting of the tail (63). While integration of the tactile cues with other exterosensory cues is crucial for the display of lordosis, this behavior depends heavily on the appropriate timing of the release of ovarian hormones and the subsequent priming of neural circuits by these hormones. The role of intracellular and membrane-bound ERs, as well as the interaction between mERs and mGluRs, have all been shown to be important components in driving lordosis (64, 65).

A core circuit controlling lordosis is within the hypothalamus. Here, signaling between the arcuate nucleus (ARH), the medial preoptic nucleus (MPN), and the ventromedial nucleus of the hypothalamus (VMH) have been shown to be fundamental in the expression of lordosis (64, 66–70) (Figure 2). Within this circuit, estradiol first acts on ER α -containing neuropeptide Y (NPY) neurons in the ARH (21, 68, 71), allowing for the release of NPY onto NPY-Y1 receptors in ARH proopiomelanocortin (POMC) neurons. The subset of POMC neurons that are involved in reproduction project

further to the MPN where they release β -End onto neurons that express μ -opioid receptors (MORs). The estradiol-induced activation, and subsequent internalization, of MOR depends upon ER α activity (65). Throughout the estrous cycle the activation/internalization of this receptor is out of phase with the ability to express lordosis (69). That is, when MOR is internalized, a measure of activation, the display of lordosis is prohibited. As the cycle progresses, increasing progesterone levels ultimately result in the restoration of MOR to the membrane, a measure indicating that the receptors are not stimulated (72), and the behavior can be expressed. While counterintuitive, this estradiol inhibition of lordosis is necessary for its later full expression. While many neural changes must occur to result in the production of lordosis, recent work has shown that much of the machinery involved in this behavior utilizes fast-acting mER signaling cascades, and particularly those signaling through mGluRs.

Within the ARH, a subset of the NPY neurons express both ER α and mGluR1a, which have been shown to interact at the membrane to initiate signaling (21). The mER-mGluR signaling in the ARH has been shown to be crucial in both the internalization of MOR and the subsequent display of lordosis. The level of estradiol determines the expression of the mER α -mGluR1 complex in the ARH. When estradiol is low, the mER α -mGluR1 complex is present, but as estradiol levels rise the



expression of mER α -mGluR1 is reduced (73). In the ARH, antagonizing mGluR1a activity before estradiol treatment is sufficient to attenuate the internalization of MOR in the MPN as well as the ensuing expression of lordosis (20, 21). Activating mGluR1a in the ARH to circumvent the necessity of estradiol is sufficient to result in the internalization of MOR and lordosis (20). Both *in vitro* and *in vivo* activating mGluR1a through estradiol-induced mER α activity increases many important phosphoproteins, including PKC and CREB (20, 21, 23), and the internalization of MOR appears to depend at least in part upon PKC signaling. Downstream from mGluR1a activity, activating PKC signaling in the ARH in the absence of estradiol was sufficient to result in the internalization of MOR, and the amount of this internalization was comparable to that seen following estradiol treatment alone (21).

Another key component in the production of lordosis regulated by fast mER-mGluR activity is morphological changes to neuronal structure. Estradiol affects both the generation and pruning of dendritic spines, though this is not unique to the hypothalamic lordosis circuit but occurs throughout the brain (64, 74, 75) and appears due to retrograde signaling by endogenous opioids (76–78). Within this circuit, important morphological changes can be induced rapidly through mER α -mGluR1a signaling in the ARH. Within 4 hours, estradiol activation of mGluR1a results in an increase in the total number of dendritic spines, which remains for at least 48 hours. By 20 hours these spines display mushroom-shaped morphology, suggesting that these synapses are functional. Blocking mGluR1a activity prevented this spinogenesis, as well as attenuated the display of lordosis (64). Importantly, this time course of changes in morphology lines up with that of the display of lordosis.

Motivation

Estrogen membrane-receptor signaling has also been found to play a role in motivation. Though the long-term consequence of reproduction is the survival of the species through the production of offspring, the short-term motivation of reproduction is often the immediate drive for the rewarding aspects of the behavior, in females as much as in males (79). While work has focused on the physiology of ovulation and the rodent's reflexive response to mounting by a male, female sexual behavior is indicative of a motivational drive. Female rats placed in a modified operant chamber, in which the female can choose if and when she wants to copulate, will seek the male for copulation timed to maximize reward (80, 81). Additionally, other pre-copulatory behaviors from female rodents, such as hopping or darting (82, 83), further indicate a level of control over the mating process. This pre-copulatory activity, which is called "pacing," contributes to a robust dopamine response in the female nucleus accumbens (NAc) in response to mating

(84–87). This dopamine response, as well as further structural changes, in the NAc is regulated at least in part by estradiol signaling at the membrane.

The NAc is a key region in reward and incentive salience, and the limbic control of behavioral motivation, and inputs here affect structural morphology and subsequently behavioral output. The limbic system is important in the motivation to engage in reproductive behaviors in both males and females (67, 79), and projections from the hypothalamic nuclei robustly innervate this circuit. The reproductive limbic circuit consists of the MPN, the ventral tegmental area (VTA), and the NAc. A key node connecting the hypothalamus to the limbic component includes the MPN (88). Projections from here reciprocally innervate the mesolimbic dopamine system, including the VTA (89). The VTA projections to the NAc arise from cells that contain ERs (90) and are sensitive to fluctuations in estradiol levels (91), as well as estradiol-mediated signals arising from the MPN (90). These estradiol-mediated changes in VTA signaling have been shown to further affect the subsequent release of DA in the NAc (90).

The predominant output neuron in the NAc is the medium spiny neuron (MSN) - named due to the density of spines it possesses (92). The MSNs in the NAc receive both DAergic and glutamatergic inputs (93), and it has been shown that estradiol plays an important role in modulating both inputs (94–97). MSNs contain few nuclear ERs, suggesting that estradiol acts primarily through membrane-bound receptors ERs (98–103). As in the hypothalamus, estradiol modulates spine density in the NAc and the estradiol-induced morphological changes in the MSNs of the NAc are dramatic in terms of functional circuitry and neuronal morphology (104, 105). In female rodents, sexual experience modulates future sexual behavior through estradiol-mediated morphological changes within the limbic circuit (79). While the complete mechanisms of estradiol modulation on motivational circuitry have yet to be fully elucidated, it is likely that mER-mGluR signaling plays a role.

The role of membrane estradiol signaling, and particularly the interaction between mERs and mGluRs, in reproductive motivational drive can be further extrapolated from studies investigating when motivational drive becomes maladaptive, such as in drug addiction. In comparison with men, women tend to show heightened vulnerability to developing a drug addiction (106, 107). Additionally, subjective effects of a drug can vary across the menstrual cycle, as has been reported in response to cocaine. When estrogen levels are high, women report the greatest effects of the drug (108). Interest in the interaction between membrane ERs and group I mGluRs has been taken in understanding the influence of estradiol on drug addiction. In ovx rats, estradiol activation of mGluR5 has been shown to facilitate self-administration of cocaine, while inhibiting this signaling through an mGluR5 antagonist before estradiol administration is sufficient to attenuate this intake of the drug (109). Estradiol activation of mGluR5 in MSNs also

results in an increase in the phosphorylation of CREB (100), and a decrease in dendritic spine density in both the core region of the NAc (27, 104, 105). Conversely, estradiol activation of mGluR1 can result in an increase in spine density in the shell region of NAc (27). Taken together, the data suggest that mER-mGluR signaling are important in the drive to seek reward generally, as is apparent in drug taking behaviors, and in the reinforcement of reproductive behaviors.

Discussion

A great deal of progress has been made in understanding the physiology of rapid estradiol signaling, including the relationship between mERs and mGluRs. Rapid estradiol signaling has been found throughout the brain and the body. In the central nervous system, the signaling cascades initiated by the mER/mGluR complex has been shown to be involved in many physiological functions in both sexes, but particularly in females. In females, the diverse signaling cascades initiated by the interaction of mERs with mGluRs have been shown to play important roles in mediating key aspects of reproduction and motivation, among other crucial functions. While uncovering the roles of CAV and palmitoylation has led to further understanding of this complex signaling cascade, current and future research will inevitably expand our knowledge of mER/mGluR signaling and its physiological and behavioral outcomes.

References

- McKenna NJ, Lanz RB, O'Malley BW. Nuclear receptor coregulators: Cellular and molecular biology. *Endocr Rev* (1999) 20(3):321–44. doi: 10.1210/edrv.20.3.0366
- Evans RM. The steroid and thyroid hormone receptor superfamily. *Science* (1988) 240(4854):889–95. doi: 10.1126/science.3283939
- Giguère V, Yang N, Segui P, Evans RM. Identification of a new class of steroid hormone receptors. *Nature* (1988) 331(6151):91–4. doi: 10.1038/331091a0
- Jensen EV, Jacobsonk HI. Basic guides to the mechanism of estrogen action. *Recent Progr Hormone Res* (1962) 18:387–414.
- Toft D, Gorski J. A receptor molecule for estrogens: Isolation from the rat uterus and preliminary characterization. *Proc Natl Acad Sci* (1966) 55(6):1574–81. doi: 10.1073/pnas.55.6.1574
- Shughrue PJ, Lane MV, Merchenthaler I. Comparative distribution of estrogen receptor- α and - β mRNA in the rat central nervous system. *J Comp Neurol* (1997) 388(4):507–25. doi: 10.1002/(SICI)1096-9861(19971201)388:4<507::AID-CNE1>3.0.CO;2-6
- Simerly RB, Swanson LW, Chang C, Muramatsu M. Distribution of androgen and estrogen receptor mRNA-containing cells in the rat brain: An *in situ* hybridization study. *J Comp Neurol* (1990) 294(1):76–95. doi: 10.1002/cne.902940107
- O'Malley BW, Tsai MJ. Molecular pathways of steroid receptor action. *Biol Reprod* (1992) 46(2):163–7. doi: 10.1095/biolreprod46.2.163
- Marino M, Galluzzo P, Ascenzi P. Estrogen signaling multiple pathways to impact gene transcription. *Curr Genomics* (2006) 7(8):497–508. doi: 10.2174/138920206779315737
- Vrtačnik P, Ostanek B, Mencej-Bedrač S, Marc J. The many faces of estrogen signaling. *Biochem Medica* (2014) 24(3):329–42. doi: 10.11613/BM.2014.035
- Aranda A, Pascual A. Nuclear hormone receptors and gene expression. *Physiol Rev* (2001) 81(3):1269–304. doi: 10.1152/physrev.2001.81.3.1269
- Göttlicher M, Heck S, Herrlich P. Transcriptional cross-talk, the second mode of steroid hormone receptor action. *J Mol Med* (1998) 76(7):480–9. doi: 10.1007/s001090050242
- Szego CM, Davis JS. Adenosine 3',5'-monophosphate in rat uterus: Acute elevation by estrogen. *Proc Natl Acad Sci* (1967) 58(4):1711–8. doi: 10.1073/pnas.58.4.1711
- Kelly MJ, Moss RL, Dudley CA. Differential sensitivity of preoptic-septal neurons to microelectrophoresed estrogen during the estrous cycle. *Brain Res* (1976) 114(1):152–7. doi: 10.1016/0006-8993(76)91017-9
- Zheng J, Ramirez VD. Demonstration of membrane estrogen binding proteins in rat brain by ligand blotting using a 17 β -estradiol-[125I]bovine serum albumin conjugate. *J Steroid Biochem Mol Biol* (1997) 62(4):327–36. doi: 10.1016/S0960-0760(97)00037-X
- Stevis PE, Deecher DC, Suhadolnik L, Mallis LM, Frail DE. Differential effects of estradiol and estradiol-BSA conjugates. *Endocrinology* (1999) 140(11):5455–8. doi: 10.1210/endo.140.11.7247
- Harrington WR, Kim SH, Funk CC, Madak-Erdogan Z, Schiff R, Katzenellenbogen JA, et al. Estrogen dendrimer conjugates that preferentially activate extranuclear, nongenomic versus genomic pathways of estrogen action. *Mol Endocrinol* (2006) 20(3):491–502. doi: 10.1210/me.2005-0186
- Filardo EJ, Quinn JA, Bland KI, Frackelton AR. Estrogen-induced activation of erk-1 and erk-2 requires the G protein-coupled receptor homolog, GPR30, and occurs via trans-activation of the epidermal growth factor receptor through release of HB-EGF. *Mol Endocrinol* (2000) 14(10):1649–60. doi: 10.1210/mend.14.10.0532
- Razandi M, Pedram A, Greene G, Levin E. Cell membrane and nuclear estrogen receptors (ERs) originate from a single transcript: Studies of ER α and ER β

Author contributions

CJ wrote the first draft of the manuscript, and revised subsequent drafts. PMe, PMi contributed to and revised the manuscript. All authors read and approved the submitted version.

Funding

This work was supported by DA041808 and HD10007 to PGM and HD098284 to PEM. CSJ was supported by DA007234.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- expressed in Chinese hamster ovary cells. *Mol Endocrinol* (1999) 13(2):307–19. doi: 10.1210/me.13.2.307
20. Dewing P, Boulware MI, Sinchak K, Christensen A, Mermelstein PG, Micevych P. Membrane estrogen receptor- α interactions with metabotropic glutamate receptor 1a modulate female sexual receptivity in rats. *J Neurosci* (2007) 27(35):9294–300. doi: 10.1523/JNEUROSCI.0592-07.2007
 21. Dewing P, Christensen A, Bondar G, Micevych P. Protein kinase c signaling in the hypothalamic arcuate nucleus regulates sexual receptivity in female rats. *Endocrinology* (2008) 149(12):5934–42. doi: 10.1210/en.2008-0847
 22. Aibrahaim IM, Todman MG, Korach KS, Herbison AE. Critical *in vivo* roles for classical estrogen receptors in rapid estrogen actions on intracellular signaling in mouse brain. *Endocrinology* (2004) 145(7):3055–61. doi: 10.1210/en.2003-1676
 23. Boulware MI, Weick JP, Becklund BR, Kuo SP, Groth RD, Mermelstein PG. Estradiol activates group I and II metabotropic glutamate receptor signaling, leading to opposing influences on cAMP response element-binding protein. *J Neurosci* (2005) 25(20):5066–78. doi: 10.1523/JNEUROSCI.1427-05.2005
 24. Gu G, Rojo AA, Zee MC, Yu J, Simerly RB. Hormonal regulation of CREB phosphorylation in the anteroventral periventricular nucleus. *J Neurosci* (1996) 16(9):3035–44. doi: 10.1523/JNEUROSCI.16-09-03035.1996
 25. Szego EM, Barabás K, Balog J, Szilágyi N, Korach KS, Juhász G, et al. Estrogen induces estrogen receptor α -dependent cAMP response element-binding protein phosphorylation *via* mitogen activated protein kinase pathway in basal forebrain cholinergic neurons *in vivo*. *J Neurosci* (2006) 26(15):4104–10. doi: 10.1523/JNEUROSCI.0222-06.2006
 26. Wade CB, Dorsa DM. Estrogen activation of cyclic adenosine 5'-monophosphate response element-mediated transcription requires the extracellularly regulated kinase/mitogen-activated protein kinase pathway. *Endocrinology* (2003) 144(3):832–8. doi: 10.1210/en.2002-220899
 27. Gross KS, Brandner DD, Martinez LA, Olive MF, Meisel RL, Mermelstein PG. Opposite effects of mGluR1a and mGluR5 activation on nucleus accumbens medium spiny neuron dendritic spine density. *PLoS One* (2016) 11(9):e0162755. doi: 10.1371/journal.pone.0162755
 28. Boulware MI, Kordasiewicz H, Mermelstein PG. Caveolin proteins are essential for distinct effects of membrane estrogen receptors in neurons. *J Neurosci* (2007) 27(37):9941–50. doi: 10.1523/JNEUROSCI.1647-07.2007
 29. Christensen A, Micevych P. CAV1 siRNA reduces membrane estrogen receptor- α levels and attenuates sexual receptivity. *Endocrinology* (2012) 153(8):3872–7. doi: 10.1210/en.2012-1312
 30. Razandi M, Oh P, Pedram A, Schnitzer J, Levin ER. Ers associate with and regulate the production of caveolin: Implications for signaling and cellular actions. *Mol Endocrinol* (2002) 16(1):100–15. doi: 10.1210/mend.16.1.0757
 31. Anderson RGW. The caveolae membrane system. *Annu Rev Biochem* (1998) 67(1):199–225. doi: 10.1146/annurev.biochem.67.1.199
 32. Eisinger KRT, Gross KS, Head BP, Mermelstein PG. Interactions between estrogen receptors and metabotropic glutamate receptors and their impact on drug addiction in females. *Horm Behav* (2018) 104:130–7. doi: 10.1016/j.yhbeh.2018.03.001
 33. Bondar G, Kuo J, Hamid N, Micevych P. Estradiol-induced estrogen receptor- α trafficking. *J Neurosci* (2009) 29(48):15323–30. doi: 10.1523/JNEUROSCI.2107-09.2009
 34. Wong AM, Scott AK, Johnson CS, Mohr MA, Mittelman-Smith M, Micevych PE. ER α Δ4, an ER α splice variant missing exon4, interacts with caveolin-3 and mGluR2/3. *J Neuroendocrinol* (2019) 31(6):e12725. doi: 10.1111/jne.12725
 35. Fukata Y, Fukata M. Protein palmitoylation in neuronal development and synaptic plasticity. *Nat Rev Neurosci* (2010) 11(3):161–75. doi: 10.1038/nrn2788
 36. Meitzen J, Luoma JJ, Boulware MI, Hedges VL, Peterson BM, Tuomela K, et al. Palmitoylation of estrogen receptors is essential for neuronal membrane signaling. *Endocrinology* (2013) 154(11):4293–304. doi: 10.1210/en.2013-1172
 37. Pedram A, Razandi M, Deschenes RJ, Levin ER. DHHC-7 and -21 are palmitoyltransferases for sex steroid receptors. *Mol Biol Cell* (2012) 23(1):188–99. doi: 10.1091/mbc.e11-07-0638
 38. Eisinger KRT, Woolfrey KM, Swanson SP, Schnell SA, Meitzen J, Dell'Acqua M, et al. Palmitoylation of caveolin-1 is regulated by the same DHHC acyltransferases that modify steroid hormone receptors. *J Biol Chem* (2018) 293(41):15901–11. doi: 10.1074/jbc.RA118.004167
 39. Schwarz JM, McCarthy MM. Cellular mechanisms of estradiol-mediated masculinization of the brain. *J Steroid Biochem Mol Biol* (2008) 109(3–5):300–6. doi: 10.1016/j.jsbmb.2008.03.012
 40. Cornil CA, Ball GF, Balthazart J. Rapid control of male typical behaviors by brain-derived estrogens. *Front Neuroendocrin* (2012) 33(4):425–46. doi: 10.1016/j.yfrne.2012.08.003
 41. Seredynski AL, Balthazart J, Ball GF, Cornil CA. Estrogen receptor β activation rapidly modulates male sexual motivation through the transactivation of metabotropic glutamate receptor 1a. *J Neurosci* (2015) 35(38):13110–23. doi: 10.1523/JNEUROSCI.2056-15.2015
 42. Hedges VL, Chen G, Yu L, Krentzel AA, Starrett JR, Zhu JN, et al. Local estrogen synthesis regulates parallel fiber-purkinje cell neurotransmission within the cerebellar cortex. *Endocrinology* (2018) 159(3):1328–38. doi: 10.1210/en.2018-00039
 43. Herbison AE. The gonadotropin-releasing hormone pulse generator. *Endocrinology* (2018) 159(11):3723–36. doi: 10.1210/en.2018-00653
 44. Kauffman AS, Smith JT eds. *Kisspeptin signaling in reproductive biology* Vol. 342). New York: Springer (2013).
 45. Wintermantel TM, Campbell RE, Porteous R, Bock D, Gröne HJ, Todman MG, et al. Definition of estrogen receptor pathway critical for estrogen positive feedback to gonadotropin-releasing hormone neurons and fertility. *Neuron* (2006) 52(2):271–80. doi: 10.1016/j.neuron.2006.07.023
 46. Ferin M, Tempone A, Zimmering PE, Wiele RLV. Effect of antibodies to 17 β -estradiol and progesterone on the estrous cycle of the rat. *Endocrinology* (1969) 85(6):1070–8. doi: 10.1210/endo-85-6-1070
 47. Labhsetwar AP. Role of estrogens in ovulation: A study using the estrogen-antagonist, I.C.I. 46,474. *Endocrinology* (1970) 87(3):542–51. doi: 10.1210/endo-87-3-542
 48. Chappell PE. Stimulation of gonadotropin-releasing hormone surges by estrogen. I. role of hypothalamic progesterone receptors. *Endocrinology* (2000) 141(4):1477–85.
 49. Micevych P, Sinchak K. Estradiol regulation of progesterone synthesis in the brain. *Mol Cell Endocrinol* (2008) 290(1–2):44–50. doi: 10.1016/j.mce.2008.04.016
 50. Herbison AE, Theodosis DT. Localization of oestrogen receptors in preoptic neurons containing neurotensin but not tyrosine hydroxylase, cholecystokinin or luteinizing hormone-releasing hormone in the male and female rat. *Neuroscience* (1992) 50(2):283–98. doi: 10.1016/0306-4522(92)90423-Y
 51. Shivers BD, Harlan RE, Morrell JL, Pfaff DW. Absence of oestradiol concentration in cell nuclei of LHRH-immunoreactive neurones. *Nature* (1983) 304(5924):345–7. doi: 10.1038/304345a0
 52. Clarkson J, de Tassigny Xd'Anglemont, Moreno AS, Colledge WH, Herbison AE. Kisspeptin-GPR54 signaling is essential for preovulatory gonadotropin-releasing hormone neuron activation and the luteinizing hormone surge. *J Neurosci* (2008) 28(35):8691–7. doi: 10.1523/JNEUROSCI.1775-08.2008
 53. Navarro VM, Gottsch ML, Chavkin C, Okamura H, Clifton DK, Steiner RA. Regulation of gonadotropin-releasing hormone secretion by kisspeptin/dynorphin/neurokinin b neurons in the arcuate nucleus of the mouse. *J Neurosci* (2009) 29(38):11859–66. doi: 10.1523/JNEUROSCI.1569-09.2009
 54. Wakabayashi Y, Nakada T, Murata K, Ohkura S, Mogi K, Navarro VM, et al. Neurokinin b and dynorphin a in kisspeptin neurons of the arcuate nucleus participate in generation of periodic oscillation of neural activity driving pulsatile gonadotropin-releasing hormone secretion in the goat. *J Neurosci* (2010) 30(8):3124–32. doi: 10.1523/JNEUROSCI.5848-09.2010
 55. Kuo J, Hamid N, Bondar G, Prossnitz ER, Micevych P. Membrane estrogen receptors stimulate intracellular calcium release and progesterone synthesis in hypothalamic astrocytes. *J Neurosci* (2010) 30(39):12950–7. doi: 10.1523/JNEUROSCI.1158-10.2010
 56. Micevych PE, Chaban V, Ogi J, Dewing P, Lu JKH, Sinchak K. Estradiol stimulates progesterone synthesis in hypothalamic astrocyte cultures. *Endocrinology* (2007) 148(2):782–9. doi: 10.1210/en.2006-0774
 57. Sinchak K, Mills RH, Tao L, LaPolt P, Lu JKH, Micevych P. Estrogen induces *de novo* progesterone synthesis in astrocytes. *Dev Neurosci-basel* (2003) 25(5):343–8. doi: 10.1159/000073511
 58. Micevych P, Sinchak K. The neurosteroid progesterone underlies estrogen positive feedback of the LH surge. *Front Endocrinol* (2011) 2:90. doi: 10.3389/fendo.2011.00090
 59. Kuo J, Hariri OR, Bondar G, Ogi J, Micevych P. Membrane estrogen receptor- α interacts with metabotropic glutamate receptor type 1a to mobilize intracellular calcium in hypothalamic astrocytes. *Endocrinology* (2009) 150(3):1369–76. doi: 10.1210/en.2008-0994
 60. Chen C, Kuo J, Wong A, Micevych P. Estradiol modulates translocator protein (TSPO) and steroid acute regulatory protein (StAR) *via* protein kinase A (PKA) signaling in hypothalamic astrocytes. *Endocrinology* (2014) 155(8):2976–85. doi: 10.1210/en.2013-1844
 61. Micevych P, Sinchak K, Mills RH, Tao L, LaPolt P, Lu JKH. The luteinizing hormone surge is preceded by an estrogen-induced increase of hypothalamic progesterone in ovariectomized and adrenalectomized rats. *Neuroendocrinology* (2003) 78(1):29–35. doi: 10.1159/000071703
 62. Mittelman-Smith MA, Wong AM, Kathiresan ASQ, Micevych PE. Classical and membrane-initiated estrogen signaling in an *in vitro* model of anterior hypothalamic kisspeptin neurons. *Endocrinology* (2015) 156(6):2162–73. doi: 10.1210/en.2014-1803

63. Beach FA. Sexual attractivity, proceptivity, and receptivity in female mammals. *Horm Behav* (1976) 7(1):105–38. doi: 10.1016/0018-506X(76)90008-8
64. Christensen A, Dewing P, Micevych P. Membrane-initiated estradiol signaling induces spinogenesis required for female sexual receptivity. *J Neurosci* (2011) 31(48):17583–9. doi: 10.1523/JNEUROSCI.3030-11.2011
65. Micevych PE, Rissman EF, Gustafsson J, Sinchak K. Estrogen receptor- α is required for estrogen-induced μ -opioid receptor internalization. *J Neurosci Res* (2003) 71(6):802–10. doi: 10.1002/jnr.10526
66. Johnson C, Hong W, Micevych P. Opiogenetic activation of β -endorphin terminals in the medial preoptic nucleus regulates sexual receptivity. *Eneuro* (2020) 7(1):ENEURO.0315–19.2019. doi: 10.1523/ENEURO.0315-19.2019
67. Micevych PE, Meisel RL. Integrating neural circuits controlling female sexual behavior. *Front Syst Neurosci* (2017) 11:42. doi: 10.3389/fnsys.2017.00042
68. Mills RH, Sohn RK, Micevych PE. Estrogen-induced μ -opioid receptor internalization in the medial preoptic nucleus is mediated via neuropeptide y-Y1 receptor activation in the arcuate nucleus of female rats. *J Neurosci* (2004) 24(4):947–55. doi: 10.1523/JNEUROSCI.1366-03.2004
69. Sinchak K, Micevych P. Visualizing activation of opioid circuits by internalization of G protein-coupled receptors. *Mol Neurobiol* (2003) 27(2):197–222. doi: 10.1385/MN:27:2:197
70. Sinchak K, Dewing P, Ponce L, Gomez L, Christensen A, Berger M, et al. Modulation of the arcuate nucleus–medial preoptic nucleus lordosis regulating circuit: A role for GABAB receptors. *Horm Behav* (2013) 64(1):136–43. doi: 10.1016/j.yhbeh.2013.06.001
71. Sar M, Sahu A, Wr C. Kalra sp. localization of neuropeptide-y immunoreactivity in estradiol-concentrating cells in the hypothalamus. *Endocrinology* (1990) 127(6):2752–6. doi: 10.1210/endo-127-6-2752
72. Sinchak K, Micevych PE. Progesterone blockade of estrogen activation of μ -opioid receptors regulates reproductive behavior. *J Neurosci* (2001) 21(15):5723–9. doi: 10.1523/JNEUROSCI.21-15-05723.2001
73. Mahavongtrakul M, Kanjiya MP, Maciel M, Kanjiya S, Sinchak K. Estradiol dose-dependent regulation of membrane estrogen receptor- α , metabotropic glutamate receptor-1a, and their complexes in the arcuate nucleus of the hypothalamus in female rats. *Endocrinology* (2013) 154(9):3251–60. doi: 10.1210/en.2013-1235
74. Gross KS, Mermelstein PG. Estrogen receptor signaling through metabotropic glutamate receptors. *Vitam Horm* (2020) 114:211–32. doi: 10.1016/bs.vh.2020.06.003
75. Woolley C, McEwen B. Estradiol mediates fluctuation in hippocampal synapse density during the estrous cycle in the adult rat [published erratum appears in *J Neurosci* 1992 Oct;12(10):following table of contents]. *J Neurosci* (1992) 12(7):2549–54. doi: 10.1523/JNEUROSCI.12-07-02549.1992
76. Torres-Reveron A, Khalid S, Williams TJ, Waters EM, Jacome L, Luine VN, et al. Hippocampal dynorphin immunoreactivity increases in response to gonadal steroids and is positioned for direct modulation by ovarian steroid receptors. *Neuroscience* (2009) 159(1):204–16. doi: 10.1016/j.neuroscience.2008.12.023
77. Cholanian M, Krajewski-Hall SJ, McMullen NT, Rance NE. Chronic oestradiol reduces the dendritic spine density of KNDy (kisspeptin/neurokinin b/dynorphin) neurones in the arcuate nucleus of ovariectomised Tac2-enhanced green fluorescent protein transgenic mice. *J Neuroendocrinol* (2015) 27(4):253–63. doi: 10.1111/jne.12263
78. Moore AM, Coolen LM, Porter DT, Goodman RL, Lehman MN. KNDy cells revisited. *Endocrinology* (2018) 159(9):3219–34. doi: 10.1210/en.2018-00389
79. Meisel RL, Mullins AJ. Sexual experience in female rodents: Cellular mechanisms and functional consequences. *Brain Res* (2006) 1126(1):56–65. doi: 10.1016/j.brainres.2006.08.050
80. Mermelstein PG, Becker JB. Increased extracellular dopamine in the nucleus accumbens and striatum of the female rat during paced copulatory behavior. *Behav Neurosci* (1995) 109(2):354–65. doi: 10.1037/0735-7044.109.2.354
81. Xiao L, Becker JB. Hormonal activation of the striatum and the nucleus accumbens modulates paced mating behavior in the female rat. *Horm Behav* (1997) 32(2):114–24. doi: 10.1006/hbeh.1997.1412
82. Erskine MS. Solicitation behavior in the estrous female rat: A review. *Horm Behav* (1989) 23(4):473–502. doi: 10.1016/0018-506X(89)90037-8
83. McClintock MK, Adler NT. The role of the female during copulation in wild and domestic norway rats (*Rattus norvegicus*). *Behaviour* (1978) 67(1–2):67–95. doi: 10.1163/156853978X00260
84. Becker JB, Rudick CN, Jenkins WJ. The role of dopamine in the nucleus accumbens and striatum during sexual behavior in the female rat. *J Neurosci* (2001) 21(9):3236–41. doi: 10.1523/JNEUROSCI.21-09-03236.2001
85. Guarraci FA, Megroz AB, Clark AS. Paced mating behavior in the female rat following lesions of three regions responsive to vaginocervical stimulation. *Brain Res* (2004) 999(1):40–52. doi: 10.1016/j.brainres.2003.10.056
86. Jenkins WJ, Becker JB. Role of the striatum and nucleus accumbens in paced copulatory behavior in the female rat. *Behav Brain Res* (2001) 121(1–2):119–28. doi: 10.1016/S0166-4328(00)00394-6
87. Jenkins WJ, Becker JB. Dynamic increases in dopamine during paced copulation in the female rat. *Eur J Neurosci* (2003) 18(7):1997–2001. doi: 10.1046/j.1460-9568.2003.02923.x
88. Simerly RB, Swanson LW. Projections of the medial preoptic nucleus: A phaseolus vulgaris leucoagglutinin anterograde tract-tracing study in the rat. *J Comp Neurol* (1988) 270(2):209–42. doi: 10.1002/cne.902700205
89. Kokane SS, Perrotti LI. Sex differences and the role of estradiol in mesolimbic reward circuits and vulnerability to cocaine and opiate addiction. *Front Behav Neurosci* (2020) 14:74. doi: 10.3389/fnbeh.2020.00074
90. Tobiansky DJ, Will RG, Lominac KD, Turner JM, Hattori T, Krishnan K, et al. Estradiol in the preoptic area regulates the dopaminergic response to cocaine in the nucleus accumbens. *Neuropsychopharmacol* (2016) 41(7):1897–906. doi: 10.1038/npp.2015.360
91. Zhang D, Yang S, Yang C, Jin G, Zhen X. Estrogen regulates responses of dopamine neurons in the ventral tegmental area to cocaine. *Psychopharmacology* (2008) 199(4):625–35. doi: 10.1007/s00213-008-1188-6
92. Yager LM, Garcia AF, Wunsch AM, Ferguson SM. The ins and outs of the striatum: Role in drug addiction. *Neuroscience* (2015) 301:529–41. doi: 10.1016/j.neuroscience.2015.06.033
93. Russo SJ, Nestler EJ. The brain reward circuitry in mood disorders. *Nat Rev Neurosci* (2013) 14(9):609–25. doi: 10.1038/nrn3381
94. Becker JB. Direct effect of 17 β -estradiol on striatum: Sex differences in dopamine release. *Synapse [Internet]* (1990) 5(2):157–64. doi: 10.1002/syn.890050211
95. Mermelstein PG, Becker JB, Surmeier DJ. Estradiol reduces calcium currents in rat neostriatal neurons via a membrane receptor. *J Neurosci Off J Soc Neurosci* (1996) 16(2):595–604. doi: 10.1523/JNEUROSCI.16-02-00595.1996
96. Surmeier DJ, Ding J, Day M, Wang Z, Shen W. D1 and D2 dopamine-receptor modulation of striatal glutamatergic signaling in striatal medium spiny neurons. *Trends Neurosci* (2007) 30(5):228–35. doi: 10.1016/j.tins.2007.03.008
97. Thompson TL, Moss RL. Estrogen regulation of dopamine release in the nucleus accumbens: Genomic and nongenomic-mediated effects. *J Neurochem* (1994) 62(5):1750–6. doi: 10.1046/j.1471-4159.1994.62051750.x
98. Almey A, Filardo EJ, Milner TA, Brake WG. Estrogen receptors are found in glia and at extranuclear neuronal sites in the dorsal striatum of female rats: Evidence for cholinergic but not dopaminergic colocalization. *Endocrinology* (2012) 153(11):5373–83. doi: 10.1210/en.2012-1458
99. Almey A, Milner TA, Brake WG. Estrogen receptor α and G-protein coupled estrogen receptor 1 are localized to GABAergic neurons in the dorsal striatum. *Neurosci Lett* (2016) 622:118–23. doi: 10.1016/j.neulet.2016.04.023
100. Grove-Strawser D, Boulware MI, Mermelstein PG. Membrane estrogen receptors activate the metabotropic glutamate receptors mGluR5 and mGluR3 to bidirectionally regulate CREB phosphorylation in female rat striatal neurons. *Neuroscience* (2010) 170(4):1045–55. doi: 10.1016/j.neuroscience.2010.08.012
101. Küppers E, Beyer C. Expression of estrogen receptor- α and β mRNA in the developing and adult mouse striatum. *Neurosci Lett* (1999) 276(2):95–8. doi: 10.1016/S0304-3940(99)00815-0
102. Schultz KN, von Esenwein SA, Hu M, Bennett AL, Kennedy RT, Musatov S, et al. Viral vector-mediated overexpression of estrogen receptor- α in striatum enhances the estradiol-induced motor activity in female rats and estradiol-modulated GABA release. *J Neurosci* (2009) 29(6):1897–903. doi: 10.1523/JNEUROSCI.4647-08.2009
103. Stanić D, Dubois S, Chua HK, Tonge B, Rinehart N, Horne MK, et al. Characterization of aromatase expression in the adult male and female mouse brain. i. coexistence with oestrogen receptors α and β , and androgen receptors. *PLoS One* (2014) 9(3):e90451. doi: 10.1371/journal.pone.0090451
104. Peterson BM, Mermelstein PG, Meisel RL. Estradiol mediates dendritic spine plasticity in the nucleus accumbens core through activation of mGluR5. *Brain Struct Funct* (2015) 220(4):2415–22. doi: 10.1007/s00429-014-0794-9
105. Staffend NA, Loftus CM, Meisel RL. Estradiol reduces dendritic spine density in the ventral striatum of female Syrian hamsters. *Brain Struct Funct* (2011) 215(3–4):187–94. doi: 10.1007/s00429-010-0284-7
106. Brady KT, Randall CL. Gender differences in substance use disorders. *Psychiat Clin N Am* (1999) 22(2):241–52. doi: 10.1016/S0193-953X(05)70074-5
107. Hernandez-Avila CA, Rounsaville BJ, Kranzler HR. Opioid-, cannabis- and alcohol-dependent women show more rapid progression to substance abuse treatment. *Drug Alcohol Depend* (2004) 74(3):265–72. doi: 10.1016/j.drugalcdep.2004.02.001
108. Evans SM, Haney M, Foltin RW. The effects of smoked cocaine during the follicular and luteal phases of the menstrual cycle in women. *Psychopharmacology* (2002) 159(4):397–406. doi: 10.1007/s00213-001-0944-7
109. Martinez LA, Gross KS, Himmler BT, Emmitt NL, Peterson BM, Zlebnik NE, et al. Estradiol facilitation of cocaine self-administration in female rats requires activation of mGluR5. *Eneuro* (2016) 3(5):ENEURO.0140–16.2016. doi: 10.1523/ENEURO.0140-16.2016