Lawrence Berkeley National Laboratory

Recent Work

Title

LIFETIMES OF ROTATIONAL STATES FROM HEAVY-ION REACTIONS

Permalink

<https://escholarship.org/uc/item/0hm6q2w7>

Authors

Diamond, R.M.
Stephens, F.S.
Kelly, W.H.
et al.

Publication Date

1969

cy. 2

RECEIVED
JAN 17 1969
LIBRARY AND
DOCUMENTS SECTION

LIFETIMES OF ROTATIONAL STATES
FROM HEAVY-ION REACTIONS

R. M. Diamond, F. S. Stephens, W. H. Kelly,
and D. Ward

January 1969

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545

34
LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Physical Review Letters

UCRL-18702
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California 94720

AEC Contract No. W-7405-eng-48

LIFETIMES OF ROTATIONAL STATES FROM HEAVY-ION REACTIONS

R. M. Diamond, F. S. Stephens, W. H. Kelly, and D. Ward

January 1969

LIFETIMES OF ROTATIONAL STATES FROM HEAVY-ION REACTIONS*

R. M. Diamond, F. S. Stephens, W. H. Kelly,[†] and D. Ward^{††}Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

January 1969

The half-lives of several transitions in the ground-state collective bands of ^{160}Er , ^{158}Er , and ^{156}Er , produced by (^{40}Ar , 4n) reactions have been measured by a recoil-distance Doppler-shift method. These lifetimes are compared with those of several models. The mean time interval between the reaction and the population of the ground band was also determined.

The measurement of transition moments is an important method of testing nuclear models. In particular, the E2 moments of the $2 \rightarrow 0$ transitions in the ground-state collective bands of even-even nuclei have yielded much information on the nature of these bands. In the present work we have measured the $2 \rightarrow 0$ and several higher ground-band E2 moments (lifetimes) in each of three even-even Er nuclei, using a recoil-distance method. These nuclei were produced in the reactions $^{120,122,124}\text{Sn}(\text{Ar}, 4n)^{156,158,160}\text{Er}$.

The recoil-distance Doppler-shift method¹ is well suited to measure half-lives in the range 10^{-9} to 10^{-12} sec.²⁻⁵ Basically the method consists of stopping part of the excited nuclei recoiling from a thin target with a movable plunger placed closely behind the target. The fraction of the nuclei that live long enough to reach the plunger before emitting their radiation

will yield a normal gamma-ray line, whereas the rest will decay in flight and yield a Doppler-shifted line. By varying the distance of the plunger from the target, the fraction of unshifted transitions can be changed, and one can obtain the half-life of the transition if the velocity of the recoiling nucleus is also known.

The $(\text{Ar}, 4n)$ reaction produces a tightly collimated beam of product nuclei recoiling along the beam direction. Using a $\text{Ge}(\text{Li})$ detector at 0° to the beam direction, we were able to resolve completely the shifted and unshifted lines for all transitions over ~ 100 keV. The average recoil velocity could be obtained directly from the fractional energy difference of the shifted and unshifted lines, after a correction was made for the finite solid angle of the detector (the effective angle is different from 0° for large solid angles). The lead-covered plunger was attached to a precision micrometer with which its position could be adjusted to ± 0.002 mm. The targets were 1 mg/cm^2 thick metallic foils of separated tin isotopes which were supported over seven closely-spaced holes, 1.5 mm in diameter, in a tantalum disk. The foils were examined under a microscope and found to be flat within about 0.01 mm.

Some typical spectra for ^{160}Er are shown in Fig. 1. We have integrated the areas under the shifted and unshifted peaks, and calculated the fraction of each line that is unshifted at each distance. The analysis of these data is rather complex as account must be taken of the preceding rotational transitions, and also of the 10-20 unobserved transitions that precede entry into the ground band. The $(\text{Ar}, 4n)$ reactions are advantageous in that almost all the feeding of the ground band occurs at high spin values. Thus the formation

and decay of a given level are marked by the ground-band transitions which populate and depopulate that level. In the analysis we have used a computer program which considers three consecutive transitions at a time, e.g., $8 \rightarrow 6 \rightarrow 4 \rightarrow 2$. The program makes a least-square fit to the first set of data ($8 \rightarrow 6$ above) varying the target location and a sequence of three arbitrary transitions to mock up whatever cascade precedes this first observed transition. Simultaneous least-square fits to the following two sets of data ($6 \rightarrow 4$ and $4 \rightarrow 2$ above) are made with their two lifetimes as the only additional variables. A modified program was also used in which two components of different intensity and half-life were allowed to feed the highest state for which data were being considered. Addition of a second (slower) component turned out to be of significance in the case of ^{158}Er , but of little importance for $^{156,160}\text{Er}$.

A number of corrections had to be considered.⁶ An important one was that due to attenuation of the angular distribution of the gamma radiation by the large hyperfine field (~ 40 Mgauss) acting at the nucleus when the highly stripped target ion recoils into vacuum.^{7,8} This caused a maximum correction of 6% for half-lives of the order of 3×10^{-11} seconds. Small corrections for differences in absorption, efficiency, and effective solid angle between the shifted and unshifted lines were also made.

There are several sources of error. Although we do not know the absolute distance between the target and plunger to better than ± 0.02 mm, this does not contribute any error to the lifetime determinations of the individual rotational transitions, as each of these is measured relative to the preceding transition. The target and plunger are neither flat nor parallel to each other within ± 0.02 mm, which gives rise to instrumental limiting slopes corresponding to a mean life of $\sim 3 \times 10^{-12}$ sec. However, the rotational transitions

are determined appreciably better than this by considering not the slope of a transition, but its separation from that of the preceding transition. In addition, there are possible errors in deducing the average velocity of the recoiling nuclei from the difference in energy of the Doppler-shifted and unshifted peaks. Seven determinations of this difference show a maximum range of $\pm 2\%$ from the average value, and this error cancels out completely in the ratio of two half-lives measured in the same experiment. It should be noted that the $2 \rightarrow 0$ transitions in $^{158,160}\text{Er}$ were determined in separate experiments using a longer recoil chamber and somewhat thinner targets. The most significant error for the fast transitions is in the peak area integrations. The spectra are not particularly clean, and a small extraneous peak could cause a systematic error in the integration of a particular line. The uncertainty introduced by the integrations is estimated to be $\pm 0.47 \times 10^{-12}$ sec for $^{160,158}\text{Er}$ and $\pm 0.66 \times 10^{-12}$ sec for ^{156}Er which has somewhat poorer spectra. An error of $\pm 5\%$ has been set for the best transitions to account for other systematic errors and for the fact that a number of corrections (mentioned above) ranging up to a few percent in magnitude had to be made to the raw data.

Figure 2 shows a comparison of the experimental points and the computer-calculated curves for ^{160}Er . The lowest points of the fastest transitions tend systematically to be high, which probably indicates small amounts of somewhat slower feeding. The half-life values obtained are shown in Table I together with the energies of the transitions and the ratio of v/c for the recoils. Also listed are the experimental $B(E2)$ values, and these are compared with three sets of theoretical $B(E2)$ values.

Several qualitative conclusions can be drawn from the $B(E2)$ values in Table I. Within our limits of error, all these values are nearly consistent

with the rigid-rotor limit; however, those for ^{160}Er probably fall below this limit for high spin values and may even show an absolute decrease for the highest spin values, whereas those for the other two nuclei probably go above this limit. In fact, the $B(E2)$ values for ^{156}Er are fit about as well by the vibrator limit, as might be expected from the energy-level spacings in this nucleus. Since the more vibrational nuclei start with smaller $B(E2; 2 \rightarrow 0)$ values than the rotors, but increase more rapidly with increasing spin, there is a tendency for the $B(E2)$ values to become the same at higher spins. Thus for the three nuclei studied here the spread between the $B(E2; 2 \rightarrow 0)$ values is a factor of 2.5, but between the $B(E2; 6 \rightarrow 4)$ values it is only a factor of 1.3. This type of behavior has already been noted for the transition energies of even-even nuclei at high spin,⁹ and suggests that the difference between the rotational- and vibrational-type nuclei near the ground state may be disappearing with increasing angular momentum.

It is also of interest to compare the $B(E2)$ values with the predictions of the centrifugal stretching model.¹⁰ The last column in Table I makes this comparison using the calculations of Davydov and Ovcharenko.¹¹ The values of μ are taken from fits to the energy levels. The $B(E2)$ values do not increase as much as predicted by this model for ^{160}Er . Thus, in this nucleus centrifugal stretching is not the only change taking place with increasing angular momentum. This conclusion has previously been reached in other similar cases based on 1) the mixing of the ground- and β -bands,^{12,13} 2) the Mössbauer isomer-shift work,¹⁴⁻¹⁸ 3) the μ -mesic X-ray studies,¹⁹ and 4) theoretical calculations.^{20,21} It is generally supposed that the other important change occurring with increasing angular momentum is a reduction of the pairing correlations due to the

Coriolis force. It is not clear how this reduction would affect the $B(E2)$ values. For ^{158}Er , however, the $B(E2)$ values from the centrifugal stretching model fit very well and it will be of interest to see if such nuclei between rotors and vibrators do indeed stretch. For ^{156}Er , the centrifugal stretching and vibrator models give very nearly the same prediction, and in this case our data cannot really distinguish these values from those of the rigid rotor.

The decay curves, such as those in Fig. 2, also indicate the mean time interval between the reaction ($d=0$) and the population of the ground band ($1/e$ point on the curves for the fastest rotational transitions). These mean feeding times are $(6\pm3)\times 10^{-12}$, $(1.1\pm 0.3)\times 10^{-11}$, and $(1.6\pm 0.3)\times 10^{-11}$ sec, respectively, for ^{160}Er , ^{158}Er , and ^{156}Er . Considering that 10-20 transitions must be emitted in this time interval, these feeding times seem rather short, especially for the rotor, ^{160}Er . This subject is discussed more fully in another publication.²² The trend between ^{160}Er and ^{156}Er is rather striking, and it will be of interest to see if this reflects a general difference between rotors and vibrators, or is just something peculiar to these three cases.

This type of half-life measurement has been shown to give sufficient accuracy to distinguish among alternative models for the collective band of even-even nuclei. It can also obviously be used to measure many other lifetimes in the range 10^{-9} - 10^{-12} sec, where, in general, high precision will be less critical. We have made measurements on the reaction, $^{152}\text{Sm}(^{20}\text{Ne}, 4n)^{168}\text{Hf}$, and these studies appear to be feasible on reactions of this type, although the problem of side feeding has not been fully evaluated in this case. It seems clear that this method can produce much systematic information on lifetimes and feeding times following (HI, xn) reactions.

ACKNOWLEDGMENTS

The authors would like to express their appreciation to Mr. G. Steers and Mr. T. Gee for preparing the targets, and to Mr. D. Lebeck and Mrs. R. Garrett for help with the analyses.

FIGURE CAPTIONS

Fig. 1 Spectra from $^{124}\text{Sn}(^{40}\text{Ar}, 4n)^{160}\text{Er}$ taken with the plunger set at the indicated distances from the target. The positions of the unshifted (shifted) lines are given at the top (bottom) of the figure.

Fig. 2 The fraction of each transition which is unshifted in energy vs. the separation distance of the target and plunger. The symbols represent experimental points for the ground-band transitions in ^{160}Er . The solid lines are the computer-calculated curves, and the dashed line for the $8 \rightarrow 6$ transition shows the effect of using the program which allows two components to feed the ground-state band.

FOOTNOTES AND REFERENCES

*This work performed under the auspices of the United States Atomic Energy Commission.

[†]Permanent address: Department of Physics, Michigan State University, East Lansing, Michigan 48823.

^{††}Present address: Nuclear Physics Division, Chalk River Nuclear Laboratories, Chalk River, Ontario, Canada.

¹S. Devons, G. Manning, St.P. Bunbury, Proc. Phys. Soc. (London) A68, 18 (1955).

²I. F. Wright, Bull. Am. Phys. Soc. 6, 285 (1961).

³T. K. Alexander and K. W. Allen, Can. J. Phys. 43, 1563 (1965).

⁴T. K. Alexander, K. W. Allen, and D. C. Healey, Phys. Letters 20, 402 (1965); K. W. Allen, T. K. Alexander, and D. C. Healey, Phys. Letters 22, 193 (1966).

⁵P. G. Bizzetti, A. M. Bizzetti-Sona, S. Kalbitzer, and B. Povh, Z. für Physik 201, 295 (1967).

⁶K. W. Jones, A. Z. Schwarzschild, and E. K. Warburton, to be published.

⁷D. Ashery, N. Bahcall, G. Goldring, A. Spinzak, and Y. Wolfson, Nucl. Physics A101, 51 (1967).

⁸I. Ben Zvi, P. Gilad, M. Goldberg, G. Goldring, A. Schwarzschild, A. Spinzak, and Z. Vager, Nucl. Physics, to be published.

⁹F. S. Stephens, N. Lark, and R. M. Diamond, Phys. Rev. Letters 12, 225 (1964); R. M. Diamond, F. S. Stephens, and W. J. Swiatecki, Phys. Letters 11, 315 (1964).

- ¹⁰A. S. Davydov and A. A. Chaban, Nucl. Physics 20, 499 (1960).
- ¹¹A. S. Davydov and V. I. Ovcharenko, Soviet J. Nucl. Physics 3, 740 (1966).
- ¹²Y. Yoshizawa, B. Elbek, B. Herskind, and M. C. Olesen, Nucl. Physics 73, 273 (1965).
- ¹³G. C. Seaman, J. G. Greenberg, D. A. Bromley, and F. K. McGowan, Phys. Rev. 149, 925 (1966).
- ¹⁴J. Fink and P. Kienle, Phys. Letters 17, 326 (1965).
- ¹⁵D. Yeboah-Amankwah, L. Grodzins, and R. B. Frankel, Phys. Rev. Letters 18, 791 (1967).
- ¹⁶P. Steiner, E. Gerdau, P. Kienle, and H. J. Körner, Phys. Letters 24, B515 (1967).
- ¹⁷U. Atzmony, E. R. Bauminger, J. Hess, A. Mustachi, and S. Ofer, Phys. Rev. Letters 18, 1061 (1967).
- ¹⁸P. Kienle, W. Henning, G. Kaendl, H. J. Körner, H. Schaller, and F. Wagner, Proc. Int. Conf. on Nuclear Structure, Tokyo, 1967, edited by J. Sanada; J. Phys. Soc. of Japan, 24, 207 (1968).
- ¹⁹S. Bernow, S. Devons, I. Duerdoth, D. Hitlin, J. W. Kast, E. R. Macagno, J. Rainwater, K. Runge, and C. S. Wu, Phys. Rev. Letters 18, 787 (1967).
- ²⁰T. Udagawa and R. K. Sheline, Phys. Rev. Letters 16, 325 (1966).
- ²¹E. R. Marshalek, Phys. Rev. Letters 20, 214 (1968).
- ²²F. S. Stephens, R. M. Diamond, W. H. Kelly, J. O. Newton, and D. Ward, to be published.

Table I.

Nucleus	Transition	Energy (keV)	Recoil v/c(%)	$T_{1/2}$ (psec)	$\alpha_T \times 10^2$ ($\alpha_K + 1.3\alpha_L$)	$B(E2; I \rightarrow I-2)$ ($e^2 \times 10^{-48} \text{ cm}^4$)	$B(E2)_{\text{rot}}^a$	$B(E2)_{\text{vib}}^a$	$B(E2)_{\text{c.s.}}^a$
^{160}Er	$2 \rightarrow 0$	126.2	1.96	919 ± 46	1.30	$.83 \pm .04$	(.83)	(.83)	(.83) ^{b)}
	$4 \rightarrow 2$	264.3	1.80	34.5 ± 1.7	.089	$1.16 \pm .06$	1.19	1.66	1.25
	$6 \rightarrow 4$	376.3	1.80	$5.39 \pm .47$	.037	$1.34 \pm .12$	1.31	2.49	1.48
	$8 \rightarrow 6$	464.6	1.80	$2.16 \pm .47$	.020	$1.18 \pm .27$	1.37	3.32	1.70
	$10 \rightarrow 8$	532.1	1.80	$1.24 \pm .47$	.014	$1.05 \pm .40$	1.40	4.15	1.91
^{158}Er	$2 \rightarrow 0$	192.7	1.96	300 ± 15	.283	$.55 \pm .03$	(.55)	(.55)	(.55) ^{c)}
	$4 \rightarrow 2$	335.7	1.87	$14.4 \pm .72$	.050	$.87 \pm .04$	.79	1.10	.87
	$6 \rightarrow 4$	443.8	1.87	$2.80 \pm .47$	.026	$1.14 \pm .19$	.87	1.65	1.11
	$8 \rightarrow 6$	523.8	1.87	$1.21 \pm .47$	.015	$1.16 \pm .45$	.91	2.20	1.35
^{156}Er	$2 \rightarrow 0$	344.4	2.20	33.2 ± 1.7	.047	$.33 \pm .02$	(.33)	(.33)	(.33) ^{d)}
	$4 \rightarrow 2$	452.9	2.20	$5.42 \pm .66$	.022	$.53 \pm .06$	.47	.66	.61
	$6 \rightarrow 4$	543.2	2.20	$1.14 \pm .66$	.013	$1.03 \pm .60$	.52	.99	.91

a) These values have been normalized to the experimental $2 \rightarrow 0$ values.

b) These values have been calculated for $\mu = 0.3$ and $\gamma = 10^\circ$.

c) These values have been calculated for $\mu = 0.4$ and $\gamma = 10^\circ$.

d) These values have been calculated for $\mu = 0.8$ and $\gamma = 10^\circ$.

XBL683-2083

Fig. 1

XBL 6812-7511

Fig. 2

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720