

## **UC Irvine**

### **Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health**

#### **Title**

Emergency Medicine Resident RVU Trends at an Academic Medical Center

#### **Permalink**

<https://escholarship.org/uc/item/0j67p1bc>

#### **Journal**

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 23(4.1)

#### **ISSN**

1936-900X

#### **Authors**

Von Dohlen, Meredith  
Evans, Lauren  
Bouldin, Meryll  
[et al.](#)

#### **Publication Date**

2022

#### **Copyright Information**

Copyright 2022 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

## 2 Don't Break Their Hearts - Discharging Patients with Moderate Risk HEART Scores from the Emergency Department

Nadia Lehtihet, Allison Becker, Jessica Waters, Rory Spiegel, Rahul Bhat

**Learning Objectives:** To undertake an ED focused quality improvement initiative to allow for resident education in the fundamentals of research inquiry and study design and to provide experience in drafting research proposals, IRB submissions, medical abstracts and final manuscripts.

**Background:** Identifying patients at low risk for clinically relevant adverse cardiac events (CRACE) has the potential to decrease unnecessary hospitalizations. Earlier studies have shown that patients discharged with non-ischemic ECG and modified HEART (History, ECG, Age, Risk Factors, Troponin) score  $\leq 3$  had no deaths, acute myocardial infarctions (MI), or coronary revascularization events at 30 days. New observational data suggest the rate of CRACE for patients classified as moderate-risk by HEART score of 4-6 may be similarly low.

**Objective:** To determine the incidence of CRACE in patients discharged with a moderate-risk HEART score.

**Methods:** We reviewed ED visits involving adults presenting to an urban teaching hospital in Washington, DC. A prior quality improvement project determined the low overall rate of CRACE in patients with 2 negative 4th generation troponins, allowing clinicians to discharge patients with a HEART score of 4-6. This study was an analysis of patients with inclusion criteria of: (1) primary diagnosis code of chest pain for either ED reason for visit or discharge, (2) HEART score of 4-6, and (3) discharge from December 1, 2019 to March 15th, 2020.

**Outcomes:** The primary outcome was the 30 day rate of CRACE, defined as all-cause mortality, STEMI, fatal arrhythmia, and cardiopulmonary arrest. Secondary outcome was the rate of NSTEMI within 30 days of ED visit.

**Results:** 298 patients with HEART score 4-6 were discharged. 296 of 298 patients were alive with no recorded events at 30 days. Two patients had no available follow up data. Overall, the CRACE rate was 0% and NSTEMI rate was 0.34% within 30 days of discharge. One discharged patient followed up with cardiology and prior to arranged stress testing, re-presented with worsening chest pain and was found to have NSTEMI.

**Conclusion:** CRACE and NSTEMI are exceedingly rare within 30 days of ED discharge for patients with a moderate risk HEART score.

**Table 1.** Class average RVUs/hour.

	Number of resident data points	Mean	SD
PGY1	10	1.97	0.26
PGY2	20	2.67	0.77
PGY3	30	3.35	0.36

**Table 2.** Longitudinal mean RVUs/hour per class.

	2020 – 2021	2021 – 2022 (YTD)
PGY 3 - will graduate 2022	1.85	2.28
PGY 2 - will graduate 2023	1.23	2.89

## 3 Emergency Medicine Resident RVU Trends at an Academic Medical Center

Meredith Von Dohlen, Lauren Evans, Meryll Bouldin, Amanda Young, Sarah Greenberger, Rachael Freeze-Ramsey, Travis Eastin, Carly Eastin

**Learning Objectives:** To determine average RVUs per hour for emergency medicine residents at a tertiary-care, university-based academic medical center and to characterize change in mean RVUs per hour as residents advance in training.

**Background:** Physician productivity is often reported in relative value units (RVU). However, RVUs are infrequently reported during residency. Studying RVUs in varied training settings may better define benchmarks for progression of resident productivity.

**Methods:** This was a retrospective, observational study of PGY 1-3 residents at an academic, tertiary-care center. PGY2s and PGY3s were considered equivalent in shift scheduling and responsibilities. From 07/2019 to 09/2021, RVUs were extracted from the electronic health record (Epic) using E&M billing codes (excluding procedures, which were not tied to specific residents). In the PGY1 year and continuing longitudinally, residents received individual productivity reports. Individual metrics were de-identified, coded, and analyzed.

**Results:** The primary outcome was the measure of mean RVUs/hr per resident overall and by class. Secondary outcomes were the change in RVUs/hr between classes at the end of each year, as well as the change in RVUs/hr for the same class year to year. Descriptive statistics were reported in mean with standard deviations. One-way ANOVA was used to determine if PGY-level had a significant effect on RVUs generated; the remainder of comparisons were made with student's t-test. 60 RVU data points were obtained, representing 40 residents. Two classes were followed longitudinally (Table 2). Overall mean RVU/hr per resident was 2.89 RVU/hr (SD 0.89). Mean RVU/hr per resident for PGY1s, PGY2s, and PGY3s were 1.97 RVU/hr (SD 0.26), 2.67 (SD 0.77), and 3.35 (SD 0.36) respectively. Class year was predictive of RVUs generated ( $p < 0.001$ ). There was no significant difference in RVUs within a single class from PGY2 to PGY3 ( $p = 0.528$ ), but there was a significant increase from PGY1 to PGY2 ( $p < 0.001$ ).

**Conclusion:** Resident RVUs in our academic ED were associated with training year, but longitudinally, the only statistically significant increase was from PGY1 to PGY2.

Limitations include ongoing effects of the COVID-19 pandemic on productivity.

## 4 Exposure of Emergency Medicine Clerkship Students to Psychiatric Emergencies

Connor Knowles

**Learning Objectives:** To determine if a dedicated didactic and clinical curriculum in psychiatric emergency care would improve medical student comfort and perceived ability to assess psychiatric emergencies. It is believed these parameters would increase with this intervention.

**Background:** Medical students rotating in the emergency department (ED) are often discouraged or otherwise limited from seeing patients with psychiatric emergencies. Little is known about students' perceived preparedness and comfort levels in the assessment of individuals with acute mental health related complaints.

**Objectives:** To determine if a dedicated didactic and clinical curriculum in psychiatric emergency care would improve medical student comfort and perceived ability to assess psychiatric emergencies. It is believed these parameters would increase with this intervention.

**Methods:** Over a 10-month study period, students rotating in an Emergency Medicine (EM) clerkship at a large community-based ED received 1-2 hours of EM attending-led psychiatric case-based discussions in addition to an 8-hour clinical experience with the ED behavioral health team, assessing a wide scope of behavioral health emergencies. An identical pre-and post-rotation 10 question survey (1-10 Likert scale) was given regarding their comfort level and perceived abilities to assess aspects of this patient population. Statistical assessment was performed utilizing the Wilcoxon signed rank test.

**Results:** Seventeen students completed this voluntary study to completion seeing a mean of 4.8 patients during their 8-hour clinical experience, about 60% presenting with acute suicidal ideation. A mean improvement in the Likert scale was noted in "comfort in evaluation and interviewing" [+1.82 (p<0.01)], "decision-making for appropriate disposition" [+2.41(p<0.01)], and "screening for acute suicidal risk" [+1.76 (p<0.01)]. Similar improvement patterns of these parameters were also shown for complaints of acute mania and acute psychosis.

**Conclusion:** Focused education and exposure to patients with psychiatric emergencies had a positive impact on medical students' perception of their ability to assess and manage this population. This may lead to greater preparedness among incoming EM residents.

## 5 High sensitivity troponin - 6 hours is the magic number

O moyemen Blue, Michael Nguyen, Huyen-Trang Thai, Hayley Harvey, Maria Cacciapuoti, Rory Spiegel, Rahul Bhat

**Learning Objectives:** We wanted to assess whether emergency department (ED) patients presenting with chest pain with a high sensitivity troponin under the 99th percentile drawn at 6 hours or greater after symptom onset could safely be discharged from the ED.

**Background:** High sensitivity troponin (hsT) assays have become widespread for emergency department (ED) evaluation of acute chest pain. Several studies have demonstrated safety of a "rapid rule-out" strategy, evaluating initial hsT with repeat testing at 1-2 hours. A small number of patients can be discharged using this strategy while the most patients have a detectable hsT level and require a prolonged workup.

**Objective:** We assessed if a hsT under the 99th percentile drawn at 6 hours or greater from symptom onset could safely rule out patients.

**Methods:** We conducted a multicenter retrospective study examining ED patients with chest pain who did not meet rapid-rule out criteria and were admitted for further evaluation. Among these admitted patients, we assessed the rate of clinically relevant acute cardiac events (CRACE) and NSTEMI in patients with hsT less than 99th percentile (34 ng/mL in females, 53 ng/mL in males) obtained after at least 6 hours of chest pain. CRACE was defined as death, cardiac arrest, STEMI, or life-threatening arrhythmia. A manual chart review was conducted with 10% of randomly selected charts to estimate the number of patients admitted with nonischemic ECG and no other compelling reason for admission (unstable vitals or additional diagnoses).

**Results:** Out of 1189 patients admitted, we found 30 CRACE, all of which occurred in patients admitted for another compelling reason or ischemic ECG. 36 patients had an NSTEMI, of which 33 were identified with hsT above 99th percentile within 6 hours of chest pain onset. This left 0 CRACE and 3 NSTEMI among the 430 patients with a negative hsT at 6 hours and nonischemic ECG and no other compelling reason for admission.

**Conclusion:** Patients who have hsT values under the 99th percentile after 6 hours of chest pain have a low rate of CRACE and NSTEMI. Future prospective studies to evaluate the safety of outpatient management are warranted.