

Lawrence Berkeley National Laboratory

Recent Work

Title

Tests of Quantum Mechanics at a Phi-Factory

Permalink

<https://escholarship.org/uc/item/0k60t2t4>

Journal

The DaPHIne Physics Handbook, III

Author

Eberhard, P.H.

Publication Date

1994-08-09



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA


Physics Division

To be published as a chapter in *The DaPHne Physics Handbook*,
L. Maiani, G. Pancheri, and N. Paver, Eds., Vol. III, Servizio
Documentazione dei Laboratori Nazionali di Frascati, Frascati,
Italy, 1994

Tests of Quantum Mechanics at a ϕ -Factory

P.H. Eberhard

August 1994



REFERENCE COPY |
Does Not |
Circulate |

Bldg. 50 Library.

Copy 1

LBL-35983

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

LBL-35983

August 9, 1994

**Tests of Quantum Mechanics
at a ϕ -Factory***

Philippe H. Eberhard

Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

*This work was supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098

LBL-35983

August 9, 1994

Tests of Quantum Mechanics at a ϕ -Factory.*

Philippe H. Eberhard
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

ABSTRACT: Unique tests of quantum mechanics, which can only be performed at a ϕ -factory, are proposed for $D_s\phi$ ne. Each of these tests consists of measuring the difference between the predicted and the actual amount of interference between two processes leading from a single pure initial state to a single pure final state of a kaon system.

Estimates are made of the upper limits that will be set for the amount of violation if the predictions of quantum mechanics turn out to be correct. They are of the order a fraction of one percent. For the case where, on the contrary, a significant violation is found, several decoherence mechanisms are considered.

*This work is supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

Contents

1	Introduction.	1
2	Test Performed in the Past.	2
3	Two-Kaon Systems.	3
3.1	Predictions and Errors.	5
3.2	Performance at $D\phi ne$	6
3.3	$K_L K_L$ events	7
4	Alternate Theories.	8
4.1	Completeness.	8
4.2	Examples of Decoherence Mechanisms.	9
5	Conclusions.	11
	References	12

1 Introduction.

Opportunities for testing quantum mechanics are unique at a ϕ -factory. Some of those that are suitable for Da ϕ ne are described here. In the past, few checks of quantum mechanics have been performed in neutral kaon physics and none with a two-kaon system. At a ϕ -factory, several such tests are possible. The property tested is the predicted amount of interference between two different physical processes.

When there are two different mechanisms, 1 and 2, by which a system can transit from one state to another, the resulting probability of transition P in classical physics is the sum of the transition probabilities, P_1 and P_2 , due to the individual processes,

$$P = P_1 + P_2 . \quad (1)$$

In quantum mechanics, when the initial and the final states are pure single states and when the system does not interact with the environment, what one is supposed to add are not probabilities, but probability amplitudes, ψ_1 and ψ_2

$$\psi = \psi_1 + \psi_2 . \quad (2)$$

The interference between the two processes results from that property.

For the tests, one can express the transition probability P as

$$P = |\psi_1|^2 + |\psi_2|^2 + 2(1 - \zeta)Re\{\psi_1^*\psi_2\} . \quad (3)$$

where $|\psi_1|^2$ and $|\psi_2|^2$ are the individual transition probabilities of each process, the last term represents the amplitude of the interference effect, and ζ is a parameter which we call the “decoherence parameter”. If there is no decoherence mechanism, quantum mechanics predicts $\zeta = 0$; thus ζ measures the amount of violation of the theoretical prediction. $1-\zeta$ is analogous to what is called “visibility” in optics. ζ can be measured by measuring P . The tests proposed here are designed to yield a low upper limit for ζ if the predictions of quantum mechanics are 100% correct.

If the initial or the final states included in the sample of events are not pure, the interference effect is reduced. Background can fake a violation. The advantage of a ϕ -factory is the possibility to have initial kaon systems in quite pure states, final decay modes that are theoretically equivalent to pure states, and practically no decoherence effect.

2 Test Performed in the Past.

Tests of quantum mechanics using $K_L \rightarrow K_S$ regeneration have been proposed long ago, [1]. They consist of transporting a neutral kaon beam over a distance long enough for all its K_S -component to decay; letting it impinge on a piece of material called the regenerator; and recording the number of decays into two charged pions as a function of distance to the regenerator.

There are two processes that produce the $\pi^+\pi^-$ -decay of a kaon in the very forward direction after the regenerator :

- 1) plain transmission of a K_L and decay via the CP violating mode;
- 2) $K_L \rightarrow K_S$ regeneration followed by a CP conserving decay.

The initial K_L can be considered as a pure state in each momentum interval, if that interval is small enough. Because the kaon has spin zero, all configurations of its $\pi^+\pi^-$ -decays can be considered as two pions in an S-wave state, which is a pure state. Thus, at each momentum, the two-pion state from the decay of a kaon in the very forward direction is a pure state. Whether the kaon is transmitted or coherently regenerated, the state of the regenerator is the same, before and also after the kaon traversed it. Therefore the two processes interfere between one another. As a function of proper time t , the decay rate can be written as

$$\Gamma(t) = c_S e^{-\Gamma_S t} + c_L e^{-\Gamma_L t} + 2c_I e^{-(\Gamma_S + \Gamma_L)\frac{t}{2}} \cos(\Delta m t - \phi) , \quad (4)$$

where Γ_S and Γ_L are the total decay rates of the K_S and of the K_L mesons, respectively; Δm is the $K_L - K_S$ mass difference; and c_S , c_L , c_I , and ϕ are parameters that can be measured by the shape of the time distribution $\Gamma(t)$.

The third term in Eq. (4) is the interference term between the two processes. Quantum theory predicts $c_i = \sqrt{c_S c_L}$. In accordance with Eq. (3), we define

$$\zeta_{\text{regen}} = 1 - \frac{c_I}{\sqrt{c_S c_L}} , \quad (5)$$

which measures the amount of violation in the regeneration process.

A test of this kind was performed with neutral kaons of momentum ranging from 4 to 10 GeV/c and a carbon regenerator, [2]. The result was

$$\zeta_{\text{regen}} = 3\% \pm 2\% . \quad (6)$$

The error was essentially statistical.

The same type of test could be performed at DaΦne. The accuracy would be improved. For instance, one may surround one of the two intersection

points with a regenerator of a cylindrical form, with its axis along the e^+e^- beam line, with a radius of 20 cm and a height of 20 cm too. The regenerator would intercept K_L emitted within a solid angle of about 5 sr. The thickness of the cylinder should be adjusted to produce, just behind the regenerator, more $\pi^+\pi^-$ -decays by regenerated K_S than by transmitted K_L . Then one is sure that the K_S - K_L interference pattern visible in the $\pi^+\pi^-$ -mode is as long as possible. For instance, a beryllium cylinder 2.4 cm thick would be an adequate regenerator. It would regenerate one K_L in 10^4 according to Ref. [3], i.e. an appropriate amount of regenerated K_S .

At a luminosity $\mathcal{L} = 10^{33}$, in one month calendar time with an efficiency of 30%, i.e. an integrated luminosity of 1000 pb^{-1} , one may expect more than 10^9 K_L through the regenerator according to Refs. [4] and [5]. Then the decoherence parameter ζ would be measured with a statistical error of less 1%. If the angle of the kaon emerging from the regenerator can be determined to better than 20 mr, the effect of the incoherent regeneration and elastic scattering should introduce an uncertainty of less than 1%. Finally one should be able to eliminate background in $\pi^+\pi^-$ -decays down to less than 10^{-3} times the K_L decay rate into $\pi^+\pi^-$ at a machine like Da ϕ ne designed to measure the CP violation parameter called ϵ'/ϵ . Therefore combining statistical and systematic uncertainties, the final error should be of the order of 1%.

Since the value of ζ of Ref. [2] is 1.5 standard deviation away from zero, it would be significant to repeat that test with an error of, let us say, 1%. Of course, this test is not one that can be considered as unique for a ϕ -factory. However it would establish a limit for spontaneous decoherence in regeneration of kaons comparable to the limit of 0.6% obtained for decoherence in neutron interferometry, [6].

3 Two-Kaon Systems.

The really unique tests of quantum mechanics at Da ϕ ne are tests with quantum states made of two kaons. In e^+e^- -collisions of this kind of energy, only objects odd under charge conjugation are formed. The two-kaon system created via ϕ production and decay in a given direction is in a pure quantum state. Its state vector can be expressed as

$$\Psi_0 = \frac{1}{\sqrt{2}}(K^0\bar{K}^0 - \bar{K}^0K^0) \quad (7)$$

$$\approx \frac{1}{\sqrt{2}}(K_LK_S - K_SK_L) . \quad (8)$$

In the terms $K^0\bar{K}^0$, \bar{K}^0K^0 , K_LK_S , and K_SK_L , the first symbol refers to the state of the particle emitted on the left and the second symbol to the particle on the right. The sign \approx is used in Eq. (8) instead of the sign = because of a slight inaccuracy in the normalization factor.

As stated above, kaons living long enough to be reduced to their K_L component are pure states and $\pi^+\pi^-$ -decay states are equivalent to pure $\pi^+\pi^-$ S-wave states. If we accept the $\Delta S/\Delta Q$ rule, each of the semileptonic decay states $\ell^+\nu\pi^-$ and $\ell^-\bar{\nu}\pi^+$ can also be considered as a pure state in the context of this paper. As to multiple-particle states where the kaon on the left has decayed into a pure state f_a and the kaon on the right into a pure state f_b , they are pure states themselves. All processes leading from the initial ϕ -meson to any one of these pure multiparticle final states $f_a f_b$ interfere between each other.

In particular, consider the possibility of having the same decay state f , for instance $\pi^+\pi^-$, or $\ell^+\nu\pi^-$, or $\ell^-\bar{\nu}\pi^+$, as the final decay product on the left as well as on the right. This may happen via two processes,

- 1) a K_L emitted to the left and a K_S to the right; or
- 2) a K_S to the left and a K_L to the right;

and, in either case, both the K_L and the K_S decaying into the same state f . These two processes interfere between one another.

Let B_L and B_S be the branching ratios of K_L and K_S , respectively, into the decay state f . From Eq. (8), one gets the rate $\Gamma(t_a, t_b)$ of a kaon on the left decaying into f at proper time t_a and the one on the right decaying also into f but at proper time t_b :

$$\Gamma(t_a, t_b) \approx \frac{B_L\Gamma_L B_S\Gamma_S}{2} \left(e^{-\Gamma_S t_a} + e^{-\Gamma_S t_b} - 2e^{-\Gamma_S \frac{t_a+t_b}{2}} \cos(\Delta m(t_a - t_b)) \right) . \quad (9)$$

The third term in the largest bracket of Eq. (9) is the interference term. For $t_a = t_b$, that term is negative and large enough to make $\Gamma(t_a, t_b) = 0$. However, if, for any reason, the interference term is reduced by a factor $1-\zeta$, the rate for $t_a = t_b$ becomes non-zero. These conditions are particularly favorable for a test of quantum mechanics because, then, the test for $\zeta = 0$ is a null experiment.

3.1 Predictions and Errors.

Let us use the variable z , which we define in each event as the difference between t_a and t_b measured in units of the K_S average lifetime,

$$z = \Gamma_S(t_a - t_b) . \quad (10)$$

One can histogram events where both kaons of the pair decay by the same decay mode as a function of the variable z , integrating over the variable $t_a + t_b$ all events corresponding to the same $t_a - t_b$, i.e. corresponding to the same z . In each bin of width dz , the number of events $\frac{dn}{dz} dz$ is a function of z that can be developed up to second order in z around the point $z = 0$. Because of the symmetry between the kaon on the left and the kaon on the right, the term in z to the first power is zero.

$$\frac{dn}{dz} = C_0 + C_2 z^2 . \quad (11)$$

C_0 and C_2 are two parameters that can be obtained from a fit to the data.

Let N be the number of pairs emitted in the fiducial volume, B_L , B_S the K_L and the K_S branching ratios, Γ_L and Γ_S the K_L and the K_S decay rates, as above. Let us define the constants A and γ :

$$A = NB_L B_S \frac{\Gamma_L}{\Gamma_S} , \quad (12)$$

$$\gamma = \frac{1 + 4 \left(\frac{\Delta m}{\Gamma_S} \right)^2}{8} = 0.239 , \quad (13)$$

using Ref. [7]. In absence of background, quantum mechanics predicts

$$C_0 = 0 , \quad (14)$$

$$C_2 = A\gamma . \quad (15)$$

However, if, because of a violation of quantum mechanics, the interference term is reduced by a factor $1 - \zeta$, and if there is background,

$$C_0 = A\zeta + \frac{dn}{dz}(\text{Background}) . \quad (16)$$

Measuring C_0 permits us to determine the decoherence parameter

$$\zeta = \frac{C_0}{A} . \quad (17)$$

At the limit of large N , the particular form of Eq. (11) permits us to derive special expressions for the error δC_0 on C_0 :

$$\delta C_0 = \left(\frac{C_0 C_2}{\pi^2} \right)^{\frac{1}{4}} \quad \text{if } C_0 > \left(\frac{C_2}{\pi^2} \right)^{\frac{1}{3}} , \quad (18)$$

$$\delta C_0 = \left(\frac{C_2}{\pi^2} \right)^{\frac{1}{3}} \quad \text{if } C_0 < \left(\frac{C_2}{\pi^2} \right)^{\frac{1}{3}} . \quad (19)$$

The proper form to be used depends on the size of C_0 . The appropriate formula is always the one that gives the largest error. In any case, the error $\delta\zeta$ on ζ is given by

$$\delta\zeta = \frac{\delta C_0}{A} . \quad (20)$$

3.2 Performance at Da ϕ ne.

To predict how good the tests could be at Da ϕ ne, we consider an experiment lasting one year with 30% efficiency, (i.e. duration 10^7 s.) at a luminosity $\mathcal{L} = 10^{33}$ cm $^{-2}$. The integrated luminosity is 10 000 pb $^{-1}$. In that time, $N \approx 10^{10}$ neutral-kaon pairs are expected to be produced.

For both kaons decaying into $\pi^+\pi^-$, using the kaon data of Ref. [7], we compute

$$A = 24000 , \quad (21)$$

$$C_2 = 5700 . \quad (22)$$

We assume the background due to misidentified events in the $\pi^+\pi^-$ sample to be no more than what can be tolerated in a measurement of ϵ'/ϵ , i.e. let us say 10^{-3} of the K_L rate into $\pi^+\pi^-$. Another effect, equivalent to a contribution of background, is due to the measurement error on the vertex positions the two kaon decays. That effect generates uncertainties δt_a and δt_b in the determinations of the proper times t_a and t_b , thus δz in z , and it populates the region around $z = 0$. We will assume that the sample of events can be limited to events where the vertex uncertainty is 1 mm in average. Then the uncertainty in z is $\delta z = 0.2$.

$$\frac{dn}{dz}(\text{Background}) = A 10^{-3} + C_2(\delta z)^2 \approx 250 . \quad (23)$$

Assuming that the contribution of these backgrounds is known within 10% and can be subtracted off with that 10% accuracy, we find that the systematic error is about equal to the statistical error, given by Eq. (16) with a

small ζ , Eqs. (18), and (20).

$$\delta\zeta \approx 10^{-3} . \quad (24)$$

It follows that this test would explore possible values of $\zeta_{\pi^+\pi^-}$ as small as one or two 10^{-3} . This value is better than the one given in Ref. [6] in the context of neutron interferometry. Even smaller values of ζ can be detected if the vertex accuracy can be improved.

A similar calculation can be made for the case where both kaons decay into $\ell^- \bar{\nu} \pi^+$ or both into $\ell^+ \nu \pi^-$. Predictions of the performance at DaΦne are less reliable than for the $\pi^+\pi^-$ decays because the background is more difficult to estimate at this point in time. However, assuming that, as for the $\pi^+\pi^-$ -mode, the background is dominated by the effect of the vertex position uncertainty, we get

$$A = 3300 , \quad (25)$$

$$C_2 = 800 , \quad (26)$$

$$C_0 \approx 30 , \quad (27)$$

$$\delta\zeta = 2 \cdot 10^{-3} \quad \text{for } \zeta = 0 , \quad (28)$$

for each of the two modes $\ell^- \bar{\nu} \pi^+$ and $\ell^+ \nu \pi^-$. This value of $\delta\zeta$ would permit exploring values of $\zeta_{\ell^- \bar{\nu} \pi^+}$ and of $\zeta_{\ell^+ \nu \pi^-}$ comparable to the value estimated for the $\pi^+\pi^-$ -mode, i.e. a few 10^{-3} .

3.3 $K_L K_L$ events

Another possible test consists of measuring the number of pairs of K_L in the same event. As obviously shown by Eq. (8), quantum mechanics predicts no such pair in any event. That same prediction can be obtained using Eq. (7) instead, as the result of a 100% destructive interference effect between the process where the ϕ decays into K^0 on the left and \bar{K}^0 on the right and the process where it is the other way around. If the interference term were reduced by a factor $1 - \zeta_{LL}$, there would be some $K_L K_L$ events and their number would be

$$n_{LL} = \frac{N \zeta_{LL}}{4} . \quad (29)$$

To identify two- K_L events in the Kloe detector, the signature can be two particles decaying or interacting at a large distance from the interaction point. Background is going to be generated by $K_L K_S$ or $K_S K_L$ events

where the K_S decays via the two- π^0 mode and the γ 's from π^0 decays get confused with a K_L decay or interaction. It is beyond the scope of this paper to estimate this background. All we can say is that, if the statistical error was the only limitation, one could detect values of ζ_{LL} as low as 10^{-8} .

4 Alternate Theories.

The tests above are designed to detect effects described by alternate theories to quantum mechanics, more specifically by those which do not use hidden variables. The maximum information that can characterize a quantum system is still given by a density matrix ρ . In quantum mechanics, the time evolution of ρ is described by a unitary transformation,

$$\rho(t) = U(t)\rho(0)U^\dagger(t) \quad , \quad (30)$$

from which one can derive a hamiltonian,

$$H(t) = i\frac{dU}{dt}U^\dagger \quad . \quad (31)$$

Eq. (30) conserves the rank of ρ , thus insures that a pure state cannot evolve into a mixed state. That is the property tested by our tests of interferences. In these tests, the initial state is one that quantum mechanics considers pure and the measurements check that, within error, the final density matrix is of rank 1, i.e. has the mathematical property that describes a pure state.

4.1 Completeness.

The test of Sect. 2 involves a one-kaon system, consisting of a pure K_L initially. The density matrix is a two-by-two matrix with only two eigenvectors ψ and ψ' . The time distribution of $\pi^+\pi^-$ decays is of the form of Eq. (4) with

$$c_S = B_S\Gamma_S \left(|K_S^\dagger\psi|^2 + |K_S^\dagger\psi'|^2 \right) \quad , \quad (32)$$

$$c_L = B_L\Gamma_L \left(|K_L^\dagger\psi|^2 + |K_L^\dagger\psi'|^2 \right) \quad , \quad (33)$$

$$c_I = \sqrt{B_S\Gamma_S B_L\Gamma_L} \left(|(K_S^\dagger\psi)(\psi^\dagger K_L) + (K_S^\dagger\psi')(\psi'^\dagger K_L)| \right) \quad . \quad (34)$$

Using Schwarz' inequality, one proves that, to have $c_I = \sqrt{c_S c_L}$, vectors ψ and ψ' have to be collinear. Then the rank of ρ is 1. If the test had infinite

precision and if it was found that ζ_{regen} of Eq. (5) was equal to zero, the density matrix would have been shown to correspond to a pure state. In that sense, the test is complete.

The tests of Sect. 3 involve a two-kaon system, made initially of a pure state of two neutral kaons from ϕ -decay in one direction. The density matrix ρ is a four-by-four matrix, with four eigenvectors. To prove that ρ is of rank 1 (pure state), it is enough to show that the product of ρ by three linearly independent vectors give zero. This is equivalent, experimentally, to demonstrating that decay rates in three different modes corresponding to three amplitudes represented by three linearly independent vectors in the two-kaon Hilbert space are zero. The tests of Sect. 3.2 consist of measuring the rates of decay of the two-kaon system with the kaon on the left decaying into the same thing at the same time as the kaon on the right; and of checking that these rates are zero. There are three of these tests, involving either the $\pi^+\pi^-$, $\ell^-\bar{\nu}\pi^+$, or $\ell^+\nu\pi^-$ decay mode. The test of Sect. 3.3 is a fourth test which measures the probability of the system evolving into two K_L . The vectors corresponding to any three of the four amplitudes measured in these four tests are linearly independent. Therefore the set of any three of these tests is complete.

4.2 Examples of Decoherence Mechanisms.

1) $K_L \rightarrow K_S$ evolution. Processes that imply such an evolution affect the $\pi^+\pi^-$ decay time distribution. Let $\Gamma_{K_L \rightarrow K_S}$ be the rate of incoherent evolution of K_L into K_S . In the experimental setup of Sect. 2, a fraction $\Gamma_{K_L \rightarrow K_S}/\Gamma_S$ of incoherent K_S would be present all the time alongside with the K_L mesons, in equilibrium, i.e. with as many K_S being formed and decaying per unit of time. Each one of these K_S would decay into $\pi^+\pi^-$ at a rate equal to $B_S\Gamma_S$ instead of $B_L\Gamma_L$ for K_L . Thus the coefficient c_L of $e^{-\Gamma_L t}$ in Eq. (4) would be increased, without c_S and c_I being appreciably changed. Thus ζ_{regen} of Eq. (5) is not zero anymore, and one can deduce

$$\Gamma_{K_L \rightarrow K_S} = 2\zeta_{\text{regen}} \frac{B_L\Gamma_L}{B_S} = \frac{\zeta_{\text{regen}}}{10 \mu\text{s}} . \quad (35)$$

The test of Ref. [2] yields

$$\Gamma_{K_L \rightarrow K_S} = (2 \pm 1.5) 10^{-21} \text{ GeV} . \quad (36)$$

Note that $\Gamma_{K_L \rightarrow K_S}$ is equivalent to the parameter γ of Refs. [8] and [9] in the context of their theory.¹

In the test of Sect. 3.2 that relies on the $\pi^+\pi^-$ decay distribution, the incoherent $K_L \rightarrow K_S$ process would act on the K_L of the $K_L K_S$ and $K_S K_L$ systems and produce $K_S K_S$ states after some time. As an average in the sample of events

$$\frac{dn}{dz} = \frac{N}{2} B_S^2 \frac{\Gamma_{K_L \rightarrow K_S}}{\Gamma_S} . \quad (37)$$

$\Gamma_{K_L \rightarrow K_S}$ can be derived from $\zeta_{\pi^+\pi^-}$ by the same relation as Eq. (35) for ζ_{regen} . At DaΦne, we can hope to get an upper limit of the order of 1/10 ms, i.e. 10^{-22} GeV.²

2) $K_S \rightarrow K_L$ evolution. Such process would produce $K_L K_L$ events in the test of Sect. 3.3.

$$n_{LL} = N \frac{\Gamma_{K_S \rightarrow K_L}}{\Gamma_S} , \quad (38)$$

$$\Gamma_{K_S \rightarrow K_L} = \frac{\zeta_{LL} \Gamma_S}{4} . \quad (39)$$

Suppose that the uncertainty in the background of the test of Sect. 3.3 can be as low as 10^{-6} , then values of $\Gamma_{K_S \rightarrow K_L}$ as low as 1/100 μs or 10^{-20} GeV could be detected.

3) Evolution $K_L K_S - K_S K_L$ into $K_L K_S + K_S K_L$. This evolution can be expressed also as

$$(K^0 \bar{K}^0 - \bar{K}^0 K^0) \rightarrow (K^0 K^0 - \bar{K}^0 \bar{K}^0) . \quad (40)$$

It will produce states where both kaons can decay into $\ell^- \bar{\nu} \pi^+$ or both into $\ell^+ \nu \pi^-$ at the same time. Let us call $\Gamma_{(-) \rightarrow (+)}$ the rate of this process. For the tests of Sect. 3.2 involving $\ell^- \bar{\nu} \pi^+$ or $\ell^+ \nu \pi^-$,

$$\frac{dn}{dz} = 2 N B_S^2 \frac{\Gamma_{(-) \rightarrow (+)}}{\Gamma_S} \quad (41)$$

$$\Gamma_{(-) \rightarrow (+)} = \zeta_{\ell^- \bar{\nu} \pi^+} \frac{\Gamma_S}{2} = 2 \zeta_{\ell^+ \nu \pi^-} \frac{\Gamma_S}{2} . \quad (42)$$

¹The error on $\Gamma_{K_L \rightarrow K_S}$ given by the measurement of Ref. [2] is twice as small as the one quoted in Refs. [8] and [9] because, in those references, only the CPT -violating effect of that parameter was taken into consideration. Another advantage of the test of Ref. [2] is that one can be sure that no constraint derived from quantum coherence was assumed anywhere in the analysis of the events.

²This is not yet comparable to the result obtained in neutron interferometry, 10^{-25} GeV, [6], but it would be the best result relevant to kaon physics.

Using the estimate of Eq. (28), we could explore values of $\Gamma_{(-)\rightarrow(+)}$ lower than $1/0.1$ ms, i.e. 10^{-17} GeV.

If a violation of quantum mechanics is ever revealed by one of the tests above, comparing the amount of violation found in each test will bring information about the decoherence mechanism responsible for the violation. Also questions will be raised about the background due to $K_S K_S$ and $K_L K_L$ events produced by ϕ decay into $K^0 \bar{K}^0 \gamma$. However, the effect of this background can be distinguished from the decoherence effects described in this section if we separate the sample of events into kaon decays near the intersection point and kaon decays far from it. The $K^0 \bar{K}^0 \gamma$ background is present immediately, starting at the time of the ϕ decay, while the effect of the decoherence mechanisms take time to build up. In case of a decoherence effect, the two samples will yield a different ζ . They should not in the case of $K^0 \bar{K}^0 \gamma$ background.

5 Conclusions.

Significant tests of quantum mechanics that can be performed at DaΦne are proposed. They consist of measuring the amplitude of interference effects and comparing it to the predicted value. They may reach the relative accuracy of 10^{-3} . The tests of Sect. 3.2 and, maybe, the one of Sect. 3.3 can be conducted with the KLOE detector as is. They only require a specific analysis of data that will be collected anyhow. These tests of Sect. 3.2 are valid regardless of the Weisskopf-Wigner approximation, [10], because the probability for the two kaons to decay at the same time into the same mode can be shown to be zero just regardless of the time evolution operator $U(t)$, [11].

Tests of quantum mechanics in kaon physics were proposed already in Ref. [1]. One of the motivations was to look for possible spontaneous wave function collapses not involving an observer, because such effects would induce decoherence. That motivation is still valid. Differential equations for the time evolution of the density matrix in presence of a decoherence mechanism were developed in Ref. [12]. In Ref. [13], that kind of evolution was considered relevant to particles in the vicinity of black holes, therefore to string theory. Guesses at how strong this effect could be in kaon physics are given in Ref. [14]. A more detailed analysis is made in Ref. [8] and expression relevant to two-kaon states are given in Ref. [9]. The latest theory assumes strangeness conservation in the decoherence mechanism and does not require “completeness”. However, it is our opinion that it is better to

keep as broad a perspective as possible and to perform all the tests proposed here, which test all the above theories, rather than only those that have the best chance to reveal a violation of quantum mechanics in the context of the theory of Refs. [8], [9], and [14].

Acknowledgments The author is indebted to N.E. Mavromatos for several very useful discussions in Frascati in April 1994.

References

- [1] P.H. Eberhard *Should Unitarity Be Tested Experimentally ?* CERN-72-1 (1972) (unpublished)
- [2] W.C. Carithers et al., *Phys. Rev.* **14 D** (1976) 290 .
- [3] R. Baldini and A. Michetti, private communication, (1993) .
- [4] *Proceeding of the Workshop on Physics and Detectors for Daφne*, G. Pancheri, ed., Frascati, It, (1991) 737 ;
- [5] Particle Data Group, *Review of Particle Properties*, *Phys. Rev. D* **45** (1992) p. II.7 .
- [6] A. Zeilinger, M. A. Horne, and C. G. Shull, *Proc. Int. Symp. Foundations of Quantum Mechanics*, Tokyo (1983) 389 .
- [7] Particle Data Group, *Review of Particle Properties*, *Phys. Rev. D* **45** (1992) p. II.10 .
- [8] J. Ellis, N.E. Mavropoulos, and D.V. Nanopoulos, *Phys. Lett. B* 267 (1991) 465, *Phys. Lett. B* **293**, (1992) 142 , CERN-TH. 6897/93 .
- [9] P. Huet and M. Peskin, SLAC-PUB-6454 (1994), SLAC preprint, Stanford, CA, 94309 .
- [10] See T.D. Lee, R. Ohme, and C.N. Yang, *Phys. Rev.* **106** (1957) 340 .
- [11] Discussions with L.A. Khalfin and N.E Mavromatos at the Meeting of the Eurodaφne Collaboration, Frascati, Italy, (April 1994).
- [12] M.S. Marinov, *Zh. Eksp. Teor. Fiz Pis'ma Red.* **15** (1972) 671 , *JETP Lett.* **15** (1972) 677 .
- [13] S. Hawking, *Comm. Math. Phys.* **43** (1975) 199 , J. Bekenstein, *Phys. Rev. D* **12** (1975) 3077 .
- [14] J. Ellis et al, *Nucl. Phys. B* **241** (1984) 395 .

LAWRENCE BERKELEY NATIONAL LABORATORY
UNIVERSITY OF CALIFORNIA
TECHNICAL & ELECTRONIC INFORMATION DEPARTMENT
BERKELEY, CALIFORNIA 94720