

UCSF

UC San Francisco Previously Published Works

Title

Surgical Debridement for Acute and Chronic Osteomyelitis in Children.

Permalink

<https://escholarship.org/uc/item/0kx104c6>

Journal

JBJS Essential Surgical Techniques, 13(1)

ISSN

2160-2204

Authors

Ekunseitan, Ernest

Sabatini, Coleen

Swarup, Ishaan

Publication Date

2023

DOI

10.2106/JBJS.ST.21.00039

Peer reviewed

KEY PROCEDURES

SURGICAL DEBRIDEMENT FOR ACUTE AND CHRONIC OSTEOMYELITIS IN CHILDREN

Ernest Ekunseitan, BS, Coleen S. Sabatini, MD, MPH, Ishaan Swarup, MD

Published outcomes of this procedure can be found at: *J Pediatr Orthop*. 1996 Mar-Apr; 16(2):220-3, *Afr J Paediatr Surg*. 2014 Oct-Dec;11(4):297-303, and *J Orthop Surg* (Hong Kong). 2016 Aug;24(2):250-2.

Investigation performed at UCSF Benioff Children's Hospital Oakland, Oakland, California

COPYRIGHT © 2023 BY THE JOURNAL OF BONE AND JOINT SURGERY, INCORPORATED

Step 5: Irrigation



Click the arrow above or go to surgicaltechniques.jbjs.org to view the video article described in this summary.

Abstract

Background: Osteomyelitis is an infection of the bone that commonly occurs in pediatric populations. First-line treatment most often involves a course of antibiotics. In recent studies, surgical debridement, in addition to antibiotics, has been shown to provide positive clinical and functional outcomes in children. Debridement is most often indicated in patients with an abscess or in those who do not respond to empiric antibiotic therapy; however, there are limited video resources describing this technique in pediatric patients.

Description: The key steps of the procedure, which are demonstrated in the present video article, are (1) preoperative planning, (2) positioning, (3) subperiosteal exposure and debridement, (4) cortical window creation, (5) irrigation, (6) adjunctive treatment, (7) drain placement, (8) wound closure, (9) dressing and immobilization, and (10) wound check and drain removal.

Alternatives: Nonoperative treatment is usually indicated for acute osteomyelitis in which patients present with little to no necrotic tissue or abscess formation. In these cases, a course of broad-spectrum antibiotics may be sufficient for a cure.

Rationale: This procedure allows for the removal of necrotic bone and soft tissue, thus facilitating the recovery process. It also allows for the retrieval of tissue samples that may be used to guide selection of the appropriate antibiotic therapy. Surgical debridement is a safe and reliable technique that has been associated with positive long-term outcomes.

Expected Outcomes: We expect that some patients will require repeat surgical debridement procedures to decrease pathogen burden and prevent future complications. However, we expect that the majority of patients who undergo surgical debridement for uncomplicated osteomyelitis will recover full functionality of the affected limb with no associated long-term sequelae¹⁰.

Important Tips:

- Understand preoperative imaging to identify areas of infection, localize critical structures and the physis, and plan surgical approaches.
- Use extensile approaches and preserve vascularity during the approach.
- Perform subperiosteal dissection and create a cortical window to debride areas of infection, but avoid excessive periosteal stripping.

Disclosure: The Disclosure of Potential Conflicts of Interest forms are provided with the online version of the article (<http://links.lww.com/JBJSST/A404>).

- Close the dead space and wound in a layered manner.

Acronyms and Abbreviations:

- MRI = magnetic resonance imaging
- K-wire = Kirschner wire
- MRSA = methicillin-resistant *Staphylococcus aureus*
- PDS = polydioxanone

Ernest Ekunseitan, BS¹

Coleen S. Sabatini, MD, MPH¹

Ishaan Swarup, MD¹

¹Division of Pediatric Orthopaedic Surgery, UCSF Benioff Children's Hospital Oakland, Oakland, California

Email for corresponding author: Ishaan.Swarup@ucsf.edu

References

1. Hamdy RC, Lawton L, Carey T, Wiley J, Marton D. Subacute hematogenous osteomyelitis: are biopsy and surgery always indicated? *J Pediatr Orthop*. 1996 Mar-Apr;16(2):220-3.
2. Wirbel R, Hermans K. Surgical treatment of chronic osteomyelitis in children admitted from developing countries. *Afr J Paediatr Surg*. 2014 Oct-Dec;11(4):297-303.
3. Chou AC, Mahadev A. Acute bacterial osteomyelitis in children. *J Orthop Surg (Hong Kong)*. 2016 Aug;24(2):250-2.
4. Jaramillo D, Dormans JP, Delgado J, Laor T, St Geme JW 3rd. Hematogenous Osteomyelitis in Infants and Children: Imaging of a Changing Disease. *Radiology*. 2017 Jun;283(3):629-43.
5. Calhoun JH, Manring MM, Shirliff M. Osteomyelitis of the long bones. *Semin Plast Surg*. 2009 May;23(2):59-72.
6. Tröbs R, Möritz R, Bühligen U, Bennek J, Handrick W, Hörmann D, Meier T. Changing pattern of osteomyelitis in infants and children. *Pediatr Surg Int*. 1999 Jul;15(5-6):363-72.
7. Canavese F, Corradin M, Khan A, Mansour M, Rousset M, Samba A. Successful treatment of chronic osteomyelitis in children with debridement, antibiotic-laden cement spacer and bone graft substitute. *Eur J Orthop Surg Traumatol*. 2017 Feb;27(2):221-8.
8. Gogia JS, Meehan JP, Di Cesare PE, Jamali AA. Local antibiotic therapy in osteomyelitis. *Semin Plast Surg*. 2009 May;23(2):100-7.
9. Murphy RF, Plumblee L, Barfield WB, Murphy JS, Fuerstenau N, Spence DD, Kelly DM, Dow MA, Mooney JF 3rd. Septic Arthritis of the Hip-Risk Factors Associated With Secondary Surgery. *J Am Acad Orthop Surg*. 2019 May 1;27(9):321-6.
10. Gornitzky AL, Kim AE, O'Donnell JM, Swarup I. Diagnosis and Management of Osteomyelitis in Children: A Critical Analysis Review. *JBJS Rev*. 2020 Jun;8(6):e1900202.
11. Vander Have KL, Karmazyn B, Verma M, Caird MS, Hensinger RN, Farley FA, Lubicky JP. Community-associated methicillin-resistant *Staphylococcus aureus* in acute musculoskeletal infection in children: a game changer. *J Pediatr Orthop*. 2009 Dec;29(8):927-31.