

UCLA

UCLA Previously Published Works

Title

Stress, Psychological Resources, and HPA and Inflammatory Reactivity During Late Adolescence

Permalink

<https://escholarship.org/uc/item/0m08h434>

Journal

Development and Psychopathology, 31(2)

ISSN

0954-5794

Authors

Chiang, Jessica J
Ko, Ahra
Bower, Julianne E
[et al.](#)

Publication Date

2019-05-01

DOI

10.1017/s0954579418000287

Peer reviewed



Published in final edited form as:

Dev Psychopathol. 2019 May ; 31(2): 699–712. doi:10.1017/S0954579418000287.

Stress, Psychological Resources, and HPA and Inflammatory Reactivity During Late Adolescence

Jessica J. Chiang, Ph.D.^a, Ahra Ko, M.A.^b, Julienne E. Bower, Ph.D.^{c,d,e}, Shelley E. Taylor, Ph.D.^c, Michael R. Irwin, M.D.^{c,d,e}, and Andrew J. Fuligni, Ph.D.^{c,d,e}

^aInstitute for Policy Research, Northwestern University

^bLos Angeles County Department for Mental Health

^cDepartment of Psychology, University of California, Los Angeles

^dDepartment of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles

^eCousins Center for Psychoneuroimmunology, University of California, Los Angeles

Abstract

Psychosocial stress during childhood and adolescence is associated with alterations in the hypothalamic-pituitary-adrenal (HPA) axis and with heightened inflammation, both of which are implicated in poor health. However, factors that may protect against these effects relatively early in life are not well understood. Thus, we examined whether psychosocial resources protect against stress-related alterations in the HPA-axis and heightened inflammation in a sample of 91 late adolescents. Participants completed measures of various stressors (major life events, daily interpersonal stress, early adversity), and psychosocial resources (mastery, optimism, self-esteem, and positive reappraisal). They also completed the Trier Social Stress Test and provided saliva and blood samples for the assessment of cortisol and IL-6 reactivity. Each of the stressors was associated with lower cortisol reactivity. Additionally, associations with major life events and daily stress were moderated by psychological resources, such that more life events and daily stress were associated with decreased HPA reactivity among adolescents with lower levels of psychological resources, but not among those with higher levels of psychological resources. This pattern of findings was observed only for cortisol reactivity and not for IL-6 reactivity. Findings suggest that psychological resources may counteract the effects of certain adversity-related decreases in cortisol reactivity.

Keywords

major life events; daily stress; early adversity; cortisol; IL-6

Psychosocial stress during childhood and adolescence can have immediate and long-term impacts on both mental and physical health, increasing risk for numerous conditions that

Address correspondence to Jessica J. Chiang, Ph.D., Northwestern University, 1801 Maple Avenue, Suite 2450, Evanston, IL 60201. jessica.chiang@northwestern.edu.

Conflicts of interest: none

range from mood and anxiety disorders to upper respiratory infections to premature mortality (Brown et al., 2009; Cohen, Doyle, Turner, Alper, & Skoner, 2004; Demakakos, Pillas, Marmot, & Steptoe, 2016; Miller, Chen, & Parker, 2011; Repetti, Taylor, & Seeman, 2002; Syed & Nemeroff, 2017). These associations are thought to be driven in part by gradual changes in the stress response system, specifically in the hypothalamic-pituitary-adrenal (HPA) axis and in inflammatory processes (Ehrlich, Miller, & Chen, 2016; Heim & Binder, 2012; Slavich & Irwin, 2014). Stress activates the HPA axis, resulting in secretion of the hormone cortisol, and increases inflammation. These biological responses to stress promote short-term adaptation to challenging circumstances. However, repeated or chronic activation of these responses can interfere with later HPA axis functioning and inflammatory processes in ways that compromise long-term health (McEwen & Seeman, 1999).

A substantial body of work has linked various childhood and adolescent stressors, including socioeconomic disadvantage, maltreatment, harsh parenting, and parental separation or loss, to alterations in the HPA axis and to heightened inflammation. Outcomes of HPA axis functioning include the components of the diurnal rhythm of cortisol (i.e., cortisol awakening response, diurnal slope, and waking and bedtime cortisol levels) and responses to acute psychosocial stress. Previous studies have linked childhood and adolescent stress to both heightened and dampened levels of cortisol, which may be due to differences in stressor characteristics and HPA-axis measurement (Chiang, Taylor, & Bower, 2015; Repetti, Robles, & Reynolds, 2011). Seemingly discrepant findings may also reflect the notion that because excessive exposure to glucocorticoids can be detrimental, hypocortisolism may emerge after sustained periods of hypercortisolism as a counter-regulatory mechanism (Fries, Hesse, Hellhammer, & Hellhammer, 2005; Miller, Chen, & Zhou, 2007). Indeed, with respect to HPA responses to acute stress, past studies have consistently linked chronic or repeated stress to dampened cortisol responses (Bunea, Szentágotai-Tóth, & Miu, 2017; Cserneci, Crișan, Vulturar, Opre, & Miu, 2015; Carpenter et al., 2007; Carpenter, Shattuck, Tyrka, Geraciotti, & Price, 2011; Elzinga et al., 2008; Engert et al., 2010; Janusek, Tell, Gaylord-Harden, & Mathews, 2017; Kraft & Luecken, 2009; Lovallo, 2013; MacMillan et al., 2009), a pattern that has also been observed in individuals with post-traumatic stress disorder, chronic fatigue syndrome, and fibromyalgia (Heim, Ehlert, & Hellhammer, 2000). Dampened cortisol reactivity to stress has also been associated with poorer lung function, decreased anti-body responses to vaccination, and greater adiposity (Carroll et al., 2012; de Rooij, 2013; Phillips, Carroll, Burns, & Drayson, 2005). In addition to its links to HPA axis alterations, childhood and adolescent stress has also been related to heightened levels of circulating inflammatory markers in adulthood (Baumeister, Akhtar, Ciufolini, Pariante, & Mondelli, 2016; Coelho, Viola, Walss-Bass, Brietzke, & Grassi-Oliveira, 2014). Studies of youth have yielded more mixed results, suggesting potential delayed effects on systemic inflammation (Slopen, Koenen, & Kubzansky, 2012; Slopen, Kubzansky, McLaughlin, & Koenen, 2013). Importantly, heightened inflammation is implicated in numerous age-related chronic conditions, including heart disease and certain cancers (Brydon & Steptoe, 2005; Choy, 2012; Elinav et al., 2013; Libby, 2006; Wellen & Hotamisligil, 2005).

Although much research has linked childhood and adolescent stress to HPA alterations and heightened inflammation, factors that protect against these are not completely understood. Previous research has pointed to warm and supportive relationships as important protective

factors (Brody, Yu, & Beach, 2016; Carroll et al., 2013; Chen, Miller, Kobor, & Cole, 2011), but less is known about individual psychological characteristics that may similarly counteract the biological effects of early stress. Theoretical models purport several individual psychological characteristics, collectively known as psychological resources, that promote resilience in the face of adversity; these include constructs such as mastery or a sense of control over one's life, purpose in life, positive affect, optimism, self-esteem, and the ability to positively reappraise negative emotions and situations (Chen & Miller, 2013; Gallo, de los Monteros, & Shivpuri, 2009; Taylor & Seeman, 1999). These individual characteristics have independently and together been associated with lower risk for clinical health outcomes such as mortality and cardiovascular disease (Carver, Scheier, & Segerstrom, 2010; Lundgren, Garvin, Jonasson, Andersson, & Kristenson, 2015; Stamatakis et al., 2004; Surtees, Wainwright, Luben, Khaw, & Day, 2006; Taylor & Broffman, 2011). They have also been shown to buffer against chronic stress-related increases in inflammation and delayed blood pressure recovery from subsequent acute stress (Boylan, Jennings, & Matthews, 2016; Elliot & Chapman, 2016).

The majority of these studies have focused on adults, raising the question of whether psychological resources can similarly buffer against the adverse effects of chronic or repeated stress during earlier life stages. Adolescence may be an especially important focal point for several reasons. First, the adolescent years are marked by an increase in stress. Demands across domains (school, peers, and family) are known to increase during this time, and adolescents have reported similar or higher levels of psychosocial stress than adults (American Psychological Association, 2014). Second, in addition to facing increasing stress, research suggests that sensitivity to the environment increases during adolescence, potentially rendering adolescents more vulnerable to the effects of stress (Blakemore & Mills, 2014; Somerville, 2013; Spear, 2009). Evidence for this notion comes from studies showing that adolescents exhibit greater negative affective and neurobiological responses to social threat compared to children and adults (Gunnar, Wewerka, Frenn, Long, & Griggs, 2009; Sebastian, Viding, Williams, & Blakemore, 2010; Stroud et al., 2009). Lastly, the foundations of future adult health may be set during adolescence. The idea here is that with rapid neurobiological and psychosocial development that facilitates the transition into an independent, productive adult, adolescent experiences can not only directly shape biological patterns of functioning, but also modify health trajectories initiated in childhood and contribute to the adoption of long-term health behaviors (Del Giudice, Ellis, & Shirtcliff, 2011; Tottenham & Galván, 2016; Viner et al., 2012).

Past studies examining whether psychological resources buffer against the negative health effects of chronic stress during adolescence have found that optimism in combination with meaning in life attenuated the association between socioeconomic disadvantage and obesity, cardio-metabolic risk, and inflammatory processes (Chen, Lee, Cavey, & Ho, 2013; Chen, McLean, & Miller, 2015; Kallem et al., 2013). However, such studies are few, and buffering effects on the HPA axis have not been examined. Yet, as described above, the HPA axis is thought to be an important pathway linking early stress to poor health. Additionally, past studies have focused on only a single form of adversity, namely socioeconomic disadvantage. However, youth encounter a variety of stressors in their lives. Furthermore, more mundane and less severe stressors occurring in everyday life (e.g., conflict, work

deadlines), in addition to other various chronic stressors (e.g., childhood maltreatment, major life events), have been linked to poor health outcomes and to compromised biological functioning (Chiang, Eisenberger, Seeman, & Taylor, 2012; DeLongis, Folkman, & Lazarus, 1988; Dietz et al., 2013; Luecken & Appelhans, 2006; Miller et al., 2011; Repetti et al., 2011; Stawski, Cichy, Piazza, & Almeida, 2013; Stone et al., 1994). Whether psychological resources protect against the health-related effects of these other stressors is not entirely clear.

The overall purpose of the present study was to examine whether various stressors were associated with HPA axis alterations and heightened inflammation during late adolescence, and whether psychological resources (mastery, self-esteem, optimism, and the tendency to positively reappraise negative situations) mitigated these associations. We focused on cortisol and interleukin (IL)-6 reactivity to a standard laboratory stress task as indices of HPA axis functioning and inflammatory processes, respectively. We examined several types of stressors, including major life events in the past year, daily negative social interactions across a 15-day period, and the early psychosocial family climate. Given that prior work has tied each of these stressors to health outcomes and biological functioning in both youth and adults (Carroll et al., 2013; Chiang et al., 2016; Fuligni et al., 2009; Kershaw et al., 2014; Loucks, Almeida, Taylor, & Matthews, 2011; Miller & Chen, 2010; Pyykkönen et al., 2010; Stroud, Chen, Doane, & Granger, 2016), we did not hypothesize any differential effects according to stressor type. Instead, we hypothesized that all three types of stress would be associated with sub-optimal biological profiles, as evidenced by dampened cortisol and heightened IL-6 responses to acute stress. We further hypothesized significant interactions between each of the three types of stress and psychological resources, such that those reporting greater levels of stress and lower levels of psychological resources would exhibit reduced cortisol and elevated IL-6 reactivity. These associations were expected to be attenuated among those who reported stress but had higher levels of psychological resources. Given that our sample included Latino and European-American adolescents, exploratory analyses examined whether there were any ethnic differences in stress associations with cortisol and IL-6 responses and in the potential buffering role of psychological resources. Socioeconomic status, family values, and rates of racial discrimination are known to differ between these two ethnic groups (Fuligni, Tseng, & Lam, 1999; Greene, Way, & Pahl, 2006; Sabogal, Marín, Otero-Sabogal, Marín, & Perez-Stable, 1987; Semega, Frontenot, Kollar, & Bureau, 2017), which may translate to differential effects of stress and its interactions with psychological resources on biological responsivity.

Method

Participants

Participants were 18 to 20-year-old adolescents ($n = 91$; $M_{age} = 18.37$, $SD = .51$; 57% female) drawn from a larger sample of 248 adolescents ($M_{age} = 18.31$, $SD = .77$) partaking in the second wave of a three-wave longitudinal study on the psychosocial contributions to the development of early health risk from mid-adolescence to early adulthood. During the first wave of the larger parent study, 316 tenth and eleventh graders ($M_{age} = 16.40$, $SD = .74$) were recruited, 214 of whom participated in the second wave two years later. During the

second wave, 34 new participants were added to refresh the sample. Two years after the second wave, 180 participants ($M_{age} = 20.29$, $SD = .74$) from Wave 1 and/or Wave 2 provided data during the third and final wave.

Data collection of the larger study involved a home visit during which participants completed a series of questionnaires and a 15-day daily diary protocol. After completing the Wave 2 protocol, participants were pre-screened to ensure that they identified as Latino or European-American (other ethnicities were excluded due to insufficient numbers in the larger study) and were at least 18 years of age. Pre-screening was based on responses to demographic questionnaires collected in the larger parent study. Eligible individuals were contacted via telephone, and those who responded ($n = 169$) were given information on the current experimental study and invited to participate. Individuals expressing interest were scheduled for a laboratory visit and instructed to refrain from eating or drinking anything (except water) the hour prior to the visit. Ninety-one participants, 79 of whom were from the original Wave 1 sample and 12 of whom were from the pool of newly added participants at Wave 2, completed the laboratory visit an average of 5.5 ($SD = 2.7$, range = 1.4 to 12.2) months after they completed Wave 2 of the parent study. Participants in the present study did not differ from the larger Wave 2 sample on age, gender, parent education, and ethnicity ($p > .156$). They also did not differ on major life events and daily interpersonal stress ($p > .173$), which were assessed as part of the larger parent study.

The subsample in the current study were from European-American (37.36%) and Latino backgrounds (62.64%). Primary caregivers reported on annual household income as well as their own and spouse's highest level of education on an 11-point scale ($1 = \text{some elementary school}$, $11 = \text{graduated from medical law, or graduate school}$). Education across parents was averaged. Across the study sample, median household income was \$79,000 (range = \$11,000 – \$350,000). Approximately 14.44% of participants had parents with less than a high school diploma, 7.78% had at least one parent with a high school diploma, 38.89% had at least one parent who completed vocational trade school or some college, 23.33% had at least one parent with a college degree, and 15.57% had at least one parent who completed some or all of graduate or professional school.

Procedure

Participants arrived at the University of California, Los Angeles (UCLA) Clinical and Translational Research Center (CTRC) and provided informed written consent and anthropometric measures. A nurse then inserted an indwelling intravenous catheter in the antecubital vein of the nondominant arm, after which participants viewed a neutral-content video for 20 minutes to facilitate acclimation to the testing environment. The first saliva and blood samples were collected after this baseline period. Participants then completed the Trier Social Stress Task (TSST), a well-established acute laboratory-based social stressor that activates the HPA axis and increases inflammation (Kirschbaum, Pirke, & Hellhammer, 1993; Steptoe, Hamer, & Chida, 2007). Participants delivered a 5-minute impromptu speech on why they were the best candidates for their ideal jobs in front of an evaluative panel of expert interviewers consisting of research assistants trained to provide nonverbal negative feedback. Participants were given five minutes to prepare for their speech. Immediately after

delivering their speech, participants performed a five-minute mental arithmetic task involving subtracting by 13's from 2,935 while the evaluative panel urged them to go more quickly and perform more accurately. After the TSST, participants provided five additional saliva samples (immediately and 30, 45, 60, and 75 minutes post-TSST onset) and three additional blood samples (30, 60, and 90 minutes post-TSST onset). After the final blood sample was collected, participants were debriefed and compensated. All visits were conducted between the hours of 12pm and 6pm, and the UCLA Institutional Review Board approved all procedures.

Measures

Major life events—During the second wave visit of the parent study, participants completed a major life event checklist. They reported on whether they had experienced any of 21 events across the domains of family, friends, and school within the past 12 months. Items were adapted from previously used measures of stressful events that have been linked to negative outcomes (Conger et al., 2002; Hammen, 1991). Example items included parents divorced or separated, family member became seriously ill, a close friend moved quite far away, you had a serious falling out or ended a friendship with a close friend, you were suspended or expelled in school, and your grades in school went down a lot. Affirmative responses were summed across items.

Daily interpersonal stress—Participants also completed a daily diary protocol during the second wave visit of the parent study. Each night for 15 consecutive evenings, participants reported on whether they had experienced any of eight negative social interactions across the contexts of family, peers, and school. Items were argued with a parent, argued with another family member, argued with a friend, punished by a parent, parents argued, something bad happened to a family member, had an argument or was punished by an adult at school, and was insulted, threatened, or made fun of by someone at school. These items were selected because they have previously been shown to be stressful for adolescents (Chung, Flook, & Fuligni, 2009; Fuligni et al., 2009; Nishina & Juvonen, 2005). Consistent with previous research on daily stress (Chiang et al., 2016; Grzywacz, Almeida, & McDonald, 2002; Sin, Graham-Engeland, Ong, & Almeida, 2015), affirmative responses were summed and recoded as 0 or 1 for each day to indicate whether any one of the stressors occurred that day. The average of recoded scores across days was subsequently computed to indicate the proportion of days that at least one stressor occurred.

Early adversity—The Risky Families questionnaire (Taylor, Lerner, Sage, Lehman, & Seeman, 2004) was used to measure early adversity and was administered during the laboratory visit. Specifically, the Risky Families questionnaire assesses the family psychosocial climate in which participants grew up when they were 5 to 15 years old. On a 5-point scale (*1=not at all to 5=very often*), participants indicated how frequently they experienced conflict, violence, corporal punishment, parental affection, neglect, and chaos. The Risky Families questionnaire has been widely used in previous research and aligns with clinical interviews assessing early life stress. Items framed positively were reverse-coded, and scores across items were averaged. The scale demonstrated good internal reliability in the present sample ($\alpha = .84$).

HPA reactivity—Six saliva samples were collected using oral swabs (Salimetrics) throughout the laboratory visit. They were collected immediately after the neutral-content video (baseline) and TSST, and 30, 45, 60, and 75 minutes after TSST onset. Samples were stored at -80°C until transported on ice to the Laboratory of Biological Psychology at the Technical University of Dresden, Germany where they were assayed for cortisol in duplicate using high-sensitivity chemiluminescence-immunoassays (IBL, International, Hamburg, Germany). Intra- and inter-assay coefficients of variations were below 10%. Outlier screening revealed a total of 10 samples across time points that had cortisol values greater than three standard deviations (SDs) from their respective means. These values were Winsorized and replaced with the value at three SDs. Cortisol values were then natural log-transformed to correct for their skewed distributions. We previously reported that the TSST activated the HPA axis, as evidenced by a significant increase in cortisol from baseline to 30 and 45 minutes post TSST (Chiang, Bower, Irwin, Taylor, & Fuligni, 2017).

Inflammatory reactivity—Four blood samples of 6mL were collected at baseline, and 30, 60, and 90 minutes after TSST onset into EDTA lavender-top tubes. Samples were placed on ice immediately after collection, after which they were centrifuged, aliquoted into plasma samples, and stored at -80°C . After data collection was completed, samples were assayed in duplicate for interleukin (IL)-6 using the Quantikine high sensitivity human IL-6 ELISA kits (R&D Systems, Inc., Minneapolis, MN) by the UCLA Cousins Center Inflammatory Biology Core. Intra- and inter-assay coefficients of variability were below 7%. Outlier screening revealed a total of 11 samples across time points that had IL-6 values greater than 3 SDs from their respective means. These values were Winsorized and replaced with the value at three SDs. All values were natural log-transformed given the skewed distribution. As reported previously (Chiang et al., 2017), the TSST was effective in eliciting an inflammatory response, with participants exhibiting a 34% average increase in IL-6 from baseline to 90 minutes after TSST onset.

Psychological resources—Measures of psychological resources were completed during the laboratory visit and assessed perceived control, optimism, self-esteem, and positive reappraisal. Perceived control was assessed using the 7-item Pearlin Mastery Scale (Pearlin & Schooler, 1978). Participants indicated the extent to which they agreed with statements such as: “There is little I can do to change many of the important things in my life (reverse coded),” and “What happens to me in the future mostly depends on me,” using a 4-point scale (*1=strongly disagree 4=strongly agree*). In the current study, internal consistency was adequate ($\alpha = .74$).

Dispositional optimism was assessed using the six-item Life Orientation Test (LOT-R) (Scheier, Carver, & Bridges, 1994). Using a 5-point scale (*0=strongly disagree, 4=strongly agree*), participants rated the degree to which they agreed with each item. Example items include “In uncertain times, I usually expect the best” and “Overall, I expect more good things to happen to me than bad.” This scale demonstrated good internal reliability in the present study ($\alpha = .83$).

Self-esteem was assessed using the ten-item Rosenberg Self-Esteem Scale (Rosenberg, 1965). Participants rated items such as, “I feel that I have a number of good qualities” and

“On the whole, I am satisfied with myself,” on a 4-point scale (0 =strongly agree, 3 =strongly disagree). Internal consistency of this scale was high ($\alpha = .93$).

Positive reappraisal was assessed using the cognitive reappraisal subscale from the Emotion Regulation Questionnaire (ERQ) (Gross & John, 2003). Using a 7-point scale (1 = not at all, 7 = a lot), participants indicated the extent to which they agreed with statements such as “To feel less negative emotion, I change what I think about” and “When I’m faced with a stressful situation, I make myself think about it in a way that helps me stay calm.” The ERQ has been well-validated and internal consistency in the present sample was high ($\alpha = .88$).

Negatively worded items on each scale were reverse-coded, such that higher scores indicated greater mastery, optimism, self-esteem, and positive reappraisal. Items were then summed for each scale. Principal-components analysis indicated that a single component explained 61.2% of the variance. Factor loadings were .83 for mastery, .83 for optimism, .73 for self-esteem, and .72 for reappraisal. Thus, we averaged standardized scores of the four scales to create a composite score indexing psychological resources.

Potential covariates—Sociodemographic characteristics (age, gender, ethnicity, socioeconomic status), health behaviors (exercise, caffeine consumption, cigarette smoking), adiposity (waist circumference), use of oral contraceptives, and time of day were considered as potential covariates given that these factors have been previously related to levels of cortisol and/or IL-6 (DeSantis et al., 2007; Dowd, Simanek, & Aiello, 2009; Jacks, Sowash, Anning, Mcgloughlin, & Andres, 2002; Lovallo et al., 2005; O’Connor et al., 2009; Uhart, Chong, Oswald, Lin, & Wand, 2006). During the second wave visit, participants reported their own ethnicity and gender. Their parents reported on participants’ date of birth, from which age was computed, and their own and spouse’s educational attainment. Educational attainment was averaged across parents to index family socioeconomic status. Waist circumference was assessed twice at the mid-point between the iliac crest and lower rib and averaged. During the laboratory visit, study experimenters recorded participants’ time of arrival, and participants reported on the number of caffeinated beverages consumed that day, whether they smoked cigarettes during the past week (yes/no), and whether they currently use oral contraceptives (yes/no). Additionally, for three consecutive mornings prior to the lab visit, participants reported on whether they engaged in physical activity the previous day and the total number of days exercised was computed.

Analytic Approach

Selection of covariates—Bivariate correlations among potential covariates and cortisol and IL-6 at each time point were examined to identify the panel of covariates to be included in primary analyses. Gender, parental education, time of day, exercise, smoking, and use of oral contraceptives were significantly or marginally associated with at least one of the cortisol measures across time points ($r_s = -.40$ - $.30$, $p_s < .06$). Thus, these were included as covariates in all models with cortisol as the outcome. For IL-6, age, gender, parent education, ethnicity, and waist circumference showed marginal or significant correlations with at least one IL-6 measure across time points ($r_s = -.21$ - $.47$, $p_s < .06$), and were

therefore included as a covariate in models with IL-6 as the outcome. Models with and without covariates yielded similar results.

Primary analyses—Multi-level modeling was used to conduct analyses given repeated measurements within individuals. Level 1 modeled within-person changes in cortisol and IL-6 as a function of time while Level 2 modeled between-person differences in cortisol and IL-6 responses based on experiences of stress (i.e., major life events, daily interpersonal stress, and early adversity), psychological resources, and their interactions. The intercepts in models represent baseline levels of cortisol and IL-6. Models with cortisol as the outcome included both a linear and a quadratic term for time given that sampling times captured cortisol reactivity as well as recovery (Chiang et al., 2017; Dickerson & Kemeny, 2004). The linear term for time represents the rate of change in cortisol whereas the quadratic term for time represents how the rate of change increases or decreases across time, or in other words, the curvature of cortisol responses. Thus, increased cortisol reactivity is reflected as a positive value for the linear term, and a slowing rate of cortisol increases (i.e., cortisol peaks then declines) is reflected as a negative value for the quadratic term. Models with IL-6 as the outcome only included a linear term for time given that sampling times captured only reactivity to the TSST, as IL-6 requires a longer period of time to return to baseline (Chiang et al., 2017; Marsland, Walsh, Lockwood, & John-Henderson, 2017).

We first tested the main effects of stress and psychological resources on cortisol and IL-6 response trajectories by examining the cross-level interactions between stress and time and between psychological resources and time. As reported below, significant stress by time interactions emerged. These interactions were followed-up with tests of simple slopes in which the effect of time on responses to the TSST were examined among individuals above and below half a standard deviation from the mean of major life events (low: $n = 21$; high: $n = 21$), daily interpersonal stress (low: $n = 29$; high: $n = 30$), and early adversity (low: $n = 32$; high: $n = 22$).

We then tested whether psychological resources protected against the effects of stress by examining the three-way interactions between stress, psychological resources, and time. Significant interactions indicate that biological responses varied as a function of stress at varying levels of psychological resources. Observed interactions were followed-up with tests of simple interactions in which two-way stress by time interactions were examined separately in individuals above ($n = 33$) and below ($n = 23$) half a standard deviation from the mean of psychological resources.

In all analyses, interaction effects were probed at half a standard deviation above and below the mean of the moderator of interest because the number of individuals above and below one standard deviation were much fewer. All continuous predictor and covariate variables were grand mean-centered and separate models were tested for each measure of stress. Because major life events and daily interpersonal stress were assessed at varying times prior to the laboratory visit, we also assessed whether the duration between the second-wave visit of the parent study and the laboratory visit altered results. Results were unchanged when time between parent study and laboratory visit was added as a covariate. Therefore, results from models without this variable are reported.

Missing data—Rates of missing data were low for variables included in analyses.

Variables that had missing data included IL-6 at all time points, parent education, and daily family stress. For IL-6, only 7 out of the 364 (1.9%) samples had missing data. Only one (1.1%) participant had missing data on parent education, and only three (3.3%) participants had missing data on daily interpersonal stress. In all models, full information maximum likelihood estimation was used.

Results

Descriptive statistics are presented in Table 1 and bivariate correlations are displayed in Table 2. Overall, adolescents experienced a few major life events, had negative social interactions on 24% of days (i.e., 3.6 days), and experienced fairly low levels of early adversity. As expected, more major life events, daily interpersonal stress, and early adversity were positively correlated with one another and negatively correlated with psychological resources. Of the different stressors, only greater early adversity was associated with lower cortisol. Psychological resources were also not associated with cortisol. Neither stress nor psychological resources was correlated with IL-6.

HPA Reactivity

Major life events—As shown in Table 3 (column 1), neither major life events ($p = .623$) nor psychological resources ($p = .870$) were associated with baseline levels of cortisol. Cortisol reactivity to the TSST was also not modulated by psychological resources (linear: $p = .297$; quadratic: $p = .201$). However, there was a significant interaction between major life events and time (linear: $p = .005$; quadratic: $p = .006$). Follow-up tests revealed that adolescents who experienced more major life events exhibited smaller cortisol responses (linear: $b(SE) = .07(.06)$, $p = .195$; quadratic: $b(SE) = -.03(.01)$, $p = .006$) relative to those who experienced fewer major life events (linear: $b(SE) = .21(.05)$, $p < .001$; quadratic: $b(SE) = -.05(.01)$, $p < .001$).

The association between major life events and cortisol responses were modulated by psychological resources, as evidenced by the significant interaction between major life events, psychological resources, and time (linear: $p < .001$; quadratic: $p = .001$; Table 3, column 2). At lower levels of psychological resources, cortisol responses varied as a function of major life (linear: $b(SE) = -.07(.02)$, $p = .002$; quadratic: $b(SE) = .01(.004)$, $p = .008$). By contrast, at higher levels of psychological resources, major life events were not associated with cortisol responses (linear: $b(SE) = .01(.02)$, $p = .715$; quadratic: $b(SE) = .0002(.004)$, $p = .968$). Visualization of these simple interaction effects shows that among individuals with low psychological resources (Figure 1a), those with more major life events exhibit smaller cortisol responses relative to their peers who report fewer major life events. However, among those with high psychological resources (Figure 1b), cortisol responses of those with more major life events more closely resemble those with fewer major life events.

Daily interpersonal stress—Paralleling findings for major life events, daily interpersonal stress operationalized as proportion of stressor days ($p = .624$) and psychological resources ($p = .990$) were not related to cortisol levels at baseline (Table 3, column 3). In addition, psychological resources were not associated with cortisol reactivity

to the TSST (linear: $p = .424$; quadratic: $p = .286$). However, there was a significant interaction between daily interpersonal stress and time (linear: $p = .017$; quadratic: $p = .011$). Adolescents who experienced more interpersonal stress in their daily lives had smaller cortisol responses (linear: $b(SE) = .19(.06)$, $p = .001$; quadratic: $b(SE) = -.04(.01)$, $p < .001$) compared to their peers who experienced less interpersonal stress in their everyday lives (linear: $b(SE) = .37(.04)$, $p < .001$; quadratic: $b(SE) = -.07(.01)$, $p < .001$).

Variation in cortisol responses associated with daily interpersonal stress were further modulated by psychological resources, as reflected in the significant three-way interaction between daily interpersonal stress, psychological resources, and time (linear: $p < .015$; quadratic: $p = .070$; Table 3, column 4). Tests of simple interactions indicated that daily interpersonal stress was associated with cortisol responses at lower levels of psychological resources (linear: $b(SE) = -.60(.22)$, $p = .006$; quadratic: $b(SE) = .09(.04)$, $p = .020$), but not at higher levels of psychological resources (linear: $b(SE) = -.07(.25)$, $p = .778$; quadratic: $b(SE) = .04(.04)$, $p = .342$). Visualization of the simple interaction effects revealed a similar pattern as that depicted in Figure 1¹.

Early adversity—Table 3 (column 5) shows that early adversity ($p = .099$) and psychological resources ($p = .396$) were not associated with baseline cortisol. Psychological resources were not associated with rate of change in cortisol (linear: $p = .095$), but was associated with quadratic change ($p = .036$). However, simple slope analyses revealed similar quadratic change at both high and low levels of psychological resources (quadratic: $bs(SEs) = -.06(.01)$, $ps < .001$).

Similar to major life events and daily interpersonal stress, early adversity also modulated cortisol responses to the TSST (linear: $p = .002$; quadratic: $p < .001$). Cortisol responses were smaller among those reporting higher levels of early adversity (linear: $b(SE) = .20(.06)$, $p = .001$; quadratic: $b(SE) = -.04(.01)$, $p < .001$) compared to cortisol responses among those reporting lower levels of early adversity (linear: $b(SE) = .36(.04)$, $p < .001$; quadratic: $b(SE) = -.07(.01)$, $p < .001$).

There was a trend towards a three-way interaction between early adversity, psychological resources, and time (linear: $p = .073$; quadratic: $p = .310$; Table 3, column 6). At low levels of psychological resources, early adversity was related to cortisol responses (linear: $b(SE) = -.26(.08)$, $p = .001$; quadratic: $b(SE) = .04(.01)$, $p = .005$). At high levels of psychological resources, early adversity was not related to cortisol responses (linear: $b(SE) = -.07(.08)$, $p = .406$; quadratic: $b(SE) = .03(.01)$, $p = .072$). Visualization of the simple interaction effects revealed a similar pattern as that depicted in Figure 1.

¹We also examined whether summing the number of daily stressors across days yielded similar findings given that the cumulative exposure to stress across days might be distinct from the proportion of stressor days. These two measures of daily stress were highly correlated ($r = .89$, $p < .001$). Similar to proportion of stressor days, cumulative stress exposure modulated cortisol responses to the TSST (linear: $b(SE) = -.01(.01)$, $p = .028$; quadratic: $b(SE) = .002(.001)$, $p = .027$). Adolescents who reported a greater number of stressors across days had smaller cortisol responses (linear: $b(SE) = .23(.07)$, $p < .001$; quadratic: $b(SE) = -.04(.01)$, $p < .001$) compared to those who reported a smaller number of stressors across days (linear: $b(SE) = .36(.04)$, $p < .001$; quadratic: $b(SE) = -.07(.01)$, $p < .001$). Contrasting findings for proportion of stressor days, however, there was no evidence that the variation in cortisol responses related to cumulative stress exposure across days was further modulated by psychological resources (linear: $b(SE) = .01(.01)$, $p = .119$; quadratic: $b(SE) = -.001(.001)$, $p = .459$).

Inflammatory Reactivity

Major life events—As shown in Table 4 (column 1), neither major life events nor psychological resources were associated with baseline levels of IL-6 (major life events: $p = .812$; psychological resources: $p = .332$) and IL-6 reactivity (major life events: $p = .669$; psychological resources: $p = .609$). There was also no major life events by psychological resources interaction effect on IL-6 reactivity ($p = .098$; Table 4, column 2).

Daily interpersonal stress—Table 4 also shows that daily interpersonal stress operationalized as proportion of stressor days and psychological resources were not related to IL-6 at baseline (daily stress: $p = .176$; psychological resources: $p = .452$) and in response to the TSST (daily stress: $p = .869$; psychological resources: $p = .408$). Daily interpersonal stress and psychological resources also did not interact to influence IL-6 reactivity ($p = .261$)².

Early adversity—Again, as shown in Table 4, early adversity and psychological resources were not associated with baseline IL-6 (early adversity: $p = .455$; psychological resources: $p = .494$) and IL-6 reactivity (early adversity: $p = .283$; psychological resources: $p = .248$). There was also no interaction effect between early family stress and psychological resources on inflammatory reactivity ($p = .10$).

Sensitivity Analyses

We tested whether higher psychological resources may be a proxy for better overall mental health by statistically adjusting for depressive symptoms and by testing whether depressive symptoms interacted with the various stressors to predict HPA reactivity. Results remained the same when including depressive symptoms as a covariate. There was also no evidence that daily interpersonal stress or early adversity interacted with depressive symptoms to influence cortisol responses (linear terms: $\beta s = .644 - .796$; quadratic terms: $\beta s = .564 - .885$). Depressive symptoms did, however, interact with major life events to predict cortisol responses (linear term: $b(SE) = -.004(.001)$, $p = .003$; quadratic term: $b(SE) = .001(.0003)$, $p = .023$). Major life events were related to cortisol reactivity only among those with higher depressive symptoms ($n = 21$; linear term: $b(SE) = -.06(.03)$, $p = .019$; quadratic term: $b(SE) = .01(.005)$, $p = .074$), but not among those with lower depressive symptoms ($n = 33$; linear term: $b(SE) = .01(.03)$, $p = .803$; quadratic term: $b(SE) = -.002(.005)$, $p = .643$).

Exploratory Analyses

Compared to European-Americans, Latinos face higher rates of poverty and socioeconomic and racial discrimination, and have different family values, which may lead to differential effects of stress and psychological resources on biological responsivity. Therefore, we examined whether ethnicity moderated the effects of stress and the effects of the stress by psychological resources interactions on cortisol and IL-6 responses. Overall there was no evidence that stress associations with cortisol (linear: $\beta s > .352$; quadratic: $\beta s > .155$) and IL-6 ($\beta s > .110$) responses were different across ethnicity. The only exception to this pattern

²The cumulative number of stressors across the days similarly did not modulate IL-6 responses ($b(SE) = .001(.003)$, $p = .619$). This association was also not further modulated by psychological resources ($b(SE) = .003(.003)$, $p = .306$).

of findings was the association between early adversity and cortisol responses (linear term: $b(SE) = -.31 (.12)$, $p = .009$; quadratic term: $b(SE) = .06 (.02)$, $p = .003$). Early adversity was associated with decreased cortisol responses among Latinos (linear term: $b(SE) = -.25 (.07)$, $p < .001$; quadratic term: $b(SE) = .05 (.01)$, $p < .001$), but not among European-Americans (linear term: $b(SE) = .01 (.09)$, $p = .947$; quadratic term: $b(SE) = -.01 (.01)$, $p = .699$). There was no evidence of ethnic differences in stress associations with cortisol (linear: $ps > .590$; quadratic: $ps > .255$) and IL-6 ($ps > .065$) responses moderated by psychological resources.

Discussion

The present study aimed to deepen our understanding of factors that may protect against the negative biological effects of psychosocial stress during adolescence, a developmental period important to the establishment of future adult health. Consistent with past work on childhood and adolescent stress and HPA axis responses to subsequent acute stress (e.g., Crum et al., 2015; Carpenter et al., 2007; Elzinga et al., 2008; Kraft & Luecken, 2009), we found that experiences of various stressors, including major life events, daily interpersonal stress, and early adversity, were associated with smaller cortisol responses to an acute laboratory-based stressor. Individual psychological resources moderated the links from major life events and daily stress to cortisol reactivity, such that they were evident only among adolescents who reported lower levels of psychological resources. By comparison, these stress-cortisol reactivity associations were not apparent among those reporting higher levels of psychological resources. These findings were specific to the HPA axis and were not evident for inflammatory processes.

The present findings converge with past work showing that among socioeconomically disadvantaged youth, positive reappraisal tendencies and finding meaning in life were associated with reduced risk of poor health-related outcomes, such as obesity and cardiovascular risk (Chen et al., 2013; Chen et al., 2015; Kalleem et al., 2013). We extend this work by demonstrating that individual psychological resources may also protect young individuals from the effects of other stressors, including major life events, daily interpersonal stress, and to some extent early adversity, on reduced HPA responses to acute stress, which is thought to be one key mechanism linking stress to health and psychopathology (Ehrlich et al., 2016; Heim & Binder, 2012). The current findings raise the question of how psychological resources may counter the effects of chronic or repeated stress on HPA axis reactivity to subsequent acute stress. One explanation is that psychological resources foster effective coping strategies (Taylor & Broffman, 2011; Taylor & Stanton, 2007). Having higher levels of optimism, self-esteem, self-mastery, and a tendency to positively reappraise negative or threatening situations likely biases one towards regulation of stress-evoked negative emotions, active problem-solving, and/or seeking support or other resources, over maladaptive ways of coping, such as rumination. In the face of major life events or repeated daily stress, these coping strategies may attenuate HPA reactivity and/or facilitate quicker recovery. To the extent that hypocortisolism results after extended periods of high levels of cortisol as a counter-regulatory mechanism (Fries et al., 2005; Miller et al., 2007), attenuated reactivity to and/or quicker recovery from previous stress may reduce the amount of overall exposure to cortisol and prevent or delay hypocortisolism from emerging.

Psychological resources seemed to have stronger protective effects for major life events and daily stress relative to early adversity. Whereas significant interactions with psychological resources emerged for major life events and daily stress, there was only a marginally significant interaction between early adversity and psychological resources on linear cortisol changes, and there was no interaction for quadratic cortisol changes. Why we observed this pattern is not entirely clear, but may have to do with temporal distance between exposure to the different stressors and the assessment of cortisol responses. In the current study, major life events tapped into stressful events within the last year and daily stress assessed the everyday interpersonal in participants' current lives. By contrast, early adversity assessed participants' family climate from ages 5 to 15. Given the longer period of time between early adversity and assessment of HPA reactivity, the effects of early adversity may have had more sufficient time to "get under the skin" and solidify. As such, psychological resources may be particularly less effective for more temporally distal stressors relative to more temporally proximal stressors. Of course, this remains speculative, and requires empirical examination.

One caveat to the observed pattern of findings is that there was some evidence that psychological resources per se did not buffer the effects of major life events on cortisol responses. Psychological resources and depressive symptoms were correlated fairly strongly ($r = -.60$), and major life events interacted with depressive symptoms, such that major life events were related to cortisol responses only among those with higher depressive symptoms, but not among individuals with lower depressive symptoms. It may be that for the effects of major life events, psychological resources served as a proxy for overall adjustment, and lower mental health may be the important moderator that accentuates relations between major life events and HPA axis functioning. By contrast, psychological resources may be effective in counteracting the adverse health-related effects not only of socioeconomic disadvantage, as shown in previous studies (Chen et al., 2013; Chen et al., 2015; Kalleem et al., 2013), but also of other chronic stressors such as more mundane daily stressors that may accumulate over time.

Interestingly, exploratory analyses revealed ethnic differences in the associations between early adversity and cortisol responses, but not in the buffering role of psychological resources. Specifically, the association between early adversity and cortisol reactivity was evident only in Latino adolescents, but not in their European-American counterparts. These findings may reflect ethnic differences in family values. Our measure of early adversity assessed overall family environment, and compared to European-American families, Latino families are characterized by higher levels of familism, or the view that the family constitutes the self and is the primary source of support; family duties, interconnectedness and unity among family members, and family needs rather than individual needs are also emphasized (Fuligni et al., 1999; Sabogal et al., 1987). As such, early family-related stress, may be more distressing for Latino adolescents. We found no ethnic differences in the protective role of psychological resources, which might suggest that psychological resources do not operate as a protective factor for Latino youth exposed to a harsh family climate. However, the current sample size may be a limiting factor. Future work should further probe psychological resources as a possible protective factor, particularly for Latino youth experiencing high levels of family-related stress.

Findings for stress and its interaction with psychological resources were specific to the HPA axis, and not observed for inflammatory reactivity. At first, this may seem somewhat surprising given that the HPA axis regulates inflammatory activity (Irwin & Cole, 2011; Rohleder, 2014). In particular, cortisol down-regulates the production of cytokines when bound to its corresponding receptors in immune cells, and as such, decreased cortisol reactivity might be expected to result in increases in circulating levels of IL-6 among those with higher levels of stress and lower levels of psychological resources. However, in order for cortisol to bind to its receptor in immune cells, cortisol must be unbound to carrier proteins. Yet, only a small fraction of cortisol (approximately 10%) remains unbound and biologically active (El-Farhan, Rees, & Evans, 2017). Furthermore, before modifying other biological systems, including the immune system, unbound, free cortisol, must avoid being excreted and converted to its inactive form by counter-regulatory mechanisms (Ehrlich et al., 2016; Miller et al., 2009). Once bound to its receptors, it must translocate to the nucleus of the cell where it can then exert its effects. Measures of salivary cortisol, as used in the present study, do not offer any insight into these processes, and therefore, the downstream effects of the smaller cortisol responses on inflammatory processes in adolescents with higher levels of stress and lower psychological resources cannot be determined in the present study.

The fact that we observed no stress and IL-6 associations or moderation by psychological resources seems to diverge from prior studies showing significant interactions between socioeconomic status and psychological resources on inflammatory processes (Chen et al., 2015; Elliot & Chapman, 2016). However, these studies examined production of inflammatory cytokines in response to a bacterial stimulus in youth and circulating markers of systemic inflammation under tonic conditions in midlife adults. In the present study, we focused on circulating IL-6 in response to an acute psychosocial stressor in late adolescents. The lack of findings for IL-6 reactivity may have to do with the fact that systemic inflammation remains low in earlier life stages whereas the HPA axis is known to be highly sensitive to the environment across the lifespan, even early in life when physical health is relatively intact (Gunnar & Quevedo, 2007; Hostinar, Sullivan, & Gunnar, 2014). As such, there may be delayed effects of stress on circulating markers of inflammation, and links may emerge in later in life. Consistent with this notion, associations between stress and circulating markers of inflammation are more consistently observed in adults compared to youth (Baumeister et al., 2016; Coelho et al., 2014; Slopen et al., 2012). The fact that circulating inflammatory cytokines stem from both immune and non-immune sources, such as adipocytes (Black, 2003), may also make it difficult to detect associations among stress, psychological resources, and inflammatory reactivity in young individuals. To the extent that stress biases immune cells specifically towards a pro-inflammatory phenotype (Miller et al., 2011) and that psychological resources counteract the impact of stress (as described above), effects in young persons may be better detected when stimulating immune cells, as in the study by Chen et al. (2015).

The current investigation was not without limitations. First, the observational nature of the study precludes any conclusion about causal relations and makes alternative explanations plausible. Although we adjusted for potential confounds and considered depressive symptoms as an alternative explanation, other unmeasured factors associated with stress and

HPA axis functioning may have contributed to the observed findings. Second, given the one-time assessment of HPA axis reactivity to stress, we were unable to determine whether the effects of the various stressors on cortisol reactivity and the attenuation of these associations by psychological resources persist across time. As development continues, later experiences may alter the health trajectories that early experiences initially set in motion (Chiang, Chen, & Miller, 2018; Gouin, Caldwell, Woods, & Malarkey, 2017), making it important for future longitudinal studies to elucidate whether the current findings persist across time. Third, the clinical relevance of the study findings cannot be determined. We did not measure any physical health outcomes, and the clinical significance of the magnitude of cortisol decreases we observed remain unknown. Lastly, participants were European- and Latin-Americans from the Los Angeles area and of a narrow age range. Thus, more work is needed to determine generalizability of our findings.

Despite these limitations, the current study nevertheless contributes to previous literature. It builds on prior research on psychological resources by extending findings of the protective role of psychological resources to an adolescent sample. It extends past work on adolescent stress biology by highlighting psychological resources as potential protective factors that may mitigate the association between stress and dampened cortisol reactivity. Results raise the possibility that intervention programs aimed at enhancing psychological resources among young individuals may ultimately help counteract the effects of certain stressors on HPA axis functioning, though the long-term health implications remain unclear.

Acknowledgments

This research was supported by funding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (R01-HD062547), UCLA California Center for Population Research funded by the National Institute of Child Health and Human Development (R24-HD041022), UCLA Older Americans Independence Center funded by the National Institute of Aging (P30-AG028748), UCLA Cousins Center for Psychoneuroimmunology, University of California Institute for Mexico and the US, American Psychological Association, and Division 38 of the American Psychological Association. Authors' efforts were supported by the National Heart, Lung, and Blood Institute (F32-HL134276 to J.J.C.).

References

- American Psychological Association A. P. Stress in America: Are teens adopting adults' stress habits. Stress in America Surveys. 2014. (<http://www.apa.org/news/press/releases/stress/2013/stress-report.pdf>)
- Baumeister D, Akhtar R, Ciufofini S, Pariante CM, Mondelli V. 2016; Childhood trauma and adulthood inflammation: a meta-analysis of peripheral C-reactive protein, interleukin-6 and tumour necrosis factor- α . *Molecular Psychiatry*. 21(5):642–649. DOI: 10.1038/mp.2015.67 [PubMed: 26033244]
- Black PH. 2003; The inflammatory response is an integral part of the stress response: Implications for atherosclerosis, insulin resistance, type II diabetes and metabolic syndrome X. *Brain, Behavior, and Immunity*. 17(5):350–364. DOI: 10.1016/S0889-1591(03)00048-5
- Blakemore SJ, Mills KL. 2014; Is adolescence a sensitive period for sociocultural processing? *Annual Review of Psychology*. 65:187–207. DOI: 10.1146/annurev-psych-010213-115202
- Boylan JM, Jennings JR, Matthews KA. 2016; Childhood socioeconomic status and cardiovascular reactivity and recovery among Black and White men: Mitigating effects of psychological resources. *Health Psychology*. 35(9):957–966. DOI: 10.1037/hea0000355 [PubMed: 27054298]

- Brody GH, Yu T, Beach SRH. 2016; Resilience to adversity and the early origins of disease. *Development and Psychopathology*. 28(4pt2):1347–1365. DOI: 10.1017/S0954579416000894 [PubMed: 27692007]
- Brown DW, Anda RF, Tiemeier H, Felitti VJ, Edwards VJ, Croft JB, Giles WH. 2009; Adverse childhood experiences and the risk of premature mortality. *American Journal of Preventive Medicine*. 37(5):389–396. DOI: 10.1016/j.amepre.2009.06.021 [PubMed: 19840693]
- Brydon L, Steptoe A. 2005; Stress-induced increases in interleukin-6 and fibrinogen predict ambulatory blood pressure at 3-year follow-up. *Journal of Hypertension*. 23(5):1001–1007. DOI: 10.1097/01.hjh.0000166841.57474.d0 [PubMed: 15834286]
- Bunea IM, Szentágotai-Tar A, Miu AC. 2017; Early-life adversity and cortisol response to social stress: a meta-analysis. *Translational Psychiatry*. 7(12):1274.doi: 10.1038/s41398-017-0032-3 [PubMed: 29225338]
- Cimneci M, Crișan LG, Vulturar R, Opre A, Miu AC. 2015; Emotional non-acceptance links early life stress and blunted cortisol reactivity to social threat. *Psychoneuroendocrinology*. 51:176–187. DOI: 10.1016/j.psyneuen.2014.09.026 [PubMed: 25462891]
- Carpenter LL, Carvalho JP, Tyrka AR, Wier LM, Mello AF, Mello MF, Price LH. 2007; Decreased adrenocorticotropic hormone and cortisol responses to stress in healthy adults reporting significant childhood maltreatment. *Biological Psychiatry*. 62(10):1080–1087. DOI: 10.1016/j.biopsych.2007.05.002 [PubMed: 17662255]
- Carpenter LL, Shattuck TT, Tyrka AR, Geraciotti TD, Price LH. 2011; Effect of childhood physical abuse on cortisol stress response. *Psychopharmacology*. 214(1):367–375. DOI: 10.1007/s00213-010-2007-4 [PubMed: 20838776]
- Carroll D, Bibbey A, Roseboom TJ, Phillips AC, Ginty AT, Rooij SR. 2012; Forced expiratory volume is associated with cardiovascular and cortisol reactions to acute psychological stress. *Psychophysiology*. 49(6):866–872. DOI: 10.1111/j.1469-8986.2012.01361.x [PubMed: 22416939]
- Carroll JE, Gruenewald TL, Taylor SE, Janicki-Deverts D, Matthews KA, Seeman TE. 2013; Childhood abuse, parental warmth, and adult multisystem biological risk in the Coronary Artery Risk Development in Young Adults study. *Proceedings of the National Academy of Sciences*. 110(42):17149–17153. DOI: 10.1073/pnas.1315458110
- Carver CS, Scheier MF, Segerstrom SC. 2010; Optimism. *Clinical Psychology Review*. 30(7):879–889. DOI: 10.1016/j.cpr.2010.01.006 [PubMed: 20170998]
- Chen E, Lee WK, Cavey L, Ho A. 2013; Role models and the psychological characteristics that buffer low-socioeconomic-status youth from cardiovascular risk. *Child Development*. 84(4):1241–1252. DOI: 10.1111/cdev.12037 [PubMed: 23278857]
- Chen E, McLean KC, Miller GE. 2015; Shift-and-persist strategies: associations with socioeconomic status and the regulation of inflammation among adolescents and their parents. *Psychosomatic Medicine*. 77(4):371–382. DOI: 10.1097/PSY.0000000000000157 [PubMed: 26167560]
- Chen E, Miller GE. 2013; Socioeconomic status and health: mediating and moderating factors. *Annual Review of Clinical Psychology*. 9:723–749. DOI: 10.1146/annurev-clinpsy-050212-185634
- Chen E, Miller GE, Kobor MS, Cole SW. 2011; Maternal warmth buffers the effects of low early-life socioeconomic status on pro-inflammatory signaling in adulthood. *Molecular Psychiatry*. 16(7):729–737. DOI: 10.1038/mp.2010.53 [PubMed: 20479762]
- Chiang JJ, Bower JE, Irwin MR, Taylor SE, Fuligni AJ. 2017; Adiposity moderates links from early adversity and depressive symptoms to inflammatory reactivity to acute stress during late adolescence. *Brain, Behavior, and Immunity*. 66:146–155. DOI: 10.1016/j.bbi.2017.06.015
- Chiang, JJ; Chen, E; Miller, GE. Midlife self-reported social support as a buffer against premature mortality risks associated with childhood abuse. *Nature Human Behaviour*. 2018.
- Chiang JJ, Eisenberger NI, Seeman TE, Taylor SE. 2012; Negative and competitive social interactions are related to heightened proinflammatory cytokine activity. *Proceedings of the National Academy of Sciences*. 109(6):1878–1882. DOI: 10.1073/pnas.1120972109
- Chiang JJ, Taylor SE, Bower JE. 2015; Early adversity, neural development, and inflammation. *Developmental Psychobiology*. 57(8):887–907. DOI: 10.1002/dev.21329 [PubMed: 26369600]
- Chiang JJ, Tsai KM, Park H, Bower JE, Almeida DM, Dahl RE, Fuligni AJ. 2016; Daily family stress and HPA axis functioning during adolescence: The moderating role of sleep.

Psychoneuroendocrinology. 71:43–53. DOI: 10.1016/j.psyneuen.2016.05.009 [PubMed: 27235639]

- Choy E. 2012; Understanding the dynamics: pathways involved in the pathogenesis of rheumatoid arthritis. *Rheumatology*. 51(suppl_5):v3–v11. [PubMed: 22718924]
- Chung GH, Flook L, Fuligni AJ. 2009; Daily family conflict and emotional distress among adolescents from Latin American, Asian, and European backgrounds. *Developmental Psychology*. 45(5):1406–1415. DOI: 10.1037/a0014163 [PubMed: 19702401]
- Coelho R, Viola TW, Walss-Bass C, Brietzke E, Grassi-Oliveira R. 2014; Childhood maltreatment and inflammatory markers: a systematic review. *Acta Psychiatrica Scandinavica*. 129(3):180–192. DOI: 10.1111/acps.12217 [PubMed: 24205846]
- Cohen S, Doyle WJ, Turner RB, Alper CM, Skoner DP. 2004; Childhood socioeconomic status and host resistance to infectious illness in adulthood. *Psychosomatic Medicine*. 66(4):553–558. [PubMed: 15272102]
- Conger RD, Wallace LE, Sun Y, Simons RL, McLoyd VC, Brody GH. 2002; Economic pressure in African American families: a replication and extension of the family stress model. *Developmental Psychology*. 38(2):179–193. DOI: 10.1037/0012-1649.38.2.179 [PubMed: 11881755]
- de Rooij SR. 2013; Blunted cardiovascular and cortisol reactivity to acute psychological stress: A summary of results from the Dutch Famine Birth Cohort Study. *International Journal of Psychophysiology*. 90(1):21–27. DOI: 10.1016/j.ijpsycho.2012.09.011 [PubMed: 23046585]
- Del Giudice M, Ellis BJ, Shirtcliff EA. 2011; The adaptive calibration model of stress responsivity. *Neuroscience & Biobehavioral Reviews*. 35(7):1562–1592. DOI: 10.1016/j.neubiorev.2010.11.007 [PubMed: 21145350]
- DeLongis A, Folkman S, Lazarus RS. 1988; The impact of daily stress on health and mood: psychological and social resources as mediators. *Journal of Personality and Social Psychology*. 54(3):486. [PubMed: 3361420]
- Demakakos P, Pillas D, Marmot M, Steptoe A. 2016; Parenting style in childhood and mortality risk at older ages: a longitudinal cohort study. *The British Journal of Psychiatry*. 209(2):135–141. DOI: 10.1192/bjp.bp.115.163543 [PubMed: 26941265]
- DeSantis AS, Adam EK, Doane LD, Mineka S, Zinbarg RE, Craske MG. 2007; Racial/ethnic differences in cortisol diurnal rhythms in a community sample of adolescents. *Journal of Adolescent Health*. 41(1):3–13. DOI: 10.1016/j.jadohealth.2007.03.006 [PubMed: 17577528]
- Dickerson SS, Kemeny ME. 2004; Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research. *Psychological Bulletin*. 130(3):355–391. DOI: 10.1037/0033-2909.130.3.355 [PubMed: 15122924]
- Dietz LJ, Stoyak S, Melhem N, Porta G, Matthews KA, Payne MW, Brent DA. 2013; Cortisol response to social stress in parentally bereaved youth. *Biological Psychiatry*. 73(4):379–387. DOI: 10.1016/j.biopsycho.2012.08.016 [PubMed: 23021533]
- Dowd JB, Simanek AM, Aiello AE. 2009; Socio-economic status, cortisol and allostatic load: a review of the literature. *International Journal of Epidemiology*. 38(5):1297–1309. DOI: 10.1093/ije/dyp277 [PubMed: 19720725]
- Ehrlich, KB, Miller, GE, Chen, E. *Childhood adversity and adult physical health* *Developmental Psychopathology*. Vol. 4. Hoboken, NJ: Wiley; 2016. 1–42.
- El-Farhan N, Rees DA, Evans C. 2017; Measuring cortisol in serum, urine and saliva—are our assays good enough? *Annals of Clinical Biochemistry*. 54(3):308–322. DOI: 10.1177/0004563216687335 [PubMed: 28068807]
- Elinav E, Nowarski R, Thaiss CA, Hu B, Jin C, Flavell RA. 2013; Inflammation-induced cancer: crosstalk between tumours, immune cells and microorganisms. *Nature Reviews Cancer*. 13(11): 759–771. DOI: 10.1038/nrc3611 [PubMed: 24154716]
- Elliot AJ, Chapman BP. 2016; Socioeconomic status, psychological resources, and inflammatory markers: Results from the MIDUS study. *Health Psychology*. 35(11):1205–1213. DOI: 10.1037/hea0000392 [PubMed: 27280368]
- Elzinga BM, Roelofs K, Tollenaar MS, Bakvis P, van Pelt J, Spinhoven P. 2008; Diminished cortisol responses to psychosocial stress associated with lifetime adverse events: a study among healthy

- young subjects. *Psychoneuroendocrinology*. 33(2):227–237. DOI: 10.1016/j.psyneuen.2007.11.004 [PubMed: 18096322]
- Engert V, Efanov SI, Dedovic K, Duchesne A, Dagher A, Pruessner JC. 2010; Perceived early-life maternal care and the cortisol response to repeated psychosocial stress. *Journal of Psychiatry & Neuroscience*. 35(6):370–377. DOI: 10.1503/jpn.100022 [PubMed: 20964960]
- Fries E, Hesse J, Hellhammer J, Hellhammer DH. 2005; A new view on hypocortisolism. *Psychoneuroendocrinology*. 30(10):1010–1016. DOI: 10.1016/j.psyneuen.2005.04.006 [PubMed: 15950390]
- Fuligni AJ, Telzer EH, Bower J, Cole SW, Kiang L, Irwin MR. 2009; A preliminary study of daily interpersonal stress and C-reactive protein levels among adolescents from Latin American and European backgrounds. *Psychosomatic Medicine*. 71(3):329–333. DOI: 10.1097/PSY.0b013e3181921b1f [PubMed: 19196810]
- Fuligni AJ, Tseng V, Lam M. 1999; Attitudes toward family obligations among American adolescents with Asian, Latin American, and European backgrounds. *Child Development*. 70(4):1030–1044. DOI: 10.1111/1467-8624.00075
- Gallo LC, de los Monteros KE, Shivpuri S. 2009; Socioeconomic status and health: what is the role of reserve capacity? *Current Directions in Psychological Science*. 18(5):269–274. DOI: 10.1111/j.1467-8721.2009.01650.x [PubMed: 22210579]
- Gouin JP, Caldwell W, Woods R, Malarkey WB. 2017; Resilience resources moderate the association of adverse childhood experiences with adulthood inflammation. *Annals of Behavioral Medicine*. 51(4):782–786. DOI: 10.1007/s12160-017-9891-3 [PubMed: 28281135]
- Greene ML, Way N, Pahl K. 2006; Trajectories of perceived adult and peer discrimination among Black, Latino, and Asian American adolescents: patterns and psychological correlates. *Developmental Psychology*. 42(2):218–236. DOI: 10.1037/0012-1649.42.2.218 [PubMed: 16569162]
- Gross JJ, John OP. 2003; Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. *Journal of Personality and Social Psychology*. 85(2):348–362. DOI: 10.1037/0022-3514.85.2.348 [PubMed: 12916575]
- Grzywacz JG, Almeida DM, McDonald DA. 2002; Work–family spillover and daily reports of work and family stress in the adult labor force. *Family Relations*. 51(1):28–36. DOI: 10.1111/j.1741-3729.2002.00028.x
- Gunnar M, Quevedo K. 2007; The neurobiology of stress and development. *Annual Review of Psychology*. 58:145–173. DOI: 10.1146/annurev.psych.58.110405.085605
- Gunnar MR, Wewerka S, Frenn K, Long JD, Griggs C. 2009; Developmental changes in hypothalamus–pituitary–adrenal activity over the transition to adolescence: normative changes and associations with puberty. *Development and Psychopathology*. 21(1):69–85. DOI: 10.1017/S0954579409000054 [PubMed: 19144223]
- Hammen C. 1991; Generation of stress in the course of unipolar depression. *Journal of Abnormal Psychology*. 100(4):555–561. DOI: 10.1037/0021-843X.100.4.555 [PubMed: 1757669]
- Heim C, Binder EB. 2012; Current research trends in early life stress and depression: Review of human studies on sensitive periods, gene–environment interactions, and epigenetics. *Experimental Neurology*. 233(1):102–111. DOI: 10.1016/j.expneurol.2011.10.032 [PubMed: 22101006]
- Heim C, Ehlert U, Hellhammer DH. 2000; The potential role of hypocortisolism in the pathophysiology of stress-related bodily disorders. *Psychoneuroendocrinology*. 25(1):1–35. DOI: 10.1016/S0306-4530(99)00035-9 [PubMed: 10633533]
- Hostinar CE, Sullivan RM, Gunnar MR. 2014; Psychobiological mechanisms underlying the social buffering of the hypothalamic–pituitary–adrenocortical axis: A review of animal models and human studies across development. *Psychological Bulletin*. 140(1):256–282. DOI: 10.1037/a0032671 [PubMed: 23607429]
- Irwin MR, Cole SW. 2011; Reciprocal regulation of the neural and innate immune systems. *Nature Reviews Immunology*. 11(9):625–632. DOI: 10.1038/nri3042
- Jacks DE, Sowash J, Anning J, Mcgloughlin T, Andres F. 2002; Effect of exercise at three exercise intensities on salivary cortisol. *The Journal of Strength & Conditioning Research*. 16(2):286–289. [PubMed: 11991783]

- Janusek LW, Tell D, Gaylord-Harden N, Mathews HL. 2017; Relationship of childhood adversity and neighborhood violence to a proinflammatory phenotype in emerging adult African American men: an epigenetic link. *Brain, Behavior, and Immunity*. 60:126–135. DOI: 10.1016/j.bbi.2016.10.006
- Kallem S, Carroll-Scott A, Rosenthal L, Chen E, Peters SM, McCaslin C, Ickovics JR. 2013; Shift-and-persist: A protective factor for elevated BMI among low-socioeconomic-status children. *Obesity*. 21(9):1759–1763. DOI: 10.1002/oby.20195 [PubMed: 23671041]
- Kershaw KN, Brenes GA, Charles LE, Coday M, Daviglius ML, Denburg NL, Tindle HA. 2014; Associations of stressful life events and social strain with incident cardiovascular disease in the Women’s Health Initiative. *Journal of the American Heart Association*. 3(3):e000687.doi: 10.1161/JAHA.113.000687 [PubMed: 24973226]
- Kirschbaum C, Pirke KM, Hellhammer DH. 1993; The ‘Trier Social Stress Test’—a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*. 28(1–2):76–81. DOI: 10.1159/000119004 [PubMed: 8255414]
- Kraft AJ, Luecken LJ. 2009; Childhood parental divorce and cortisol in young adulthood: evidence for mediation by family income. *Psychoneuroendocrinology*. 34(9):1363–1369. DOI: 10.1016/j.psyneuen.2009.04.008 [PubMed: 19442451]
- Libby P. 2006; Inflammation and cardiovascular disease mechanisms. *The American Journal of Clinical Nutrition*. 83(2):456S–460S. DOI: 10.1093/ajcn/83.2.456S [PubMed: 16470012]
- Loucks EB, Almeida ND, Taylor SE, Matthews KA. 2011; Childhood family psychosocial environment and coronary heart disease risk. *Psychosomatic Medicine*. 73(7):563–571. DOI: 10.1097/PSY.0b013e318228c820 [PubMed: 21810898]
- Lovallo WR. 2013; Early life adversity reduces stress reactivity and enhances impulsive behavior: Implications for health behaviors. *International Journal of Psychophysiology*. 90(1):8–16. DOI: 10.1016/j.ijpsycho.2012.10.006 [PubMed: 23085387]
- Lovallo WR, Whitsett TL, al’Absi M, Sung BH, Vincent AS, Wilson MF. 2005; Caffeine stimulation of cortisol secretion across the waking hours in relation to caffeine intake levels. *Psychosomatic Medicine*. 67(5):734–739. DOI: 10.1097/01.psy.0000181270.20036.06 [PubMed: 16204431]
- Luecken LJ, Appelhans BM. 2006; Early parental loss and salivary cortisol in young adulthood: the moderating role of family environment. *Development and Psychopathology*. 18(1):295–308. [PubMed: 16478564]
- Lundgren O, Garvin P, Jonasson L, Andersson G, Kristenson M. 2015; Psychological resources are associated with reduced incidence of coronary heart disease. An 8-year follow-up of a community-based Swedish sample. *International Journal of Behavioral Medicine*. 22(1):77–84. DOI: 10.1007/s12529-014-9387-5 [PubMed: 24430130]
- MacMillan HL, Georgiades K, Duku EK, Shea A, Steiner M, Niec A, Vella E. 2009; Cortisol response to stress in female youths exposed to childhood maltreatment: results of the youth mood project. *Biological Psychiatry*. 66(1):62–68. DOI: 10.1016/j.biopsych.2008.12.014 [PubMed: 19217075]
- Marsland AL, Walsh C, Lockwood K, John-Henderson NA. 2017; The effects of acute psychological stress on circulating and stimulated inflammatory markers: a systematic review and meta-analysis. *Brain, Behavior, and Immunity*. 64:208–219. DOI: 10.1016/j.bbi.2017.01.011
- McEwen BS, Seeman T. 1999; Protective and damaging effects of mediators of stress: elaborating and testing the concepts of allostasis and allostatic load. *Annals of the New York Academy of Sciences*. 896(1):30–47. DOI: 10.1111/j.1749-6632.1999.tb08103.x [PubMed: 10681886]
- Miller GE, Chen E. 2010; Harsh family climate in early life presages the emergence of a proinflammatory phenotype in adolescence. *Psychological Science*. 21(6):848–856. DOI: 10.1177/0956797610370161 [PubMed: 20431047]
- Miller GE, Chen E, Fok AK, Walker H, Lim A, Nicholls EF, Kobor MS. 2009; Low early-life social class leaves a biological residue manifested by decreased glucocorticoid and increased proinflammatory signaling. *Proceedings of the National Academy of Sciences*. 106(34):14716–14721. DOI: 10.1073/pnas.0902971106
- Miller GE, Chen E, Parker KJ. 2011; Psychological stress in childhood and susceptibility to the chronic diseases of aging: moving toward a model of behavioral and biological mechanisms. *Psychological Bulletin*. 137(6):959–997. DOI: 10.1037/a0024768 [PubMed: 21787044]

- Miller GE, Chen E, Zhou ES. 2007; If it goes up, must it come down? Chronic stress and the hypothalamic-pituitary-adrenocortical axis in humans. *Psychological Bulletin*. 133(1):25–45. DOI: 10.1037/0033-2909.133.1.25 [PubMed: 17201569]
- Nishina A, Juvonen J. 2005; Daily reports of witnessing and experiencing peer harassment in middle school. *Child Development*. 76(2):435–450. DOI: 10.1111/j.1467-8624.2005.00855.x [PubMed: 15784092]
- O'Connor MF, Bower JE, Cho HJ, Creswell JD, Dimitrov S, Hamby ME, Sloan EK. 2009; To assess, to control, to exclude: effects of biobehavioral factors on circulating inflammatory markers. *Brain, Behavior, and Immunity*. 23(7):887–897. DOI: 10.1016/j.bbi.2009.04.005
- Pearlin LI, Schooler C. 1978; The structure of coping. *Journal of Health and Social Behavior*. 19(1):2–21. [PubMed: 649936]
- Phillips AC, Carroll D, Burns VE, Drayson M. 2005; Neuroticism, cortisol reactivity, and antibody response to vaccination. *Psychophysiology*. 42(2):232–238. DOI: 10.1111/j.1469-8986.2005.00281.x [PubMed: 15787860]
- Pyykkönen AJ, Rääkkönen K, Tuomi T, Eriksson JG, Groop L, Isomaa B. 2010; Stressful life events and the metabolic syndrome. *Diabetes Care*. 33(2):378–384. DOI: 10.2337/dc09-1027 [PubMed: 19880581]
- Repetti RL, Robles TF, Reynolds B. 2011; Allostatic processes in the family. *Development and Psychopathology*. 22:921–938. DOI: 10.1017/S095457941100040X
- Repetti RL, Taylor SE, Seeman TE. 2002; Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*. 128(2):330–366. DOI: 10.1037/0033-2909.128.2.330 [PubMed: 11931522]
- Rohleder N. 2014; Stimulation of systemic low-grade inflammation by psychosocial stress. *Psychosomatic Medicine*. 76(3):181–189. DOI: 10.1097/PSY.0000000000000049 [PubMed: 24608036]
- Rosenberg, M. *Society and the adolescent self-image*. Princeton, NJ: Princeton University Press; 1965.
- Sabogal F, Marín G, Otero-Sabogal R, Marín BV, Perez-Stable EJ. 1987; Hispanic familism and acculturation: What changes and what doesn't? *Hispanic Journal of Behavioral Sciences*. 9(4): 397–412. DOI: 10.1177/07399863870094003
- Scheier MF, Carver CS, Bridges MW. 1994; Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): a reevaluation of the Life Orientation Test. *Journal of Personality and Social Psychology*. 67(6):1063–1078. DOI: 10.1037/0022-3514.67.6.1063 [PubMed: 7815302]
- Sebastian C, Viding E, Williams KD, Blakemore SJ. 2010; Social brain development and the affective consequences of ostracism in adolescence. *Brain and Cognition*. 72(1):134–145. DOI: 10.1016/j.bandc.2009.06.008 [PubMed: 19628323]
- Semega, JL, Frontenot, KR, Kollar, MA, Bureau, USC. *Income and poverty in the United States: 2016*. Washington DC: U.S Government Printing Office; 2017.
- Sin NL, Graham-Engeland JE, Ong AD, Almeida DM. 2015; Affective reactivity to daily stressors is associated with elevated inflammation. *Health Psychology*. 34(12):1154–1165. DOI: 10.1037/hea0000240 [PubMed: 26030309]
- Slavich GM, Irwin MR. 2014; From stress to inflammation and major depressive disorder: A social signal transduction theory of depression. *Psychological Bulletin*. 140(3):774–815. DOI: 10.1037/a0035302 [PubMed: 24417575]
- Slopen N, Koenen KC, Kubzansky LD. 2012; Childhood adversity and immune and inflammatory biomarkers associated with cardiovascular risk in youth: a systematic review. *Brain, Behavior, and Immunity*. 26(2):239–250. DOI: 10.1016/j.bbi.2011.11.003
- Slopen N, Kubzansky LD, McLaughlin KA, Koenen KC. 2013; Childhood adversity and inflammatory processes in youth: a prospective study. *Psychoneuroendocrinology*. 38(2):188–200. DOI: 10.1016/j.psyneuen.2012.05.013 [PubMed: 22727478]
- Somerville LH. 2013; The teenage brain: Sensitivity to social evaluation. *Current Directions in Psychological Science*. 22(2):121–127. DOI: 10.1177/0963721413476512 [PubMed: 24761055]

- Spear LP. 2009; Heightened stress responsivity and emotional reactivity during pubertal maturation: Implications for psychopathology. *Development and Psychopathology*. 21(01):87–97. DOI: 10.1017/S0954579409000066 [PubMed: 19144224]
- Stamatakis KA, Lynch JW, Everson SA, Raghunathan T, Salonen JT, Kaplan GA. 2004; Self-esteem and mortality: prospective evidence from a population-based study. *Annals of Epidemiology*. 14(1):58–65. DOI: 10.1016/S1047-2797(03)00078-4 [PubMed: 14664781]
- Stawski RS, Cichy KE, Piazza JR, Almeida DM. 2013; Associations among daily stressors and salivary cortisol: Findings from the National Study of Daily Experiences. *Psychoneuroendocrinology*. 38(11):2654–2665. DOI: 10.1016/j.psyneuen.2013.06.023 [PubMed: 23856186]
- Steptoe A, Hamer M, Chida Y. 2007; The effects of acute psychological stress on circulating inflammatory factors in humans: a review and meta-analysis. *Brain, Behavior, and Immunity*. 21(7):901–912. DOI: 10.1016/j.bbi.2007.03.011
- Stone AA, Neale JM, Cox DS, Napoli A, Valdimarsdottir H, Kennedy-Moore E. 1994; Daily events are associated with a secretory immune response to an oral antigen in men. *Health Psychology*. 13(5):440–446. [PubMed: 7805639]
- Stroud CB, Chen FR, Doane LD, Granger DA. 2016; Individual differences in early adolescents' latent trait cortisol (LTC): Relation to early adversity. *Developmental Psychobiology*. 58(6):700–713. DOI: 10.1002/dev.2141010.1002/dev.21410 [PubMed: 26990202]
- Stroud LR, Foster E, Papandonatos GD, Handwerker K, Granger DA, Kivlighan KT, Niaura R. 2009; Stress response and the adolescent transition: Performance versus peer rejection stressors. *Development and Psychopathology*. 21(1):47–68. DOI: 10.1017/S0954579409000042 [PubMed: 19144222]
- Surtees PG, Wainwright NWJ, Luben R, Khaw KT, Day NE. 2006; Mastery, sense of coherence, and mortality: evidence of independent associations from the EPIC-Norfolk Prospective Cohort Study. *Health Psychology*. 25(1):102–110. [PubMed: 16448303]
- Syed SA, Nemeroff CB. 2017; Early life stress, mood, and anxiety disorders. *Chronic Stress*. 1:1–16. DOI: 10.1177/2470547017694461
- Taylor SE, Broffman, JI. Psychosocial resources: Functions, origins, and links to mental and physical health. In: Olson, JM, Zanna, MP, editors. *Advances in Experimental Social Psychology*. Vol. 1–57. New York: Academic Press; 2011. 1
- Taylor SE, Lerner JS, Sage RM, Lehman BJ, Seeman TE. 2004; Early environment, emotions, responses to stress, and health. *Journal of Personality*. 72(6):1365–1394. DOI: 10.1111/j.1467-6494.2004.00300.x [PubMed: 15509286]
- Taylor SE, Seeman TE. 1999; Psychosocial resources and the SES-health relationship. *Annals of the New York Academy of Sciences*. 896(1):210–225. DOI: 10.1111/j.1749-6632.1999.tb08117.x [PubMed: 10681899]
- Taylor SE, Stanton AL. 2007; Coping resources, coping processes, and mental health. *Annual Review of Clinical Psychology*. 3:377–401. DOI: 10.1146/annurev.clinpsy.3.022806.091520
- Tottenham N, Galván A. 2016; Stress and the adolescent brain: Amygdala-prefrontal cortex circuitry and ventral striatum as developmental targets. *Neuroscience & Biobehavioral Reviews*. 70:217–227. DOI: 10.1016/j.neubiorev.2016.07.030 [PubMed: 27473936]
- Uhart M, Chong RY, Oswald L, Lin P, Wand GS. 2006; Gender differences in hypothalamic–pituitary–adrenal (HPA) axis reactivity. *Psychoneuroendocrinology*. 31(5):642–652. DOI: 10.1016/j.psyneuen.2006.02.003 [PubMed: 16616815]
- Viner RM, Ozer EM, Denny S, Marmot M, Resnick M, Fatusi A, Currie C. 2012; Adolescence and the social determinants of health. *The Lancet*. 379(9826):1641–1652. DOI: 10.1016/S0140-6736(12)60149-4
- Wellen KE, Hotamisligil GS. 2005; Inflammation, stress, and diabetes. *Journal of Clinical Investigation*. 115(5):1111–1119. DOI: 10.1172/JCI200525102 [PubMed: 15864338]




Figure 1.

(a) At lower levels of psychological resources, major life events are associated with decreased cortisol reactivity. (b) At higher levels of psychological resources, major life events are not associated with cortisol reactivity. Error bars reflect 95% confidence intervals.

Table 1

Sample characteristics and descriptive statistics of study variables.

Variable	Mean (SD)	Range
Age	18.37 (.51)	18.00 – 20.00
Female (%)	57%	
Latino (%)	63%	
Parent education	7.44 (2.01)	2.00 – 10.50
Major life events	3.15 (2.04)	0.00 – 12.00
Daily interpersonal stress (proportion of days)	0.24 (0.21)	0.00 – 0.93
Early adversity	1.92 (0.60)	1.00 – 3.92
Psychological resources	0.00 (0.78)	–2.47 – 1.26
Cortisol	13.03 (39.06)	–74.24 – 94.67
Baseline	16.84 (15.69)	3.51 – 111.16
Post-TSST	17.76 (13.81)	4.69 – 109.53
+30 minutes	24.60 (15.40)	5.68 – 90.47
+45 minutes	19.71 (11.95)	4.24 – 71.57
+60 minutes	15.82 (8.11)	4.09 – 45.41
+75 minutes	14.20 (6.98)	2.10 – 41.86
IL-6		
Baseline	1.47 (1.39)	0.34 – 7.53
+30 minutes	1.44 (1.29)	0.27 – 7.54
+60 minutes	1.70 (1.49)	0.35 – 9.28
+90 minutes	1.97 (1.65)	0.41 – 8.36

Note. Cortisol and IL-6 reflect raw, unwinsorized values.

Table 2

Bivariate correlations among primary study variables.

Variable	Major Life Events	Daily Interpersonal Stress	Early Adversity	Psychological Resources
Major life events				
Daily stress	0.28 ^{***}			
Early adversity	0.39 ^{***}	0.28 ^{**}		
Psychological resources	-0.33 ^{**}	-0.33 ^{**}	-0.49 ^{***}	
Cortisol: baseline	-0.01	-0.06	-0.10	0.11
Cortisol: post TSST	-0.04	-0.08	-0.28 ^{**}	0.11
Cortisol: +30 min	-0.20 [†]	-0.20 [†]	-0.28 ^{**}	0.08
Cortisol: +45 min	-0.19 [†]	-0.18	-0.31 ^{**}	0.13
Cortisol: +60 min	-0.17	-0.18	-0.26 [*]	0.15
Cortisol +90 min	-0.10	-0.08	-0.21 [*]	0.16
IL-6: baseline	-0.01	-0.14	0.01	0.05
IL-6: +30 min	-0.02	-0.12	0.002	-0.01
IL-6: +45 min	-0.02	-0.14	-0.03	-0.003
IL-6: +90 min	0.03	-0.12	-0.02	-0.02

Note.

† $p = .06 - .09$ * $p < .05$ ** $p < .01$ *** $p < .001$

Table 3

Results of models predicting cortisol from stress, psychological resources, and their interactions.

	Major Life Events		Daily Interpersonal Stress		Early Adversity	
	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2
	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)
Intercept	6.25 (0.92)***	6.31 (0.90)***	5.76 (0.94)***	5.99 (0.92)***	5.50 (0.88)***	5.67 (0.87)***
Gender	-0.17 (0.10) [†]	-0.20 (0.10)*	-0.16 (0.10)	-0.18 (0.10) [†]	-0.20 (0.09)*	-0.17 (0.09) [†]
Parent education	-0.004 (0.02)	-0.01 (0.02)	-0.003 (0.02)	-0.02 (0.02)	-0.01 (0.02)	-0.01 (0.02)
Time of day	-0.28 (0.07)***	-0.28 (0.07)***	-0.24 (0.07)***	-0.26 (0.07)***	-0.22 (0.06)***	-0.24 (0.06)***
Exercise	0.09 (0.04)*	0.09 (0.04)*	0.09 (0.04)*	0.09 (0.04)*	0.11 (0.04)*	0.12 (0.04)**
Smoking	0.23 (0.18)	0.23 (0.17)	0.20 (0.16)	0.22 (0.17)	0.27 (0.17)	0.23 (0.17)
Oral contraceptives	0.08 (0.11)	0.13 (0.12)	0.0002 (0.11)	0.04 (0.11)	-0.05 (0.10)	-0.07 (0.10)
Time	0.27 (0.03)***	0.31 (0.03)***	0.27 (0.03)***	0.27 (0.03)***	0.27 (0.03)***	0.29 (0.03)***
Time ²	-0.06 (0.01)***	-0.06 (0.01)***	-0.06 (0.01)***	-0.06 (0.01)***	-0.06 (0.004)***	-0.06 (0.01)***
Stress	-0.02 (0.03)	-0.02 (0.03)	-0.15 (0.31)	-0.19 (0.31)	-0.19 (0.12)	-0.20 (0.12)
Stress × Time	-0.04 (0.01)**	-0.04 (0.01)**	-0.34 (0.14)*	-0.29 (0.14)*	-0.17 (0.05)**	-0.16 (0.05)**
Stress × Time ²	0.01 (0.003)**	0.01 (0.003)**	0.06 (0.03)*	0.06 (0.03)*	0.03 (0.01)***	0.03 (0.01)**
Psychological resources	0.01 (0.09)	0.02 (0.09)	0.001 (0.09)	-0.001 (0.09)	-0.08 (0.09)	-0.08 (0.10)
Psychological resources × Time	-0.04 (0.04)	-0.06 (0.04)	-0.03 (0.04)	-0.06 (0.04)	-0.07 (0.04)	-0.09 (0.04)*
Psychological Resources × Time ²	0.01 (0.01)	0.01 (0.01) [†]	0.01 (0.01)	0.01 (0.01)	0.02 (0.01)*	0.02 (0.01)*
Stress × Psychological Resources		-0.04 (0.04)		-0.05 (0.37)		-0.03 (0.10)
Stress × Psychological Resources × Time		0.07 (0.02)***		0.41 (0.17)*		0.08 (0.05) [†]
Stress × Psychological Resources × Time ²		-0.01 (0.003)***		-0.05 (0.03) [†]		-0.01 (0.01)

Note.

Gender was coded as 0=male 1=female and ethnicity was coded as 0=European American, 1=Latino.

[†] $p = .06 - .08$ * $p < .05$

100' d

10' d
**

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4

Results of models predicting IL-6 from stress, psychological resources, and their interactions.

	Major Life Events		Daily Interpersonal Stress		Early Adversity	
	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2
	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)	<i>b</i> (<i>SE</i>)
Intercept	-0.51 (0.12)***	-0.50 (0.12)***	-0.50 (0.12)***	-0.49 (0.12)***	-0.49 (0.12)***	-0.51 (0.12)***
Age	-0.02 (0.11)	-0.001 (0.11)	-0.04 (0.11)	-0.02 (0.11)	-0.002 (0.11)	0.001 (0.11)
Gender	0.77 (0.11)***	0.76 (0.11)***	0.74 (0.11)***	0.74 (0.11)***	0.73 (0.11)***	0.73 (0.11)***
Ethnicity	0.21 (0.12) [†]	0.21 (0.12) [†]	0.22 (0.12) [†]	0.22 (0.12) [†]	0.22 (0.12) [†]	0.22 (0.12) [†]
Parent education	-0.02 (0.03)	-0.02 (0.03)	-0.03 (0.03)	-0.03 (0.03)	-0.02 (0.03)	-0.02 (0.03)
Waist circumference	0.02 (0.004)***	0.02 (0.004)***	0.02 (0.004)***	0.02 (0.004)***	0.02 (0.004)***	0.02 (0.004)***
Time	0.12 (0.01)***	0.12 (0.02)***	0.11 (0.01)***	0.11 (0.01)***	0.12 (0.01)***	0.12 (0.02)***
Stress	-0.03 (0.03)	-0.03 (0.03)	-0.37 (0.29)	-0.37 (0.29)	-0.09 (0.12)	-0.10 (0.12)
Stress × Time	0.003 (0.01)	0.003 (0.01)	-0.01 (0.07)	-0.001 (0.07)	-0.03 (0.03)	-0.03 (0.03)
Psychological resources	0.08 (0.08)	0.08 (0.08)	0.06 (0.08)	0.05 (0.08)	0.06 (0.09)	0.07 (0.09)
Psychological resources × Time	-0.01 (0.02)	-0.02 (0.02)	-0.02 (0.02)	-0.02 (0.02)	-0.03 (0.02)	-0.03 (0.02)
Stress × Psychological Resources		-0.003 (0.04)		0.16 (0.34)		-0.06 (0.09)
Stress × Psychological Resources × Time		0.02 (0.01)		0.09 (0.08)		0.03 (0.02)

Note.

Gender was coded as 0=male 1=female and ethnicity was coded as 0=European American, 1=Latino.

[†] *p* = .06 - .08* *p* .05** *p* .01*** *p* .001