

UCSF

UC San Francisco Previously Published Works

Title

Coping with oil spills: oil exposure and anxiety among residents of Gulf Coast states after the Deepwater Horizon Oil Spill.

Permalink

<https://escholarship.org/uc/item/0m7215c3>

Authors

Goldman, Zachary

Kaufman, John

Sharpe, J

et al.

Publication Date

2022





DOI

10.14324/111.444/ucloe.000035

Peer reviewed

RESEARCH ARTICLE

Coping with oil spills: oil exposure and anxiety among residents of Gulf Coast states after the Deepwater Horizon Oil Spill

Zachary E. Goldman¹, John A. Kaufman², J. Danielle Sharpe², Amy F. Wolkin³
and Matthew O. Gribble^{4*}

How to cite

Goldman ZE, Kaufman JA, Sharpe JD, Wolkin AF, Gribble MO. Coping with oil spills: oil exposure and anxiety among residents of Gulf Coast states after the Deepwater Horizon Oil Spill. *UCL Open: Environment*. 2022;(4):06. Available from: <https://doi.org/10.14324/111.444/ucloe.000035>

Submission date: 18 March 2021; Acceptance date: 12 April 2022; Publication date: 27 May 2022

Peer review

UCL Open: Environment is an open scholarship publication, all previous versions and open peer review reports can be found online in the *UCL Open: Environment* Preprint server at ucl.scienceopen.com

Copyright and open access

©2022 The Authors. Creative Commons Attribution Licence (CC BY) 4.0 International licence
<https://creativecommons.org/licenses/by/4.0/>

Open access

This is an open access article distributed under the terms of the Creative Commons Attribution Licence (CC BY) 4.0 <https://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.



*Corresponding author

Email: mgribble@uab.edu

¹Department of Environmental Health, Emory University Rollins School of Public Health, Atlanta, GA 30322, USA

²Department of Epidemiology, Emory University Rollins School of Public Health, Atlanta, GA 30322, USA

³National Center for Injury Prevention and Control, Centers for Disease Control and Prevention, Atlanta, GA 30345, USA

Abstract

In April 2010, a fatal explosion on the Deepwater Horizon drilling rig in the Gulf of Mexico resulted in the largest marine oil spill in history. This research describes the association of oil exposure with anxiety after the Deepwater Horizon Oil Spill and evaluates effect modification by self-mastery, emotional support and cleanup participation. To assess the impacts of the Deepwater Horizon Oil Spill, the Centers for Disease Control and Prevention (CDC) conducted the Gulf States Population Survey (GSPS), a random-digit-dial telephone cross-sectional survey completed between December 2010 and December 2011 with 38,361 responses in four different Gulf Coast states: Louisiana, Florida, Alabama and Mississippi. Anxiety severity was measured using the Generalised Anxiety Disorder (GAD) symptom inventory. We used Tobit regression to model underlying anxiety as a function of oil exposure and hypothesised effect modifiers, adjusting for socio-demographics. Latent anxiety was higher among those with direct contact with oil than among those who did not have direct contact with oil in confounder-adjusted models [$\beta = 2.84$, 95% confidence interval (CI): 0.78, 4.91]. Among individuals with direct contact with oil, there was no significant interaction between participating in cleanup activities and emotional support for anxiety ($p = 0.20$). However, among those with direct contact with oil, in confounder-adjusted models, participation in oil spill cleanup activities was associated with lower latent anxiety ($\beta = -3.55$, 95% CI: -6.15 , -0.95).

⁴Department of Epidemiology,
University of Alabama at
Birmingham School of Public
Health, Birmingham, AL 35294,
USA

Oil contact was associated with greater anxiety, but this association appeared to be mitigated by cleanup participation.

Keywords: generalised anxiety, disaster recovery, mental health, emergency response, Gulf States Population Survey (GSPS)

Introduction

In April 2010, the explosion of the Deepwater Horizon drilling rig in the Gulf of Mexico resulted in massive ecological and community harms leading to a \$4 billion plea agreement with BP Exploration & Production, a \$400 million criminal plea agreement with Transocean, a \$14.9 billion civil settlement with BP Exploration & Production and a \$159 million civil penalty ruling against Anadarko Petroleum Corporation [1]. Two thousand kilometres of coastline were impacted by the pollution from the spill [2], which was a complex mixture that often included oil as well as dispersants (chemicals that contain surfactants and/or solvent compounds) used in cleanup efforts [3,4]. There has been extensive research on the environmental impacts of the spill [5–7], as well as several public health impact assessments of nearby communities [8–11]. It was anticipated by public health authorities that there was potential for adverse mental health impacts on the affected community based on the experience of previous oil spills (e.g., smaller spills such as the *Hebei Spirit* tanker disaster in Korea [12] and *Exxon Valdez* in Alaska [13]), so to collect data on experiences pertaining to the oil spill and the mental health of the community in the wake of the Deepwater Horizon Oil Spill, the Substance Abuse and Mental Health Services Administration (SAMHSA) and the Centers for Disease Control and Prevention (CDC) developed the Gulf States Population Survey (GSPS) as a telephone survey administered to more than 38,000 adults of age ≥18 years old in affected Gulf Coast states (Louisiana, Mississippi, Alabama and Florida) from December 2010 to December 2011 [14].

The Deepwater Horizon Oil Spill had a large number of people involved in cleanup efforts [15–17]. There have been several investigations specifically focused on the mental health of participants in oil spill cleanup efforts following the Deepwater Horizon Spill, in particular evidence from the National Institute for Environmental Health Science's Gulf Long-Term Follow-Up Study (GuLF) [16,18,19] that found complex patterns of association that speak to multiple mediating pathways that might act in different directions on health outcomes and sometimes lead to net harms of cleanup participation [20,21]. However, a previous analysis of GSPS data focused on depression outcomes of cleanup participation among persons exposed to oil and found a potentially ameliorative association [22]. Several studies have focused on the associations between mental health and oil spill cleanup participation by comparing cleanup participants to populations not involved with the disaster [18,23,24]. Few studies have compared cleanup participants to non-cleanup participants among populations impacted by the disaster. A recent study on the associations of oil contact and depression found that respondents with oil contact had lower depression severity if they participated in cleanup activities, compared to exposed individuals who did not participate in cleanup activities [22].

The primary objective of this cross-sectional study was to evaluate whether oil exposure was associated with the severity of anxiety after the Deepwater Horizon Oil Spill in the GSPS, and assess effect modification of that association by self-mastery (coping ability), emotional support and cleanup participation. A secondary objective was to contribute to the evidence base regarding oil spill cleanup participation impacts on mental health among persons exposed to oil.

Methods

Study population

Following the Deepwater Horizon Oil Spill, the Substance Abuse and Mental Health Services Administration (SAMHSA) and the Centers for Disease Control and Prevention (CDC) published a report titled, *Behavioral Health in the Gulf Coast Region Following the Deepwater Horizon Oil Spill*, which described the behavioural health of residents impacted by the Deepwater Horizon Oil Spill [25]. Included in the report was the GSPS, a random-digit-dial telephone survey conducted between December 2010 and December 2011 to assess the impacts of the Deepwater Horizon Oil Spill [26]. The survey was created to provide information on the mental health status of the coastal

population in areas affected by the oil spill [14]. The GSPS, a publicly available, coded dataset, included 38,361 responses from individuals aged ≥ 18 years old residing in Louisiana, Florida, Alabama and Mississippi [14]. The majority of the interviews were conducted in 25 coastal counties or parishes within 32 miles (approximately 50 km) of fishing locations closed due to the oil spill [14]. To compare the results from coastal counties to non-coastal counties, 27,947 interviews were conducted in coastal counties and 10,414 interviews were conducted in non-coastal counties [14].

Data description

There were 16 questions found in the GSPS that measured contact with oil from the Deepwater Horizon Oil Spill. Direct oil contact was assessed by the question, 'Did you have direct contact with the oil from the Gulf oil spill? (yes/no)'. Questions from the 7-item Generalised Anxiety Disorder (GAD-7) questionnaire were also used to self-screen for moderate-to-severe cases of generalised anxiety [14]. Of the individuals contacted for the GSPS, 44.2% completed the interview [27]. The data for the GSPS were weighted to adjust for United States Census population estimates, and the survey design, sampling methods and weights were based on methods used for the Behavioral Risk Factor Surveillance System [25]. Analysis of this resource does not qualify as Human Subjects Research under 45 CFR 46.102 because this is a publicly available deidentified dataset.


Analysis

In the GSPS, the observed GAD-7 scores of survey respondents could range from 0 to 21 for anxiety symptom severity [14]. We conceptualised mental health as being on a continuous gradient and therefore assumed there to be left-censoring of 'negative' anxiety (e.g., calmness, confidence) and right-censoring of extreme anxiety by the GAD-7 psychometric instrument. Tobit regression is a tool for modelling a latent dependent variable assumed to be normally distributed whose true values are censored by the measurement instrument [28]. Therefore, under the assumption that latent anxiety in the population was normally distributed rather than bounded between 0 and 21, Tobit regression was used to model underlying latent anxiety as a normally distributed latent variable. Figure 1 presents a histogram of observed GAD-7 scores along with modelled Tobit latent anxiety severity.

We adjusted models for age (continuous), gender (male/female), race (White/Black/other), Hispanic ethnicity (yes/no), smoking status (never/former/current), binge drinking (yes if over previous 30 days had consumed ≥ 5 drinks on one occasion if male or ≥ 4 drinks on one occasion if female), participating in any form of exercise (yes/no), marital status (married/not currently married) and employment status [employed (employed for wages, self-employed)/not employed (out of work,

Figure 1

Latent anxiety and observed anxiety symptom scores in the GSPS.



unable to work)/other (homemaker, student, retired)]. We further added interaction terms with how often people felt they had the emotional support they needed (always/usually/sometimes/rarely or never) and perceived self-mastery, a confirmatory factor analysis score derived from five items (strength of agreement with 'I have little control over the things that happen to me', 'What happens to me in the future mostly depends on me', 'I can do just about anything I really set my mind to do', 'I am confident in my ability to handle unexpected problems' and 'When I need suggestions about how to deal with a personal problem, I know there is someone I can turn to') [29].

In fully adjusted models, we tested for an interaction between direct contact with oil and participation in oil spill cleanup efforts. We also examined the association of cleanup participation with anxiety, controlling for the same confounders among the subpopulation with direct oil exposure, and tested for interactions with emotional support and self-mastery.

Survey estimation methods were used to account for the survey design with singleton primary sampling units treated as certainty units [30,31]. Missing data were handled by multiple imputation by chained equations with 80 imputations using all modelled variables (GAD-7 score, race, age, Hispanic ethnicity, direct oil contact, marital status, employment status, exercise, binge drinking, self-mastery, oil cleanup participation, smoking status and level of emotional support) [32]. All analyses were conducted using Stata/IC 16.1 (StataCorp, College Station, TX, USA).

Results

Population and sample demographics

Characteristics of the population are provided in Table 1, according to whether they had direct contact with oil from the Deepwater Horizon Oil Spill and whether they participated in cleanup activities. These estimates incorporate multiple imputation and survey estimation for population inference. Characteristics of the sample population according to direct oil contact and participation in cleanup activities during the Deepwater Horizon Oil Spill are provided in Table 2.

Oil contact with anxiety

In fully adjusted models, latent anxiety was higher ($\beta = 2.84$, 95% confidence interval (CI): 0.78, 4.91, t -test p value = 0.01) among individuals with directly contact with oil compared to individuals not with direct contact to oil (Table 3). There was no heterogeneity in the oil-anxiety association across levels of emotional support (always, usually, sometimes and rarely or never) (3-degrees-of-freedom F test: p value = 0.38), with the lowest β -coefficient of 0.96 (95% CI: -0.84, 2.76) for 'usually supported' and the highest β -coefficient of 4.72 (95% CI: 0.53, 8.92) for 'always supported' (Table 4). There was also no significant modification of the association between oil exposure and anxiety by self-mastery (t -test p value = 0.12). Among those with direct contact to oil, individuals with lower self-mastery scores (minimum in sample: -4.04) had higher latent anxiety ($\beta = 9.15$, 95% CI: 0.03, 18.28) compared to individuals with higher self-mastery scores (maximum in sample: 3.31) ($\beta = -3.13$, 95% CI: -9.72, 3.45).

Cleanup participation and anxiety among those with oil contact

There was, however, a significant antagonistic interaction between direct oil exposure and participation in cleanup activities for latent anxiety (t -test p value < 0.01). Among individuals with direct contact with oil in fully adjusted models, cleanup participation was associated with lower latent anxiety ($\beta = -3.55$, 95% CI: -6.15, -0.95) (Table 5). Among individuals with direct contact with oil, there was no significant interaction between participating in cleanup activities and levels of emotional support (always, usually, sometimes, rarely or never) (3-degrees-of-freedom F test: p value = 0.20), with the lowest β -coefficient of -4.08 (-8.53, 0.37) for 'usually' and the highest β -coefficient of 1.55 (-2.59, 5.68) for 'sometimes' (Table 6).

Among those with direct contact to oil and who participated in cleanup activities, individuals with lower self-mastery scores (minimum in sample: -4.04) had less latent anxiety ($\beta = -12.87$, 95% CI: -24.98, -0.76) compared to individuals with higher self-mastery scores (maximum in sample: 3.31) ($\beta = 4.69$, 95% CI: -5.66, 15.05). Thus, although the interaction term between cleanup participation

Table 1. Estimated characteristics (point estimates and 95% CI) of the population according to direct oil contact and participation in cleanup activities during the Deepwater Horizon oil spill. These estimates incorporate MI and survey estimation for population inference

Characteristic	Direct oil contact	No direct oil contact	Participated in cleanup activities	Did not participate in cleanup activities
Gender (%)				
Female	46 (38, 53)	52 (50, 54)	44 (33, 54)	52 (50, 54)
Male	54 (47, 62)	48 (46, 50)	56 (46, 67)	48 (46, 50)
Mean age (years)	43 (40, 46)	49 (48, 49)	39 (35, 42)	49 (48, 50)
Mean GAD-7 score	6 (5, 7)	4 (3, 4)	3 (3, 4)	4 (3, 4)
Race (%)				
White	76 (67, 85)	69 (67, 72)	72 (61, 83)	70 (67, 72)
Black	11 (7, 14)	20 (19, 22)	18 (8, 28)	20 (18, 22)
Other	13 (4, 22)	10 (8, 12)	10 (4, 17)	10 (9, 12)
Hispanic ethnicity (%)				
Yes	13 (4, 22)	11 (9, 13)	8 (1, 15)	12 (10, 13)
No	87 (78, 96)	89 (87, 91)	92 (85, 99)	88 (87, 90)
Employment (%)				
Employed	60 (52, 68)	52 (50, 54)	64 (53, 75)	52 (50, 54)
Not employed	16 (10, 21)	15 (14, 17)	12 (6, 18)	16 (14, 17)
Other	24 (18, 31)	32 (30, 35)	24 (13, 35)	32 (30, 34)
Marital status (%)				
Married	51 (43, 59)	52 (50, 54)	43 (32, 53)	52 (50, 55)
Not married	49 (41, 57)	48 (46, 50)	57 (47, 68)	48 (45, 50)
Exercise (%)				
Yes	82 (76, 88)	73 (71, 75)	89 (83, 95)	73 (71, 75)
No	18 (12, 24)	27 (25, 29)	11 (5, 17)	27 (25, 29)
Binge drinking (%)				
Yes	23 (17, 28)	15 (13, 16)	29 (19, 38)	15 (13, 16)
No	77 (72, 83)	85 (84, 87)	71 (62, 81)	85 (84, 87)
Smoking (%)				
Never	52 (44, 60)	55 (52, 57)	50 (39, 60)	55 (52, 57)
Former	21 (15, 27)	24 (22, 26)	22 (13, 32)	24 (22, 25)
Current	27 (20, 33)	22 (20, 24)	28 (19, 37)	22 (20, 24)
Emotional support (%)				
Always	40 (32, 48)	50 (48, 52)	53 (42, 63)	49 (47, 51)
Usually	33 (26, 40)	27 (25, 29)	32 (23, 41)	27 (25, 29)
Sometimes	16 (11, 21)	14 (13, 16)	6 (2, 10)	15 (13, 16)
Rarely or never	11 (6, 16)	9 (7, 10)	9 (3, 14)	9 (7, 10)

Abbreviations: CI: confidence intervals; MI, multiple imputation.

with self-mastery among those with direct contact with oil was not statistically significant (t -test p value = 0.12), there is suggestive evidence that participation may have had a larger benefit among individuals with lower self-mastery.

Discussion

Oil contact and increased anxiety

This study found positive associations between oil contact and anxiety among residents of Gulf States after the Deepwater Horizon Oil Spill. These results are consistent with previous findings from studies examining associations between oil exposure and acute mental health impacts [11,18,22,33,34]. The implications for longer-term mental health are unclear; one recent study found that physical exposure to the Deepwater Horizon Oil Spill was not associated with measures of psychological resiliency for several years among adults with children [35], while another study found that later mental health in the community might be a function of cumulative exposure to other disasters as well (e.g., Hurricane Katrina) not only the Deepwater Horizon Oil Spill [36]. Our findings that exposed individuals with lower self-mastery scores had greater mental health impacts also agree with studies demonstrating the positive associations between high levels of self-mastery and improved mental health [37].

Table 2. Characteristics of the sample stratified by direct oil contact and participation in cleanup activities during the Deepwater Horizon oil spill. The numbers in rows that show (%) are N and percent, the numbers without (%) are mean and SD

Characteristic	Direct oil contact	No direct oil contact	Participated in cleanup activities	Did not participate in cleanup activities
Gender (%)				
Female	1,985 (55%)	21,300 (63%)	740 (47%)	22,698 (63%)
Male	1,640 (45%)	12,578 (37%)	835 (53%)	13,459 (37%)
Mean age (years)	3,625 (50)	33,878 (56)	1,575 (47)	36,157 (56)
Mean GAD-7 score	3,539 (5)	33,108 (3)	1,548 (4)	35,318 (3)
Race (%)				
White	2,983 (82%)	26,160 (77%)	1,289 (82%)	28,005 (78%)
Black	428 (12%)	5,910 (17%)	172 (11%)	6,234 (17%)
Other	210 (6%)	1,751 (5%)	112 (7%)	1,860 (5%)
Hispanic ethnicity (%)				
Yes	3,496 (97%)	32,749 (97%)	1,518 (97%)	34,948 (97%)
No	111 (3%)	962 (3%)	52 (3%)	1,024 (3%)
Employment (%)				
Employed	2,063 (57%)	15,510 (46%)	1,033 (66%)	16,654 (46%)
Not employed	615 (17%)	4,423 (13%)	215 (14%)	4,879 (14%)
Other	938 (26%)	13,856 (41%)	323 (21%)	14,531 (40%)
Marital status (%)				
Married	2,114 (59%)	17,709 (52%)	882 (56%)	19,045 (53%)
Not married	1,496 (41%)	16,027 (48%)	683 (44%)	16,963 (47%)
Exercise (%)				
Yes	2,868 (80%)	24,251 (72%)	1,299 (83%)	25,992 (72%)
No	732 (20%)	9,400 (28%)	265 (17%)	9,925 (28%)
Binge drinking (%)				
Yes	2,857 (80%)	29,550 (88%)	1,229 (80%)	31,366 (88%)
No	706 (20%)	3,889 (12%)	314 (20%)	4,322 (12%)
Smoking (%)				
Never	1,754 (49%)	17,927 (54%)	777 (50%)	19,018 (53%)
Former	939 (26%)	9,003 (27%)	368 (24%)	9,627 (27%)
Current	882 (25%)	6,513 (19%)	405 (26%)	7,053 (20%)
Emotional support (%)				
Always	1,570 (44%)	16,622 (50%)	743 (48%)	17,556 (49%)
Usually	1,114 (31%)	9,526 (28%)	494 (32%)	10,203 (29%)
Sometimes	543 (15%)	4,864 (15%)	208 (13%)	5,237 (15%)
Rarely or never	35 (10%)	2,490 (7%)	119 (8%)	2,760 (8%)

Table 3. Differences in anxiety (β) with direct contact with oil in communities participating in the GSPS

Model	β (95% CI)
Model 1: Unadjusted association	3.51 (1.79, 5.24)
Model 2: Adjusted for age, race, Hispanic ethnicity, gender	3.35 (1.60, 5.09)
Model 3: Further adjusted for exercise, smoking, and binge drinking	3.42 (1.61, 5.24)
Model 4: Further adjusted for employment and marital status	3.32 (1.50, 5.16)
Model 5: Further adjusted for self-mastery and emotional support	2.84 (0.78, 4.91)

Abbreviations: CI, confidence interval; GSPS, Gulf States Population Survey.

Table 4. Differences in anxiety (β) with direct contact with oil, by frequency of emotional support, in communities participating in the GSPS

Emotional support	Proportion of population (95% CI)	β (95% CI)
Always	0.49 (0.47, 0.52)	4.72 (0.53, 8.92)
Usually	0.27 (0.26, 0.29)	0.96 (-0.84, 2.76)
Sometimes	0.15 (0.13, 0.16)	1.89 (-0.46, 4.23)
Rarely or never	0.09 (0.07, 0.10)	3.10 (-0.51, 6.71)

Abbreviations: CI, confidence interval; GSPS, Gulf States Population Survey.

Table 5. Differences in anxiety (β) between individuals participating in oil spill cleanup activities compared to individuals not participating in oil spill cleanup activities, among individuals with direct contact with oil, in communities participating in the GSPS

Model	β (95% CI) among persons with direct oil contact	β (95% CI) among persons without direct oil contact
Model 1: Unadjusted association	-3.48 (-6.81, -0.15)	-0.35 (-2.30, 1.60)
Model 2: Adjusted for age, race, Hispanic ethnicity, gender	-3.18 (-6.16, -0.19)	-0.83 (-2.91, 1.24)
Model 3: Further adjusted for exercise, smoking, and binge drinking	-3.25 (-6.16, -0.33)	-0.56 (-2.49, 1.37)
Model 4: Further adjusted for employment and marital status	-3.47 (-6.12, -0.82)	-0.30 (-2.19, 1.60)
Model 5: Further adjusted for self-mastery and emotional support	-3.55 (-6.15, -0.95)	0.30 (-1.56, 2.17)

Abbreviations: CI, confidence interval; GSPS, Gulf States Population Survey.

Table 6. Associations between oil spill cleanup participation and anxiety, by frequency of emotional support, in communities participating in the GSPS

Emotional support	Proportion of population (95% CI)	β (95% CI)
Always	0.49 (0.47, 0.52)	-3.83 (-7.47, -0.19)
Usually	0.27 (0.26, 0.29)	-4.08 (-8.53, 0.37)
Sometimes	0.15 (0.13, 0.16)	1.55 (-2.59, 5.68)
Rarely or never	0.09 (0.07, 0.10)	-3.31 (-9.03, 2.40)

Abbreviations: CI, confidence interval; GSPS, Gulf States Population Survey.

Oil contact, cleanup participation and anxiety

We found that the association between oil exposure and latent anxiety is diminished among participants in oil cleanup activities. To the best of our knowledge, this is the first study to examine associations between anxiety and participation in oil spill cleanup activities within populations directly affected by the oil spill. These findings agree with a recent study examining the association between depression and participating in oil spill cleanup activities compared to not participating in these activities within populations directly affected by the oil spill [22]. This previous study also found associations between oil contact and mental health effects that were attenuated among the persons who participated in oil spill cleanup efforts [22].

Volunteering, cleanup participation and mental health; similar findings in literature

Although this paper is the first to examine the relationship between oil exposure and anxiety by cleanup participation, prior studies have shown associations between general volunteering and mental health. In a longitudinal study conducted in the United Kingdom, individuals who engaged in volunteering regularly appeared to have higher levels of mental well-being compared to those who never volunteered [38]. A 2017 cross-sectional study found that combat veterans who volunteered in disaster relief social service organisations reported positive mental health responses as a result of helping those in need [39]. A previous study examining public servants working in the area affected by the Great East Japan Earthquake of 2011 also reported that engaging in disaster-related work may reduce mental health distress among workers [40].

Study strengths and limitations

The main strengths of this paper were the use of a large, representative sample and the ability to compare different populations affected by the oil spill. However, because this was a cross-sectional survey, there was potential for reverse causality between cleanup participation and anxiety symptoms, both assessed after the oil spill [20]. It is also unknown whether respondents who participated in oil spill cleanup activities were healthier at baseline and had fewer mental health conditions. As the GSPS was a random-digit-dial telephone survey, the associations may introduce bias due to potential effects of recall bias or social desirability bias among the study participants [41].

Although the associations between cleanup participation and anxiety were adjusted for several factors related to anxiety and socio-demographics, confounding by unmeasured variables is possible. A recent study examining determinants of participating in oil spill cleanup activities

found positive associations between cleanup participation and proximity to affected areas [15]. Approximately 4.7% of residents in affected Gulf communities participated in the oil spill cleanup activities, and most were young men in excellent physical health [15]. These individuals residing in affected communities may have been easily mobilised and motivated to help their affected communities by participating in cleanup activities [15].

Implications and conclusions

It is possible that volunteering in disaster relief efforts could serve as an ameliorative intervention for mental health impacts after disasters occur. By responding to major disasters and working as a community, a sense of belonging or contribution may create a protective response for mental health [40]. Additionally, it has been shown that individuals who experience a disaster may benefit from the resources of their social networks [42]. We think it is plausible that participation in disaster-related response activities may give people a feeling of partial control over the consequences of an event that was otherwise out of their control. Increasing participation in disaster-related cleanup activities could also increase resilience and recovery of individuals and the surrounding community [15].

Although this cross-sectional study is limited in its ability to draw firm causal inferences, the findings from this paper raise the possibility that encouraging affected individuals to volunteer during oil spill disasters may mitigate the harmful effects of oil exposure. Further research, such as prospective evidence on the impacts of post-oil exposure cleanup participation on mental health outcomes, is needed.

Acknowledgements

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.

Declarations and conflicts of interest

Research ethics statement

This was a deidentified public dataset and so according to United States Department of Health and Human Services Regulation 45 CFR 46.102(f) this is Research Not Involving Human Subjects.

Consent for publication statement

Not applicable to this article.

Conflicts of interest statement

Dr. Gribble is a current Editor for this journal. All authors declare no conflict of interest with the content of this work.

Data availability statement

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.


Funding

Dr. Gribble's efforts were supported in part by funding from the National Institute of Environmental Health Sciences (P30 ES019776).

References

- [1] United States Environmental Protection Agency. Enforcement. Deepwater Horizon – BP Gulf of Mexico Oil Spill. [Accessed 23 October 2021]. Available from: <https://www.epa.gov/enforcement/deepwater-horizon-bp-gulf-mexico-oil-spill#settlement>
- [2] Kujawinski EB, Reddy CM, Rodgers RP, Thrash JC, Valentine DL, White HK. The first decade of scientific insights from the Deepwater Horizon oil release. *Nat Rev Earth Environ*. 2020;1(5):237–50.
- [3] Allan SE, Smith BW, Anderson KA. Impact of the Deepwater Horizon oil spill on bioavailable polycyclic aromatic hydrocarbons in gulf of mexico coastal waters. *Environ Sci Technol*. 2012;46:2033–9.

- [4] EPA. Dispersing Agents. [Accessed 10 October 2021]. Available from: <https://www.epa.gov/emergency-response/dispersing-agents>
- [5] Beyer J, Trannum HC, Bakke T, Hodson PV, Collier TK. Environmental effects of the Deepwater Horizon oil spill: a review. *Mar Pollut Bull.* 2016;110(1):28–51.
- [6] Zengel S, Weaver J, Mendelssohn IA, Graham SA, Lin Q, Hester MW, et al. Meta-analysis of salt marsh vegetation impacts and recovery: a synthesis following the Deepwater Horizon oil spill. *Ecol Appl.* 2022;32:e02489.
- [7] Ward CP, Overton EB. How the 2010 Deepwater Horizon spill reshaped our understanding of crude oil photochemical weathering at sea: a past, present, and future perspective. *Environ Sci Process Impacts.* 2020;22:1125–38.
- [8] Buttke D, Vagi S, Schnall A, Bayleyegn T, Morrison M, Allen M, et al. Community assessment for public health emergency response (CASPER) one year following the Gulf Coast oil spill: Alabama and Mississippi, 2011. *Prehosp Disaster Med.* 2012;27(6):496–502.
- [9] Buttke D, Vagi S, Bayleyegn T, Sircar K, Strine T, Morrison M, et al. Mental health needs assessment after the Gulf Coast oil spill—Alabama and Mississippi, 2010. *Prehosp Disaster Med.* 2012;27(5):401–408.
- [10] Peres LC, Trapido E, Rung AL, Harrington DJ, Oral E, Fang Z, et al. The Deepwater Horizon Oil Spill and physical health among adult women in southern Louisiana: The Women and Their Children's Health (WaTCH) study. *Environ Health Perspect.* 2016;124(8):1208–13.
- [11] Osofsky HJ, Osofsky JD, Hansel TC. Deepwater Horizon oil spill: mental health effects on residents in heavily affected areas. *Disaster Med Public Health Prep.* 2011;5(4):280–6.
- [12] Song M, Hong YC, Cheong HK, Ha M, Kwon H, Ha EH, et al. [Psychological health in residents participating in clean-up works of Hebei Spirit oil spill]. *J Prev Med Public Health.* 2009;42(2):82–8.
- [13] Palinkas LA. A conceptual framework for understanding the mental health impacts of oil spills: lessons from the Exxon Valdez oil spill. *Psychiatry.* 2012;75:203–22.
- [14] Substance Abuse and Mental Health Services Administration (SAMHSA) and the Centers for Disease Control and Prevention. *Behavioral Health in the Gulf Coast Region Following the Deepwater Horizon Oil Spill.* [Accessed 6 February 2020]. Available from: <https://www.samhsa.gov/data/sites/default/files/NSDUH-GSPS-GulfCoast-Apps-2012/NSDUH-GSPS-GulfCoast-2012.pdf>
- [15] Sharpe JD, Kaufman JA, Goldman ZE, Wolkin A, Gribble MO. Determinants of oil-spill cleanup participation following the Deepwater Horizon oil spill. *Environ Res.* 2019;170:472–80.
- [16] Kwok RK, Engel LS, Miller AK, Blair A, Curry MD, Jackson WB, et al. The GuLF STUDY: a prospective study of persons involved in the Deepwater Horizon oil spill response and clean-up. *Environ Health Perspect.* 2017;125(4):570–8.
- [17] Rusiecki J, Alexander M, Schwartz EG, Wang L, Weems L, Barrett J, et al. The Deepwater Horizon Oil Spill Coast Guard Cohort Study. *Occup Environ Med.* 2018;75(3):165–75.
- [18] Kwok RK, McGrath JA, Lowe SR, Engel LS, Jackson WB 2nd, Curry MD, et al. Mental health indicators associated with oil spill response and clean-up: cross-sectional analysis of the GuLF STUDY cohort. *Lancet Public Health.* 2017;2(12):e560–7.
- [19] Lowe SR, Kwok RK, Payne J, Engel LS, Galea S, Sandler DP. Mental health service use by cleanup workers in the aftermath of the Deepwater Horizon oil spill. *Soc Sci Med.* 2015;130:125–34.
- [20] Lowe SR, Kwok RK, Payne J, Engel LS, Galea S, Sandler DP. Why does disaster recovery work influence mental health?: Pathways through physical health and household income. *Am J Community Psychol.* 2016;58(3–4):354–64.
- [21] O'Rourke HP, MacKinnon DP. Reasons for testing mediation in the absence of an intervention effect: a research imperative in prevention and intervention research. *J Stud Alcohol Drugs.* 2018;79:171–81.
- [22] Kaufman JA, Goldman ZE, Sharpe JD, Wolkin AF, Gribble MO. Mechanisms of resiliency against depression following the Deepwater Horizon oil spill. *J Environ Psychol.* 2019;65:101329.
- [23] Brooks SK, Dunn R, Amlôt R, Greenberg N, Rubin GJ. Social and occupational factors associated with psychological distress and disorder among disaster responders: a systematic review. *BMC Psychol.* 2016;4(1):18.
- [24] Benedek DM, Fullerton C, Ursano RJ. First responders: mental health consequences of natural and human-made disasters for public health and public safety workers. *Annu Rev Public Health.* 2007;28:55–68.
- [25] Gould DW, Teich JL, Pemberton MR, Pierannunzi C, Larson S. Behavioral health in the Gulf Coast region following the Deepwater Horizon oil spill: findings from two federal surveys. *J Behav Health Serv Res.* 2015;42(1):6–22.
- [26] United States Centers for Disease Control and Prevention. CDC Mental Illness Surveillance. Gulf States Population Survey. Retrieved via Wayback Machine. [Accessed 23 October 2021]. Archived copy available from: <http://web.archive.org/web/20150907184930/www.cdc.gov/mentalhealthsurveillance/gsp-data.html>
- [27] Fan AZ, Prescott MR, Zhao G, Gotway CA, Galea S. Individual and community-level determinants of mental and physical health after the Deepwater Horizon oil spill: findings from the Gulf States Population Survey. *J Behav Health Serv Res.* 2015;42(1):23–41.
- [28] Tobin J. Estimation of relationships for limited dependent variables. *Econometrica.* 1958;26(1):24–36.
- [29] Pearlin LI, Schooler C. The structure of coping. *J Health Soc Behav.* 1978;19(1):2–21.
- [30] Korn EL, Graubard BI. Simultaneous testing of regression coefficients with complex survey data: use of Bonferroni t statistics. *Am Stat.* 1990;44(4):270–6.
- [31] West BT, Berglund P, Heeringa SG. A closer examination of subpopulation analysis of complex-sample survey data. *Stata J.* 2008;8:520–31.
- [32] Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple imputation by chained equations: what is it and how does it work? *Int J Methods Psychiatr Res.* 2011;20(1):40–9.
- [33] Grattan LM, Roberts S, Mahan WT Jr, McLaughlin PK, Otwell WS, Morris JG Jr. The early psychological impacts of the Deepwater Horizon oil spill on Florida and Alabama communities. *Environ Health Perspect.* 2011;119(6):838–43.
- [34] Gill DA, Picou JS, Ritchie LA. The Exxon Valdez and BP oil spills: a comparison of initial social and psychological impacts. *Am Behav Sci.* 2012;56(1):3–23.
- [35] Slack T, Kroeger RA, Stroope S, Keating KS, Beedasy J, Chandler T, et al. Disparate effects of BP Deepwater Horizon oil spill exposure on psychological resilience. *Traumatology.* 2021. [Accessed 18 March 2021]. Available from: <https://psycnet.apa.org/record/2021-98452-001?doi=1>
- [36] Lowe SR, McGrath JA, Young MN, Kwok RK, Engel LS, Galea S, et al. Cumulative disaster exposure and



Extra information

UCL Open: Environment is an open scholarship publication, all previous versions and open peer review reports can be found online in the *UCL Open: Environment Preprint* server at ucl.scienceopen.com

- mental and physical health symptoms among a large sample of Gulf Coast residents. *J Trauma Stress*. 2019;32(2):196–205.
- [37] Rabani Bavojdan M, Towhidi A, Rahmati A. The relationship between mental health and general self-efficacy beliefs, coping strategies and locus of control in male drug abusers. *Addict Health*. 2011;3(3–4):111–8.
- [38] Tabassum F, Mohan J, Smith P. Association of volunteering with mental well-being: a lifecourse analysis of a national population-based longitudinal study in the UK. *BMJ Open*. 2016;6:e011327.
- [39] Kranke D, Weiss EL, Heslin KC, Dobalian A. 'We are disaster response experts': a qualitative study on the mental health impact of volunteering in disaster settings among combat veterans. *Soc Work Public Health*. 2017;32:500–9.
- [40] Fukasawa M, Suzuki Y, Obara A, Kim Y. Relationships between mental health distress and work-related factors among prefectural public servants two months after the Great East Japan Earthquake. *Int J Behav Med*. 2015;22:1–10.
- [41] Mazor KM, Clauser BE, Field T, Yood RA, Gurwitz JH. A demonstration of the impact of response bias on the results of patient satisfaction surveys. *Health Serv Res*. 2002;37(5):1403–17.
- [42] Parks V, Ayer L, Ramchand R, Finucane ML. Disaster experience, social capitals, and behavioral health. *Natural Hazards*. 2020;104(1):959–77.