

Lawrence Berkeley National Laboratory

Recent Work

Title

MEASUREMENT OF THE PANOFSKY RATIO IN ^3He

Permalink

<https://escholarship.org/uc/item/0m87x2n0>

Authors

Truol, Peter
Baer, Helmet W.
Bistirlich, James A.
et al.

Publication Date

1973-10-01

MEASUREMENT OF THE PANOFSKY RATIO IN ^3He


Peter Truöl, Helmut W. Baer, James A. Bistirlich,
Kenneth M. Crowe, Nico de Botton and Jerome A. Helland

October 1973

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48

For Reference

Not to be taken from this room



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

MEASUREMENT OF THE PANOFSKY RATIO IN ${}^3\text{He}^\dagger$

Peter Truöl*

Physik-Institut der Universität Zürich
Zürich, Switzerland

and

Helmut W. Baer, James A. Bistirlich, Kenneth M. Crowe,
Nico de Botton,[‡] and Jerome A. Helland[§]

Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

ABSTRACT

The photon spectrum from radiative and charge-exchange capture of pions in ${}^3\text{He}$ was measured in a high-resolution pair-spectrometer yielding the new value of the Panofsky ratio $P_3 = (\pi^- + {}^3\text{He} \rightarrow {}^3\text{H} + \pi^0) / (\pi^- + {}^3\text{He} \rightarrow {}^3\text{H} + \gamma) = 2.68 \pm 0.13$. An impulse approximation analysis is presented which gives a value $P_3 = 2.6$. In addition, the branching ratios for the ${}^3\text{He} + \pi^0$, ${}^3\text{H} + \gamma$, and (${}^2\text{H} + n + \gamma$ and $p + n + n + \gamma$) channels are measured to be 17.8 ± 2.3 , 6.6 ± 0.8 , and $7.4 \pm 1.0\%$ respectively.

The absorption of negative pions from atomic orbits around free protons proceeds almost exclusively via the charge-exchange reaction $\pi^- + p \rightarrow n + \pi^0$ and the radiative capture reaction $\pi^- + p \rightarrow \gamma + n$. The ratio of the transition rates for these two processes, the so-called Panofsky ratio,¹ links pion-photoproduction at threshold to pion-nucleon scattering and provides a determination of the pion-nucleon coupling strength. The equivalent ratio for protons bound in nuclei has been observed only in ${}^3\text{He}$, where an earlier measurement by Zaïmidoroga et al.,² yielded $P_3 = (\pi^- + {}^3\text{He} \rightarrow \pi^0 + {}^3\text{H}) / (\pi^- + {}^3\text{He} \rightarrow \gamma + {}^3\text{H}) = 2.28 \pm 0.18$. Some authors^{3,4} have regarded this quantity as a test case in the application of the PCAC hypothesis to soft-pion problems involving complex nuclei. Other authors employ the impulse approximation (IA) directly and relate the Panofsky ratio for ${}^3\text{He}$ to ${}^1\text{H}$. By making this assumption it has been shown^{5,6} that P_3 depends primarily one parameter, viz., the ${}^3\text{He}$ - ${}^3\text{H}$ rms transition radius. The value $\langle r^2 \rangle_{3\text{He}}^{1/2} = 1.4 \pm 0.2 \text{ F}$ extracted from $P_3 = 2.28 \pm 0.18$ disagrees with the value $1.88 \pm 0.05 \text{ F}$ determined⁷ by electron scattering. In view of the importance of this quantity both to the study of the elementary particle approach to nuclei as well as the structure of the mass-3 system and possible 3-body forces it was thought desirable to remeasure this quantity and to study directly the radiative breakup reactions $\pi^- + {}^3\text{He} \rightarrow d + n + \gamma$ and $p + n + n + \gamma$.

The experiment was performed in the stopped- π beam of the Lawrence Berkeley Laboratory 184-inch cyclotron. Details of the experimental setup are given in Ref. 8. A π^- beam is brought to rest in a 9.5-cm-diameter, 12.7-cm-long Mylar flask (0.02 cm wall thickness) filled with liquid ${}^3\text{He}$ at 1.9°K. A typical rate was 3×10^4 π^- /sec stopping

in the helium content of the target. The photons were detected in a 180° pair spectrometer employing three wire spark chambers. The energy resolution is 2 MeV (FWHM) at 130 MeV; the acceptance (Fig. 1a) ($\text{conversion efficiency} \times \Delta\Omega/4\pi$) was determined in a Monte Carlo calculation.⁸ An independent calibration of the instrument was achieved with a hydrogen target. Since the relative efficiency at low energy was found to depend to some extent on the spark chamber performance and could vary by as much as 15-20%,⁹ a total of 15 hydrogen runs were taken during the experiment. Despite the changes in the relative efficiency we find a constant value for the Panofsky ratio in ^3He when compared with hydrogen.

The $^3\text{He}(\pi^-, \gamma)$ spectrum (Fig. 1b) exhibits the expected four photon channels: $t\gamma$ with $E_\gamma = 135.8$ MeV; $dn\gamma$ and $pnn\gamma$ with endpoint energies of 129.8 and 127.7 MeV respectively; and $t\pi^0, \pi^0 \rightarrow 2\gamma$ with a uniform distribution between 53.1 and 85.7 MeV. There is a suggestion of a broad peak corresponding to 10- to 15 MeV excitation in the ^3H system (Fig. 1c), although the statistical evidence for the state proposed by Chang *et al.*¹¹ is inconclusive. The two breakup channels cannot be separated from each other, but their separation from the $t\gamma$ reaction can be achieved reliably by shifting the hydrogen line by 6.35 MeV and normalizing to ^3He events above 130 MeV. Separation of the small contribution ($\sim 4\%$) which the breakup reactions make to the charge-exchange peak ($E_\gamma < 90$ MeV) was performed with a pole-model^{8,10} calculation (Fig. 1c). Comparing (Table I) results with ones obtained in Ref. 2, we find that the difference in the Panofsky ratio stems mainly from difference in the charge-exchange yields, since the radiative yields agree very well. This seems under-

standable, since the small kinetic energy of the recoil triton (190 keV) may cause difficulties in observing them in diffusion chambers. This point is discussed by the authors in Ref. 2.

Analyses of radiative π^- capture in light nuclei are in general complicated by the fact that a large fraction of pions gets captured from the 2p Bohr orbit. The Panofsky ratio in ^3He , however, appears to be very nearly independent of 2p-state capture. Estimates³ for the fraction (pions captured)/(pions making 2p \rightarrow 1s x-ray transition) range up to 55%. However Ericson and Figureau³ estimate that only 0.1% and 0.03% of pions captured from the 2p orbit undergo charge exchange (CEX) and radiative (REX) capture, respectively. Thus the measured Panofsky ratio should be given quite accurately by the relative 1s-capture CEX/REX matrix elements.

The transition rates in the IA are given for radiative π^- capture¹² by

$$\Lambda_{\gamma}(1s) = \frac{1}{4\pi} \frac{k}{m_{\pi}} C^2 \left(1 - \frac{k}{m_3 + m_{\pi}}\right) \left(1 + \frac{m_{\pi}}{m_n}\right)^2 |\phi_{\pi}(0)|^2 |M|^2 \quad \text{with}$$

$$|M|^2 = \frac{1}{2J_i + 1} \sum_{m_i, m_f, \lambda} \int \frac{d\Omega_{\hat{k}}}{4\pi} |\langle J_f M_f | \sum_{j=1}^3 (\hat{\epsilon}_{\lambda} \cdot \vec{\sigma}_j) \tau_j^{(-)} e^{-i\vec{k} \cdot \vec{r}_j} | J_i M_i \rangle|^2$$

and for charge-exchange³ by

$$\Lambda_{\pi^0}(1s) = \frac{1}{4\pi} \frac{q_0}{m_{\pi}} A^2 \left(1 - \frac{q_0}{m_3 + m_{\pi}}\right) \left(1 + \frac{m_{\pi}}{m_n}\right)^2 |\phi_{\pi}(0)|^2 |M_0|^2 \quad \text{with}$$

$$|M_0|^2 = \frac{1}{2J_i + 1} \sum_{m_i, m_f} \int \frac{d\Omega_{\hat{q}}}{4\pi} |\langle J_f M_f | \sum_{j=1}^3 \tau_j^{(-)} e^{-i\vec{q}_0 \cdot \vec{r}_j} | J_i M_i \rangle|^2$$

[$\hbar = c = 1$; $m_3 = \text{mass}(^3\text{He})$; $m_n = \text{mass}(^1\text{H})$; $(\omega_0, \vec{q}_0) = \pi_0$ four momentum; $\vec{k}, \hat{\epsilon}$ = photon momentum, polarization]. It is assumed that the pion wave function may be taken out of the matrix element and replaced by its value at the origin with a small correction³ for the extended charge distribution $|\phi_\pi(0)|^2 = (0.97/\pi)(Z\alpha m_\pi)^3 (1 + m_\pi/m_n)^{-3}$, where $\alpha = 1/137$. The value of C is determined by pion photoproduction cross sections at threshold and has the value¹³ $C = 4\pi |E_0^{(\pi^-)}| = 4\pi (3.15 \pm 0.06) \times 10^{-2}/m_\pi$. A is related to the πN isospin singlet and triplet scattering lengths¹³ by $A = (4\pi\sqrt{2}/3) |a_1 - a_3|$.

For radiative capture, $|M|^2$ is related to the axial form factors of the mass-3 system and the nucleon and the Gamow-Teller matrix element $M_{GT} = \langle ^3\text{H} | \sum_{j=1}^3 \tau_j^{(-)} \sigma_j | ^3\text{He} \rangle$ by¹⁴

$$|M|^2 = \frac{2}{3} |M_{GT}|^2 \left(\frac{F_A^{^3\text{He} \rightarrow ^3\text{H}}(q^2)}{F_A^{^3\text{He} \rightarrow ^3\text{H}(0)}} \right)^2 \left(\frac{F_A^{n \rightarrow p(0)}}{F_A^{n \rightarrow p(q^2)}} \right)^2$$

The GT matrix element has the value $|M_{GT}|^2 = 3$ if ^3He and ^3H are exact mirror states with totally space-symmetric wave functions, whereas the experimental value is $|M_{GT}|^2 = 2.84 \pm 0.06$ measured¹⁵ in β^- decay. A value of $F_A^{^3\text{He} \rightarrow ^3\text{H}}(0.474)/F_A^{^3\text{He} \rightarrow ^3\text{H}(0)} = 0.80$ [Ref. 3] is found from linear extrapolation of μ -capture (4-momentum transfer, $q^2 = 0.27 F^{-2}$) to π -capture in good agreement with 0.776 ± 0.016 based on the assumption¹⁴ $F_A(q^2)/F_A(0) = F_M(q^2)/F_M(0)$, where $F_M(q^2)$ is the $^3\text{He} \rightarrow ^3\text{H}$ magnetic form factor determined using ^3He and ^3H electron scattering data.⁷ For the nucleon¹³ the latter procedure gives

$F_A^{n \rightarrow p}(0.474)/F_A^{n \rightarrow p}(0) = 0.948$. Combining the last two results with the experimental value for $|M_{GT}|^2$ gives $\lambda_Y(1s) = 3.41 \times 10^{15} \text{ sec}^{-1}$.

The charge-exchange matrix element $|M_O|^2$ is related to the vector-form factors and Fermi matrix element by¹⁴

$$|M_O|^2 = |M_F|^2 \times \left(\frac{F_V^{3\text{He} \rightarrow 3\text{H}(q^2)}}{F_V^{3\text{He} \rightarrow 3\text{H}(0)}} \right)^2 \left(\frac{F_V^{n \rightarrow p(0)}}{F_V^{n \rightarrow p(q^2)}} \right)^2 = 1 \times .974,$$

applying the CVC hypothesis to both nucleon and mass-3 form factors.

The Panofsky ratio for ^3He expressed in terms of the same quantity for hydrogen is

$$P_3 = 2P_1 \frac{q_{03}}{k_3} \frac{k_1}{q_{01}} \frac{m_3 + m_\pi - \omega_{03}}{m_3 + m_\pi - k_3} \frac{m_n + m_\pi - k_1}{m_n + m_\pi - \omega_{01}} \frac{|M_O|^2}{|M|^2}.$$

Inserting the experimental value¹ $P_1 = 1.533 \pm 0.021$ yields $P_3 (= 3.42 |M_O|^2 / |M|^2) = 2.63$, in good agreement with our measured value $P_3 = 2.68 \pm 0.13$.

Since P_1 appears in the evaluation of the experimental value for P_3 as well as in the theoretical expression, our result is independent of the particular value for P_1 chosen, and can therefore be considered a direct test of the IA in s-wave pion-nucleus interactions. The good agreement with the IA calculation has to be contrasted with the poor agreement with recent values for P_3 obtained by current-algebra methods. Ericson and Figureau³ obtain values between 1.9 and 2.1, depending on whether the CEX cross section is calculated in IA or in the soft-pion technique. In this calculation the electric-dipole amplitude

$|E_{0+}(\pi^-)|$ in the nucleon case gets replaced by the soft-pion value $\sqrt{\alpha/4\pi} (1/f_\pi) (g_A/g_V)$. When the elementary amplitude is applied to the nuclear case,⁴ corrections for ρ -meson exchange, incoherent rescattering, and nuclear intermediate states are included. The corrections have the effect of increasing the radiative rate to $4.43 \times 10^{15} \text{ sec}^{-1}$ and thereby reducing P_3 . It would appear therefore that these corrections are smaller than estimated. Other calculations along these lines,¹³ where, however, terms first order in m_π/m_n are neglected, give values for the radiative rate around $2.3 \times 10^{15} \text{ sec}^{-1}$.^{18,19}

We wish to thank Professor N. Straumann and Dr. G. Nixon for clarifying discussions of previous theoretical work.

Footnotes and References

† Work done under the auspices of the U. S. Atomic Energy Commission.

* Part of the work was done while at the Lawrence Berkeley Laboratory.

‡ Permanent address: DPHN/HE, C.E.N. Saclay, BP n^o2, 91-Gif sur Yvette, France.

§ Present address: Los Alamos Scientific Laboratory, University of California, Los Alamos, New Mexico.

1. W. Panofsky et al., Phys. Rev. 81, 565 (1951); V. Cocconi et al., Nuovo Cimento 22, 494 (1961).
2. O. Zaimidoroga et al., Sov. Phys. JETP 21, 848 (1965) and 24, 1111 (1967).
3. M. Ericson and A. Figureau, Nucl. Phys. B3, 609 (1967) and B11, 621 (1969).
4. M. Ericson, M. Rho, Phys. Rep. 5C, 59 (1972).
5. A. Fujii, D. Hall, Nucl. Phys. 32, 102 (1962).
6. B. Struminsky, Int. Conf. High-Energy Phys., CERN(1962), p.17.
7. J. M. McCarthy, Phys. Rev. Lett. 25, 884 (1970);
H. Collard et al., Phys. Rev. B138, 57 (1965).
8. H. Baer et al., Lawrence Berkeley Report LBL-1737 (Phys. Rev. C, in press).
9. This effect was not fully realized in the first part of the run. This is the reason that our preliminary analysis based on 25% of our data yielded the higher value $P_3 = 3.5 \pm 0.4$ [H. Baer et al., Int. Conf. Few Part. Prob. in Nucl. Interaction, Los Angeles, 1972 (North-Holland, Amsterdam, 1973), p. 877.]

10. L. Dakhno and Yu. Prokoshkin, Sov. J. Nucl. Phys. 7, 351 (1968).
11. C. Chang et al., Phys. Rev. Lett. 29, 307 (1972).
12. J. Delorme and T. Ericson, Phys. Lett. 21, 98 (1966).
13. G. Ebel et al., Nucl. Phys. B33, 317 (1971).
14. C. Kim and H. Primakoff, Phys. Rev. 140B, 566 (1965).
15. R. Salgo and H. Staub, Nucl. Phys. A138, 417 (1969).
16. P. Divakaran, Phys. Rev. 139, B387 (1965).
17. Previous calculations yielded values $8.32 \times 10^{15} \text{ sec}^{-1}$ (Ref. 5) and $0.97 \times 10^{15} \text{ sec}^{-1}$ (Ref. 16.) Differences with our value are due to missing factors of $m_{\pi}/2k$ in Ref. 5, 4 in Ref. 16, and the older value $r_{\text{rms}}(^3\text{He}) = 1.55 \text{ F}$.
18. D. Griffiths and C. Kim, Phys. Rev. 173, 1584 (1968);
L. Fulcher and J. Eisenberg, Nucl. Phys. B18, 271 (1970).
19. P. Pascual and A. Fujii, Nuovo Cimento 65, 411 (1970);
these authors obtain $\lambda_{\gamma}(1s) = 3.37 \times 10^{15} \text{ sec}^{-1}$ and $P_3 = 2.2$.
After introducing a missing factor of 2/3 in the expression for $\lambda_{\gamma}(1s)$, one obtains $\lambda_{\gamma}(1s) = 2.25 \times 10^{15} \text{ sec}^{-1}$ and $P_3 = 3.3$.

TABLE I. Results for stopped- π^- absorption on ^3He and ^1H .

Final state	N_γ^a	R^b (%)	R^c (%)
$^3\text{H } \pi^0$	6273 ± 82	17.8 ± 2.3	15.8 ± 0.8^2
$^3\text{H } \gamma$	5580 ± 157	6.6 ± 0.8	6.9 ± 0.5^2
$\text{dn}\gamma + \text{pnny}$	5331 ± 137	7.4 ± 1.0	$[3.6 \pm 1.2]^{d,2}$
$n \pi^0 (^1\text{H})$	2355 ± 49	65.6 ± 11.1	60.5 ± 0.3^1
$n \gamma (^1\text{H})$	3860 ± 62	42.4 ± 4.4	39.5 ± 0.3^1
dn		$[68.2 \pm 2.6]$	15.9 ± 2.3^2
pnn			57.8 ± 5.4^2
$P_3(^3\text{He})^e$		2.68 ± 0.13^f	2.28 ± 0.18^2
$P_1(^1\text{H})^e$		1.54 ± 0.26	1.533 ± 0.021^1
B_3^g		1.12 ± 0.05	
C_3^h		10.3 ± 1.3	10.7 ± 1.2^2

^a Raw number of events in spectrum. ^b This experiment.


^c Previous experiment. ^d $\text{dn}\gamma$ only. ^e Panofsky ratio. ^f $P_3 = P_1$

$$\begin{aligned}
 & (= 1.533 \pm 0.021) \times \frac{N_\gamma(^3\text{H}\pi^0)}{N_\gamma(^3\text{H}\gamma)} \times \frac{N_\gamma(n\gamma)}{N_\gamma(n\pi^0)} \times (1-f), \text{ where } f = (5.3 \pm 2.0) \times 10^{-2}
 \end{aligned}$$

is small correction for difference in photon detection efficiency for ^3He and ^1H . ^g $\sigma[\pi^- + ^3\text{He} \rightarrow (\text{dn}\gamma + \text{pnny})] / \sigma[\pi^- + ^3\text{He} \rightarrow ^3\text{H} + \gamma]$. ^h Ratio of nucleon ejection modes to radiative absorption.

FIGURE CAPTION

- Fig. 1. (a) Hydrogen spectrum and pair spectrometer acceptance.
(b) ${}^3\text{He}$ spectrum, 50-150 MeV.
(c) ${}^3\text{He}$ spectrum in region where the breakup channels dominate. The curve is a pole model calculation^{8,10} ($\Delta = 6.8$ MeV) with complete kinematics incorporated.



LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720