

# UCLA

## UCLA Previously Published Works

### Title

A multi-stage genome-wide association study of uterine fibroids in African Americans

### Permalink

<https://escholarship.org/uc/item/0mc5r0xh>

### Journal

Human Genetics, 136(10)

### ISSN

0340-6717

### Authors

Hellwege, Jacklyn N

Jeff, Janina M

Wise, Lauren A

et al.

### Publication Date

2017-10-01

### DOI

10.1007/s00439-017-1836-1

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# A multi-stage genome-wide association study of uterine fibroids in African Americans

Jacklyn N. Hellwege<sup>1,2,3</sup> · Janina M. Jeff<sup>4</sup> · Lauren A. Wise<sup>5,6</sup> · C. Scott Gallagher<sup>7</sup> · Melissa Wellons<sup>8,9</sup> · Katherine E. Hartmann<sup>3,9</sup> · Sarah F. Jones<sup>1,3</sup> · Eric S. Torstenson<sup>1,2</sup> · Scott Dickinson<sup>10</sup> · Edward A. Ruiz-Narváez<sup>6</sup> · Nadin Rohland<sup>7</sup> · Alexander Allen<sup>7</sup> · David Reich<sup>7,11,12</sup> · Arti Tandon<sup>7</sup> · Bogdan Pasaniuc<sup>13,14</sup> · Nicholas Mancuso<sup>13</sup> · Hae Kyung Im<sup>10</sup> · David A. Hinds<sup>15</sup> · Julie R. Palmer<sup>6</sup> · Lynn Rosenberg<sup>6</sup> · Joshua C. Denny<sup>16,17</sup> · Dan M. Roden<sup>2,16,17,18</sup> · Elizabeth A. Stewart<sup>19</sup> · Cynthia C. Morton<sup>12,20,21,22</sup> · Eimear E. Kenny<sup>4</sup> · Todd L. Edwards<sup>1,2,3</sup> · Digna R. Velez Edwards<sup>2,3,9</sup>

Received: 12 April 2017 / Accepted: 16 August 2017 / Published online: 23 August 2017  
© Springer-Verlag GmbH Germany 2017

**Abstract** Uterine fibroids are benign tumors of the uterus affecting up to 77% of women by menopause. They are the leading indication for hysterectomy, and account for \$34 billion annually in the United States. Race/ethnicity and age are the strongest known risk factors. African American (AA) women have higher prevalence, earlier onset, and larger and more numerous fibroids than European American women. We conducted a multi-stage genome-wide association study (GWAS) of fibroid risk among AA women followed by in silico genetically predicted gene expression profiling of top hits. In Stage 1, cases and controls were confirmed by pelvic

imaging, genotyped and imputed to 1000 Genomes. Stage 2 used self-reported fibroid and GWAS data from 23andMe, Inc. and the Black Women's Health Study. Associations with fibroid risk were modeled using logistic regression adjusted for principal components, followed by meta-analysis of results. We observed a significant association among 3399 AA cases and 4764 AA controls at rs739187 (risk-allele frequency = 0.27) in *CYTH4* (OR (95% confidence interval) = 1.23 (1.16–1.30),  $p$  value =  $7.82 \times 10^{-9}$ ). Evaluation of the genetic association results with MetaXcan identified lower predicted gene expression of *CYTH4* in thyroid tissue as significantly associated with fibroid risk ( $p$  value =  $5.86 \times 10^{-8}$ ). In this first multi-stage GWAS for fibroids among AA women, we identified a novel risk locus

**Electronic supplementary material** The online version of this article (doi:10.1007/s00439-017-1836-1) contains supplementary material, which is available to authorized users.

✉ Digna R. Velez Edwards  
digna.r.velez.edwards@vanderbilt.edu

- <sup>1</sup> Division of Epidemiology, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA
- <sup>2</sup> Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Nashville, TN, USA
- <sup>3</sup> Institute for Medicine and Public Health, Vanderbilt University Medical Center, Nashville, TN, USA
- <sup>4</sup> Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA
- <sup>5</sup> Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA
- <sup>6</sup> Slone Epidemiology Center at Boston University, Boston, MA, USA
- <sup>7</sup> Department of Genetics, Harvard Medical School, Boston, MA, USA
- <sup>8</sup> Division of Diabetes, Endocrinology and Metabolism, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

- <sup>9</sup> Department of Obstetrics and Gynecology, Vanderbilt University Medical Center, Nashville, TN, USA
- <sup>10</sup> Section of Genetic Medicine, The University of Chicago, Chicago, IL, USA
- <sup>11</sup> Howard Hughes Medical Institute, Chevy Chase, MD, USA
- <sup>12</sup> Broad Institute of Harvard and MIT, Cambridge, MA, USA
- <sup>13</sup> Department of Pathology and Laboratory Medicine, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, USA
- <sup>14</sup> Department of Human Genetics, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, USA
- <sup>15</sup> 23andMe, Inc., Mountain View, CA, USA
- <sup>16</sup> Department of Biomedical Informatics, Vanderbilt University School of Medicine, Nashville, TN, USA
- <sup>17</sup> Department of Medicine, Vanderbilt University School of Medicine, Nashville, TN, USA
- <sup>18</sup> Department of Pharmacology, Vanderbilt University School of Medicine, Nashville, TN, USA

for fibroids within *CYTH4* that impacts gene expression in thyroid and has potential biological relevance for fibroids.

## Introduction

Uterine fibroids or leiomyomata are common, benign tumors of the uterus with an estimated lifetime risk of 77% by menopause (Baird et al. 2003). African Americans (AA) are more likely to have fibroids than women of European ancestry (EA), with AA having greater than 80% incidence of fibroids by menopause compared to nearly 70% for EA (Baird et al. 2003). AA women also have larger and more numerous fibroids as well as a younger age-of-onset on average (Baird et al. 2003). In addition to race/ethnicity (Baird et al. 2003; Faerstein et al. 2001; Marshall et al. 1997), there are other well-characterized risk factors for fibroids, including early age at menarche (Baird and Dunson 2003; Cha et al. 2011; Faerstein et al. 2001; Luoto et al. 2000; Moore et al. 2008), being overweight (BMI 25–29 kg/m<sup>2</sup>) (Baird et al. 2007; Moore et al. 2008; Terry et al. 2007; Wise et al. 2005a), and older premenopausal age (Baird et al. 2003; Moore et al. 2008). In addition, higher parity is associated with reduced fibroid risk, likely due to pregnancy-related hormonal and uterine changes (Baird and Dunson 2003). Symptoms of uterine fibroids may include pelvic pain and abnormal or heavy menses, though many fibroids are asymptomatic (Baird et al. 2003; Borah et al. 2013; Vollenhoven 1998). The lack of overt symptoms makes imaging crucial for classification of case/control status, as up to 51% of women may be misclassified by self-report (Baird et al. 2003; Myers et al. 2012). We have developed and validated a phenotyping algorithm to classify fibroid case/control status using electronic health records (Feingold-Link et al. 2014). This algorithm requires pelvic imaging for identification of cases and controls, which reduces misclassification of both cases and controls.

Several lines of evidence suggest genetic risk factors influence fibroid development. Estimates of the heritability of fibroids from familial aggregation and twin studies range from 26 to 69% in European populations (Luoto et al. 2000). Racial disparities in age at onset, number, size, and lifetime

incidence of fibroids by menopause (Baird et al. 2003) also strongly support a role for genetic factors. There has been one published genome-wide association study (GWAS) of uterine fibroids, which was performed among Japanese subjects (Cha et al. 2011). Genome-wide linkage and follow-up association studies in a meta-analysis of EA women implicated an additional locus for risk of fibroid diagnosis (Egger et al. 2012). The loci implicated in these previous studies (*SLK*, *BET1*, *TNRC6B*, and *FASN/CCDC57*) have been replicated among EAs (Aissani et al. 2015a; Edwards et al. 2013b), which have been evaluated but failed to replicate in AAs (Aissani et al. 2015b; Wise et al. 2012). The predominant studies conducted among AA subjects have involved admixture mapping, which has shown significant regions of increased African ancestry in cases, particularly at 1q42.2 (Zhang et al. 2015), 4p16, and 10q26 (Wise et al. 2012). The aim of this work was to perform the first GWAS for fibroids risk among AA women using image-verified fibroids for discovery, with replication in cohorts that collected data on self-reported fibroids.

## Methods

### Study populations

Individuals with imaging-confirmed uterine fibroids and genome-wide genotype data were included from the following studies in Stage 1: Vanderbilt University BioVU, Mt Sinai, BioME and the Coronary Artery Risk Development in Young Adults (CARDIA) Women's Study (CARDIA-WS) for a total of 1273 cases and 1379 controls. All studies received Institutional Review Board approval at their respective institutions and written informed consent was obtained for all participants.

BioVU is an electronic health record (EHR) biorepository at Vanderbilt University Medical Center, Nashville, TN and was designed to link clinical data available from de-identified EHRs to DNA specimens. Methods have been previously described (Pulley et al. 2010). BioME is the ongoing, consented EHR-linked biobank at the Institute for Personalized Medicine at the Icahn School of Medicine at Mt. Sinai. A subset of the available BioME samples were genotyped as part of the Electronic Medical Records and Genetics (eMERGE) Network and are referred to in this instance as Mt. Sinai, while genotyped samples acquired through other means were analyzed separately and will be henceforth referred to as BioME. The phenotyping algorithms used to identify cases and controls in BioVU, Mt. Sinai, and BioME have been previously published (Feingold-Link et al. 2014). Briefly, the phenotyping algorithm required at least one instance of pelvic imaging with a diagnosis code for fibroids to define cases among women aged

<sup>19</sup> Departments of Obstetrics and Gynecology and Surgery, Mayo Clinic and Mayo Clinic School of Medicine, Rochester, MN, USA

<sup>20</sup> University of Manchester Academic Health Science Centre, Manchester, England, UK

<sup>21</sup> Department of Obstetrics, Gynecology and Reproductive Biology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

<sup>22</sup> Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

18 and over. For controls, at least two instances of pelvic imaging were required, on different dates, with no diagnosis or procedure codes indicating a fibroid, fibroid treatment, or hysterectomy. The CARDIA study is a prospective multi-center study with 5115 adult European ancestry and AA participants of the age group 18 to 30 years, recruited from four centers. Details of the CARDIA study design have been previously published (Friedman et al. 1988). The CARDIA Women's Study performed standardized study ultrasounds on women from CARDIA to detect the presence or absence of fibroids, as previously described (Wellons et al. 2008).

Stage 2 data were provided by the direct-to-consumer genetic testing company 23andMe, which used a survey for determining case or control status. Specifically, cases were defined as females responding positively to either of the following questions: "Has a doctor ever told you that you have uterine fibroids?", or "Have you ever been diagnosed with uterine fibroids?". Controls were defined as females responding with a 'no' to either of the questions above. Age at enrollment with 23andMe was also collected (mean ages of 56 in cases and 47 in controls).

In addition, in Stage 2, GWAS samples were provided by the Black Women's Health Study (BWHS), a U.S. prospective cohort study of premenopausal women (aged 22–50 years) in which uterine fibroid diagnoses were ascertained by self-report, with medical record validation among a subset of cases (Wise et al. 2004, 2005b). BWHS participants reported whether they had been diagnosed with


fibroids on the follow-up questionnaires every 2 years (1997–2009). The women reported the calendar year of their first diagnosis, and whether the diagnosis was confirmed by ultrasound or surgery. Controls were premenopausal women who reported no diagnosis of fibroids through 2009. Among those who released their medical records, 96% of self-reported fibroid cases were confirmed (Wise et al. 2005b). The study workflow is diagrammed in Fig. 1.

## Genotyping

BioVU samples were genotyped using both the Affymetrix BioBank array and the Axiom World array 2 (Affymetrix Inc., Santa Clara, CA, USA) to attain better coverage for African-derived variants. BioME and the Mt. Sinai eMERGE site used the Illumina 1 M (Illumina Inc., San Diego, CA, USA) platform, with an exome chip backbone included for BioME samples. CARDIA-WS was genotyped on the Affymetrix 6.0 array (Affymetrix Inc., Santa Clara, CA, USA) as part of the Candidate-gene Association Resource which has been previously described in detail (Lettre et al. 2011).

The Stage 2 23andMe samples (1744 cases and 2906 controls) were genotyped on a custom GWAS panel across four versions. Stage 2 participants from BWHS (382 cases and 392 controls) were genotyped on the Illumina Infinium Expanded Multi-Ethnic Genotyping Array (MEGA; Illumina, Inc., San Diego, CA, USA) at Vanderbilt University VANTAGE Core Genotyping facility.

**Fig. 1** Study workflow diagram



## Quality control

Genotype quality control was performed within each study population, and a uniform protocol was implemented for all Stage 1 studies using PLINK (Purcell et al. 2007), including a 95% SNP and individual call rate threshold, removal of first-degree related individuals, sex checks, alignment of alleles to the genomic ‘+’ strand, and visualization of ancestry by principal components analysis using the Eigenstrat software (Price et al. 2006). No samples were excluded from analyses based on principal components.

For 23andMe in Stage 2, SNPs with Hardy–Weinberg equilibrium  $p$  value  $<10^{-20}$  were excluded, as were those with call rate  $<95\%$ , or with large allele frequency discrepancies compared to European 1000 Genomes reference data. For BWHS in Stage 2, genotype quality control was also performed in PLINK and consisted of 95% SNP and 90% individual call rate thresholds, removal of first-degree related individuals, sex checks, alignment of alleles to the genomic ‘+’ strand, and visualization of ancestry by principal components analysis using the Eigenstrat software (Price et al. 2006). No samples were excluded from analyses based on principal components.

## Statistical analysis

All demographic data were summarized and evaluated with Stata 14.0 (College Station, TX, USA).

The Stage 1 data were imputed to the 1000 Genomes phase 3 reference panel using SHAPEIT2 (Delaneau et al. 2012) for haplotype phasing and IMPUTE2 (Duan et al. 2013) for genotype imputation, with phasing and imputation performed separately for each data set. SNPTEST2 (Marchini et al. 2007) was used to perform single-variant association analyses for each cohort independently. Analyses were adjusted for the first five principal components in each sample to account for population substructure (Supplemental Fig. 1). Additional analysis adjusting for age was also performed, as was an analysis to avoid misclassification errors by restricting controls to peri-menopausal women (age  $\geq 45$ ) to remove women who were at increased risk of developing fibroids.

The Stage 2 data from 23andMe were imputed to the March 2012 “v3” release of 1000 Genomes reference haplotypes. Phasing and imputation were performed for each genotyping platform version separately, using Beagle4 (version 3.3.1) to phase batches of 8000–9000 individuals across chromosomal segments of no more than 10,000 genotyped single nucleotide polymorphisms (SNPs), with overlaps of 200 SNPs, and a high-performance version of Minimac5 for imputation of each phased segment against all-ethnicity 1000 Genomes haplotypes, using five rounds and 200 states for parameter estimation. Logistic regression was modeled

using an additive genetic model adjusting for age and five principal components, reporting likelihood ratio test  $p$  values for association with genotypes.

Stage 2 also utilized 776 samples from BWHS which were imputed to the 1000 Genomes phase 3 reference panel using SHAPEIT2 (Delaneau et al. 2012) for haplotype phasing and IMPUTE2 (Duan et al. 2013) for genotype imputation, with phasing and imputation performed separately for each data set. SNPTEST2 (Marchini et al. 2007) was used to perform single-variant association analyses. Association analysis was performed using logistic regression of the imputed genotype data. Analyses were adjusted for ten principal components.

All association results for Stage 1 and across Stages 1–2 were meta-analyzed using fixed effects inverse-variance weighted meta-analysis in METAL (Willer et al. 2010). The final meta-analysis of all variants including all stages consisted of 3399 cases and 4764 controls.

To further evaluate the genetic association results in the context of gene expression, we employed the novel method MetaXcan (Barbeira et al. 2016), an extension of the PrediXcan method (Gamazon et al. 2015). PrediXcan conducts a test of association between phenotypes and gene expression levels predicted by genetic variants in a library of tissues from the Genotype-Tissue Expression (GTEx) project (2015; Carithers et al. 2015; Mele et al. 2015). MetaXcan is a meta-analysis approach that conducts the PrediXcan test using genotype association summary statistics, rather than performing the tests in individual-level data. For the purposes of this study, we utilized covariance matrices built for nine relevant tissues from GTEx (i.e., uterus, ovary, breast, vagina, subcutaneous adipose, visceral omentum adipose, thyroid, whole blood, and transformed fibroblasts) to annotate SNP association signals as well as to provide information about likely tissue expression patterns and relevant biological information.

## Results

We used our previously validated EHR algorithm to define 999 image-confirmed cases and 1233 image-confirmed controls for fibroids from BioVU and Mt. Sinai (including the Mt. Sinai eMERGE subset as well as other cases and controls from the remainder of the BioME resource) sites. We also included additional samples from the CARDIA Women’s Study, which incorporated a standardized research ultrasound, for a total of 1273 case and 1379 control samples for analysis in Stage 1 discovery analyses (Fig. 1). Descriptive characteristics of each study are presented in Table 1. Briefly, cases and controls were of similar ages within each individual study in Stage 1 (no more than 4 year difference in means), though the Mt. Sinai sample was substantially

**Table 1** Demographic characteristics of study populations

Study	Cases (N)	Controls (N)	Genotyping platform	Age <sup>a</sup> Mean (SD) Cases	Age <sup>b</sup> Mean (SD) Controls
BioVU AA	578	804	Affymetrix Biobank Array and World Array 3	40 (11)	41 (15)
Mt Sinai (eMERGE)	74	78	Illumina Omni 1M	59 (10)	63 (12)
BioME	347	351	Illumina Omni 1M	48 (10)	45 (16)
CARDIA-WS	274	146	Affymetrix 6.0	40 (4) <sup>c</sup>	39 (4)
Total Stage 1	1273	1379			
23andMe	1744	2903	Custom Array	56 (12) <sup>b</sup>	47 (15)
BWHS	382	392	Illumina MEGA	36 (6)	32 (6)
Total Stage 2	2126	3295			
Total Stage 1 + 2	3399	4764			

<sup>a</sup>Age at diagnosis unless otherwise specified<sup>b</sup>Age at enrollment<sup>c</sup>Age at CARDIA-WS standardized study ultrasound

older on average than the other three studies. However, comparing ages across studies is challenging due to the differing study designs (case–control vs. cohort) and incident periods.

All data sets were genotyped on study-specific genome-wide genotyping arrays, imputed to the 1000 Genomes reference panel, and analyzed for association with fibroids, adjusting for principal components of ancestry (Supplemental Table 1). Meta-analysis of both Stages (3399 cases and 4764 controls) revealed a single genome-wide significant locus near *CYTH4* (cytohesin 4) on chromosome 22 (Table 2; bottom panel of Fig. 2). The lead SNP at this locus, rs739187 [ $p$  value =  $7.82 \times 10^{-9}$ , OR [95% confidence

interval (CI)] = 1.23 (1.16–1.30)] had low heterogeneity across cohorts.

To further evaluate the genetic association results in the context of gene expression, we employed the novel method MetaXcan which conducts a test of association between phenotypes and gene expression levels predicted by genetic variants in a library of tissues from the GTEx project (GTEx Consortium et al. 2015; Carithers et al. 2015; Mele et al. 2015). Analysis with MetaXcan to evaluate association with genetically predicted gene expression (GPGE) levels resulted in 1542 results across 9 GTEx tissues with  $p$  value <0.05, from a total of 44,577 comparisons (Table 3; upper

**Table 2** Summary of genome-wide significant and suggestive ( $<5 \times 10^{-7}$ ) associations in GWAS meta-analysis

Chr. <sup>a</sup>	Position	Nearest gene	SNP	EA/RA <sup>b</sup>	EAF <sup>c</sup>	Stage 1 + 2 (N = 3399/4764)				
						OR (95% CI) <sup>d</sup>	P value	Direction	$P_{\text{het}}$	$I^2$
22	37,728,254	<i>CYTH4</i>	rs739187	T/C	0.27	1.23 (1.16–1.30)	<b>7.83E–09</b>	++++++	0.24	26.0
22	37,722,301	<i>CYTH4</i>	rs4821628	A/G	0.75	0.81 (0.74–0.88)	<b>8.86E–09</b>	-----	0.22	29.0
22	37,726,660	<i>CYTH4</i>	rs713939	T/C	0.28	1.22 (1.15–1.29)	<b>1.37E–08</b>	++++++	0.30	17.8
22	37,719,004	<i>CYTH4</i>	rs5995416	T/C	0.29	1.21 (1.14–1.28)	<b>3.44E–08</b>	++++++	0.40	1.3
22	37,717,946	<i>CYTH4</i>	rs4821627	T/C	0.29	1.21 (1.14–1.27)	5.86E–08	++++++	0.33	13.1
3	5,348,595	<i>EDEMI</i>	rs55768811	A/T	0.91	0.76 (0.66–0.87)	2.62E–07	-----	0.72	0
8	75,119,342	<i>JPH1</i>	rs6472827	T/C	0.07	0.76 (0.65–0.86)	3.89E–07	-----	0.99	0
3	5,346,825	<i>EDEMI</i>	rs6804817	T/G	0.09	1.31 (1.20–1.41)	4.18E–07	++++++	0.64	0
3	5,348,095	<i>EDEMI</i>	rs62255982	T/C	0.09	1.31 (1.21–1.42)	4.39E–07	++++++	0.70	0
22	37,736,406	<i>CYTH4</i>	rs34377599	G/GC	0.70	0.78 (0.69–0.88)	4.62E–07	---?--	0.54	0
8	4,441,780	<i>CSMD1</i>	rs11987640	C/G	0.20	0.81 (0.73–0.89)	4.85E–07	?----++	0.01	68.8
8	4,441,757	<i>CSMD1</i>	rs11987636	A/G	0.20	0.81 (0.73–0.89)	4.94E–07	?----++	0.01	68.8

Boldface indicates genome-wide significant  $p$  values

<sup>a</sup>Chromosome<sup>b</sup>Effect allele/reference allele<sup>c</sup>Effect allele frequency<sup>d</sup>Odds ratio per additional effect allele (95% confidence interval)





**Table 3** Genetically predicted gene expression results from GWAS summary statistics

Tissue	Gene	Chr	Z score	P value	R <sup>2a</sup>	Variance <sup>b</sup>	N SNPs <sup>c</sup>	Model N <sup>d</sup>
Thyroid	<i>CYTH4</i> *	22	−5.42	5.86E−08	0.030	0.0003	1	17
Breast	<b><i>ZNF391</i></b>	6	4.40	1.07E−05	0.157	0.008	6	17
Thyroid	<i>AGT</i>	1	−4.22	2.49E−05	0.010	0.005	6	50
Whole blood	<i>SLC26A8</i>	6	4.08	4.43E−05	0.046	0.001	4	14
Fibroblasts	<b><i>ZNF391</i></b>	<b>6</b>	4.04	5.34E−05	0.519	0.123	34	77
Ovary	<b><i>ZNF391</i></b>	6	3.97	7.17E−05	0.315	0.110	14	26
Whole blood	<i>ALDH2</i>	12	3.97	7.30E−05	0.008	0.002	8	42
Breast	<i>LY6D</i> *	8	−3.79	1.52E−04	0.046	0.001	6	35
Fibroblasts	<i>MOB3A</i>	19	3.77	1.62E−04	0.008	0.0002	2	11
Subcutaneous adipose	<i>GLIS1</i>	1	−3.75	1.75E−04	0.034	0.0002	2	6
Whole blood	<b><i>ZNF391</i></b>	6	3.70	2.13E−04	0.103	0.001	12	32
Subcutaneous adipose	<b><i>ZNF391</i></b>	6	3.69	2.20E−04	0.225	0.038	6	33
Fibroblasts	<i>FAM174A</i>	5	−3.66	2.51E−04	0.040	0.004	8	48
Vagina	<b><i>ZNF391</i></b>	6	3.65	2.63E−04	0.193	0.037	23	39
Breast	<i>OAZ1</i>	19	3.61	3.11E−04	0.019	0.013	11	42
Thyroid	<b><i>ZNF391</i></b>	6	3.57	3.60E−04	0.317	0.023	26	57
Thyroid	<i>RAB30</i>	11	3.53	4.21E−04	0.062	0.047	33	89
Vagina	<i>UCKL1</i>	20	3.53	4.22E−04	0.100	0.123	24	113
Whole blood	<i>TMEM91</i>	19	3.52	4.28E−04	0.180	0.0009	2	10

Boldface highlights suggestive predicted *ZNF391* expression across multiple tissues

\* represents genes which localize to top GWAS signals (Table 2)

<sup>a</sup>Performance prediction R<sup>2</sup>

<sup>b</sup>Variance of the gene's predicted expression, calculated as  $W' \times G \times W$  (where  $W$  is the vector of SNP weights in a gene's model,  $W'$  is its transpose, and  $G$  is the covariance matrix)

<sup>c</sup>Number of SNPs included in the prediction model for that gene available in the summary statistics

<sup>d</sup>Number of SNPs used to construct the prediction model for the gene in the tissue of interest using the GTEx data

and more interestingly, is expressed in several reproductive-related tissues in GTEx. The highest expression of *ZNF391* is in testis and ovary, with uterine expression falling in the top third of tissues. Furthermore, a SNP in this gene (rs4713108) is suggestively associated with expression of *ZNF391* in uterine tissue ( $p$  value =  $1.1 \times 10^{-7}$ ). This particular SNP was not associated with fibroid risk in our meta-analysis ( $p$  value = 0.08); however, there was a broad base of suggestive SNPs in the region with  $p$  values as low as  $2.01 \times 10^{-6}$  (Supplemental Fig. 2).


Another gene target identified through GPGE, *ALDH2* ( $p$  value =  $7.30 \times 10^{-5}$ ), encodes a mitochondrial isoform of alcohol dehydrogenase, which is involved in alcohol metabolism. Five studies have shown a consistent relationship between increased alcohol consumption and fibroid risk (OR = 1.25–2.78) (D'Aloisio and Baird 2004; Marshall et al. 1997; Nagata et al. 2009; Templeman et al. 2009; Wise et al. 2004). Predicted decreased levels of *AGT* in thyroid tissue were also implicated as being associated with fibroid risk ( $p$  value =  $2.49 \times 10^{-5}$ ). *AGT* encodes angiotensinogen, which is involved in renal homeostasis, and given the observed link between hypertension and

fibroids, may support a plausible biological connection between the two.

Previous GWAS for fibroids in Japanese subjects has implicated variants near the *BETIL*, *TNRC6B*, and *OBFC1-SLK* genes (Cha et al. 2011). We did not observe suggestive associations in our AA subjects at the prior GWAS-associated SNPs near *BETIL* and *OBFC1-SLK* which were reported by Cha et al. (Supplemental Figs. 3–8). The best result at these loci was near *TNRC6B* (rs59426214;  $p$  value =  $2.42 \times 10^{-5}$ ; Supplemental Fig. 3), although this variant was not in LD with the index SNP (rs12484776) detected by Cha et al. This lack of strong association is largely consistent with prior published studies that have attempted to replicate the associations within cohorts of African ancestry.

A potential consideration of this study was that the Stage 2 studies relied upon self-reported fibroids information, which is known to be subject to misclassification bias. However, the top result at *CYTH4* was suggestive in the imaged Stage 1 analysis with a  $p$  value of  $6.46 \times 10^{-7}$ , and effect sizes were consistent between Stages 1 and 2. Additional points to consider are that the control group was not





**Fig. 3** Regional association plot for *CYTH4*. There is substantial LD between the lead SNP rs739187 and other SNPs in the region

restricted to women beyond menopause, when they would have a negligible likelihood of developing new fibroids. We did explore this as a secondary analysis in the studies from Stage 1, albeit with reduced power due to the smaller sample size (data not shown). Restricting the controls to women peri-menopausal and younger (age <45) did not change the overall associations, as many of the upper tier signals were of similar effect size ( $r^2 = 0.87$ ) despite the lack of significance due to reduced power. In addition, adjustment for age in Stage 1 did not markedly alter the results seen in the primary analysis, with several of the top overall signals remaining suggestively significant (6/12 variants with  $p$  value <  $1e-4$ ) and having consistent effect sizes ( $r^2 = 0.96$ ). Further considerations are the small numbers and heterogeneity among tissues included in the GTEx models. Uterus in particular had a very small number of tissue samples ( $N = 70$ ), and is known to be a heterogeneous mix of specific cell/tissue types.

In conclusion, in this first multi-stage GWAS of uterine fibroids among AA women, we identified a novel genome-wide significant result in *CYTH4* with biological support from the significant GPGE showing reduced *CYTH4* expression in thyroid tissue. Further studies are needed to confirm the SNP and target loci and elucidate the biological role of

this gene and others identified through GWAS and GPGE in relation with fibroid development (Edwards et al. 2013a). Overall, we have identified novel genetic signals, highlighting the importance of this work for understanding fibroids in AA women.

**Acknowledgements** This work was supported by the National Institutes of Health Grants R01HD074711 (DRVE), R03HD078567 (DRVE), U01HG08672-01 (DMR), and R01HD060530 (CCM), as well as the Vanderbilt Molecular and Genetic Epidemiology of Cancer (MAGEC) training program, funded by R25CA160056 (PI: X.-O. Shu). One of the data sets used for the analyses described was obtained from Vanderbilt University Medical Center's BioVU which is supported by institutional funding, the 1S10RR025141-01 instrumentation award, and by the Vanderbilt CTSA Grant UL1TR000445 from NCATS/NIH. The BioMe Biobank program is supported by The Andrea and Charles Bronfman Philanthropies. The Electronic Medical Records and Genomics Network was initiated and funded by the National Human Genome Research Institute through by eMERGE Network Grant U01HG006380 to Mt. Sinai. The Coronary Artery Risk Development in Young Adults Study (CARDIA) is supported by contracts HHSN268201300025C, HHSN268201300026C, HHSN268201300027C, HHSN268201300028C, HHSN268201300029C, and HHSN268200900041C from the National Heart, Lung, and Blood Institute (NHLBI), the Intramural Research Program of the National Institute on Aging (NIA), and an intra-agency agreement between NIA and NHLBI (AG0005). Genotyping was funded as part of the NHLBI Candidate-gene Association Resource (N01-HC-65226). Uterine fibroid-related work from

the Black Women's Health Study (BWHS) was supported by Grants R01CA58420 (LR), UM1CA164974 (LR), and R01CA098663 (JP) from the National Cancer Institute, and Grant R01HD057966 (LAW). The content is solely the responsibility of the authors and does not necessarily represent the official view of the National Institutes of Health. We thank the research participants and staff from all studies, as well as the employees of 23andMe, for making this work possible.

### Compliance with ethical standards

**Conflict of interest** DAH is an employee of and owns stock options in 23andMe, Inc. EAS declares the following financial competing interests: consultancy related to uterine fibroids: AbbVie, Allergan, Astellas, Bayer, and Gynesonics, Viteava. Patent applications (pending or actual), including individual applications or those belonging to the institution to which the authors, are affiliated and from which the authors may benefit. US 6440445 (With Nowak, RA, issued 2002, no commercial activity). Research Grants: NIH. R01HD060503, PI. P50 HS023418, Site PI for 3 component parts. R01HD074711, Consultant. K12HD065987, Board Member for Mayo Clinic Site. Other: Focused Ultrasound Surgery Foundation Borah(PI) 11/15/11–11/14/12 Costs of Uterine Fibroid Treatments Including Focused Ultrasound Surgery. Insightec Inc, PI Patient care costs in conjunction with R01HD060530. In addition, the following non-financial competing interests: relationship (paid or unpaid) with organizations and funding bodies including nongovernmental organizations, research institutions, or charities. Registry Steering Committee, REgistry for Leiomyoma International Efficacy of Focused Ultrasound (RELIEF). Writing or consulting for an educational company. Honoraria from: UpToDate, Massachusetts Medical Society.

### References

- Aissani B, Zhang K, Wiener H (2015a) Evaluation of GWAS candidate susceptibility loci for uterine leiomyoma in the multi-ethnic NIEHS uterine fibroid study. *Front Genet* 6:241. doi:[10.3389/fgene.2015.00241](https://doi.org/10.3389/fgene.2015.00241)
- Aissani B, Zhang K, Wiener H (2015b) Follow-up to genome-wide linkage and admixture mapping studies implicates components of the extracellular matrix in susceptibility to and size of uterine fibroids. *Fertil Steril* 103(528–34):e13. doi:[10.1016/j.fertnstert.2014.10.025](https://doi.org/10.1016/j.fertnstert.2014.10.025)
- Baird DD, Dunson DB (2003) Why is parity protective for uterine fibroids? *Epidemiology* 14:247–250. doi:[10.1097/01.ede.0000054360.61254.27](https://doi.org/10.1097/01.ede.0000054360.61254.27)
- Baird DD, Dunson DB, Hill MC, Cousins D, Schectman JM (2003) High cumulative incidence of uterine leiomyoma in black and white women: ultrasound evidence. *Am J Obstet Gynecol* 188:100–107
- Baird DD, Dunson DB, Hill MC, Cousins D, Schectman JM (2007) Association of physical activity with development of uterine leiomyoma. *Am J Epidemiol* 165:157–163. doi:[10.1093/aje/kwj363](https://doi.org/10.1093/aje/kwj363)
- Barbeira A, Dickinson SP, Torres JM, Torstenson ES, Zheng J, Wheeler HE, Shah KP, Edwards T, Consortium G, Nicolae D, Cox NJ, Im HK (2016) Integrating tissue specific mechanisms into GWAS summary results. *bioRxiv*. doi:[10.1101/045260](https://doi.org/10.1101/045260)
- Borah BJ, Nicholson WK, Bradley L, Stewart EA (2013) The impact of uterine leiomyomas: a national survey of affected women. *Am J Obstet Gynecol* 209:319.e1–319.e20. doi:[10.1016/j.ajog.2013.07.017](https://doi.org/10.1016/j.ajog.2013.07.017)
- Braganza MZ, de Gonzalez AB, Schonfeld SJ, Wentzensen N, Brenner AV, Kitahara CM (2014) Benign breast and gynecologic conditions, reproductive and hormonal factors, and risk of thyroid cancer. *Cancer Prev Res (Phila)* 7:418–425. doi:[10.1158/1940-6207.capr-13-0367](https://doi.org/10.1158/1940-6207.capr-13-0367)
- Carithers LJ, Ardlie K, Barcus M, Branton PA, Britton A, Buia SA, Compton CC, DeLuca DS, Peter-Demchok J, Gelfand ET, Guan P, Korzeniewski GE, Lockhart NC, Rabiner CA, Rao AK, Robinson KL, Roche NV, Sawyer SJ, Segre AV, Shive CE, Smith AM, Sobin LH, Undale AH, Valentino KM, Vaught J, Young TR, Moore HM (2015) A novel approach to high-quality postmortem tissue procurement: the GTEx project. *Biopreserv Biobank* 13:311–319. doi:[10.1089/bio.2015.0032](https://doi.org/10.1089/bio.2015.0032)
- Cha PC, Takahashi A, Hosono N, Low SK, Kamatani N, Kubo M, Nakamura Y (2011) A genome-wide association study identifies three loci associated with susceptibility to uterine fibroids. *Nat Genet* 43:447–450. doi:[10.1038/ng.805](https://doi.org/10.1038/ng.805)
- Cobb J, Cule E, Moncrieffe H, Hinks A, Ursu S, Patrick F, Kassoumeri L, Flynn E, Bulatovic M, Wulffraat N, van Zelst B, de Jonge R, Bohm M, Dolezalova P, Hirani S, Newman S, Whitworth P, Southwood TR, De Iorio M, Wedderburn LR, Thomson W (2014) Genome-wide data reveal novel genes for methotrexate response in a large cohort of juvenile idiopathic arthritis cases. *Pharmacogenomics J* 14:356–364. doi:[10.1038/tpj.2014.3](https://doi.org/10.1038/tpj.2014.3)
- D'Aloisio AA, Baird D (2004) Variation in the association of alcohol intake with uterine fibroids by race and tumor size. In: *American College of Epidemiology*, vol 14. *Annals of epidemiology*, Boston, p 622
- Delaneau O, Marchini J, Zagury JF (2012) A linear complexity phasing method for thousands of genomes. *Nat Methods* 9:179–181. doi:[10.1038/nmeth.1785](https://doi.org/10.1038/nmeth.1785)
- Duan Q, Liu EY, Auer PL, Zhang G, Lange EM, Jun G, Bizon C, Jiao S, Buyske S, Franceschini N, Carlson CS, Hsu L, Reiner AP, Peters U, Haessler J, Curtis K, Wassel CL, Robinson JG, Martin LW, Haiman CA, Le Marchand L, Matisse TC, Hindorff LA, Crawford DC, Assimes TL, Kang HM, Heiss G, Jackson RD, Kooperberg C, Wilson JG, Abecasis GR, North KE, Nickerson DA, Lange LA, Li Y (2013) Imputation of coding variants in African Americans: better performance using data from the exome sequencing project. *Bioinformatics* 29:2744–2749. doi:[10.1093/bioinformatics/btt477](https://doi.org/10.1093/bioinformatics/btt477)
- Edwards SL, Beesley J, French JD, Dunning AM (2013a) Beyond GWASs: illuminating the dark road from association to function. *Am J Hum Genet* 93:779–797. doi:[10.1016/j.ajhg.2013.10.012](https://doi.org/10.1016/j.ajhg.2013.10.012)
- Edwards TL, Michels KA, Hartmann KE, Velez Edwards DR (2013b) BET1L and TNRC6B associate with uterine fibroid risk among European Americans. *Hum Genet* 132:943–953. doi:[10.1007/s00439-013-1306-3](https://doi.org/10.1007/s00439-013-1306-3)
- Eggert SL, Huyck KL, Somasundaram P, Kavalla R, Stewart EA, Lu AT, Painter JN, Montgomery GW, Medland SE, Nyholt DR, Treloar SA, Zondervan KT, Heath AC, Madden PA, Rose L, Buring JE, Ridker PM, Chasman DI, Martin NG, Cantor RM, Morton CC (2012) Genome-wide linkage and association analyses implicate FASN in predisposition to Uterine Leiomyomata. *Am J Hum Genet* 91:621–628. doi:[10.1016/j.ajhg.2012.08.009](https://doi.org/10.1016/j.ajhg.2012.08.009)
- Faerstein E, Szklo M, Rosenshein N (2001) Risk factors for uterine leiomyoma: a practice-based case-control study. I. African-American heritage, reproductive history, body size, and smoking. *Am J Epidemiol* 153:1–10
- Feingold-Link L, Edwards TL, Jones S, Hartmann KE, Velez Edwards DR (2014) Enhancing uterine fibroid research through utilization of biorepositories linked to electronic medical record data. *J Womens Health (Larchmt)* 23:1027–1032. doi:[10.1089/jwh.2014.4978](https://doi.org/10.1089/jwh.2014.4978)
- Friedman GD, Cutter GR, Donahue RP, Hughes GH, Hulley SB, Jacobs DR Jr, Liu K, Savage PJ (1988) CARDIA: study design, recruitment, and some characteristics of the examined subjects. *J Clin Epidemiol* 41:1105–1116
- Gamazon ER, Wheeler HE, Shah KP, Mozaffari SV, Aquino-Michaels K, Carroll RJ, Eyer AE, Denny JC, Nicolae DL, Cox NJ, Im

- HK (2015) A gene-based association method for mapping traits using reference transcriptome data. *Nat Genet* 47:1091–1098. doi:[10.1038/ng.3367](https://doi.org/10.1038/ng.3367)
- Genotype-Tissue Expression (GTEx) (2015) Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* 348:648–660. doi:[10.1126/science.1262110](https://doi.org/10.1126/science.1262110)
- Kanamarlupudi V, Owens SE, Lartey J, Lopez Bernal A (2012) ADP-ribosylation factor 6 expression and activation are reduced in myometrium in complicated pregnancies. *PLoS One* 7:e37954. doi:[10.1371/journal.pone.0037954](https://doi.org/10.1371/journal.pone.0037954)
- Kim MH, Park YR, Lim DJ, Yoon KH, Kang MI, Cha BY, Lee KW, Son HY (2010) The relationship between thyroid nodules and uterine fibroids. *Endocr J* 57:615–621
- Klemke M, Muller MH, Wosniok W, Markowski DN, Nimzyk R, Helmke BM, Bullerdiek J (2014) Correlated expression of HMGA2 and PLAG1 in thyroid tumors, uterine leiomyomas and experimental models. *PLoS One* 9:e88126. doi:[10.1371/journal.pone.0088126](https://doi.org/10.1371/journal.pone.0088126)
- Lette G, Palmer CD, Young T, Ejebe KG, Allayee H, Benjamin EJ, Bennett F, Bowden DW, Chakravarti A, Dreisbach A, Farlow DN, Folsom AR, Fornage M, Forrester T, Fox E, Haiman CA, Hartiala J, Harris TB, Hazen SL, Heckbert SR, Henderson BE, Hirschhorn JN, Keating BJ, Kritchevsky SB, Larkin E, Li M, Rudock ME, McKenzie CA, Meigs JB, Meng YA, Mosley TH, Newman AB, Newton-Cheh CH, Paltoo DN, Papanicolaou GJ, Patterson N, Post WS, Psaty BM, Qasim AN, Qu L, Rader DJ, Redline S, Reilly MP, Reiner AP, Rich SS, Rotter JI, Liu Y, Shrader P, Siscovick DS, Tang WH, Taylor HA, Tracy RP, Vasani RS, Waters KM, Wilks R, Wilson JG, Fabsitz RR, Gabriel SB, Kathiresan S, Boerwinkle E (2011) Genome-wide association study of coronary heart disease and its risk factors in 8,090 African Americans: the NHLBI CARE Project. *PLoS Genet* 7:e1001300. doi:[10.1371/journal.pgen.1001300](https://doi.org/10.1371/journal.pgen.1001300)
- Luo L, McGarvey P, Madhavan S, Kumar R, Gusev Y, Upadhyay G (2016) Distinct lymphocyte antigens 6 (Ly6) family members Ly6D, Ly6E, Ly6K and Ly6H drive tumorigenesis and clinical outcome. *Oncotarget* 7:11165–11193. doi:[10.18632/oncotarget.7163](https://doi.org/10.18632/oncotarget.7163)
- Luoto R, Kaprio J, Rutanen EM, Taipale P, Perola M, Koskenvuo M (2000) Heritability and risk factors of uterine fibroids—the Finnish Twin Cohort study. *Maturitas* 37:15–26
- Marchini J, Howie B, Myers S, McVean G, Donnelly P (2007) A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat Genet* 39(7):906–913. doi:[10.1038/ng2088](https://doi.org/10.1038/ng2088)
- Makaridze T, Mardaleishvili K (2011) Combined development of thyroid gland and reproductive system benign diseases. *Georgian Med News* (199):20–29
- Marshall LM, Spiegelman D, Barbieri RL, Goldman MB, Manson JE, Colditz GA, Willett WC, Hunter DJ (1997) Variation in the incidence of uterine leiomyoma among premenopausal women by age and race. *Obstet Gynecol* 90:967–973
- Mele M, Ferreira PG, Reverter F, DeLuca DS, Monlong J, Sammeth M, Young TR, Goldmann JM, Pervouchine DD, Sullivan TJ, Johnson R, Segre AV, Djebali S, Niarchou A, Wright FA, Lappalainen T, Calvo M, Getz G, Dermizakis ET, Ardlie KG, Guigo R (2015) Human genomics. The human transcriptome across tissues and individuals. *Science* 348:660–665. doi:[10.1126/science.aaa0355](https://doi.org/10.1126/science.aaa0355)
- Moore AB, Flake GP, Swartz CD, Heartwell G, Cousins D, Haseman JK, Kissling GE, Sidawy MK, Dixon D (2008) Association of race, age and body mass index with gross pathology of uterine fibroids. *J Reprod Med* 53:90–96
- Myers SL, Baird DD, Olshan AF, Herring AH, Schroeder JC, Nylander-French LA, Hartmann KE (2012) Self-report versus ultrasound measurement of uterine fibroid status. *J Womens Health (Larchmt)* 21:285–293. doi:[10.1089/jwh.2011.3008](https://doi.org/10.1089/jwh.2011.3008)
- Nagata C, Nakamura K, Oba S, Hayashi M, Takeda N, Yasuda K (2009) Association of intakes of fat, dietary fibre, soya isoflavones and alcohol with uterine fibroids in Japanese women. *Br J Nutr* 101:1427–1431
- Orozco G, Barton A, Eyre S, Ding B, Worthington J, Ke X, Thomson W (2011) HLA-DPB1-COL11A2 and three additional xMHC loci are independently associated with RA in a UK cohort. *Genes Immun* 12:169–175. doi:[10.1038/gene.2010.57](https://doi.org/10.1038/gene.2010.57)
- Ott J, Kurz C, Braun R, Promberger R, Seemann R, Vytiska-Binstorfer E, Walch K (2014) Overt hypothyroidism is associated with the presence of uterine leiomyoma: a retrospective analysis. *Eur J Obstet Gynecol Reprod Biol* 177:19–22. doi:[10.1016/j.ejogrb.2014.03.003](https://doi.org/10.1016/j.ejogrb.2014.03.003)
- Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D (2006) Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet* 38:904–909. doi:[10.1038/ng1847](https://doi.org/10.1038/ng1847)
- Pulley J, Clayton E, Bernard GR, Roden DM, Masys DR (2010) Principles of human subjects protections applied in an opt-out, de-identified biobank. *Clin Transl Sci* 3:42–48. doi:[10.1111/j.1752-8062.2010.00175.x](https://doi.org/10.1111/j.1752-8062.2010.00175.x)
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 81:559–575. doi:[10.1086/519795](https://doi.org/10.1086/519795)
- Rezazadeh M, Ghahesouran J, Mirabzadeh A, Khorram Khorshid HR, Biglarian A, Ohadi M (2015) A primate-specific functional GTTT-repeat in the core promoter of CYTH4 is linked to bipolar disorder in human. *Prog Neuropsychopharmacol Biol Psychiatry* 56:161–167. doi:[10.1016/j.pnpbp.2014.09.001](https://doi.org/10.1016/j.pnpbp.2014.09.001)
- Spinou N, Terzis G, Crysanthopoulou A, Adonakis G, Markou KB, Vervita V, Koukouras D, Tsapanos V, Decavalas G, Kourounis G, Georgopoulos NA (2007) Increased frequency of thyroid nodules and breast fibroadenomas in women with uterine fibroids. *Thyroid* 17:1257–1259. doi:[10.1089/thy.2006.0330](https://doi.org/10.1089/thy.2006.0330)
- Templeman C, Marshall SF, Clarke CA, Henderson KD, Largent J, Neuhausen S, Reynolds P, Ursin G, Bernstein L (2009) Risk factors for surgically removed fibroids in a large cohort of teachers. *Fertil Steril* 92:1436–1446. doi:[10.1016/j.fertnstert.2008.08.074](https://doi.org/10.1016/j.fertnstert.2008.08.074)
- Terry KL, De Vivo I, Hankinson SE, Spiegelman D, Wise LA, Missmer SA (2007) Anthropometric characteristics and risk of uterine leiomyoma. *Epidemiology* 18:758–763. doi:[10.1097/EDE.0b013e3181567eed](https://doi.org/10.1097/EDE.0b013e3181567eed)
- Vollenhoven B (1998) Introduction: the epidemiology of uterine leiomyomas. *Baillieres Clin Obstet Gynaecol* 12:169–176
- Wellons MF, Lewis CE, Schwartz SM, Gunderson EP, Schreiner PJ, Sternfeld B, Richman J, Sites CK, Siscovick DS (2008) Racial differences in self-reported infertility and risk factors for infertility in a cohort of black and white women: the CARDIA Women's Study. *Fertil Steril* 90:1640–1648. doi:[10.1016/j.fertnstert.2007.09.056](https://doi.org/10.1016/j.fertnstert.2007.09.056)
- Willer CJ, Li Y, Abecasis GR (2010) METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 26:2190–2191. doi:[10.1093/bioinformatics/btq340](https://doi.org/10.1093/bioinformatics/btq340)
- Wise LA, Palmer JR, Harlow BL, Spiegelman D, Stewart EA, Adams-Campbell LL, Rosenberg L (2004) Risk of uterine leiomyomata in relation to tobacco, alcohol and caffeine consumption in the Black Women's Health Study. *Hum Reprod* 19:1746–1754. doi:[10.1093/humrep/deh309](https://doi.org/10.1093/humrep/deh309)
- Wise LA, Palmer JR, Spiegelman D, Harlow BL, Stewart EA, Adams-Campbell LL, Rosenberg L (2005a) Influence of body size and body fat distribution on risk of uterine leiomyomata in U.S. black women. *Epidemiology* 16:346–354

- Wise LA, Palmer JR, Stewart EA, Rosenberg L (2005b) Age-specific incidence rates for self-reported uterine leiomyomata in the Black Women's Health Study. *Obstet Gynecol* 105:563–568. doi:[10.1097/01.AOG.0000154161.03418.e3](https://doi.org/10.1097/01.AOG.0000154161.03418.e3)
- Wise LA, Ruiz-Narvaez EA, Palmer JR, Cozier YC, Tandon A, Patterson N, Radin RG, Rosenberg L, Reich D (2012) African ancestry and genetic risk for uterine leiomyomata. *Am J Epidemiol* 176:1159–1168. doi:[10.1093/aje/kws276](https://doi.org/10.1093/aje/kws276)
- Zhang K, Wiener H, Aissani B (2015) Admixture mapping of genetic variants for uterine fibroids. *J Hum Genet*. doi:[10.1038/jhg.2015.60](https://doi.org/10.1038/jhg.2015.60)