

# UCSF

## UC San Francisco Previously Published Works

### Title

A Cross-Sectional Relationship Between Social Capital, Self-Compassion, and Perceived HIV Symptoms.

### Permalink

<https://escholarship.org/uc/item/0n82f78d>

### Journal

Journal of Pain and Symptom Management, 50(1)

### Authors

Webel, Allison  
Wantland, Dean  
Rose, Carol  
[et al.](#)

### Publication Date

2015-07-01

### DOI

10.1016/j.jpainsymman.2014.12.013

Peer reviewed



Published in final edited form as:

*J Pain Symptom Manage.* 2015 July ; 50(1): 59–68. doi:10.1016/j.jpainsymman.2014.12.013.

## A Cross-Sectional Relationship Between Social Capital, Self-Compassion and Perceived HIV Symptoms

Allison Webel, PhD, RN, Dean Wantland, PhD, RN, Carol Dawson Rose, PhD, RN, Jeanne Kempainen, PhD, RN, William L. Holzemer, PhD, RN, Wei-Ti Chen, PhD, RN, Mallory O. Johnson, PhD, Patrice Nicholas, PhD, RN, Lucille Sanzero Eller, PhD, RN, Puangtip Chaiphibalsarisdi, PhD, Elizabeth Sefcik, PhD, RN, Kathleen Nokes, PhD, RN, Inge B. Corless, PhD, RN, Lynda Tyer-Viola, PhD, RN, Kenn Kirksey, PhD, RN, Joachim Voss, PhD, RN, ACRN, Kathy Sullivan, PhD, RN, Marta Rivero-Méndez, PhD, RN, John Brion, PhD, RN, Scholastika Ipinge, PhD, J. Craig Phillips, PhD, RN, and Carmen Portillo, PhD, RN

Frances Payne Bolton School of Nursing (A.W.), Case Western Reserve University, Cleveland, Ohio; Rutgers School of Nursing (D.W., W.L.H., L.S.E.), Newark, New Jersey; School of Nursing (C.D.R.), University of California at San Francisco, San Francisco, California; University of North Carolina Wilmington (J.K.), Wilmington, North Carolina; Yale University (W.-T.C.), New Haven, Connecticut; University of California at San Francisco (M.O.J.) San Francisco, California; Global Health and Academic Partnerships (P.N.), Brigham and Women's Hospital and Massachusetts General Hospital, and Massachusetts General Hospital Institute of Health Professions (P.N., I.B.C., L.T.-V.), Boston, Massachusetts, USA; Suan Sunandha Rajabhat University (P.C.), Bangkok, Thailand; Texas A&M University-Corpus Christi (E.S.), Corpus Christi, Texas; Hunter College, City University of New York, and Hunter-Bellevue School of Nursing (K.N.), New York, New York; Center for Nursing Research (K.K.), Seton Family of Hospitals, Austin, Texas; University of Washington (J.V.), Seattle, Washington; School of Nursing and Dental Hygiene (K.S.), University of Hawaii at Manoa, Honolulu, Hawaii; University of Puerto Rico (M.R.-M.), San Juan, Puerto Rico; Ohio State University College of Nursing (J.B.), Columbus, Ohio; USA; University of Namibia (S.I.), Windhoek, Namibia; University of Ottawa (J.C.P.), Ottawa, Ontario, Canada; and Community Health Systems (C.P.), University of California School of Nursing at San Francisco, San Francisco, California, USA

### Abstract

**Context**—Individual resources of social capital and self-compassion are associated with health behaviors and perceived symptoms, suggesting that both are positive resources that can be modified to improve a person's symptom experience.

---

Address correspondence to: Allison Webel, PhD, RN, Frances Payne Bolton School of Nursing, Case Western Reserve University, 10900 Euclid Avenue, Cleveland, OH 44106-4904, USA, allison.webel@case.edu.

#### Disclosures

The contents of this manuscript are solely the responsibility of the authors and do not necessarily represent the official views of the NIH or any other funders. These funding agencies had no role in the study design; in the collection, analysis and interpretation of data; in the writing of the report; or in the decision to submit the paper for publication. The authors declare no conflicts of interest.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Objectives**—To examine the relationship between self-compassion and social capital and its impact on current HIV symptom experience in adult people living with HIV (PLWH). We further explored the impact of age on this relationship.

**Methods**—We conducted a cross-sectional analysis of 2182 PLWH at 20 sites in five countries. Social capital, self-compassion, and HIV symptom experience were evaluated using valid and reliable scales. To account for inflated significance associated with a large sample size, we took a random sample of 28% of subjects ( $n=615$ ) and conducted correlation analyses and zero inflated Poisson regression, controlling for known medical and demographic variables impacting HIV symptom experience.

**Results**—Controlling for age, sex at birth, year of HIV diagnosis, comorbid health conditions, employment and income, our model significantly predicted HIV symptom experience (overall model  $z=5.77$ ,  $P<0.001$ ). Employment status and social capital were consistent, negative and significant predictors of HIV symptom experience. Self-compassion did not significantly predict HIV symptom experience. If reporting symptoms, an increase in age was significantly associated with an increase in symptoms.

**Conclusion**—Employment and social capital modestly predicted current HIV symptom experience. Social capital can be incorporated into symptom management interventions, possibly as a way to reframe a person's symptom appraisal. This may be increasingly important as PLWH age. The relationship between employment status and HIV symptom experience was significant and should be explored further.

## Keywords

Symptom; HIV; self-compassion; social capital

---

## Introduction

Over the past two decades, HIV has transitioned from an acute to a chronic health condition. The symptom experiences for people living with HIV (PLWH) also have evolved from symptoms related to illness progression and toxic antiretroviral therapies (ARTs), to symptoms consistent with long-term adherence to better tolerated ARTs and age-related comorbidities (1). Recently, investigators found that despite the remarkable advances in treatment, PLWH continue to experience prevalent and distressing symptoms, and that those symptoms have a significant impact on daily functioning across varied populations (2–6). Despite improvements in ARTs, symptoms such as depression and fatigue are a leading contributor to poor antiretroviral treatment adherence (4, 7) and poor quality of life (8), underscoring the significance of and need for effective and scalable symptom management strategies for PLWH.

Recent HIV symptom management strategies include clinician- and peer-delivered cognitive skill-building interventions (9, 10), an HIV-specific symptom management manual (11), a web-based HIV symptom management information tool (12), and mindfulness-based stress reduction interventions (13, 14). The success of mindfulness-based interventions in PLWH suggests that symptom management strategies focusing on regulating emotional responses to

symptoms is a promising approach. Strategies targeting self-compassion as a way to modulate emotional regulation may improve HIV symptom management.

Self-compassion involves feelings of “caring and kindness towards oneself in the face of personal suffering and involves the recognition that one’s suffering, failures, and inadequacies are part of the human condition” (15–17). PLWH often confront stigma and discrimination that may contribute to self-criticism, isolation and avoidance of internally satisfying experiences (18). This can lead to over-identification with negative thoughts and feelings and may subsequently lead to an increased negative appraisal of symptom intensity. Accordingly, investigators have found negative associations between self-compassion and depressive symptoms (19), symptoms of schizophrenia (20), and pain catastrophizing (21). However, the relationship between self-compassion and symptom intensity has not been explored among PLWH.

Social capital is the “aggregate of potential resources, which are linked to possession of a durable network of more or less institutionalized relationships of mutual acquaintance or recognition” (23, 24) and also may modulate self-regulation. Social capital represents the degree to which a person has access to high-quality social resources, including social support through social networks. Among PLWH, social capital has been shown to impact a number of health-related outcomes including decreases in HIV transmission (25–28), self-reported physical and psychological health (29), behaviors including HIV self-management (30), and HIV medication adherence (31–33). Available high-quality social resources have been shown to influence one’s regulation of emotions surrounding symptoms (22). Two particular social resources, social support and social networks, have been found to be important in chronic disease symptom management (34). Recently, Fierz and colleagues (34) reported on the importance of social resources and individual interpretation in managing symptoms(34). As a measure of available social resources, individual social capital may encapsulate many of the resources necessary for PLWH to adequately manage their daily symptoms. Further, one’s perception of available social resources may influence his or her appraisal of symptoms, yet this relationship is unexplored.

As PLWH age, they may experience an increase in chronic comorbid health conditions and distressing symptoms; however, older PLWH are more socially connected than younger PLWH (35). This increased social connectedness may allow older PLWH to better manage their daily symptoms. Further, Brion and colleagues demonstrated that self-compassion may increase with the time a person is living with HIV, especially as they learn how to better accept their diagnosis and manage living with HIV (18). The effect of age, or time since HIV diagnosis, on a person’s HIV symptom experience may be associated the individual’s perception of social capital and self-compassion, but this relationship has not been described.

The objective of the present study was to extend the previous literature by examining the relationship between self-compassion and social capital and their impact on the current HIV symptom experience in adult PLWH. With the rapid increase in older PLWH globally, we further explored the impact of age on this relationship hypothesizing that older PLWH would have more self-compassion and social capital, which would attenuate their symptom

intensity. Given available literature, we anticipated that social capital and self-compassion would be negatively associated with symptom intensity in PLWH. The potential influence of social capital and self-compassion on symptom experience is unexplored; it may offer novel insights into how to efficiently modify one's appraisal of his or her symptoms.

## Methods

### Setting

This was a cross-sectional study conducted by the International Nursing Network for HIV/AIDS Research. Data were collected from study sites in Vancouver, Canada; Shanghai, China; Windhoek, Namibia; Bangkok, Thailand; and from 11 sites in the United States (Boston, MA; Chicago, IL; Cleveland, OH; Corpus Christi, TX; Durham, NC; Harlingen, TX; Honolulu, HI; Newark, NJ; New York, NY; San Francisco, CA; San Juan, PR; Seattle, WA; and Wilmington, NC). Institutional Review Board (IRB) approval was obtained by the coordinating site and at each respective study site. Each site director adhered to a common data collection protocol and all participants gave informed consent prior to completing any study procedures.

### Participants and Data Collection Procedures

Each site sought to recruit a convenience sample of a minimum of 100 adults living with HIV infection. Participants included adults (over the age of 18 years), who were diagnosed with HIV infection, and were able to give informed consent in their local language. Participants were recruited through flyers and health care provider referral at local HIV clinics or HIV service organizations. Potential participants contacted study staff who screened them for study eligibility and scheduled an appointment to complete study procedures. Additional participants were recruited in HIV clinics by study staff who screened and enrolled them, and completed study procedures in a private room. After giving informed consent, participants completed a self-administered survey comprising 16 instruments. Participants who were unable to read the survey were given the opportunity to complete it by interview with a research assistant. Upon completing the survey, participants were compensated with a financial incentive consistent with local IRB approvals and standards. All data were entered into an electronic database and were de-identified. The de-identified data were sent to the coordinating site, cleaned, entered into the master database, and stored until all sites completed data collection and entry. Original data were stored at each individual site.

### Measurements

The measures used to assess our variables of interest are listed below. While participants completed a battery of 16 surveys measures, these variables were chosen a priori and based on our hypothesis that social capital and self-compassion would be negatively associated with symptom intensity among PLWH. For sites where participants did not speak English as their native language, appropriate translation procedures were followed (36).

*Demographic characteristics and HIV disease status* were measured with a 20-item demographic and illness characteristics instrument. This included age, gender, race,

ethnicity, education, income, health insurance, date first learned of HIV diagnosis, self-reported current CD4 count, viral load, HIV medications, and general health.

*HIV symptom intensity* was measured with the 72-item revised Sign and Symptom Checklist (37, 38). This scale measures the number and intensity of symptoms being experienced by PLWH on the day of data collection. This scale has been validated and repeatedly used to assess symptoms in PLWH (11, 38–40). Participants were asked to rate whether they experienced the symptom in the past 24-hours and if so, how intense it was on a 0–3 scale. Zero indicates that a participant did not experience the symptom at all in the past 24 hours, one indicates mild intensity symptoms, two indicates moderate intensity symptoms and three indicates a severe intensity symptom. Intensity was summed across symptoms and a higher score indicates higher HIV symptom intensity. The possible range for symptom intensity is 0 to 216 for women and 0 to 192 for men (because gynecological symptoms are gender specific). Reliability for all items of the revised Sign and Symptom Checklist for our study was 0.97.

*Self-compassion* was measured with the 12-item Brief Version Self-Compassion Inventory (41). This scale was developed from Neff's (16) 26-item inventory and has demonstrated acceptable psychometric properties. We chose the brief version of the scale to minimize participant burden. Participants were asked to rate how they dealt with 12 difficult situations on a 1–5 scale with a possible range of 12–60. After reverse-scoring negatively worded items, items are summed. Higher scores indicate more self-compassion. Internal consistency reliability for the Brief Self-Compassion Inventory was 0.71 and it demonstrated acceptable internal and external validity in PLWH (41).

*Social capital* was measured with 31 items from the 36-item Social Capital Scale (42–45). This widely-used instrument provides a total score and measures eight subscale scores including: participation in the local community, social agency, feelings of trust and safety, neighborhood connections, friends and family connections, tolerance of diversity, value of life, and workplace connections. We excluded three workplace connections items and two work-related questions that are part of the social agency dimension. This modification occurred with the approval and recommendation of the scale's original authors because we anticipated high unemployment in our sample. Participants were asked to rate each item on a 1–4 scale with a possible range of 31 to 124. Higher mean scores indicate more social capital. Reliability and validity of the scale have been reported as acceptable (42). Reliability for the Social Capital Scale for our study was 0.88 (29).

### Statistical Analysis

All data were entered into a data management program and data integrity and assumptions were checked. Analyses were conducted in Stata v. 13.1 (StataCorp. LP, College Station, TX) and included descriptive statistics and bivariate correlational analyses including Spearman's rank order correlation. Because of the preponderance of values of zero in 24-hour HIV symptom intensity, zero-inflated Poisson regression modeling was used to assess effects of predictor variables on HIV symptom intensity. In addition to examining our variables of interest (self-compassion, social capital and age), we controlled for the effect of

covariates known to influence the HIV symptom experience (sex, years living with HIV diagnosis, comorbid health conditions, work and income) (46).

Because of the large sample size, we wanted to minimize the risk of overestimating the significance of our analyses and conducted all analyses on a subsample comprising 28% of the overall sample. The dependent variable (HIV symptom intensity) for the overall sample was reviewed for outliers using the  $z$  score test method. We removed cases that had a mean symptom intensity score greater than three standard deviations away from the group mean from the random subsample selection process. The subsample data ( $n=620$ ) were then reviewed for symptom intensity data completeness and data from an additional five cases were dropped for incomplete symptom intensity data for a final subsample of 615 participants. Only those cases that had complete data for all variables were included in the regression models, resulting in a final sample of 528.

## Results

A randomly selected subsample of 615 surveys was selected from a total of 2183 participants completing the survey. From this subsample, approximately 70% were male, 42% were Black/African American, 22% White/Anglo and over 60% had a high school diploma or less. Most (75%) were unemployed, had health insurance (72%), and reported having at least one comorbid health condition (63%). On average, most participants had been living with HIV for approximately 12 years, were prescribed HIV ART (81%), and had a mean CD4+ T cell count of 431 cells/ $\mu$ l. The participants reported experiencing an average of 18.7 (SD 13.6) symptoms (range 0–50), with an average total symptom intensity of 30.2 (range 0–147). Participants reported an average score of 39 on the Brief Self-Compassion Inventory (range 12–60) and had an average score of 2.54 on the social capital scale (range 1.2–3.9). Additional demographics and HIV disease characteristics can be found in Table 1.

Using bivariate analyses (Table 2) of relationships among HIV symptom intensity, social capital, self-compassion and age, we observed significant negative correlations between HIV symptom intensity and social capital ( $\rho = -0.23$ ,  $P < 0.001$ ) and HIV symptom intensity and self-compassion ( $\rho = -0.28$ ,  $P < 0.001$ ). These significant negative associations indicated that as one variable increases (i.e., self-compassion), the other variable is reduced (i.e., symptom intensity). Simply stated, as the reported average level of symptom intensity increases, the average social capital score, or average self-compassion score significantly decreases. We observed no relationship between HIV symptom intensity and age. We also observed a significant moderate relationship between social capital and self-compassion ( $\rho = 0.40$ ,  $P < 0.001$ ).

Zero-inflated Poisson regression analyses were used to assess our model's predictive power to estimate HIV symptom intensity. As zero on the HIV Sign and Symptom Checklist indicates that the participant did not experience the symptom in the previous 24 hours, it was expected that some of the participants would not report any current symptoms; approximately 8% of the sample did not experience any symptoms. Zero-inflated Poisson regression analyses are used to model count data, as is the case with the number and



intensity of self-reported symptoms. that have an excess in zero counts (i.e., no symptoms). Further, zero-inflated Poisson regression models are easy to interpret, can lead to more refined data analyses by evaluating the excess zeroes that are generated by a separate process from the non-zero count values, and the excess zeroes can be modeled independently (47, 48). The zero-inflated Poisson regression model has two components, a Poisson count model and a logit model for predicting excess zeroes (representing membership in the “no symptom” group). When examining these models together, only two variables consistently predicted HIV symptom intensity, being employed and social capital. Being employed was significantly associated with lower symptom intensity ( $\beta = -0.10$ , 95% confidence interval [CI]  $-0.14, -0.06$ ), and was associated with an increased likelihood (~17%) of not reporting any symptoms ( $\beta = 1.17$ , 95% CI 0.38, 1.96). Social capital was significantly associated with lower symptom intensity ( $\beta = -0.004$ , 95% CI  $-0.005, -0.003$ ) and was associated with an increased likelihood (~5%) of not reporting any symptoms ( $\beta = 0.05$ , 95% CI 0.02, 0.07). As expected, additional variables were significant in their respective models (age, comorbidities, years since HIV diagnosis, income, race) (Table 3).

To assess the model fit characteristics, we conducted the Vuong test, which compares the zero-inflated Poisson model to a standard Poisson model (49, 50). Because the  $z$ -value is both positive and significant ( $z = 5.77, p < 0.001$ ), the Vuong test shows that the zero-inflated Poisson is a better fit than the standard Poisson regression. This is also supported by improved log likelihood fit statistics for the two models showing that the count-only data model has a log likelihood value of  $-7428.50$ , compared to the model using both non-zero and zero values for 24-hour HIV symptom intensity that had a lower (closer to zero) log likelihood of  $-6878.81$ , indicating a better fit.

## Discussion

Our analyses of the relationship between self-compassion and social capital and its impact on current HIV symptom experience in adult PLWH yielded several observations that add to existing literature. First, consistent with our hypothesis we found that social capital is a significant negative predictor of HIV symptom intensity. Our results further indicate that among covariates known to impact HIV symptom intensity, social capital is one of the strongest predictors of HIV symptom intensity. Social resources in the form of social support and social networks have long been recognized as important to symptom management, particularly in chronic disease (34, 51, 52). Social capital, while controversial, is a promising construct for better understanding the precise nature of how social resources can improve HIV symptom experience (53, 54). As a modifiable asset of PLWH's social context, social capital can be targeted as a means to improve health-related outcomes in tangible ways (29). Lewis and colleagues (55) reviewed the literature and developed a social capital framework for end-of-life palliative care among terminally ill patients. They argued that social capital was an important construct within the social context of particular populations but also that specificity is needed in its application (55). By choosing a social capital instrument that has been widely used and extensively validated, we can provide specificity about which domains of social capital were associated with HIV symptom intensity. The social capital scale we used comprises discrete measures of participation in the local community, social agency, feelings of trust and safety, neighborhood connections,



friends and family connections, tolerance of diversity, and value of life. Additional research is required because if our results are replicated, the findings can direct us to the components of an intervention to improve symptom intensity based on social capital and its subscales.

An explanation for our findings related to social capital is that the appraisal and interpretation of the HIV symptom experience is likely to be linked to how much someone feels they can readily access the necessary emotional, psychological, and material resources available to them when dealing with physical and psychological symptoms. Consistent with the collective network approach to social capital, these resources can improve health outcomes by influencing health behavior, mobilizing a network to access health services, and providing emotional and psychological support (56). Someone who believes they have less access to these resources may interpret his or her symptoms as insurmountable. This may lead them to become emotionally overwhelmed with the experience and less likely to respond to symptom management interventions. Emerging evidence in the field of cognitive neuroscience suggested that there is an antagonistic relationship between emotional neurocognitive processing and analytic neurocognitive processing that, when activated, prevents one from using the other neural system. (57). This may help explain the variable success of evidence-based, efficacious symptom management interventions (58) and is consistent with analysis suggesting that higher social capital mediates limited literacy and health outcomes (59), and improves chronic disease self-management (60). For example, a person living with HIV who has low social capital may be focused on the emotional distress surrounding their HIV symptom experience and be unable to use the skills taught in symptom management interventions including self-monitoring, knowledge acquisition, and self-regulation. But symptom management interventions that integrate both emotional and analytic neurocognitive processing may be more successful. The mindfulness-based interventions serve as a model of strategies that integrate both emotional and analytic neurocognitive processing. Considered in this context, our results suggest that screening PLWH with high symptom intensity for their perceived social capital may be a useful strategy (13, 14, 61). Social capital screening information can then be used to tailor an intervention to the patient's unique social context and may facilitate better targeted HIV symptom management interventions.

We also found a significant, moderate, and positive association between self-compassion and social capital. The relationship between how kind someone is to herself in times of distress and her perceived social resources is intriguing and deserves further study, preferably with a causal research design. However, we did not find a relationship between self-compassion and HIV symptom intensity. As the Brief Self-Compassion Scale is a new instrument, in the future it also may be advantageous to operationalize self-compassion using several tools to ensure optimal assessment. It is possible that either self-compassion or social capital is a condition for the other or they tend to co-vary together. However, we were unable to examine these relationships because of our cross-sectional study design.

Being in paid employment significantly predicted HIV symptom intensity and this evidence adds to the growing body of literature documenting the positive role of paid employment among PLWH (30, 62–65). However, the relationship between symptoms and employment status is difficult to disentangle in a cross-sectional study. If a PLWH has fewer symptoms,

they may be more likely to seek out and engage in paid employment which may explain our findings. However, it is also possible that paid employment allows someone to interpret their symptoms as less intense and bothersome. Paid employment also may afford different opportunities to access health care and other services that may reduce severity of HIV symptoms.

Finally, we did not observe a consistent relationship between age, social capital, and symptom intensity. Once a PLWH reports symptoms, age was significantly associated with their symptom intensity; however, it was not associated with the likelihood that they did not experience HIV symptoms. This unexpected finding may be related to the limited age range of our sample. With 99% of participants less than 69 years of age, we may not have enough older participants to detect associations.

### Limitations

In addition to the cross-sectional study design, several limitations should be noted. We used convenience sampling of PLWH from clinics and community organizations. It is possible that symptom intensity, self-compassion, and social capital are different in our sample compared to the global population of PLWH. However, our large sample size and random selection of 28% of that sample reduced this risk. Further, many of our study sites were only able to obtain laboratory values via self-report. For this reason, we only provide this data descriptively and did not include it in regression analyses. We also did not include the workplace-related items on the Social Capital Scale. Although only 25% of the participants were employed, this information could have provided additional evidence to help us understand the relationship between employment and symptom intensity. Future work should include these items when using the Social Capital Scale.

Despite these limitations we found, for the first time, that social capital predicted current HIV symptom intensity as well as demonstrated that individuals with greater social capital were also more likely to have lower symptom intensity or no reported symptoms. Additional research is needed to understand this relationship. A better understanding of how social capital influences HIV symptom intensity can help in the development of targeted and tailored HIV symptom management interventions.

### Acknowledgments

This project was supported in part by: NIH UL1 RR024131; NIH T32NR007081; NIH KL2RR024990; NIH R15NR011130; International Pilot Award, University of Washington Center for AIDS Research; University of British Columbia School of Nursing Helen Shore Fund; Duke University School of Nursing Office of Research Affairs; MGH Institute for Health Professions; University of Washington School of Nursing Research and Intramural Funding Program; Rutgers College of Nursing; City University of New York and the Irwin Belk Distinguished Professorship Fund-University of North Carolina Wilmington.

### References

1. Cook PF, Sousa KH, Matthews EE, Meek PM, Kwong J. Patterns of change in symptom clusters with hiv disease progression. *J Pain Symptom Manage*. 2011; 42:12–23. [PubMed: 21429701]
2. Fierz K, Engberg S, Nicca D, et al. Symptom manageability in Swiss HIV-infected patients. *Appl Nurs Res*. 2013; 26:110–115. [PubMed: 23481487]

3. Harding R, Selman L, Aquipio G, et al. Prevalence, burden, and correlates of physical and psychological symptoms among hiv palliative care patients in sub-Saharan Africa: an international multicenter study. *J Pain Symptom Manage.* 2012; 44:1–9. [PubMed: 22658471]
4. Harding R, Lampe FC, Norwood S, et al. Symptoms are highly prevalent among HIV outpatients and associated with poor adherence and unprotected sexual intercourse. *Sex Transm Infection.* 2010; 86:520–524. [PubMed: 20551235]
5. Namisango E, Powell RA, Atuhaire L, et al. Is symptom burden associated with treatment status and disease stage among adult HIV outpatients in East Africa? *J Palliat Med.* 2014; 17:304–312. [PubMed: 24359205]
6. Merlin JS, Cen L, Praestgaard A, et al. Pain and physical and psychological symptoms in ambulatory hiv patients in the current treatment era. *J Pain Symptom Manage.* 2012; 43:638–645. [PubMed: 22115794]
7. Gay C, Portillo CJ, Kelly R, et al. Self-reported medication adherence and symptom experience in adults with HIV. *J Assoc Nurses AIDS Care.* 2011; 22:257–268. [PubMed: 21377900]
8. Vyavaharkar M, Moneyham L, Murdaugh C, Tavakoli A. Factors associated with quality of life among rural women with HIV disease. *AIDS Behav.* 2012; 16:295–303. [PubMed: 21380494]
9. Johnson M, Dilworth SE, Taylor JM, Neiland TB. Improving coping skills for self-management of treatment side effects can reduce antiretroviral medication nonadherence among people living with HIV. *Ann Behav Med.* 2011; 41:83–91. [PubMed: 20922510]
10. Webel AR. Testing a peer-based symptom management intervention for women living with HIV/AIDS. *AIDS Care.* 2010; 22:1029–1040. [PubMed: 20146111]
11. Wantland DJ, Holzemer WL, Moezzi S, et al. A randomized controlled trial testing the efficacy of an HIV/AIDS symptom management manual. *J Pain Symptom Manage.* 2008; 36:235–246. [PubMed: 18400461]
12. Schnall R, Wantland D, Velez O, Cato K, Jia H. Feasibility testing of a web-based symptom self-management system for persons living with HIV. *J Assoc Nurses AIDS Care.* 2014; 25:364–371. [PubMed: 24434198]
13. Gayner B, Esplen MJ, DeRoche P, et al. A randomized controlled trial of mindfulness-based stress reduction to manage affective symptoms and improve quality of life in gay men living with HIV. *J Behav Med.* 2012; 35:272–285. [PubMed: 21597980]
14. Duncan LG, Moskowitz JT, Neilands TB, et al. Mindfulness-based stress reduction for hiv treatment side effects: a randomized, wait-list controlled trial. *J Pain Symptom Manage.* 2012; 43:161–171. [PubMed: 21925831]
15. Birnie K, Speca M, Carlson LE. Exploring self-compassion and empathy in the context of mindfulness-based stress reduction (MBSR). *Stress and Health.* 2010; 26:359–371.
16. Neff K. Self-compassion: an alternative conceptualization of a healthy attitude toward oneself. *Self and Identity.* 2003; 2:85–101.
17. Neff KD, Brabeck KM, Kearney LK. Relationship styles of self-focused autonomy, other-focused connectedness, and mutuality across multiple relationship contexts. *J Soc Psychol.* 2006; 146:568–590. [PubMed: 17042403]
18. Brion JM, Leary MR, Drabkin AS. Self-compassion and reactions to serious illness: the case of HIV. *J Health Psychol.* 2014; 19:218–229. [PubMed: 23300046]
19. Raes F. The effect of self-compassion on the development of depression symptoms in a non-clinical sample. *Mindfulness.* 2011; 2:33–36.
20. Eicher AC, Davis LW, Lysaker PH. Self-compassion: a novel link with symptoms in schizophrenia? *J Nerv Ment Dis.* 2013; 201:389–393. [PubMed: 23588223]
21. Wren AA, Somers JT, Wright MA, et al. Self-compassion in patients with persistent musculoskeletal pain: relationship of self-compassion to adjustment to persistent pain. *J Pain Symptom Manage.* 2012; 43:759–770. [PubMed: 22071165]
22. Billings AG, Moos RH. The role of coping responses and social resources in attenuating the stress of life events. *J Behav Med.* 1981; 4:139–157. [PubMed: 7321033]
23. Portes A. Social capital: its origins and applications in modern sociology. *Annu Rev Sociol.* 1998; 24:1–24.

24. Bourdieu, P. The forms of capital. In: Richardson, J., editor. Handbook of theory and research for the sociology of education. New York: Greenwood; 1985. p. 241-58.
25. Bhattacharya G. Social capital and AIDS risk among acculturating Asian Indian men in New York. *AIDS Educ Prev.* 2005; 17:555–567. [PubMed: 16398577]
26. Frumence G, Killewo J, Kwegigabo G, et al. Social capital and the decline in HIV transmission—A case study in three villages in the Kagera region of Tanzania. *SAHARA.* 2010; 7:9–20.
27. Morrison SD, Howard R, Hardy C, Stinson B. Social capital, health and HIV awareness of girls in a rural Caribbean community. *Int Electron J Health Educ.* 2005; 8:135.
28. Cené CW, Akers AY, Lloyd SW, et al. Understanding social capital and HIV risk in rural African American communities. *J Gen Intern Med.* 2011; 26:737–744. [PubMed: 21311999]
29. Webel A, Phillips JC, Rose CD, et al. A cross-sectional description of social capital in an international sample of persons living with HIV/AIDS (PLWH). *BMC Public Health.* 2012; 12:188. [PubMed: 22414342]
30. Webel AR, Cuca Y, Okonsky JG, et al. The impact of social context on self-management in women living with HIV. *Soc Sci Med.* 2013; 87:147–154. [PubMed: 23631790]
31. Phillips JC, Webel AR, Rose CD, et al. Associations between the legal context of HIV, perceived social capital, and HIV antiretroviral adherence in North America. *BMC Public Health.* 2013; 13:736. [PubMed: 23924399]
32. Ware NC, Idoko J, Kaaya S, et al. Explaining adherence success in sub-Saharan Africa: an ethnographic study. *PLoS Med.* 2009; 6:e11. [PubMed: 19175285]
33. Merten S, Kenter E, McKenzie O, et al. Patient-reported barriers and drivers of adherence to antiretrovirals in sub-Saharan Africa: a meta-ethnography. *Trop Med Int Health.* 2010; 15:16–33. [PubMed: 20586957]
34. Fierz K, Nicca D, Spirig R. Perceived HIV symptom manageability: synthesis of a new use for a known concept. *J Adv Nurs.* 2013; 69:229–241. [PubMed: 22765386]
35. Webel AR, Longenecker CT, Gripshover B, et al. Age, stress, and isolation in older adults living with HIV. *AIDS Care.* 2014; 26:523–531. [PubMed: 24116852]
36. McGorry SY. Measurement in a cross-cultural environment: survey translation issues. *Qualitative Market Research: An International Journal.* 2000; 3:74–81.
37. Holzemer WL, Henry SB, Nokes KM, et al. Validation of the Sign and Symptom CheckList for Persons with HIV Disease (SSC-HIV). *J Adv Nurs.* 1999; 30:1041–1049. [PubMed: 10564402]
38. Holzemer WL, Henry SB, Portillo CJ, Miramontes H. The Client Adherence Profiling-Intervention Tailoring (CAP-IT) intervention for enhancing adherence to hiv/aids medications: a pilot study. *J Assoc Nurses AIDS Care.* 2000; 11:36–44. [PubMed: 10670005]
39. Sousa KH, Tann SS, Kwok OM. Reconsidering the assessment of symptom status in HIV/AIDS care. *J Assoc Nurses AIDS Care.* 2006; 17:36–46. [PubMed: 16800166]
40. Raes F, Pommier E, Neff KD, Van Gucht D. Construction and factorial validation of a short form of the Self-Compassion Scale. *Clin Psychol Psychother.* 2011; 18:250–255. [PubMed: 21584907]
41. Kempainen JK, Brion JM, Leary M, et al. Use of a brief version of the self-compassion inventory with an international sample of people with HIV/AIDS. *AIDS Care.* 2013; 25:1513–1519. [PubMed: 23527887]
42. Onyx J, Bullen JP. Measuring social capital in five communities. *J Appl Behav Sci.* 2000; 36:23.
43. Buys L, Godber AM, Summerville JA, Barnett K. Building community: collaborative individualism and the challenge for building social capital. *Aust J Reg Studies.* 2007; 13:165–176.
44. Pillai VK. Democratic attitudes and social capital in Latin America. *International Social Work.* 2011; 54:767–779.
45. George BP. Local community's support for post-tsunami recovery efforts in an agrarian village and a tourist destination: a comparative analysis. *Community Development J.* 2008; 43:444–458.
46. Dodd M, Janson S, Facione N, et al. Advancing the science of symptom management. *J Adv Nurs.* 2001; 33:668–676. [PubMed: 11298204]
47. Atkins DC, Gallop RJ. Rethinking how family researchers model infrequent outcomes: a tutorial on count regression and zero-inflated models. *J Fam Psychol.* 2007; 21:726–735. [PubMed: 18179344]

48. Lambert D. Zero-inflated Poisson regression, with an application to defects in manufacturing. *Technometrics*. 1992; 34:1–14.
49. Greene, WH. NYU Working Paper No. EC-94-10. Social Science Research Network; 1994. Accounting for excess zeros and sample selection in Poisson and negative binomial regression models.
50. Vuong QH. Likelihood ratio tests for model selection and non-nested hypotheses. *Econometrica*. 1989:307–333.
51. Rogers A, Vassilev I, Sanders C, et al. Social networks, work and network-based resources for the management of long-term conditions: a framework and study protocol for developing self-care support. *Implement Sci*. 2011; 6:56. [PubMed: 21619695]
52. Miaskowski C, Cooper BA, Melisko M, et al. Disease and treatment characteristics do not predict symptom occurrence profiles in oncology outpatients receiving chemotherapy. *Cancer*. 2014; 120:2371–2378. [PubMed: 24797450]
53. Lochner K, Kawachi I, Kennedy BP. Social capital: a guide to its measurement. *Health Place*. 1999; 5:259–270. [PubMed: 10984580]
54. Derose KP, Varda DM. Social capital and health care access. *Med Care Res Rev*. 2009; 66:272–306. [PubMed: 19174538]
55. Lewis JM, DiGiacomo M, Luckett T, Davison PM, Currow DC. A Social capital framework for palliative care: supporting health and well-being for people with life-limiting illness and their carers through social relations and networks. *J Pain Symptom Manage*. 2013; 45:92–103. [PubMed: 22795051]
56. Berkman LF, Glass T. Social integration, social networks, social support, and health. *Soc Epi*. 2000:137–173.
57. Boyatzis RE, Rochford K, Jack AI. Antagonistic neural networks underlying differentiated leadership roles. *Front Hum Neurosci*. 2014; 8:114. [PubMed: 24624074]
58. Sikorskii A, Given CW, Siddigi AE, et al. Testing the differential effects of symptom management interventions in cancer. *Psychooncology*. 2014 Apr 15. Epub ahead of print.
59. Bouchard L, Gilbert A, Landry R, Deveau K. Social capital, health, and Francophone minorities. *Can J Public Health*. 2006; 97:S16–20. [PubMed: 16805156]
60. Reeves D, Blickem C, Vassilev I, et al. The contribution of social networks to the health and self-management of patients with long-term conditions: a longitudinal study. *PLoS One*. 2014; 9:e98340. [PubMed: 24887107]
61. Riley KE, Kalichman S. Mindfulness-based stress reduction for people living with HIV/AIDS: preliminary review of intervention trial methodologies and findings. *Health Psychol Rev*. 2014:1–20.
62. Rueda S, Raboud J, Mustard C, et al. Employment status is associated with both physical and mental health quality of life in people living with HIV. *AIDS Care*. 2011; 23:435–443. [PubMed: 21271396]
63. Martin DJ, Chernoff RA, Buitron M, et al. Helping people with HIV/AIDS return to work: a randomized clinical trial. *Rehabil Psychol*. 2012; 57:280–289. [PubMed: 23148715]
64. Richardson LA, Milloy MJ, Kerr TH, et al. Employment predicts decreased mortality among HIV-seropositive illicit drug users in a setting of universal HIV care. *J Epidemiol Community Health*. 2014; 68:93–96. [PubMed: 24153247]
65. Rodger AJ, Brecker N, Bhagani S, et al. Attitudes and barriers to employment in HIV-positive patients. *Occup Med*. 2010; 60:423–429.

Table 1

## Demographic and HIV Disease Information

	Entire Sample (n=2,183)		30% Subsample (n=655)	
	Frequency (%)	Mean ( $\pm$ SD)	Frequency (%)	Mean ( $\pm$ SD)
Age (yrs)		45.1 (9.5)		44.9 (9.7)
Gender				
Male	1,486 (68.7)		453 (69.9)	
Female	623 (28.8)		180 (27.8)	
Transgender	42 (1.9)		12 (1.9)	
Race				
Asian/Pacific Islander	261 (12.1)		75 (11.6)	
African American/Black	854 (39.6)		274 (42.3)	
Hispanic/Latino	425 (19.7)		119 (18.4)	
Native American	69 (3.3)		20 (3.1)	
White/Anglo	488 (22.6)		143 (22.1)	
Education Level				
11th grade or less	601 (27.7)		192 (29.5)	
High School or GED	804 (37.2)		221 (33.9)	
2 yrs college/Associates Degree	461 (21.3)		142 (21.8)	
4 yrs college/Bachelor's Degree	224 (10.4)		74 (11.4)	
Master's Degree or Doctorate	74 (3.4)		23 (3.6)	
Work for Pay	538 (24.9)		164 (25.2)	
Has Health Insurance	1,553 (71.6)		468 (71.9)	
Has Co-Morbid Health Conditions	1,323 (64.7)		386 (63.0)	
Self-Reported HIV Indicators				
Year Diagnosed with HIV <sup>3</sup>		1999 (7.6)		1998 (7.6)
Prescribed Anti-Retroviral Therapy	1,775 (83.4)		514 (80.8)	
Undetectable Viral Load <sup>a,b</sup>	798 (36.5)		373 (59.1)	
Viral Load for those with detectable values /mL <sup>c</sup>		43,032 (109,646)		48,114 (128,333)
CD4 cells/ $\mu$ l		432 (472.8)		431.5 (489.7)
HIV Sign and Symptom Checklist				
Mean frequency of symptoms		18.41 (13.21)		18.74 (13.6)
Mean intensity of Symptoms		29.29 (24.50)		30.23 (26.7)
Total Self-Compassion Score		38.7 (7.5)		38.5 (7.7)
Total Social Capital Score <sup>3</sup>		2.6 (0.55)		2.54 (0.55)

<sup>a</sup>Subjects were asked if their HIV viral load was undetectable, detectable and report the actual value, or if they do not know their HIV viral load. We calculated those who are undetectable by summing those who reported having an undetectable HIV viral load with those who reported an HIV viral load less than 200 copies per mL and divided that by the total number of participants in the respective column.

<sup>b</sup>Difference between overall sample and subsample statistically significant with a *P*-value less than 0.01.

<sup>c</sup>189 subjects in the full dataset and 373 subjects in the subsample reported an HIV viral load value greater than or equal to 200 copies per mL. The mean reported is based on those subjects.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Table 2**

Bivariate Relationship Between Social Capital, Self-Compassion, Age, and Perceived HIV Symptom Intensity ( $n=655$ )<sup>a</sup>

	HIV Symptom Intensity	Social Capital	Self-Compassion
HIV Symptom Intensity	1.00		
Social Capital	-0.23 ( $P<0.001$ )	1.0	
Self-Compassion	-0.28 ( $P<0.001$ )	0.40 ( $P<0.001$ )	1.0
Age	0.07 ( $P=0.08$ )	0.14 ( $P<0.001$ )	0.15 ( $P<0.001$ )

<sup>a</sup>The relationships between the variables were not linear; therefore, we used Spearman's rank order correlations tests to analyze these relationships.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 3**  
Zero-Inflated Poisson Regression Model Predicting HIV Symptom Intensity in the Past 24 Hours

Poisson Count Model (for non-zero values of symptom intensity)				
Variable	B Coefficient	Standard Error	Z	95% Confidence Interval
Age	0.003 <sup>a</sup>	0.001	3.37	0.001 0.005
Sex (male)	-0.140 <sup>a</sup>	0.016	-8.63	-0.171 -0.108
Year of HIV Diagnosis	-0.001 <sup>a</sup>	0.001	-0.45	-0.003 0.002
Has other comorbid health conditions	0.286 <sup>a</sup>	0.017	16.85	0.253 0.319
Work	-0.099 <sup>a</sup>	0.020	-4.96	-0.139 -0.060
Income	-0.150 <sup>a</sup>	0.011	-13.10	-0.172 -0.127
Race				
Black	-0.287 <sup>a</sup>	0.028	-10.26	-0.342 -0.232
Hispanic	-0.006	0.029	-0.21	-0.063 0.051
White	-0.172 <sup>a</sup>	0.029	-5.95	-0.229 -0.115
Total Self-Compassion Score	-0.001	0.001	-1.03	-0.003 0.001
Total Social Capital Score	-0.004 <sup>a</sup>	0.001	-8.17	-0.005 -0.003
Constant	4.180	.067	61.99	4.05 4.313

Logit Model for Predicting Membership to "No Symptom" Group				
Variable	B Coefficient	Standard Error	Z	95% Confidence Interval
Age	-0.011	0.021	-0.51	0.609 -0.053
Sex (male)	0.427	0.449	0.96	-0.445 1.299
Year of HIV Diagnosis	-0.090	0.032	-2.78	-0.154 -0.027
Has other co-morbid health conditions	0.338	0.419	0.81	-0.483 1.158
Work	1.170 <sup>a</sup>	0.405	2.89	0.376 1.964
Income	-0.335	0.279	-1.20	-0.882 0.211
Race				
Black	-0.120	0.605	-0.20	-1.306 1.066

Logit Model for Predicting Membership to “No Symptom” Group					
Variable	B Coefficient	Standard Error	Z	95% Confidence Interval	
Hispanic	-2.173 <sup>a</sup>	0.928	-2.34	-3.991	-0.355
White	-1.905 <sup>a</sup>	0.919	-2.07	-3.706	-0.104
Total Self-Compassion Score	0.006	0.026	0.24	-0.044	0.057
<b>Total Social Capital Score</b>	<b>0.048<sup>a</sup></b>	<b>0.012</b>	<b>3.92</b>	<b>0.024</b>	<b>0.071</b>
Constant	-4.851	1.802	-2.69	-8.384	-1.319
Vuong Test of Zero Inflated Poisson Regression vs. Standard Poisson Regression					
			Z		5.77
				P-value	<0.001

<sup>a</sup>Indicates a *P*-value less than 0.05.

AU: pls define meaning of bolded values.