

Lawrence Berkeley National Laboratory

LBL Publications

Title

The Rate of Aquation of Dichlorotetraaquochromic Ion as a Function of pH in Chloride Media

Permalink

<https://escholarship.org/uc/item/0pd1b6pw>

Authors

Merideth, Charles W

Mathews, Wayne

Orlemann, Edwin F

Publication Date

1963-05-01

UCRL-10787

University of California

Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY


*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

THE RATE OF AQUATION
OF DICHLOROTETRAAQUOCHROMIC ION
AS A FUNCTION OF pH IN CHLORIDE MEDIA

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.



UCRL-10787

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

THE RATE OF AQUATION OF DICHLOROTETRAAQUOCHROMIC
ION AS A FUNCTION OF pH IN CHLORIDE MEDIA

Charles W. Merideth, Wayne Mathews and Edwin F. Orlemann

May 1963

THE RATE OF AQUATION OF DICHLOROTETRAAQUOCHROMIC ION
AS A FUNCTION OF pH IN CHLORIDE MEDIA

Charles W. Merideth, Wayne Mathews and Edwin F. Orlemann

Inorganic Materials Research Division of
Lawrence Radiation Laboratory and
Department of Chemistry
University of California
Berkeley, California

Abstract

The first order rate constant for the aquation of $\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2^+$ ions has been determined in chloride media of ionic strength 0.5 as a function of pH over the range 1.0 to 3.7 and found to be

$$k(\text{min.}^{-1}) = 0.0050 + \frac{1.55 \times 10^{-5}}{[\text{H}^+]}$$

The results are discussed in relation to existing data and in terms of a proposed mechanism.

THE RATE OF AQUATION OF DICHLOROTETRAAQUOCHROMIC ION
AS A FUNCTION OF pH IN CHLORIDE MEDIA

Charles W. Merideth, Wayne Mathews and Edwin F. Orlemann

Inorganic Materials Research Division of
Lawrence Radiation Laboratory and
Department of Chemistry
University of California
Berkeley, California

Introduction

1-5 The rate of aquation of dichlorotetraaquo chromic ions, hereafter referred to as CrCl_2^+ , has been reported by several investigators.¹⁻⁵ These workers disagree either in the dependence of the rate on hydrogen ion concentration or the value of the rate constant or both.

In the present work the rate of aquation was followed spectrophotometrically over a pH range of 1.0 to 3.7 and a concentration of 0.005 M to 0.008 M in CrCl_2^+ .

Experimental Method

A Cary model 11 spectrophotometer was used to measure the change in absorbancy of the solution at 700 m μ as a function of time. Stopped Corex 10-cm cells were employed and air was used as a reference. The absorbancy of the solutions without the chromic salt was measured against air and this value was subtracted from the absorbancy of the chromium solutions. All determinations were made at $25 \pm 0.05^\circ\text{C}$. Solid $(\text{CrCl}_2(\text{H}_2\text{O})_4)\text{Cl}\cdot 2\text{H}_2\text{O}$ was prepared by the method outlined by Elving and Zemel,⁶ and characterized by analysis of percent water and chloride. The

7 salt prepared in this manner has been shown to be predominantly the trans isomer.⁷ Weighed amounts of this salt were dissolved in solutions of HCl and NaCl of suitable ionic strength and the pH was determined by a Beckman model G pH meter.

Results and Calculations

Determination of the Rate Constants.

The following equation can be written for the total absorbancy of the solution at any time t :

$$A = \ell \left\{ a_1 [\text{CrCl}_2^+] + a_2 [\text{CrCl}^{++}] + a_3 [\text{Cr}^{+++}] \right\} \quad (1)$$

Since the rate of aquation of CrCl^{++} is known to be much smaller than that of the CrCl_2^+ species, the concentration of Cr^{+++} ion at the early stages of the experiment will be very small. This factor and the fact that the absorbancy index of the Cr^{+++} ion is approximately 0.6 at 700 m μ permits us to neglect the contribution of the Cr^{+++} ion to the total absorbancy at the early stages of the experiment. Within this approximation and neglecting the back reaction, the following expression can be derived for the total absorbancy as a function of time:

$$A = \ell [\text{CrCl}_2^+]_0 \left\{ a_1 e^{-kt} + a_2 (1 - e^{-kt}) \right\} \quad (2)$$

Dividing through by A_0 and rearranging, we can write

$$\log \left(\frac{A}{A_0} - \frac{a_2}{a_1} \right) = - \frac{kt}{2.3} + \log \left(\frac{a_1 - a_2}{a_1} \right) \quad (3)$$

where A = total absorbancy at time t
 A_0 = total absorbancy at time t = 0
 k = first-order rate constant for the aquation of CrCl_2^+
 a_1 = molar absorbancy index of CrCl_2^+
 a_2 = molar absorbancy index of CrCl^{++}
 $[\text{CrCl}_2^+]_0$ = molar concentration of $[\text{CrCl}_2^+]$ at t = 0
 l = length of cell


From this equation values of k at the various pH's were calculated.

F1 Figure 1 shows typical plots of eq. 3 for the lowest and highest pH used. The rate constant, k, was evaluated from the slope of the straight portion of these curves. In all determinations the linear region covers approximately two half-lives. At times greater than two half-lives the conversion of significant amounts of CrCl^{++} , $a_2 = 4.0$, to Cr^{+++} , $a_3 = 0.6$, leads to a larger decrease in light absorption with time than that predicted by eq. 3, as shown by the data in Fig. 1.

8 In analyzing the experimental data, a value of 4.0 was obtained for
 9 a_2 as an average of the data reported by Gates and King⁸ and Taube and Myers.⁹ For a_1 , King⁷ gives 11.6 for the cis isomer and 13.2 as the value for the trans isomer. Since the rate of isomerization of these species is unknown, we chose to use an average value of 12.4 for a_1 in the evaluation of our data. This leads to an uncertainty of $\pm 6\%$ in the ratio of a_2/a_1 , which in turn leads to an uncertainty of $\pm 3\%$ in the reported values of k.


Dependence of the Rate Constant on pH.

TI Table I shows the values of k measured at the various pH's and in
 F2 Fig. 2 we have plotted k versus $1/[\text{H}^+]$, where the symbol $[\text{H}^+]$ = antilog (-pH).



MU-30489

Fig. 1



MU-30488

Fig. 2

TABLE I
RATE OF AQUATION OF CrCl_2^+ AS A FUNCTION OF pH

pH	3.69	3.66	3.47	3.37	3.28	3.21	2.80	2.50	2.39	2.35	1.75	1.0
$k \times 10^3 (\text{min.}^{-1})$	70.80	69.20	42.80	35.40	31.10	25.40	13.50	9.45	10.40	7.22	4.72	4.7

The data in Fig. 2 were analyzed graphically to obtain the following equation for the rate of conversion of CrCl_2^+ to CrCl^{++} :

$$k = 0.0050 + \frac{1.55 \times 10^{-5}}{[\text{H}^+]'} \quad (4)$$

In evaluating these data, two effects that influence the pH during an experiment need to be considered. First there is the initial hydrolysis of the added CrCl_2^{++} , which gives rise to a change in pH that is easily measured before any significant amount of reaction has occurred. It is these measured initial pH values that are reported in Table I and Fig. 2. The second factor is the hydrolysis of the CrCl^+ species formed during the reaction, which gives rise to a continuous change in pH during the reaction. In the determination at pH = 3.69, for example, the pH decreased to 3.50 over a 24 min. interval as determined in a separate experiment. This should be born in mind in regard to the reliability of the data as given in Table I and Fig. 2 at pH greater than 3.3.

Discussion

Comparison with Existing Data.

The results of this investigation are compared with the existing data in Table II.

TABLE II
 REPORTED FIRST-ORDER RATE CONSTANTS FOR THE AQUATION OF
 DICHLOROTETRAAQUOCHROMIC ION

Investigator	Method	$k(\text{min.}^{-1})$	$k(\text{min.}^{-1})$ at 0.01M
Present study	Spectrophotometrically	$0.0050 + \frac{1.55 \times 10^{-5}}{[\text{H}^+]}$	$6.55 \times 10^{-3}{}^f$
Bjerrum ^a	Conductometrically	$0.00272 + \frac{1.62 \times 10^{-5}}{[\text{H}^+]}$	4.30×10^{-3}
Lamb and Fonda ^b	Conductometrically	$0.00265 + \frac{1.42 \times 10^{-5}}{[\text{H}^+]}$	4.10×10^{-3}
Nazarenko ^c	Argentimetric titration	...	5.30×10^{-3}
Hamm and Shull ^d	Dropping mercury electrode	$\frac{5.2 \times 10^{-6}}{[\text{H}^+]}$	5.20×10^{-4}
Ibraz, Virgili and Costa ^e	Dropping mercury electrode	$0.00095 + \frac{7.6 \times 10^{-9}}{[\text{H}^+]^2}$	10.30×10^{-4}

^a Ref. 1.

^b Ref. 2.


^c Ref. 3.

^d Ref. 4.

^e Ref. 5.

^f Referred to antilog (-pH) = 0.01; all other data referred to actual $[\text{H}^+]$.

It is apparent that our results are in substantial agreement with the data reported in refs. 1, 2, and 3. The disagreement shown by the entries for refs. 4 and 5 in Table II suggests a systematic error is involved in following this reaction at the dropping mercury electrode. In fact the rate law found for aquation of CrCl^{++} as determined in this Laboratory¹⁰ is in substantial agreement with that given in Table II for ref. 5. It would thus appear that the reaction actually being followed at the dropping mercury electrode is that of the aquation of CrCl^{++} . This can be explained by assuming that the chromous ion formed at the electrode rapidly converts CrCl_2^+ to CrCl^{++} according to the following fast reaction:






which has a rate constant⁹ of roughly $10^4 \text{ M}^{-1} \text{ min.}^{-1}$.

Mechanism.

Although the starting chromic salt used in these measurements has been shown to be predominately the trans isomer⁸, our data cannot be said to refer only to the rate of aquation of this isomer. The absence of significant experimental deviation of the data from eq. 3 as shown in Fig. 1 does not rule out the possibility that part of the actual reaction path is conversion of the original trans isomer to the corresponding cis isomer and the aquation of the latter species. In view of the term 0.0050 min.^{-1} in the rate eq. 4, this value may be taken as an upper limit for the aquation of the trans CrCl_2^{++} species.

The second term in eq. 4 represents the dependence of the forward reaction on the concentration of hydrogen ion. In light of the inverse hydrogen dependence the following mechanism is assumed, in agreement with that outlined by Hamm and Shull,⁴ to account for this term in the rate law.



11 This same type of mechanism is generally accepted in regard to many related reactions.¹¹

12 The rate constant for the aquation of $\text{CrCl}_2(\text{H}_2\text{O})_3\text{OH}$ is 6.21 as compared with 0.0050 for the $\text{CrCl}_2(\text{H}_2\text{O})_4^+$ species. This ratio of approximately 1240 is in agreement with that found in many analogous systems and is subject to the same general interpretation.¹²

Acknowledgement. This research was carried out with funds supplied by the U.S. Atomic Energy Commission Contract No. W-7405-eng-48.

References

- (1) N. Bjerrum, Z. physik.chem., 59, 336, 581 (1907).
- (2) A. B. Lamb and G. R. Fonda, J. Am. Chem. Soc., 43, 1154 (1921).
- (3) Yu. P. Nazarenko, Zhur. neorg. Khim., 4, 1829 (1959).
- (4) R. E. Hamm and C. M. Shull, Jr., J. Am. Chem. Soc., 73, 1240 (1951).
- (5) J. Ibarz, J. Virgili and J. M. Costa, Anales real. Soc. espan. Fis. Quim., 57B, 489 (1961); 58B, 89 (1962).
- (6) P. J. Elving and B. Zemel, J. Am. Chem. Soc., 79, 1281 (1957).
- (7) E. L. King, Sister M. J. M. Woods, O. P. and H. S. Gates, J. Am. Chem. Soc., 80, 5015 (1958).
- (8) H. S. Gates and E. L. King, J. Am. Chem. Soc., 80, 5011 (1958).
- (9) H. Taubé and H. Myers, J. Am. Chem. Soc., 76, 2103 (1954).
- (10) Unpublished results.
- (11) R. G. Pearson and F. Basolo, J. Am. Chem. Soc., 78, 4878 (1956).
- (12) R. Pearson and F. Basolo, "Mechanisms of Inorganic Reactions,"
John Wiley & Sons, Inc., New York, N. Y., 1958, Chapter 3.

Figure Captions

Figure 1. Experimental data analyzed by equation (3) at pH = 1.01
and 3.69.

Figure 2. The dependence of k on acidity at 25°C and $\mu = 0.5$ (NaCl added).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

