Lawrence Berkeley National Laboratory

Recent Work

Title

THERMODYNAMICS OF AQUEOUS CARBONATE SOLUTIONS INCLUDING MIXTURES OF SODIUM CARBONATE, BICARBONATE AND CHLORIDE

Permalink

<https://escholarship.org/uc/item/0q17p49d>

Authors

Peiper, J.C.
Pitzer, K.S.

Publication Date

1981-05-01

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA, BERKELEY

EARTH SCIENCES DIVISION

RECEIVED
LAWRENCE
BERKELEY LABORATORY

JUN 17 1981

Submitted to the Journal of Chemical Thermodynamics
LIBRARY AND DOCUMENTATION SECTION

THERMODYNAMICS OF AQUEOUS CARBONATE SOLUTIONS INCLUDING
MIXTURES OF SODIUM CARBONATE, BICARBONATE AND CHLORIDE

J. Christopher Peiper and Kenneth S. Pitzer

May 1981

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782

LBL-12725
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THERMODYNAMICS OF AQUEOUS CARBONATE SOLUTIONS INCLUDING
MIXTURES OF SODIUM CARBONATE, BICARBONATE AND CHLORIDE

by

J. Christopher Peiper and Kenneth S. Pitzer

This work was supported by the Director, Office of Energy Research,
Office of Basic Energy Sciences, Division of Engineering, Mathematics,
and Geosciences of the U. S. Department of Energy under Contract No.
W-7405-ENG-48.

Department of Chemistry
and
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Abstract

Recently the authors examined electrochemical cell data leading to values of the activity coefficient for aqueous sodium bicarbonate. Since that preliminary analysis, new experimental measurements have been published which contribute significantly to the overall thermodynamic understanding of the sodium carbonate - sodium bicarbonate - carbonic acid system. In this more extensive examination we consider a wide variety of measurements leading to activity coefficients of Na_2CO_3 and NaHCO_3 from 273 K to 323 K and to relative molal enthalpies and heat capacities at 298 K. Tables of thermodynamic quantities at selected temperatures are included.

1. Introduction

Measured thermodynamic properties of aqueous solutions of Na_2CO_3 and NaHCO_3 cannot be interpreted in the same straightforward manner applicable to simple strong electrolytes. The presence of the equilibria

requires that even stoichiometrically single-solute solutions be treated as mixed electrolytes. In addition, the presence of doubly- and singly-charged anions implies that the electrostatic effects of asymmetric mixing be incorporated into any complete analysis of experimental data. Thus, a full interpretation of the carbonate system becomes quite complex.

While the aqueous solutions of sodium carbonate and bicarbonate themselves are of considerable importance, naturally occurring solutions include these ions among a wider array of other ions [1,2]. Chloride is especially common and is included in this study. But sulfate, magnesium, potassium and calcium are often present with smaller concentrations of other species. The parameters of this study are determined within a general formalism which is applicable to mixed electrolytes of unlimited complexity. Thus, the present results for sodium carbonate and bicarbonate may be combined with corresponding parameters for other ions to predict the properties of a wide range of natural and artificial solutions.

A further objective of this study is to demonstrate the modeling of a complex mixture of weak and strong electrolytes within the framework of Pitzer's ion interaction theory [3]. This theory was originally formulated for single-solute solutions of strong electrolytes at 298 K [4], and at

high temperature [5], but was subsequently expanded to include mixtures of strong electrolytes [6], and higher valence type electrolytes [7,8]. Aqueous H_3PO_4 [9] and H_2SO_4 [10] were successfully treated as solutions with a single ionization equilibrium. The carbonate solutions considered herein are more complex in having two reactions at equilibrium. Also, we self-consistently treat enthalpy and heat capacity data as well as measurements leading to activity and osmotic coefficients.

The experimental data here considered include the emf measurements of Harned and Davis [11] and Harned and Bonner [12] which formed the basis of our earlier report on sodium bicarbonate [13]. These concern the cell

with reaction

Recently, Roy, et. al., [14] have made additional measurements for the same electrochemical cell but at significantly higher molalities of NaHCO_3 and NaCl .

The activity coefficient of Na_2CO_3 is obtainable at low molalities from the work of Harned and Scholes [15], who measured the potentials of similar cells with solutions containing NaCl , NaHCO_3 and Na_2CO_3 and without $\text{CO}_2(\text{g})$. The cell reaction for these measurements is

Robinson and Macaskill [16] have recently measured the activity of water above sodium carbonate solutions at 298 K and to 3.0 mol kg^{-1} by an isopiestic method. Their determinations are supplemented by similar measurements on sodium carbonate - sodium chloride mixtures by White and Bates [17].

For the enthalpy of these systems we adopt the recent results of Berg and Vanderzee [18]. These authors measured the enthalpy of dilution of sodium carbonate and sodium bicarbonate solutions at 298 K, and corrected their measurements for heat effects arising from equilibria (1). Heat capacities for these solutions have been measured by Perron, Desnoyers and Millero [19] at 298 K, who report data for other systems as well.

Older data for aqueous sodium carbonate solutions are listed and evaluated by Goldberg [20] in his review. It did not appear to us that any of these data would contribute significantly to our analysis; hence for simplicity they were omitted. Goldberg's analysis of the data for sodium carbonate ignored the hydrolysis equilibrium above.

2. Equations

We adopt the ion interaction or virial coefficient equations for mixed electrolytes introduced by Pitzer and Kim [6]. These authors were able to represent quite accurately the experimental activity and osmotic coefficient information for some sixty aqueous systems with and without a common ion. Recently, Harvie and Weare [21] in an extensive study of complex seawater related mixed electrolyte systems have also found these equations to be very satisfactory.

It is convenient to begin with the excess Gibbs energy for a solution containing 1.0 kg of water; equation (5) is written in exactly the form

from a recent review [22], but it is equivalent to the expressions of Pitzer and Kim and of Harvie and Weare.

$$\begin{aligned} \frac{G^{ex}}{RT} = & f + 2 \sum_{c,a} m_c m_a [B_{c,a} + (\sum m_z) C_{c,a}] \\ & + \sum_{c,c'} m_c m_{c'} [\theta_{c,c'} + \frac{1}{2} m_a \psi_{c,c',a}] \\ & + \sum_{a,a'} m_a m_{a'} [\theta_{a,a'} + \frac{1}{2} m_c \psi_{c,a,a'}] \end{aligned} \quad (5)$$

$$f = - \frac{4I}{1.2} A_\phi \ln(1 + 1.2I^{1/2})$$

$$B_{c,a} = \beta_{c,a}^{(0)} + \frac{\beta_{c,a}^{(1)}}{2I} [1 - (1 + 2I^{1/2})e^{-2I^{1/2}}]$$

$$C_{c,a} = \frac{C_{c,a}^\phi}{2|z_c z_a|^{1/2}}$$

The quantities $B_{c,a}$ and $C_{c,a}$ are the second and third virial coefficients for the electrolyte ca and f is the appropriate Debye-Huckel function with parameter A_ϕ . The summations are over all cations (c,c') and anions (a,a') with the quantity $(\sum m_z) = \sum m_c z_c = \sum m_a |z_a|$, where m_i and z_i represent the ionic molality and charge. The $\theta_{i,j}$ are second virial coefficients representing the difference in averaged interactions of unlike ions with charges of the same sign from the appropriate average for like ions. The $\psi_{i,j,k}$ are third virial coefficients similarly defined but for three ions with charges not all the same sign. These terms are discussed more fully by Pitzer and Kim. Finally I represents the molal ionic strength and R , T retain their usual meanings.

Equation (5) defines a Gibbs energy in excess of that for an ideal solution of the same composition. Ordinarily one obtains the ideal

solution by a simple dilution process; in this case the equilibria (1) must be frozen during the dilution. Otherwise, for the real solution, this process involves the interconversion of some small amount of solute due to the composition dependence of (1). Our analysis for a carbonate solution always involves determination of the equilibrium composition and subsequent evaluation of a nonideality term related to equation (5).

In anticipation of the activity coefficient ratios required in the analysis of emf data from the laboratories of Harned and of Roy, we present equations (6) and (7). Equation (6) is merely equation (17) of Pitzer and Kim's report, rewritten here in explicit form for solutions containing Na_2CO_3 , NaHCO_3 , NaOH and NaCl . Equation (7) may be derived from equation (15) of Pitzer and Kim by appropriate combination of mean ionic activity coefficients.

$$\begin{aligned}
 \ln \left[\frac{\gamma_{\text{Cl}}}{\gamma_{\text{HCO}_3}} \right] &= 2 \ln(\gamma_{\text{NaCl}}) - 2 \ln(\gamma_{\text{NaHCO}_3}) \\
 &= 2m_{\text{Na}} (B_{\text{Na,Cl}} - B_{\text{Na,HCO}_3}) \\
 &\quad + 2m_{\text{Na}}^2 (C_{\text{Na,Cl}} - C_{\text{Na,HCO}_3}) \\
 &\quad + 2(m_{\text{HCO}_3} - m_{\text{Cl}}) \theta_{\text{HCO}_3,\text{Cl}} \\
 &\quad + m_{\text{Na}} (m_{\text{HCO}_3} - m_{\text{Cl}}) \psi_{\text{Na,HCO}_3,\text{Cl}} \\
 &\quad + 2m_{\text{CO}_3} (\theta_{\text{CO}_3,\text{Cl}} - \theta_{\text{CO}_3,\text{HCO}_3}) \\
 &\quad + m_{\text{Na}} m_{\text{CO}_3} (\psi_{\text{Na,CO}_3,\text{Cl}} - \psi_{\text{Na,CO}_3,\text{HCO}_3}) \\
 &\quad + 2m_{\text{OH}} (\theta_{\text{OH,Cl}} - \theta_{\text{HCO}_3,\text{OH}}) \\
 &\quad + m_{\text{Na}} m_{\text{OH}} (\psi_{\text{Na,OH,Cl}} - \psi_{\text{Na,HCO}_3,\text{OH}})
 \end{aligned} \tag{6}$$

$$\begin{aligned}
 \ln \left[\frac{\gamma_{\text{HCO}_3} \gamma_{\text{Cl}}}{\gamma_{\text{CO}_3}} \right] &= 2 \ln(\gamma_{\text{NaHCO}_3}) + 2 \ln(\gamma_{\text{NaCl}}) - 3 \ln(\gamma_{\text{Na}_2\text{CO}_3}) \\
 &= 2A_\phi \left[\frac{I^{1/2}}{1 + 1.2I^{1/2}} + \frac{2}{1.2} \ln(1 + 1.2I^{1/2}) \right] \\
 &\quad + 2m_{\text{Na}} (B_{\text{Na},\text{HCO}_3} + B_{\text{Na},\text{Cl}} - B_{\text{Na},\text{CO}_3}) \\
 &\quad + 2m_{\text{Na}}^2 (C_{\text{Na},\text{HCO}_3} + C_{\text{Na},\text{Cl}} - C_{\text{Na},\text{CO}_3}) \\
 &\quad - 2m_{\text{Na}} (m_{\text{CO}_3} B'_{\text{Na},\text{CO}_3} + m_{\text{HCO}_3} B'_{\text{Na},\text{HCO}_3} \\
 &\quad \quad + m_{\text{OH}} B'_{\text{Na},\text{OH}} + m_{\text{Cl}} B'_{\text{Na},\text{Cl}}) \\
 &\quad + 2m_{\text{CO}_3} (\theta_{\text{CO}_3,\text{HCO}_3} + \theta_{\text{CO}_3,\text{Cl}}) \\
 &\quad + m_{\text{Na}} m_{\text{CO}_3} (\psi_{\text{Na},\text{CO}_3,\text{HCO}_3} + \psi_{\text{Na},\text{CO}_3,\text{Cl}}) \quad (7) \\
 &\quad + 2m_{\text{HCO}_3} (\theta_{\text{HCO}_3,\text{Cl}} - \theta_{\text{CO}_3,\text{HCO}_3}) \\
 &\quad + m_{\text{Na}} m_{\text{HCO}_3} (\psi_{\text{Na},\text{HCO}_3,\text{Cl}} - \psi_{\text{Na},\text{CO}_3,\text{HCO}_3}) \\
 &\quad + 2m_{\text{OH}} (\theta_{\text{HCO}_3,\text{OH}} + \theta_{\text{OH},\text{Cl}} - \theta_{\text{CO}_3,\text{OH}}) \\
 &\quad + 2m_{\text{Na}} m_{\text{OH}} (\psi_{\text{Na},\text{HCO}_3,\text{OH}} + \psi_{\text{Na},\text{OH},\text{Cl}} - \psi_{\text{Na},\text{CO}_3,\text{OH}}) \\
 &\quad + 2m_{\text{Cl}} (\theta_{\text{HCO}_3,\text{Cl}} - \theta_{\text{CO}_3,\text{Cl}}) \\
 &\quad + m_{\text{Na}} m_{\text{Cl}} (\psi_{\text{Na},\text{HCO}_3,\text{Cl}} - \psi_{\text{Na},\text{CO}_3,\text{Cl}}) \\
 &\quad - 2m_{\text{CO}_3} (m_{\text{HCO}_3} + m_{\text{OH}} + m_{\text{Cl}})^{E_\theta}_{2,1}
 \end{aligned}$$

The charge has been omitted from all ionic species appearing as subscripts in these equations. The final term of equation (7) is explained below.

Also,

$$B'_{c,a} = - \frac{\beta_{c,a}^{(1)}}{2I^2} [1 - (1 + 2I^{1/2} + 2I^2)e^{-2I^{1/2}}]$$

The activity of water may be obtained directly from equation (11) of Pitzer and Kim [6] by the definition

$$\ln(a_w) = - \frac{M_w}{1000} (\sum m_i) \phi$$

wherein M_w represents the formula weight of water (taken as 18.01533 g mol⁻¹) and $\sum m_i$ the sum of molalities of all ionic and neutral species when the solution is at equilibrium with respect to (1). ϕ is the osmotic coefficient of the solution.

As noted in the introduction, carbonate solutions must be discussed within the framework of asymmetric mixing theory since at equilibrium the doubly-charged CO_3^{2-} species coexists with singly-charged HCO_3^- and OH^- ions. Pitzer [23] derived the additional electrostatic terms, first noted by Friedman [24], arising from asymmetric mixing. He found that inclusion of these terms is, from the standpoint of modeling experimental emf measurements, required for 3:1 and higher charge type electrolyte mixtures, but for many 2:1 and 1:2 mixtures the experimental data may be adequately represented without them. Harvie and Weare [21] found the full electrostatic formulation essential for accurate treatment of CaSO_4 solubility in aqueous NaCl. Employing the notation of Pitzer,

$$\begin{aligned} \theta_{c,c'} &= E_{\theta_{c,c'}} + S_{\theta_{c,c'}} \\ \theta'_{c,c'} &= E_{\theta'_{c,c'}} + S_{\theta'_{c,c'}} \end{aligned} \tag{8}$$

wherein $E_{\theta_{c,c'}}$ and $E_{\theta'_{c,c'}}$ represent the higher order electrostatic effects of asymmetric mixing. These terms may be calculated a priori and are complex

functions of the charges on ions c and c' , the ionic strength and temperature; if the charges on c and c' are equal, these terms vanish. $S_{\theta_{c,c'}}$ and $S_{\theta'_{c,c'}}$ represent the effect of short-range interactions between ions c and c' , and must be evaluated empirically. It has been found in previous investigations [6-10] that the $S_{\theta'_{c,c'}}$ terms are negligible, hence they are omitted in this analysis. The remaining nonvanishing terms in $E_{\theta'_{c,c'}}$ may be collected as indicated in equation (7), with the subscript indicating only the magnitude of charge on each of the ions $c = \text{CO}_3^{2-}$ and $c' = \text{HCO}_3^-$, OH^- and Cl^- .

The excess enthalpy L and heat capacity J of the solution are obtained from temperature derivatives of equation (5) in the usual manner. If all θ and ψ parameters are assumed to have vanishing temperature derivatives, one obtains

$$L = \frac{A_H I}{1.2} \ln(1 + 1.2I^{1/2}) - 2RT^2 \sum_{c,a} m_c m_a [B_{c,a}^T + (\sum m_z) C_{c,a}^T] \quad (9)$$

$$J = \frac{A_J I}{1.2} \ln(1 + 1.2I^{1/2}) - 2RT^2 \sum_{c,a} m_c m_a [B_{c,a}^{TT} + (\sum m_z) C_{c,a}^{TT}] \quad (10)$$

$$B_{c,a}^T = \left[\frac{\partial B_{c,a}}{\partial T} \right]_P \quad ; \quad B_{c,a}^{TT} = \left[\frac{\partial^2 B_{c,a}}{\partial T^2} \right]_P + \frac{2}{T} \left[\frac{\partial B_{c,a}}{\partial T} \right]_P$$

$$C_{c,a}^T = \left[\frac{\partial C_{c,a}}{\partial T} \right]_P \quad ; \quad C_{c,a}^{TT} = \left[\frac{\partial^2 C_{c,a}}{\partial T^2} \right]_P + \frac{2}{T} \left[\frac{\partial C_{c,a}}{\partial T} \right]_P$$

The Debye-Huckel slopes A_H and A_J are defined in accordance with the results of Bradley and Pitzer [25], who give values for a wide range of temperatures and pressures.

The enthalpy of dilution for a mixture of electrolytes from some set of initial ionic molalities $\underline{m}^i = (m_1^i, m_2^i, \dots)$ to a final set of ionic molalities $\underline{m}^f = (m_1^f, m_2^f, \dots)$ is

$$\Delta H_d = L(\underline{m}^f) - L(\underline{m}^i) + \overline{\Delta H^0}(\underline{m}^f, \underline{m}^i) \quad (11)$$

The last term of this formula represents the ideal enthalpy change accompanying the interconversion of the appropriate amount of material required to shift the solution composition from $\underline{m}^i$ to $\underline{m}^f$; for the carbonate system the reactions governing this interconversion are given in (1).

Berg and Vanderzee [18] discuss this effect in some detail.

The enthalpy of solution of a salt (or mixture of pure solid salts) with net composition $\underline{m}^s$ in a solution of equilibrium composition $\underline{m}^i$, giving a solution with equilibrium composition $\underline{m}^f$ is

$$\Delta H_s = \overline{\Delta H_s^0}(\underline{m}^s) + \overline{\Delta H^0}(\underline{m}^f, \underline{m}^i + \underline{m}^s) + L(\underline{m}^f) - L(\underline{m}^i) \quad (12)$$

wherein $\overline{\Delta H_s^0}$ represents the ideal enthalpy of solution.

Equations (11) and (12) are applicable only to processes in which the temperature remains constant. The more general form of (11) may be found by integrating the total differential of the enthalpy

$$dH = \left(\frac{\partial H}{\partial T}\right)_{\underline{m}} dT + \left(\frac{\partial H}{\partial \underline{m}}\right)_T d\underline{m} \quad (13)$$

from $(T_i, \underline{m}^i)$ to $(T_f, \underline{m}^f)$. The heat of solution may be formed by adding to this result the ideal enthalpy of solution, $\overline{\Delta H_s^0}(\underline{m}^s)$, and correcting the ideal enthalpy change $\overline{\Delta H^0}$ to include $\underline{m}^s$ as in (12).

The heat capacity of a solution with some or all ionic species obeying a set of equilibria such as (1) includes the heat capacity effect

due to the change of equilibrium composition with temperature. This effect arises from the last term of (13), and was overlooked by Perron, Desnoyers and Millero [19], but was included by Woolley and Hepler [26] for some other systems. It is assumed in the following discussion that the kinetics of these equilibria are fast in comparison to the time required to measure the heat capacity. For some flow microcalorimeters the latter may be less than one second. In the present case, the hydrolysis of CO_3^{2-} , equation (1), attains equilibrium essentially instantaneously. However, it is widely known that the interconversion of $\text{CO}_2(\text{g})$ and carbonic acid is slow, and thus the disproportionation of HCO_3^- , equation (1), may not attain equilibrium within the time span of an heat capacity measurement. Fortunately the heat of disproportionation is small so that the error incurred in this analysis by assuming that equilibrium has been achieved is also small. Our calculations indicated that this error contributes negligibly to the heat capacity.

Determinations of the solution heat capacity commonly measure the small but finite rise in temperature ΔT accompanying the addition, usually via an electrical resistance heater, of an amount of heat Q under rigorously controlled conditions. The experimental heat capacity is then taken as the finite difference approximation $Q/\Delta T$. For systems which obey a set of temperature-dependent interconversion equilibria one must observe that contributions arising from these reactions are included in $Q/\Delta T$. If the solution is initially at temperature T_i with equilibrium composition $\tilde{m}^i$ and during the course of an heat capacity determination the temperature rises to $T_f = T_i + \Delta T$, causing a displacement in equilibrium composition to $\tilde{m}^f$, then

$$C_P = \frac{Q}{\Delta T} = \frac{[H(T_f, \underline{m}^f) - H(T_i, \underline{m}^i)]}{\Delta T}$$

wherein H represents the total enthalpy of the solution.

Separation of the direct effect of the temperature rise upon the solution enthalpy from its indirect effect (through the change in equilibrium composition) may be achieved by forming the total differential of the enthalpy (13). Division by dT and slight rearrangement yields

$$C_P = \frac{dH}{dT} = \left(\frac{\partial H}{\partial T}\right)_{\underline{m}} + \sum_k \left(\frac{\partial H}{\partial \alpha_k}\right)_{T, \underline{\alpha}^*} \frac{d\alpha_k}{dT} \quad (14)$$

The unique set of parameters $\underline{\alpha}$ indicate the degree to which the interconversion equilibria are displaced from the initial composition $\underline{m}^i$ to yield the final composition $\underline{m}^f$. $\underline{\alpha}^*$ is written simply to indicate that all parameters with the exception of α_k are held constant; further, the units on the α_i are moles.

The first term on the right in equation (14) may be identified as the heat capacity at constant composition $C_{P, \underline{m}}(\underline{m})$ with which $Q/\Delta T$ is normally equated. This quantity may be separated into ideal and non-ideal terms

$$C_{P, \underline{m}}(\underline{m}) = C_P^0(\underline{m}) + J(\underline{m}) \quad (15)$$

where $J(\underline{m})$ is given by equation (10). The remaining partial derivatives in equation (14) may be seen as equivalent to the molar enthalpy changes of the interconversion reactions. For reaction k,

$$\left(\frac{\partial H}{\partial \alpha_k}\right)_{T, \underline{\alpha}^*} = \overline{\Delta H}_k(\underline{m})$$

As for the heat capacities, this quantity may be partitioned into ideal

and non-ideal contributions, thus

$$\Delta H_k(\underline{m}) = \overline{\Delta H}_k^0 + \overline{\Delta L}_k(\underline{m}) \quad (16)$$

Substitution of (15) and (16) into (14) yields

$$C_p(\underline{m}) = C_p^0(\underline{m}) + J(\underline{m}) + \sum_k \overline{\Delta H}_k^0 \frac{d\alpha_k}{dT} + \sum_k \overline{\Delta L}_k(\underline{m}) \frac{d\alpha_k}{dT}$$

Although entirely correct, this last expression is cumbersome since equations for the temperature derivatives of the α_k are complicated, e.g., equation (22) of Pitzer, Roy and Silvester [10]. However, if the equilibrium composition of the solution over a range of temperature ($T_i, T_i + \Delta T$) is known to sufficient accuracy, it is possible to numerically calculate the finite difference quantity $\Delta\alpha_k/\Delta T$. With this substitution, the summation in the final term may be evaluated explicitly to give

$$\sum_k [\overline{\Delta L}_k(\underline{m})][\Delta\alpha_k] = L(\underline{m}^f) - L(\underline{m}^i)$$

wherein L is the relative enthalpy given by equation (9). Note that these enthalpies derive from the constant temperature partial derivatives of the solution enthalpy; as such they are evaluated at T . Thus our final equation for the heat capacity is

$$C_p(\underline{m}) = C_p^0(\underline{m}) + J(\underline{m}) + \sum_k \overline{\Delta H}_k^0 \frac{\Delta\alpha_k}{\Delta T} + \frac{1}{\Delta T} [L(\underline{m}^f) - L(\underline{m}^i)] \quad (17)$$

Equations (9) to (17) have been formulated "per kilogram of solvent." Also, all enthalpies and heat capacities appearing in this equation are evaluated at T , which may be taken as the midpoint of the temperature interval of experimental measurement.

3. Method of Calculation

Before discussing in detail the method used in the present analysis it is convenient to explain all data manipulation and conversion performed prior to that analysis.

Harned and Davis [11], Harned and Bonner [12] and Roy, et. al., [14] measured potentials for cell (2), which are given by

$$E = E_{\text{Ag,AgCl}}^{\circ} - \frac{RT}{F} \left[\ln(K_1 K_S) + \ln\left(\frac{a_w^f \text{CO}_2}{f_{\text{H}_2}^{1/2}}\right) + \ln\left(\frac{m_{\text{Cl}} \gamma_{\text{Cl}}}{m_{\text{HCO}_3} \gamma_{\text{HCO}_3}}\right) \right] \quad (18)$$

The measurements of Harned and Scholes [15] yield potentials according to

$$E = E_{\text{Ag,AgCl}}^{\circ} - \frac{RT}{F} \left[\ln\left(\frac{K_2}{f_{\text{H}_2}^{1/2}}\right) + \ln\left(\frac{m_{\text{Cl}} m_{\text{HCO}_3} \gamma_{\text{Cl}} \gamma_{\text{HCO}_3}}{m_{\text{CO}_3} \gamma_{\text{CO}_3}}\right) \right] \quad (19)$$

In these equations: $E_{\text{Ag,AgCl}}^{\circ}$ is the standard half-cell potential for the silver - silver chloride electrode; K_1 , K_2 and K_S are the first and second ionization constants of carbonic acid and the Henry's law constant for solution of CO_2 , respectively; a_w and f_i are the activity of water and the fugacity of gaseous species i ; and m_j , γ_j represent the molality and activity coefficient of ionic species j when the solution is at equilibrium with respect to (1). R , T and F retain their usual meanings; values for R and F , as well as other physical constants used in this analysis, were taken from the work of Cohen and Taylor [27].

The emf measurements of Roy, et. al., were used without manipulation. However, the older measurements from Harned's laboratory were based upon the international volt standard in use at that time. We have converted these potentials to the modern definition of the absolute volt by dividing

Harned's measurements by the factor $1.00033 \text{ int } V (\text{abs } V)^{-1}$. Also, Bates and Macaskill [28] and Bates, et. al., [29] have found that silver - silver halide electrodes are extremely sensitive to slight variations in preparative technique, and that, if high precision is desired, a standardization of the silver - silver halide electrode should be carried out. In brief, the difference $E - E^0$ is more reproducible from laboratory to laboratory than the absolute potential E . Therefore, in this analysis we have used the Ag,AgCl standard cell potentials of Harned and Ehlers [30] in interpreting measurements from Harned's laboratory and the similar measurements included in the report of Roy, et. al., in interpreting their own emf's.

The isopiestic determinations of Robinson and Macaskill [16] and White and Bates [17] were performed using sodium chloride reference solutions. Our analysis of these data involves converting this reference molality into an activity of water using the known properties of NaCl solutions at 298 K [4,5].

The enthalpies of dilution reported by Berg and Vanderzee [18] could be incorporated into the present analysis through equation (11) in a comprehensive simultaneous treatment of all experimental data. However, these authors have already corrected their apparent molal enthalpies for the effects of equilibria (1), with a treatment based largely upon the emf work of Harned and coworkers. Although the first method is preferable in terms of self-consistency of the calculation, the additional complexity seems hardly justified since our corrections would not differ significantly from those made by Berg and Vanderzee. Therefore, we determined the Na_2CO_3 and NaHCO_3 coefficients of equation (9) via independent least squares fits to the corrected enthalpies of Berg and Vanderzee. The results of those calculations are given in Table 1 with other results.

The heat capacities of Perron, Desnoyers and Millero [19] were incorporated into the present analysis in the following manner: Perron, et. al., measured solution heat capacities using a flow microcalorimeter and vibrating tube densimeter. These instruments directly determine the difference in volumetric heat capacity and density, respectively, between an aqueous electrolyte solution and pure water. Unfortunately, their measurements were later found to be slightly in error by Desnoyers, de Visser, Perron and Picker [31], due to experimental difficulties with the microcalorimeter, but were also found to be correctable. That correction involves multiplying the volumetric heat capacity difference, $\Delta\sigma$ in Perron's formalism, by the factor 1.02. The corrected measurements of Perron, et. al., were converted to total heat capacities for an amount of solution containing 1.0 kg of water.

In calculating the activity of water it is necessary to know the molality of dissolved but unionized carbon dioxide, which requires an accurate knowledge of K_S alone. This information may be taken from the work of Harned and Davis [11]. Their measurements of CO_2 solubility as a function of temperature and ionic strength may be reproduced within experimental error by an equation of the form adopted by them

$$\ln(K_S) = \frac{q_1}{T} + q_2 + q_3T \quad (20)$$

We found that the composition dependence of each parameter could be represented by

$$q_j = a_j + b_j I + c_j \exp(d_j I^{1.3}) \quad (21)$$

The coefficients a_j , b_j , c_j , and d_j are given in Table 2.

The equilibrium constant for the self-ionization of water

is also required. Grenthe, Ots and Ginstrup [32] reported precise measurements of the enthalpy of ionization in the 273 K to 323 K range, and fitted their results to an equation which may be integrated to yield

$$\begin{aligned} \ln(K_w) = & -32.234 + 50103.2\left(\frac{T-T_r}{TT_r}\right) - 330.089\ln\left(\frac{T}{T_r}\right) \\ & + 0.8457096(T-T_r) - 3.801538 \times 10^{-4}(T^2-T_r^2) \end{aligned} \quad (22)$$

with $T_r = 298.15$ K. The value of K_w at 298 K was taken from the report of Guggenheim and Turgeon [33], who thoroughly reanalysed the emf determinations of Harned and his students [34-36]. Recently, Olofsson and Olofsson [37] and Olofsson and Hepler [38] reviewed the available information concerning the ionization of water, however these authors were primarily interested in temperatures above those of interest in this report. In the region of overlap, 298 K to 323 K, their "best" values agree very well with values of K_w and its temperature derivatives calculated from the equation above.

A non-linear least squares procedure was used to evaluate the unknown virial parameters. However, due to equilibria (1), this approach is somewhat complicated since the preceding equations require a knowledge of the detailed ionic composition. Given estimates of the unknown parameters, ionic molalities may be calculated by an iterative procedure from the usual activity expressions for (1). The activity coefficient ratios for these expressions are readily derived from the equations in references [6] or [22]. The resulting equations are similar to equations

(6) and (7) above and involve the same parameters. The calculation of ionic molalities must be performed in a manner which allows no material to be expelled from or taken into the system, with the exception of measurements made under a constant and known pressure of CO_2 . Since the higher order electrostatic contributions, E_θ , and $E_{\theta'}$ depend upon the solution ionic strength, these must also be recalculated. At this point the normal least squares calculation may proceed for that iteration, and the entire process repeated as necessary for convergence.

The quantities which were determined via the iterative least squares process include: the product $K_1 K_S$, K_2 , the heat capacities at infinite dilution of Na_2CO_3 and NaHCO_3 , $C_p^O(\text{Na}_2\text{CO}_3)$ and $C_p^O(\text{NaHCO}_3)$, and the minimal number of virial coefficients and ion interaction parameters required to reproduce measurements to within experimental error. Of the pure-electrolyte parameters ($\beta^{(0)}$, $\beta^{(1)}$, and C^ϕ) only the third virial C^ϕ for NaHCO_3 was not needed. The temperature dependence of $\beta^{(0)}$ and $\beta^{(1)}$ was recognized in all cases. For NaOH , these temperature derivatives as well as that for C^ϕ are independently available from Silvester and Pitzer [39] and from fits of Perron, Desnoyers and Millero's [19] heat capacity data; temperature derivatives of $\beta^{(0)}$, $\beta^{(1)}$ and C^ϕ for NaCl were not required as no enthalpy or heat capacity measurements were performed on solutions containing NaCl . In no case was it necessary to recognize the temperature dependency of the virial coefficients for mixing ions of the same sign, θ and ψ . In all there are 12 parameters for mixing ions of the same sign of which two were known independently, three were evaluated from the present data, and seven were not needed.

Weighting for the least squares calculation was based principally upon the uncertainty of the data; specifically, the weight was inversely proportional to the square of the uncertainty. The measurements of Harned

and Davis were observed to be very precise near room temperature, exhibiting approximately 0.2 mV of scatter at 298 K, which worsened to 0.4 mV at either temperature extreme. Those of Harned and Bonner showed similar behavior across the temperature range, but appeared to have an additional 0.1 mV uncertainty. Harned and Scholes measured emf's with an uncertainty of 0.3 mV, which showed no variation with temperature. The measurements of Roy, et al., exhibit an uncertainty of 0.4 mV, with no apparent variation with temperature. We found that the isopiestic determinations of Robinson and Macaskill yielded a standard error of 0.002 in the osmotic coefficient, which corresponds to an error of less than 0.0001 in the water activity at an ionic strength of 1.0 mol kg⁻¹. White and Bates' data appear to be somewhat less precise, exhibiting an uncertainty of 0.0002 in the activity of water at the same ionic strength. The heat capacity data of Perron, et al., exhibit a precision better than 0.00005 J g⁻¹ K⁻¹ in the specific heat capacity.

4. Results and Discussion

Table 1 presents the results of least squares calculations with inclusion of the higher order electrostatic terms due to asymmetric mixing. Neglect of these terms yields results not significantly different from those of Table 1, with the exception of $S_{\Theta}^{\text{CO}_3, \text{Cl}}$. This quantity then has the value -0.092 ± 0.002 and may be relabelled $\Theta_{\text{CO}_3, \text{Cl}}$ since it is the only nonzero term in equation (8). Inclusion of terms required in the full electrostatic treatment lowers the standard error of fit slightly, as it should, but the difference is small. Therefore, on the basis of reproducing the experimental measurements here considered, it appears that these terms may be neglected without serious error. However, in treating certain properties of complex mixed electrolyte systems such as natural waters, these terms may be more important; hence the full electrostatic

treatment should be used if maximum accuracy is desired.

The values of parameters given in Table I yield thermodynamic quantities which reproduce experimental measurements to within the above-cited errors, with the following exceptions. All e.m.f. data at 273 K and 323 K showed a small but consistent deviation which we interpret to be the result of taking the solution heat capacity to be constant for this range of temperature. Even so, the deviations at the temperature extremes were rarely in excess of 1.0 mV. The newer e.m.f. measurements of Roy, et al., [14] were satisfactorily reproduced to within 0.8 mV, even at saturation of NaHCO_3 . The most dilute measurements of Robinson and Macaskill [16] appear to conflict with Harned and Scholes' [15] e.m.f. measurements, but the discrepancy is small (approximately 0.0002 in the water activity). White and Bates' [17] isopiestic measurements were reproduced with an uncertainty of 0.0003 in the activity of water, with the exception of those measurements on solutions with highest solute mole fraction of Na_2CO_3 . These measurements, in conflict with those of Robinson and Macaskill, exhibit a worst-case deviation of 0.002 in the water activity. The heat capacity data were reproduced to within $0.0001 \text{ J K}^{-1} \text{ g}^{-1}$ at all solution compositions.

The virial parameters $\beta^{(0)}$, $\beta^{(1)}$ and C^ϕ for Na_2CO_3 have been given previously by Pitzer and Mayorga [4] and by Robinson and Macaskill [16]. In the earlier report of Pitzer and Mayorga the parameters for Na_2CO_3 are based primarily upon the cell data of Taylor [40], which we suspect may be slightly in error due to irreversibilities in or contamination of the $\text{Ag-Ag}_2\text{CO}_3$ electrode. Robinson and Macaskill's values for the virial parameters agree reasonably well with the present values, particularly in light of the close coupling between the parameters. Differences

between the two sets of values can be explained by our recognition of hydrolysis and use of Harned and Scholes data. The only published values of $\beta^{(0)}$ and $\beta^{(1)}$ for NaHCO_3 and of $\theta_{\text{HCO}_3, \text{Cl}}$ are given in our previous report [13]. The present values of $\beta^{(0)}$ and $\beta^{(1)}$ agree with those results well within the indicated uncertainties; indeed, agreement is nearly exact. The earlier value of the anion mixing parameter $\theta_{\text{HCO}_3, \text{Cl}}$ was quite uncertain because of the limited composition range of the data then available. The much wider range of data from Roy, et al. [14] makes it possible to determine $\theta_{\text{HCO}_3, \text{Cl}}$ more accurately and to obtain a value for the corresponding third virial term $\psi_{\text{Na}, \text{HCO}_3, \text{Cl}}$. It is not surprising that the old estimate of $\theta_{\text{HCO}_3, \text{Cl}}$ differs considerably from the new value.

One further cautionary note concerning the effects of asymmetric mixing: in this study temperature derivatives of the higher-order electrostatic terms were omitted, even though they are in principle known. Our calculations indicate that for the presently available measurements, contributions of these derivatives to the enthalpy and heat capacity are negligible. Therefore, it seems pointless to include the rather complex expressions required to obtain these contributions purely in the interest

of formal completeness. However, one must be cautious in using the results of Table 1 in calculating enthalpies or heat capacities for solutions with ionic strength greater than 1.5 mol kg^{-1} and approximately equal molalities of univalent and divalent anions. An expanded discussion of the temperature derivatives of the electrostatic effects of asymmetric mixing is being prepared [41].

The standard state thermodynamics of the ionization equilibria of carbonic acid were represented with equations of the form suggested originally by Clarke and Glew [42] and recommended by Bezboruah, Camoes, Covington and Dobson [43]. Due to the lack of heat capacity data at temperatures other than 298 K, we were forced to use an equation which yields a temperature independent standard state heat capacity of reaction, an assumption which Pitzer and Silvester [5] found barely tenable for sodium chloride solutions at temperatures below 323 K.

Of course, the heat capacity of reaction for an aqueous electrolyte system is related in a precise manner to the heat capacities at infinite dilution of the several electrolytes involved. The first and second ionization equilibria of carbonic acid may be written in the form

The corresponding heat capacities of reaction are then

$$\Delta C_{p,1} = C_p^{\circ}(\text{NaHCO}_3) + C_p^{\circ}(\text{HCl}) - C_p^{\circ}(\text{CO}_2) - C_p^{\circ}(\text{H}_2\text{O}) - C_p^{\circ}(\text{NaCl})$$

$$\Delta C_{p,2} = C_p^{\circ}(\text{Na}_2\text{CO}_3) + C_p^{\circ}(\text{HCl}) - C_p^{\circ}(\text{NaHCO}_3) - C_p^{\circ}(\text{NaCl})$$

At 298 K the heat capacities at infinite dilution of NaCl and HCl are

$(C_p^{\circ}/R)(\text{NaCl}) = -10.15$ and $(C_p^{\circ}/R)(\text{HCl}) = -15.27$, respectively [31]. The heat

capacities of H_2O and CO_2 were taken from Stimson [44] and Rossini [45], respectively, who give $(C_p^0/R)(\text{H}_2\text{O}) = 8.8166$ and $(C_p^0/R)(\text{CO}_2) = 4.4650$. Including the values of $C_p^0(\text{Na}_2\text{CO}_3)$ and $C_p^0(\text{NaHCO}_3)$ taken from Table 1, we find $\Delta C_{p,1}^0 = -19.99 \text{ R}$ and $\Delta C_{p,2}^0 = -31.25 \text{ R}$.

Berg and Vanderzee [18] give values for the standard state enthalpies of the above ionization reactions: $\Delta H_1^0/R = -1274 \text{ K}$ and $\Delta H_2^0/R = +1768 \text{ K}$. The equilibrium constants for ionization of carbonic acid may be written as

$$\begin{aligned} \ln(K_1 K_S)_T &= \ln(K_1 K_S)_{298} - 1274 \left(\frac{T-298}{298T} \right) \\ &\quad - 19.99 \left[\frac{298}{T} - 1 + \ln\left(\frac{T}{298}\right) \right] \end{aligned} \quad (23)$$

$$\begin{aligned} \ln(K_2)_T &= \ln(K_2)_{298} + 1768 \left(\frac{T-298}{298T} \right) \\ &\quad - 31.25 \left[\frac{298}{T} - 1 + \ln\left(\frac{T}{298}\right) \right] \end{aligned} \quad (24)$$

Table 3 presents a summary of the standard state thermodynamics of carbonic acid at 5 K intervals based upon these equations. The values of $\ln(K_1 K_S)$ and $\ln(K_2)$ agree reasonably well with the original interpretation of Harned and coworkers, who found -17.995 and -23.758 at 298 K, respectively. However, the heat capacities of reaction at 298 K differ substantially from Harned's values: this analysis relies heavily upon the measurements of Perron, et. al., whereas Harned's values were derived from the temperature dependence of their emf work.

Tables 4 and 5 present our results for the heat capacities of Na_2CO_3 and NaHCO_3 solutions at 298 K and finite molalities. In Table 4 we compare the measurements of Perron, et. al., with our optimized values, and give also a breakdown of contributions to the total heat capacity, equation (17). It may be seen that for solutions of Na_2CO_3 at low molalities, the contri-

bution from the next-to-last term of (17) is very significant. Figure 1 indicates the magnitude of corrections which must be applied to the apparent molal heat capacity, ϕC_p . The uppermost curve presents the authors' fit of experimental measurements for pure Na_2CO_3 solutions at equilibrium with respect to (1). The central curve gives contributions from the first two terms of (17), $C_p^0 + J(m)$. The remaining curve presents the heat capacity for an hypothetical non-hydrolysed solution of Na_2CO_3 . Adding a small amount of NaOH to these solutions has the effect of reducing the contributions $\Delta H^0 + \Delta L$ (equation (17)) as Perron, *et. al.*, had observed, but do not diminish them to the point of negligibility. Figure 2 shows the effect that various small amounts of NaOH have upon ϕC_p .

The apparent molal heat capacity for a mixed electrolyte is not readily defined. For single-solute solutions ϕC_p is an excess quantity per mole of solute. Extending this idea, we suggest that the apparent molal heat capacity of one component in an electrolyte mixture may be defined as the total solution heat capacity in excess of that for the solution if the component were not present, divided by the molality of the component. Unfortunately, this definition of ϕC_p may be applied only to solutions with single common cation or a single common anion. Also, the apparent molal heat capacities for the various components of a mixture are not easily combined to give a total solution heat capacity, equation (10). It appears, then, that these quantities will be of little utility, and are used here only for demonstrative purposes.

Enthalpies of dilution at 298 K are presented in Table 6. These have been calculated for a solution at an initial stoichiometric molality of 1.0 mol kg^{-1} in Na_2CO_3 or NaHCO_3 . The various contributions to equation (11) are detailed; as with the heat capacities, contributions stemming

from the last term of (11) are important for dilute solutions of Na_2CO_3 .

Lastly, Tables 7 through 11 present the activity of water and various activity coefficients for Na_2CO_3 and NaHCO_3 solutions at 278, 288, 298, 308, and 318 K. In general, all activities and activity coefficients are believed accurate to the number of figures given; however, those near the extremes of the temperature range are certainly less accurate than values at 298 K.

Also presented in these tables for nominally single-solute solutions are stoichiometric activity and osmotic coefficients. These may be defined in terms of the equilibrium quantities by the formulae

$$\phi^{\text{st}} = \frac{\sum m_i}{m^{\text{st}}} \phi \quad (25)$$

$$\gamma_{\pm}^{\text{st}} = \left[\frac{m_x}{m_{\text{st}}} \right]^{1/\nu} \gamma_{\pm} \quad (26)$$

wherein the superscript st indicates the stoichiometric quantity and ν the number of ions generated upon complete dissociation of an electrolyte, $\nu = 2$ for NaHCO_3 and $\nu = 3$ for Na_2CO_3 . For the activity coefficient only the anion molality appears since the molality of Na^+ is unaffected. It may be noted that at low molalities of Na_2CO_3

$$\phi = 1 \quad \text{but} \quad \phi^{\text{st}} \approx 1.33$$

$$\gamma_{\pm} = 1 \quad \text{but} \quad \gamma_{\pm}^{\text{st}} \approx 0.$$

For solute molalities below approximately $10^{-5} \text{ mol kg}^{-1}$, the meaning of stoichiometric properties becomes ambiguous due to the self-ionization of water.

For nominally more than single-solute solutions, there is no simple definition of such stoichiometric quantities, a severe disadvantage of that formulation. However, if one wishes to calculate activities only for single-solute solutions, these stoichiometric quantities will yield accurate Gibbs energy relationships with a minimum of effort, a distinct advantage over the complex equilibrium formulation.

The stoichiometric activity and osmotic coefficients at 25°C in Table 9 may be compared with published values. For Na_2CO_3 the values of Goldberg [20] and of Robinson and Macaskill [16] may be considered. Since the osmotic coefficients at moderate and high concentration were based in all treatments on Robinson and Macaskill's results, they agree. Also the hydrolysis equilibrium (1) has little effect in this range. In view of the characteristics of a Gibbs-Duhem integration, the activity coefficients in the same concentration range may differ only by a constant factor. For Goldberg's results that factor is significantly less than unity (0.97₄); the fact that he ignored the hydrolysis reaction (1) yet attempted to fit the freezing point data of Ender [45] for dilute solutions is a sufficient explanation. Robinson and Macaskill simply fitted the Pitzer-type of virial coefficient equation to their own data without further guidance below 0.3 molal. It is probably fortuitous that their results agree almost exactly with the present values. For NaHCO_3 our earlier results agree very well with those in Table 9.

Lastly, we note that a few additional measurements would improve thermodynamic understanding of the carbonate system. Although accurate thermodynamic information beyond 323 K is severely lacking, there is a need to measure the heat capacity of aqueous Na_2CO_3 and NaHCO_3 solutions and their mixtures with NaCl and other salts at molalities in excess of

1.0 mol kg⁻¹ at 298 K and at all molalities at temperatures other than 298 K. In combination with a few additional measurements of the enthalpy of dilution at 298 K for solutions in excess of 1.0 mol kg⁻¹, extremely accurate thermodynamics for this system will be obtained.

5. Final Comment

The success of this general procedure for bicarbonate - carbonate - chloride solutions with sodium ion suggests similar studies with other cations. The extension to potassium or other alkali ions is self-evident. But bicarbonate - chloride solutions with Ca⁺⁺, Mg⁺⁺ and other cations could also be investigated with cell (2) and equation (3) would apply. Equation (6), as appropriately modified, would yield the parameters determining the activity coefficients of Ca(HCO₃)₂, Mg(HCO₃)₂, etc. This information would be valuable in interpreting the properties of various natural waters and the solubility therein of CaCO₃, MgCO₃, etc.

Acknowledgment

This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Division of Engineering, Mathematics, and Geosciences of the U. S. Department of Energy under Contract No. W-7405-ENG-48. We thank Prof. Roy and his associates for making their data available in advance of publication.

References

1. Millero, F. J., in The Sea, vol. 5, E. D. Goldberg, ed., John Wiley and Sons, New York, 1974, 3.
2. Davison, W., and Whitfield, M., J. Electroanal. Chem., 1977, 75, 763.
3. Pitzer, K. S., J. Phys. Chem., 1973, 77, 268.
4. Pitzer, K. S., and Mayorga, G., J. Phys. Chem., 1973, 77, 2300.
5. Silvester, L. F., and Pitzer, K. S., J. Phys. Chem., 1977, 81, 1822.
6. Pitzer, K. S., and Kim, J. J., J. Am. Chem. Soc., 1974, 96, 5701.
7. Pitzer, K. S., and Mayorga, G., J. Solution Chem., 1974, 3, 539.
8. Pitzer, K. S., and Silvester, L. F., J. Phys. Chem., 1978, 82, 1239.
9. Pitzer, K. S., and Silvester, L. F., J. Solution Chem., 1976, 5, 269.
10. Pitzer, K. S., Roy, R. N., and Silvester, L. F., J. Am. Chem. Soc., 1977, 99, 4930.
11. Harned, H. S., and Davis, R., Jr., J. Amer. Chem. Soc., 1943, 65, 2030.
12. Harned, H. S., and Bonner, F. T., J. Amer. Chem. Soc., 1945, 67, 1026.
13. Pitzer, K. S., and Peiper, J. C., J. Phys. Chem., 1980, 84, 2396.
14. Roy, R. N., Gibbons, J. J., Trower, J. K., Lee, G. A., Hartley, J., and Mack, G., in preparation.
15. Harned, H. S., and Scholes, S. R., Jr., J. Amer. Chem. Soc., 1941, 63, 1706.
16. Robinson, R. A., and Macaskill, J. B., J. Solution Chem., 1979, 8, 35.
17. White, D. R., Jr., and Bates, R. G., Aust. J. Sci., 1980, 33, 1903.
18. Berg, R. L., and Vanderzee, C. E., J. Chem. Thermodyn., 1978, 10, 1049; 1978, 10, 1113.
19. Perron, G., Desnoyers, J. E., and Millero, F. J., Can. J. Chem., 1975, 53, 1134.
20. Goldberg, R. N., J. Phys. Chem. Ref. Data, in press.
21. Harvie, C. E., and Weare, J. H., Geochim. Cosmochim. Acta, 1980, 44, 981.
22. Pitzer, K. S., in Activity Coefficients in Electrolyte Solutions, vol. 1, R. M. Pytkowicz, ed., CRC Press, Boca Raton, Florida, 1979, 7.

23. Pitzer, K. S., J. Solution Chem., 1975, 4, 249.
24. Friedman, H. L., Ionic Solution Theory, Interscience, New York, 1962.
25. Bradley, D. J., and Pitzer, K. S., J. Phys. Chem., 1979, 83, 1599.
26. Woolley, E. M., and Hepler, L. G., Can. J. Chem., 1977, 55, 158.
27. Cohen, E. R., and Taylor, B. N., J. Phys. Chem. Ref. Data, 1973, 2, 663.
28. Bates, R. G., and Macaskill, J. B., Pure Appl. Chem., 1978, 50, 1703.
29. Bates, R. G., Guggenheim, E. A., Harned, H. S., Ives, D. J. G., Janz, G. J., Monk, C. B., Prue, J. E., Robinson, R. A., Stokes, R. H., and Wynne-Jones, W. F. K., J. Chem. Phys., 1956, 25, 361; 1957, 26, 222.
30. Harned, H. S., and Ehlers, R. W., J. Am. Chem. Soc., 1932, 54, 1350; 1933, 55, 3179.
31. Desnoyers, J. E., de Visser, C., Perron, G., and Picker, P., J. Solution Chem., 1976, 5, 605.
32. Grenthe, I., Ots, H., Ginstrup, O., Acta Chem. Scand., 1970, 24, 1067.
33. Guggenheim, E. A., and Turgeon, J. C., Trans. Faraday Soc., 1955, 51, 757.
34. Harned, H. S., and Geary, C. G., J. Amer. Chem. Soc., 1937, 59, 2032.
35. Harned, H. S., and Owen, B. B., The Physical Chemistry of Electrolyte Solutions, 3rd ed., Reinhold Publishing Corporation, New York, NY, 1958.
36. Robinson, R. A., and Stokes, R. H., Electrolyte Solutions, 2nd ed., revised, Butterworths Publications Ltd., 1965.
37. Olofsson, G., and Olofsson, I., J. Chem. Thermodyn., 1973, 5, 533.
38. Olofsson, G., and Hepler, L. G., J. Solution Chem., 1975, 4, 127.
39. Silvester, L. F., and Pitzer, K. S., J. Solution Chem., 1978, 7, 327.
40. Taylor, C. E., J. Am. Chem. Soc., 1955, 59, 653.
41. Peiper, J. C. and Pitzer, K. S., in preparation.
42. Clarke, E. C. W., and Glew, D. N., Trans. Faraday Soc., 1966, 62, 539.
43. Bezboruah, C. P., Camoes, M. F., Covington, A. K., and Dobson, J. V., J. Chem. Soc., Faraday I, 1973, 69, 949.
44. Stimson, H. F., Am. J. Phys., 1955, 23, 622.
45. Rossini, F. D., Selected Values of Chemical Thermodynamic Properties, U. S. National Bureau of Standards, Circular 500, available at U. S. Government Printing Office, Washington, D. C., 1952.
46. Ender, F., Z. Elektrochem., 1937, 43, 234.

Table 1. Ion Interaction Parameters

	<u>Na₂CO₃</u>	<u>NaHCO₃</u>	<u>NaOH</u>	<u>NaCl</u>
$\beta^{(0)}$	0.0362±.0008	0.028±.003	0.0864 ^a	0.0765 ^a
$\beta^{(1)}$	1.51±.03	0.044±.009	0.253 ^a	0.2664 ^a
C^ϕ	0.0052±.0002		0.0044 ^a	0.00127 ^a
$\frac{\partial \beta}{\partial T}^{(0)} \times 10^3$	1.79±.04	1.00±.03	0.7 ^b	
$\frac{\partial \beta}{\partial T}^{(1)} \times 10^3$	2.05±.07	1.10±.06	0.134 ^b	
$\frac{\partial C^\phi}{\partial T} \times 10^3$			-0.1894 ^b	
$\frac{\partial^2 \beta}{\partial T^2}^{(0)} \times 10^5$	-4.22±.03	-2.6±.2	-2.00±.008	
$\frac{\partial^2 \beta}{\partial T^2}^{(1)} \times 10^5$	-16.8±.3	-4.3±.8	-2.1±.4	
$\frac{\partial^2 C^\phi}{\partial T^2} \times 10^5$			0.29±.02	
C_P^O/R	-27.48±.08	-1.35±.10	-11.63±.07	
$\Theta_{HCO_3, Cl}$	0.0359±.0010	$\Theta_{Cl, OH} - 0.050^d$		
$\psi_{Na, HCO_3, Cl}$	-0.0143±.0010	$\psi_{Na, Cl, OH} - 0.006^d$		
$S_{\Theta_{CO_3, Cl}}$	-0.053±.002 ^c			

^aPitzer and Mayorga [4]. ^bSilvester and Pitzer [39]. ^cIf higher order electrostatic terms are omitted, $\Theta_{CO_3, Cl} = -0.092$.

^dPitzer and Kim [6].

Table 2. Parameters for CO₂ solubility, equation (21)

<u>i</u>	<u>a_i</u>	<u>b_i</u>	<u>c_i</u>	<u>d_i</u>
1	5652.1±.4	-653.3±.2	-155.3±.2	-2.504±.010
2	-33.473±.003	3.9398±.0014	1.1695±.0012	-2.276±.010
3	.037177±.000006	.006491±.000002	.001981±.000002	-2.163±.010

Table 3. Standard state thermodynamics of ionization of carbonic acid.

Temperature	$\ln K_1$	$(\Delta H_1^0)/R$	$(\Delta C_{p,1}^0)/R$	$\ln K_2$	$(\Delta H_2^0)/R$	$(\Delta C_{p,2}^0)/R$
K		K			K	
273.15	-17.694	-774		-24.468	2549	
278.15	-17.748	-874		-24.305	2393	
283.15	-17.807	-974		-24.158	2237	
288.15	-17.870	-1074		-24.026	2080	
293.15	-17.936	-1174		-23.907	1924	
298.15	-18.006	-1274	-19.99	-23.802	1768	-31.25
303.15	-18.079	-1374		-23.708	1612	
308.15	-18.156	-1474		-23.626	1456	
313.15	-18.235	-1574		-23.555	1299	
318.15	-18.316	-1674		-23.493	1143	
323.15	-18.400	-1774		-23.442	987	

Table 4. A comparison of experimental and smoothed heat capacities for solutions of sodium carbonate and bicarbonate at 298 K. A solution with nominal composition given in the first three columns has an equilibrium composition given in columns 4 through 7. The experimental results of Perron, Desnoyers, and Millero [19] are given in the final column as a total heat capacity for a quantity of solution containing 1.0 kg water in excess of the heat capacity of that mass of pure water. The quantities of columns 8 and 12 are similarly defined. Columns 8 through 12 may be identified with terms of equation (17). All quantities in columns 8 through 13 are given in units of $\text{J K}^{-1} (\text{kg H}_2\text{O})^{-1}$.

Nominal Molalities			Equilibrium Molalities				$C_p^0(\bar{m}) - C_p^0(\text{H}_2\text{O})$	$J(\bar{m})$	$\Delta H^0/\Delta T$	$\Delta L/\Delta T$	$C_p(\bar{m}) - C_p^0(\text{H}_2\text{O})$	
Na_2CO_3	NaHCO_3	NaOH	CO_3^{2-}	HCO_3^-	H_2CO_3	OH^-	$C_p^0(\bar{m}) - C_p^0(\text{H}_2\text{O})$				eq (17)	PDM
0.0	0.0081	0.0	0.0001	0.0079	0.0001	0.0	-0.09	0.03	0.01	0.00	-0.06	-0.08
0.0	0.0141	0.0	0.0002	0.0137	0.0002	0.0	-0.16	0.07	0.01	0.00	-0.08	-0.06
0.0	0.0231	0.0	0.0003	0.0226	0.0003	0.0	-0.26	0.15	0.01	0.00	-0.10	-0.08
0.0	0.0350	0.0	0.0004	0.0341	0.0004	0.0	-0.39	0.28	0.01	0.00	-0.09	0.06
0.0	0.0644	0.0	0.0008	0.0628	0.0008	0.0	-0.72	0.75	0.02	0.00	0.06	0.14
0.0	0.1272	0.0	0.0016	0.1240	0.0016	0.0	-1.42	2.26	0.04	0.01	0.89	0.80
0.0	0.1880	0.0	0.0025	0.1831	0.0025	0.0	-2.10	4.27	0.06	0.01	2.24	2.30
0.0	0.2701	0.0	0.0036	0.2629	0.0036	0.0	-3.01	7.71	0.09	0.02	4.80	4.85
0.0	0.3189	0.0	0.0043	0.3103	0.0043	0.0	-3.56	10.11	0.11	0.03	6.68	6.70
0.0	0.5285	0.0	0.0072	0.5140	0.0072	0.0	-5.89	23.10	0.19	0.06	17.45	17.43
0.0	0.7740	0.0	0.0107	0.7526	0.0107	0.0	-8.63	43.20	0.29	0.11	34.97	35.06
0.0086	0.0	0.0	0.0075	0.0011	0.0	0.0011	-1.83	0.15	1.14	-0.01	-0.55	-0.50
0.0397	0.0	0.0	0.0376	0.0022	0.0	0.0022	-8.81	1.69	2.39	-0.04	-4.78	-4.52
0.0652	0.0	0.0	0.0625	0.0027	0.0	0.0027	-14.58	3.70	2.98	-0.06	-7.97	-7.66
0.1552	0.0	0.0	0.1513	0.0039	0.0	0.0039	-34.99	14.31	4.27	-0.12	-16.53	-16.46
0.2414	0.0	0.0	0.2368	0.0046	0.0	0.0046	-54.60	28.28	5.07	-0.17	-21.42	-21.39
0.2960	0.0	0.0	0.2911	0.0049	0.0	0.0049	-67.04	38.65	5.45	-0.19	-23.12	-23.05
0.4325	0.0	0.0	0.4270	0.0056	0.0	0.0056	-98.15	68.89	6.19	-0.23	-23.30	-23.07
0.5831	0.0	0.0	0.5771	0.0061	0.0	0.0061	-132.50	108.46	6.76	-0.26	-17.54	-16.57
0.9839	0.0	0.0	0.9772	0.0069	0.0	0.0069	-224.00	240.67	7.67	-0.29	24.05	23.91
0.0354	0.0	0.0100	0.0350	0.0004	0.0	0.0104	-9.01	1.67	0.91	-0.02	-6.46	-6.83
0.0709	0.0	0.0100	0.0702	0.0007	0.0	0.0107	-17.07	4.60	1.54	-0.04	-10.97	-11.12
0.0227	0.0	0.0050	0.0221	0.0005	0.0	0.0055	-5.60	0.80	1.09	-0.02	-3.73	-3.74
0.0401	0.0	0.0050	0.0392	0.0008	0.0	0.0058	-9.54	1.86	1.62	-0.03	-6.08	-5.97
0.0628	0.0	0.0050	0.0617	0.0012	0.0	0.0062	-14.70	3.67	2.17	-0.05	-8.90	-8.79
0.0846	0.0	0.0050	0.0832	0.0014	0.0	0.0064	-19.64	5.78	2.59	-0.06	-11.32	-11.10
0.0862	0.0	0.0050	0.0847	0.0014	0.0	0.0064	-20.00	5.95	2.62	-0.06	-11.49	-11.41
0.1844	0.0	0.0050	0.1820	0.0023	0.0	0.0073	-42.32	19.03	3.93	-0.12	-19.48	-19.62
0.2399	0.0	0.0050	0.2372	0.0027	0.0	0.0077	-54.97	28.42	4.45	-0.15	-22.25	-22.67
0.3195	0.0	0.0050	0.3164	0.0031	0.0	0.0081	-73.10	43.90	5.03	-0.18	-24.34	-24.62
0.5091	0.0	0.0050	0.5053	0.0039	0.0	0.0089	-116.34	88.86	5.99	-0.23	-21.72	-21.65
0.7539	0.0	0.0050	0.7495	0.0045	0.0	0.0095	-172.20	160.99	6.75	-0.27	-4.72	-3.94
0.9682	0.0	0.0050	0.9635	0.0048	0.0	0.0098	-221.14	235.64	7.18	-0.27	21.41	21.08

Table 5. Heat capacities of sodium carbonate and bicarbonate solutions at 298 K and rounded molalities. A solution with nominal composition given in the first two columns and equilibrium composition given in columns 3 through 6 has the total heat capacity given in the final column. Columns 7 through 11 may be identified with the terms of equation (17). The heat capacities have units of $\text{J K}^{-1} (\text{kg H}_2\text{O})^{-1}$.

Nominal Molalities		Equilibrium Molalities				$C_p^0(m) - C_p^0(\text{H}_2\text{O})$	$J(m)$	$\Delta H^0/\Delta T$	$\Delta L/\Delta T$	$C_p(m) - C_p^0(\text{H}_2\text{O})$
Na_2CO_3	NaHCO_3	CO_3^{2-}	HCO_3^-	H_2CO_3	OH^-					
0.0	0.0010	0.0000	0.0010	0.0000	0.0	-0.01	0.00	0.00	0.00	-0.01
0.0	0.0020	0.0000	0.0020	0.0000	0.0	-0.02	0.00	0.01	0.00	-0.01
0.0	0.0050	0.0001	0.0049	0.0001	0.0	-0.06	0.01	0.01	0.00	-0.04
0.0	0.0100	0.0001	0.0098	0.0001	0.0	-0.11	0.04	0.01	0.00	-0.07
0.0	0.0200	0.0002	0.0195	0.0002	0.0	-0.22	0.12	0.01	0.00	-0.10
0.0	0.0500	0.0006	0.0488	0.0006	0.0	-0.56	0.50	0.02	0.00	-0.04
0.0	0.1000	0.0013	0.0975	0.0013	0.0	-1.12	1.53	0.03	0.00	0.45
0.0	0.2000	0.0026	0.1947	0.0026	0.0	-2.23	4.72	0.06	0.01	2.57
0.0	0.4000	0.0054	0.3891	0.0054	0.0	-4.46	14.64	0.14	0.04	10.35
0.0	0.6000	0.0082	0.5835	0.0082	0.0	-6.69	28.44	0.22	0.07	22.04
0.0	0.8000	0.0110	0.7778	0.0110	0.0	-8.92	45.61	0.30	0.12	37.11
0.0	1.0000	0.0139	0.9720	0.0139	0.0	-11.15	65.87	0.40	0.17	55.29
0.0010	0.0010	0.0009	0.0011	0.0	0.0001	-0.22	0.01	0.24	0.00	0.03
0.0020	0.0020	0.0019	0.0022	0.0	0.0002	-0.46	0.03	0.29	0.00	-0.15
0.0050	0.0050	0.0049	0.0052	0.0	0.0002	-1.18	0.11	0.32	0.00	-0.76
0.0100	0.0100	0.0099	0.0101	0.0	0.0001	-2.38	0.32	0.31	0.00	-1.75
0.0200	0.0200	0.0199	0.0201	0.0	0.0001	-4.78	0.96	0.29	0.00	-3.53
0.0500	0.0500	0.0499	0.0501	0.0	0.0001	-11.97	4.08	0.25	-0.01	-7.64
0.1000	0.1000	0.0999	0.1001	0.0	0.0001	-23.96	12.16	0.22	-0.01	-11.59
0.2000	0.2000	0.2000	0.2000	0.0	0.0001	-47.93	35.93	0.19	-0.01	-11.82
0.4000	0.4000	0.4000	0.4000	0.0	0.0001	-95.87	105.65	0.14	0.00	9.92
0.6000	0.6000	0.6001	0.5998	0.0	0.0001	-143.81	199.14	0.12	0.00	55.45
0.8000	0.8000	0.8001	0.7998	0.0	0.0001	-191.75	313.69	0.10	0.00	122.05
1.0000	1.0000	1.0001	0.9997	0.0	0.0000	-239.69	448.14	0.09	0.01	208.55
0.0010	0.0	0.0007	0.0004	0.0	0.0004	-0.19	0.00	0.31	0.00	0.13
0.0020	0.0	0.0015	0.0005	0.0	0.0005	-0.40	0.01	0.50	0.00	0.11
0.0050	0.0	0.0042	0.0008	0.0	0.0008	-1.04	0.06	0.86	-0.01	-0.13
0.0100	0.0	0.0088	0.0012	0.0	0.0012	-2.15	0.19	1.24	-0.01	-0.73
0.0200	0.0	0.0184	0.0016	0.0	0.0016	-4.38	0.57	1.74	-0.02	-2.09
0.0500	0.0	0.0476	0.0024	0.0	0.0024	-11.13	2.43	2.65	-0.05	-6.11
0.1000	0.0	0.0968	0.0032	0.0	0.0032	-22.46	7.22	3.57	-0.09	-11.76
0.2000	0.0	0.1958	0.0043	0.0	0.0043	-45.19	21.18	4.72	-0.15	-19.43
0.4000	0.0	0.3946	0.0054	0.0	0.0054	-90.74	61.18	6.04	-0.22	-23.75
0.6000	0.0	0.5939	0.0061	0.0	0.0061	-136.36	113.27	6.82	-0.27	-16.54
0.8000	0.0	0.7935	0.0066	0.0	0.0066	-182.01	175.43	7.33	-0.29	0.47
1.0000	0.0	0.9932	0.0069	0.0	0.0069	-227.67	246.72	7.69	-0.29	26.45

Table 6. Enthalpies of dilution at 298.15 K, calculated for a solution at an initial nominal molality of 1.0 mol kg⁻¹. A quantity of this solution containing 1.0 kg H₂O is diluted by the amount of water in column 7 to give a solution with the equilibrium composition in columns 3 through 6. Columns 8 through 10 represent terms of equation (11). Enthalpies are given in units of J (kg H₂O)⁻¹.

Nominal Molalities		Equilibrium Molalities				kg H ₂ O	ΔH ^o	ΔL	ΔH _d
Na ₂ CO ₃	NaHCO ₃	CO ₃ ⁻²	HCO ₃ ⁻	H ₂ CO ₃	OH ⁻	added			
0.0	0.001	0.0000	0.0010	0.0000	0.0	999.00	-77	678	602
0.0	0.002	0.0000	0.0020	0.0000	0.0	499.00	-77	701	624
0.0	0.005	0.0001	0.0049	0.0001	0.0	199.00	-77	741	665
0.0	0.010	0.0001	0.0098	0.0001	0.0	99.00	-76	781	705
0.0	0.020	0.0002	0.0195	0.0002	0.0	49.00	-76	826	751
0.0	0.050	0.0006	0.0488	0.0006	0.0	19.00	-74	885	812
0.0	0.100	0.0013	0.0975	0.0013	0.0	9.00	-70	906	836
0.0	0.200	0.0026	0.1947	0.0026	0.0	4.00	-62	866	804
0.0	0.400	0.0054	0.3891	0.0054	0.0	1.50	-47	691	644
0.0	0.600	0.0082	0.5835	0.0082	0.0	0.67	-31	475	443
0.0	0.800	0.0110	0.7778	0.0110	0.0	0.25	-16	241	226
0.0	1.000	0.0139	0.9720	0.0139	0.0	0.00	0	0	0
0.001	0.0	0.0007	0.0004	0.0	0.0004	999.00	-268	840	572
0.002	0.0	0.0015	0.0005	0.0	0.0005	499.00	-261	953	692
0.005	0.0	0.0042	0.0008	0.0	0.0008	199.00	-248	1163	914
0.010	0.0	0.0088	0.0012	0.0	0.0012	99.00	-235	1373	1138
0.020	0.0	0.0184	0.0016	0.0	0.0016	49.00	-216	1626	1409
0.050	0.0	0.0476	0.0024	0.0	0.0024	19.00	-183	1993	1810
0.100	0.0	0.0968	0.0032	0.0	0.0032	9.00	-150	2229	2079
0.200	0.0	0.1958	0.0043	0.0	0.0043	4.00	-108	2300	2193
0.400	0.0	0.3946	0.0054	0.0	0.0054	1.50	-59	1966	1907
0.600	0.0	0.5939	0.0061	0.0	0.0061	0.67	-31	1403	1372
0.800	0.0	0.7935	0.0066	0.0	0.0066	0.25	-12	733	721
1.000	0.0	0.9933	0.0069	0.0	0.0069	0.00	0	0	0

Table 7. Osmotic and activity coefficients for sodium carbonate and bicarbonate solutions at 278 K. A solution with nominal composition given in the first two columns has an equilibrium composition given in columns 3 through 6, and an equilibrium vapor pressure of CO_2 given in column 7. See the text for definitions of stoichiometric properties. (X = CO_3 , Y = HCO_3 , and Z = OH .)

Nominal Molalities		Equilibrium Molalities				P_{CO_2} (bar)									
Na_2X	NaY	X^{2-}	Y^-	H_2X	Z^-		a_w	ϕ	$\gamma_{\text{Na}_2\text{X}}$	γ_{NaY}	γ_{NaZ}	ϕ^{st}	$\gamma_{\text{Na}_2\text{X}}^{\text{st}}$	$\gamma_{\text{NaY}}^{\text{st}}$	
0.0	0.001	0.0000	0.0010	0.0000	0.0	0.000	0.99996	0.988	0.931	0.965	0.966	0.989			0.956
0.0	0.002	0.0000	0.0020	0.0000	0.0	0.000	0.99993	0.984	0.906	0.952	0.952	0.984			0.942
0.0	0.005	0.0001	0.0049	0.0001	0.0	0.001	0.99982	0.975	0.858	0.926	0.928	0.975			0.917
0.0	0.010	0.0001	0.0098	0.0001	0.0	0.002	0.99965	0.966	0.810	0.900	0.902	0.966			0.890
0.0	0.020	0.0002	0.0196	0.0002	0.0	0.004	0.99931	0.954	0.752	0.865	0.870	0.954			0.856
0.0	0.050	0.0006	0.0489	0.0006	0.0	0.009	0.99832	0.933	0.658	0.805	0.815	0.933			0.796
0.0	0.100	0.0011	0.0977	0.0011	0.0	0.019	0.99671	0.913	0.579	0.749	0.766	0.913			0.740
0.0	0.200	0.0024	0.1951	0.0024	0.0	0.040	0.99360	0.891	0.498	0.685	0.712	0.890			0.676
0.0	0.400	0.0050	0.3901	0.0050	0.0	0.087	0.98760	0.866	0.417	0.614	0.658	0.865			0.606
0.0	0.600	0.0075	0.5850	0.0075	0.0	0.138	0.98179	0.850	0.371	0.571	0.627	0.850			0.564
0.0	0.800	0.0100	0.7799	0.0100	0.0	0.194	0.97610	0.840	0.339	0.540	0.607	0.839			0.534
0.0	1.000	0.0125	0.9748	0.0125	0.0	0.255	0.97050	0.831	0.315	0.517	0.593	0.831			0.510
0.001	0.001	0.0010	0.0010	0.0000	0.0000	0.000	0.99991	0.966	0.876	0.934	0.935				
0.002	0.002	0.0020	0.0021	0.0000	0.0001	0.000	0.99983	0.954	0.833	0.910	0.911				
0.005	0.005	0.0050	0.0051	0.0000	0.0001	0.000	0.99958	0.932	0.761	0.867	0.870				
0.010	0.010	0.0100	0.0100	0.0000	0.0000	0.000	0.99918	0.912	0.696	0.825	0.831				
0.020	0.020	0.0200	0.0200	0.0000	0.0000	0.000	0.99840	0.890	0.623	0.775	0.785				
0.050	0.050	0.0500	0.0500	0.0000	0.0000	0.000	0.99615	0.855	0.520	0.696	0.717				
0.100	0.100	0.1000	0.1000	0.0000	0.0000	0.000	0.99259	0.826	0.441	0.629	0.662				
0.200	0.200	0.2000	0.2000	0.0000	0.0000	0.001	0.98588	0.789	0.361	0.555	0.606				
0.400	0.400	0.4000	0.3999	0.0001	0.0000	0.001	0.97365	0.741	0.282	0.477	0.552				
0.600	0.600	0.6001	0.5999	0.0001	0.0000	0.002	0.96244	0.708	0.237	0.430	0.527				
0.800	0.800	0.8001	0.7998	0.0001	0.0000	0.003	0.95184	0.685	0.207	0.399	0.516				
1.000	1.000	1.0001	0.9998	0.0001	0.0000	0.005	0.94160	0.668	0.186	0.375	0.514				
0.001	0.0	0.0008	0.0002	0.0	0.0002	0.0	0.99994	0.968	0.894	0.944	0.944	1.036	0.826		
0.002	0.0	0.0017	0.0003	0.0	0.0003	0.0	0.99989	0.955	0.855	0.922	0.923	1.003	0.810		
0.005	0.0	0.0045	0.0005	0.0	0.0005	0.0	0.99974	0.931	0.790	0.883	0.886	0.961	0.764		
0.010	0.0	0.0093	0.0007	0.0	0.0007	0.0	0.99950	0.909	0.729	0.845	0.849	0.929	0.712		
0.020	0.0	0.0191	0.0009	0.0	0.0009	0.0	0.99903	0.883	0.659	0.799	0.806	0.897	0.649		
0.050	0.0	0.0487	0.0014	0.0	0.0014	0.0	0.99770	0.846	0.559	0.725	0.740	0.853	0.554		
0.100	0.0	0.0982	0.0018	0.0	0.0018	0.0	0.99558	0.814	0.481	0.662	0.686	0.819	0.479		
0.200	0.0	0.1976	0.0024	0.0	0.0024	0.0	0.99161	0.777	0.402	0.591	0.629	0.780	0.401		
0.400	0.0	0.3970	0.0031	0.0	0.0031	0.0	0.98440	0.725	0.321	0.514	0.571	0.727	0.321		
0.600	0.0	0.5966	0.0035	0.0	0.0035	0.0	0.97792	0.687	0.275	0.466	0.538	0.688	0.274		
0.800	0.0	0.7964	0.0037	0.0	0.0037	0.0	0.97194	0.657	0.242	0.432	0.517	0.658	0.242		
1.000	0.0	0.9962	0.0039	0.0	0.0039	0.0	0.96630	0.634	0.219	0.406	0.503	0.634	0.218		

Table 8. Osmotic and activity coefficients for sodium carbonate and bicarbonate solutions at 288 K. A solution with nominal composition given in the first two columns has an equilibrium composition given in columns 3 through 6, and an equilibrium vapor pressure of CO_2 given in column 7. See the text for definitions of stoichiometric properties. (X = CO_3 , Y = HCO_3 , and Z = OH.)

Nominal Molalities		Equilibrium Molalities				P_{CO_2}								
Na_2X	NaY	X^{2-}	Y^-	H_2X	Z^-	(bar)	a_w	ϕ	$\gamma_{\text{Na}_2\text{X}}$	γ_{NaY}	γ_{NaZ}	ϕ^{st}	$\gamma_{\text{Na}_2\text{X}}^{\text{st}}$	$\gamma_{\text{NaY}}^{\text{st}}$
0.0	0.001	0.0000	0.0010	0.0000	0.0	0.000	0.99996	0.988	0.931	0.965	0.965	0.989		0.955
0.0	0.002	0.0000	0.0020	0.0000	0.0	0.001	0.99993	0.984	0.904	0.951	0.952	0.984		0.942
0.0	0.005	0.0001	0.0049	0.0001	0.0	0.001	0.99982	0.975	0.856	0.926	0.927	0.975		0.916
0.0	0.010	0.0001	0.0098	0.0001	0.0	0.002	0.99965	0.966	0.808	0.899	0.901	0.966		0.889
0.0	0.020	0.0002	0.0196	0.0002	0.0	0.005	0.99931	0.954	0.749	0.864	0.868	0.954		0.854
0.0	0.050	0.0006	0.0488	0.0006	0.0	0.013	0.99832	0.933	0.656	0.805	0.814	0.933		0.795
0.0	0.100	0.0012	0.0976	0.0012	0.0	0.027	0.99671	0.914	0.578	0.749	0.765	0.914		0.740
0.0	0.200	0.0025	0.1950	0.0025	0.0	0.058	0.99359	0.893	0.498	0.687	0.713	0.893		0.678
0.0	0.400	0.0052	0.3896	0.0052	0.0	0.125	0.98753	0.871	0.420	0.619	0.661	0.871		0.611
0.0	0.600	0.0078	0.5843	0.0078	0.0	0.200	0.98162	0.858	0.376	0.579	0.632	0.858		0.571
0.0	0.800	0.0105	0.7789	0.0105	0.0	0.281	0.97579	0.850	0.346	0.551	0.615	0.850		0.544
0.0	1.000	0.0132	0.9735	0.0132	0.0	0.369	0.97003	0.845	0.323	0.530	0.603	0.844		0.523
0.001	0.001	0.0009	0.0011	0.0000	0.0001	0.000	0.99991	0.967	0.874	0.934	0.934			
0.002	0.002	0.0019	0.0021	0.0000	0.0001	0.000	0.99983	0.954	0.831	0.909	0.911			
0.005	0.005	0.0049	0.0051	0.0000	0.0001	0.000	0.99958	0.932	0.759	0.866	0.869			
0.010	0.010	0.0099	0.0101	0.0000	0.0001	0.000	0.99918	0.912	0.693	0.824	0.830			
0.020	0.020	0.0199	0.0201	0.0000	0.0001	0.000	0.99840	0.889	0.621	0.774	0.784			
0.050	0.050	0.0499	0.0501	0.0000	0.0001	0.000	0.99615	0.857	0.519	0.697	0.717			
0.100	0.100	0.1000	0.1000	0.0000	0.0001	0.000	0.99256	0.829	0.442	0.632	0.663			
0.200	0.200	0.2000	0.2000	0.0000	0.0000	0.001	0.98575	0.797	0.365	0.562	0.610			
0.400	0.400	0.4000	0.3999	0.0001	0.0000	0.002	0.97309	0.757	0.289	0.490	0.563			
0.600	0.600	0.6001	0.5998	0.0001	0.0000	0.004	0.96118	0.733	0.247	0.449	0.543			
0.800	0.800	0.8001	0.7998	0.0001	0.0000	0.005	0.94960	0.718	0.220	0.423	0.536			
1.000	1.000	1.0001	0.9997	0.0001	0.0000	0.007	0.93810	0.709	0.201	0.404	0.539			
0.001	0.0	0.0007	0.0003	0.0	0.0003	0.0	0.99994	0.970	0.893	0.944	0.944	1.058	0.803	
0.002	0.0	0.0016	0.0004	0.0	0.0004	0.0	0.99989	0.956	0.854	0.922	0.923	1.020	0.794	
0.005	0.0	0.0044	0.0006	0.0	0.0006	0.0	0.99974	0.932	0.788	0.883	0.885	0.972	0.753	
0.010	0.0	0.0091	0.0009	0.0	0.0009	0.0	0.99949	0.910	0.727	0.844	0.848	0.937	0.704	
0.020	0.0	0.0188	0.0012	0.0	0.0012	0.0	0.99903	0.884	0.657	0.798	0.805	0.902	0.643	
0.050	0.0	0.0482	0.0018	0.0	0.0018	0.0	0.99769	0.847	0.558	0.725	0.740	0.857	0.551	
0.100	0.0	0.0976	0.0024	0.0	0.0024	0.0	0.99556	0.817	0.481	0.663	0.687	0.823	0.477	
0.200	0.0	0.1968	0.0032	0.0	0.0032	0.0	0.99154	0.782	0.403	0.596	0.632	0.786	0.401	
0.400	0.0	0.3959	0.0041	0.0	0.0041	0.0	0.98416	0.736	0.326	0.523	0.577	0.739	0.325	
0.600	0.0	0.5954	0.0047	0.0	0.0047	0.0	0.97739	0.703	0.281	0.479	0.547	0.705	0.281	
0.800	0.0	0.7951	0.0050	0.0	0.0050	0.0	0.97100	0.679	0.251	0.448	0.530	0.681	0.251	
1.000	0.0	0.9949	0.0052	0.0	0.0052	0.0	0.96481	0.662	0.229	0.426	0.519	0.663	0.229	

Table 9. Osmotic and activity coefficients for sodium carbonate and bicarbonate solutions at 298 K. A solution with nominal composition given in the first two columns has an equilibrium composition given in columns 3 through 6, and an equilibrium vapor pressure of CO_2 given in column 7. See the text for definitions of stoichiometric properties. ($X = \text{CO}_3$, $Y = \text{HCO}_3$, and $Z = \text{OH}$.)

Nominal Molalities		Equilibrium Molalities				P_{CO_2}								
Na_2X	NaY	X^{2-}	Y^-	H_2X	Z^-	(bar)	a_w	ϕ	$\gamma_{\text{Na}_2\text{X}}$	γ_{NaY}	γ_{NaZ}	ϕ^{st}	$\gamma_{\text{Na}_2\text{X}}^{\text{st}}$	$\gamma_{\text{NaY}}^{\text{st}}$
0.0	0.001	0.0000	0.0010	0.0000	0.0	0.000	0.99996	0.988	0.930	0.964	0.965	0.989		0.954
0.0	0.002	0.0000	0.0020	0.0000	0.0	0.001	0.99993	0.983	0.903	0.951	0.951	0.984		0.941
0.0	0.005	0.0001	0.0049	0.0001	0.0	0.002	0.99982	0.974	0.855	0.925	0.926	0.975		0.915
0.0	0.010	0.0001	0.0098	0.0001	0.0	0.003	0.99965	0.965	0.806	0.898	0.900	0.965		0.888
0.0	0.020	0.0002	0.0195	0.0002	0.0	0.007	0.99931	0.953	0.747	0.863	0.867	0.953		0.853
0.0	0.050	0.0006	0.0488	0.0006	0.0	0.018	0.99832	0.933	0.653	0.803	0.812	0.933		0.793
0.0	0.100	0.0013	0.0975	0.0013	0.0	0.038	0.99671	0.914	0.576	0.748	0.763	0.914		0.739
0.0	0.200	0.0026	0.1947	0.0026	0.0	0.081	0.99358	0.894	0.496	0.687	0.712	0.894		0.678
0.0	0.400	0.0054	0.3891	0.0054	0.0	0.174	0.98748	0.874	0.420	0.622	0.661	0.874		0.613
0.0	0.600	0.0082	0.5835	0.0082	0.0	0.278	0.98150	0.864	0.378	0.584	0.635	0.864		0.576
0.0	0.800	0.0110	0.7778	0.0110	0.0	0.391	0.97557	0.858	0.349	0.558	0.619	0.858		0.550
0.0	1.000	0.0139	0.9720	0.0139	0.0	0.514	0.96969	0.854	0.327	0.539	0.609	0.854		0.531
0.001	0.001	0.0009	0.0011	0.0000	0.0001	0.000	0.99991	0.967	0.873	0.933	0.934			
0.002	0.002	0.0019	0.0022	0.0000	0.0002	0.000	0.99983	0.954	0.829	0.908	0.910			
0.005	0.005	0.0049	0.0052	0.0000	0.0002	0.000	0.99958	0.932	0.756	0.865	0.868			
0.010	0.010	0.0099	0.0101	0.0000	0.0001	0.000	0.99918	0.911	0.690	0.822	0.828			
0.020	0.020	0.0199	0.0201	0.0000	0.0001	0.000	0.99840	0.889	0.617	0.773	0.782			
0.050	0.050	0.0499	0.0501	0.0000	0.0001	0.000	0.99615	0.856	0.516	0.696	0.715			
0.100	0.100	0.0999	0.1001	0.0000	0.0001	0.001	0.99254	0.831	0.440	0.633	0.663			
0.200	0.200	0.2000	0.2000	0.0000	0.0001	0.001	0.98566	0.801	0.365	0.566	0.612			
0.400	0.400	0.4000	0.3999	0.0001	0.0001	0.003	0.97269	0.768	0.293	0.500	0.569			
0.600	0.600	0.6001	0.5998	0.0001	0.0000	0.005	0.96024	0.751	0.254	0.464	0.553			
0.800	0.800	0.8001	0.7998	0.0001	0.0000	0.008	0.94789	0.743	0.229	0.441	0.550			
1.000	1.000	1.0001	0.9997	0.0002	0.0000	0.011	0.93539	0.742	0.213	0.427	0.556			

Table 9, continued. Osmotic and activity coefficients for sodium carbonate and bicarbonate solutions at 298 K.

Nominal Molalities		Equilibrium Molalities				P_{CO_2} (bar)	a_w	ϕ	$\gamma_{\text{Na}_2\text{X}}$	γ_{NaY}	γ_{NaZ}	ϕ^{st}	$\gamma_{\text{Na}_2\text{X}}^{\text{st}}$	$\gamma_{\text{NaY}}^{\text{st}}$
Na_2X	NaY	X^{2-}	Y^-	H_2X	Z^-									
0.001	0.0	0.0007	0.0004	0.0	0.0004	0.0	0.99994	0.971	0.893	0.944	0.944	1.083	0.775	
0.002	0.0	0.0015	0.0005	0.0	0.0005	0.0	0.99989	0.957	0.853	0.922	0.923	1.039	0.774	
0.005	0.0	0.0042	0.0008	0.0	0.0008	0.0	0.99973	0.933	0.786	0.882	0.884	0.985	0.740	
0.010	0.0	0.0088	0.0012	0.0	0.0012	0.0	0.99949	0.910	0.724	0.843	0.847	0.945	0.695	
0.020	0.0	0.0184	0.0016	0.0	0.0016	0.0	0.99902	0.884	0.654	0.797	0.803	0.908	0.636	
0.050	0.0	0.0476	0.0024	0.0	0.0024	0.0	0.99768	0.847	0.554	0.724	0.738	0.860	0.545	
0.100	0.0	0.0968	0.0032	0.0	0.0032	0.0	0.99554	0.817	0.478	0.663	0.685	0.826	0.473	
0.200	0.0	0.1958	0.0043	0.0	0.0043	0.0	0.99150	0.784	0.402	0.598	0.632	0.790	0.399	
0.400	0.0	0.3946	0.0054	0.0	0.0054	0.0	0.98399	0.743	0.327	0.529	0.580	0.746	0.326	
0.600	0.0	0.5939	0.0061	0.0	0.0061	0.0	0.97700	0.715	0.285	0.488	0.553	0.717	0.284	
0.800	0.0	0.7935	0.0066	0.0	0.0066	0.0	0.97029	0.696	0.256	0.460	0.538	0.698	0.256	
1.000	0.0	0.9933	0.0069	0.0	0.0069	0.0	0.96369	0.683	0.236	0.440	0.530	0.684	0.235	
1.200	0.0	1.1931	0.0071	0.0	0.0071	0.0	0.95708	0.675	0.221	0.425	0.527	0.676	0.220	
1.400	0.0	1.3930	0.0072	0.0	0.0072	0.0	0.95036	0.672	0.209	0.414	0.528	0.673	0.209	
1.600	0.0	1.5929	0.0073	0.0	0.0073	0.0	0.94348	0.672	0.200	0.405	0.533	0.673	0.200	
1.800	0.0	1.7929	0.0073	0.0	0.0073	0.0	0.93636	0.675	0.193	0.399	0.540	0.676	0.193	
2.000	0.0	1.9930	0.0073	0.0	0.0073	0.0	0.92898	0.681	0.188	0.394	0.550	0.682	0.188	
2.200	0.0	2.1930	0.0073	0.0	0.0073	0.0	0.92129	0.689	0.184	0.390	0.563	0.689	0.184	
2.400	0.0	2.3931	0.0072	0.0	0.0072	0.0	0.91327	0.699	0.181	0.388	0.578	0.699	0.180	
2.600	0.0	2.5932	0.0072	0.0	0.0072	0.0	0.90489	0.710	0.179	0.387	0.595	0.711	0.178	
2.767	0.0	2.7603	0.0071	0.0	0.0071	0.0	0.89760	0.722	0.177	0.387	0.611	0.722	0.177	
2.800	0.0	2.7933	0.0071	0.0	0.0071	0.0	0.89613	0.724	0.177	0.387	0.615	0.725	0.177	
3.000	0.0	2.9934	0.0070	0.0	0.0070	0.0	0.88699	0.739	0.177	0.387	0.637	0.740	0.176	

Table 10. Osmotic and activity coefficients for sodium carbonate and bicarbonate solutions at 308 K. A solution with nominal composition given in the first two columns has an equilibrium composition given in columns 3 through 6, and an equilibrium vapor pressure of CO_2 given in column 7. See the text for definitions of stoichiometric properties. ($X = \text{CO}_3$, $Y = \text{HCO}_3$, and $Z = \text{OH}$.)

Nominal Molalities		Equilibrium Molalities				P_{CO_2} (bar)								
Na_2X	NaY	X^{2-}	Y^-	H_2X	Z^-		a_w	ϕ	$\gamma_{\text{Na}_2\text{X}}$	γ_{NaY}	γ_{NaZ}	ϕ^{st}	$\gamma_{\text{Na}_2\text{X}}^{\text{st}}$	$\gamma_{\text{NaY}}^{\text{st}}$
0.0	0.001	0.0000	0.0010	0.0000	0.0	0.001	0.99996	0.988	0.928	0.964	0.964	0.989		0.953
0.0	0.002	0.0000	0.0020	0.0000	0.0	0.001	0.99993	0.983	0.901	0.950	0.950	0.984		0.939
0.0	0.005	0.0001	0.0049	0.0001	0.0	0.002	0.99982	0.974	0.852	0.923	0.925	0.974		0.913
0.0	0.010	0.0001	0.0098	0.0001	0.0	0.004	0.99965	0.965	0.803	0.896	0.898	0.965		0.886
0.0	0.020	0.0002	0.0195	0.0002	0.0	0.009	0.99931	0.953	0.743	0.861	0.865	0.953		0.850
0.0	0.050	0.0006	0.0487	0.0006	0.0	0.024	0.99832	0.932	0.649	0.801	0.809	0.932		0.791
0.0	0.100	0.0013	0.0974	0.0013	0.0	0.051	0.99671	0.914	0.571	0.747	0.761	0.914		0.737
0.0	0.200	0.0028	0.1945	0.0028	0.0	0.108	0.99357	0.895	0.493	0.686	0.709	0.895		0.676
0.0	0.400	0.0057	0.3886	0.0057	0.0	0.233	0.98745	0.876	0.417	0.622	0.660	0.876		0.613
0.0	0.600	0.0087	0.5826	0.0087	0.0	0.373	0.98142	0.867	0.376	0.586	0.634	0.867		0.577
0.0	0.800	0.0117	0.7765	0.0117	0.0	0.525	0.97544	0.863	0.348	0.561	0.619	0.863		0.553
0.0	1.000	0.0147	0.9704	0.0147	0.0	0.690	0.96946	0.861	0.328	0.543	0.610	0.861		0.535
0.001	0.001	0.0008	0.0012	0.0000	0.0002	0.000	0.99991	0.968	0.872	0.933	0.933			
0.002	0.002	0.0018	0.0022	0.0000	0.0002	0.000	0.99982	0.954	0.827	0.907	0.909			
0.005	0.005	0.0048	0.0052	0.0000	0.0002	0.000	0.99958	0.931	0.753	0.863	0.866			
0.010	0.010	0.0098	0.0102	0.0000	0.0002	0.000	0.99918	0.910	0.686	0.820	0.826			
0.020	0.020	0.0198	0.0202	0.0000	0.0002	0.000	0.99840	0.887	0.612	0.770	0.779			
0.050	0.050	0.0498	0.0502	0.0000	0.0002	0.000	0.99615	0.855	0.511	0.694	0.712			
0.100	0.100	0.0999	0.1001	0.0000	0.0002	0.001	0.99255	0.830	0.435	0.632	0.660			
0.200	0.200	0.1999	0.2001	0.0000	0.0001	0.002	0.98563	0.803	0.362	0.567	0.611			
0.400	0.400	0.4000	0.3999	0.0001	0.0001	0.005	0.97245	0.775	0.293	0.504	0.570			
0.600	0.600	0.6001	0.5998	0.0001	0.0001	0.008	0.95961	0.763	0.256	0.472	0.557			
0.800	0.800	0.8001	0.7997	0.0002	0.0001	0.012	0.94669	0.760	0.234	0.453	0.558			
1.000	1.000	1.0001	0.9997	0.0002	0.0001	0.016	0.93345	0.764	0.219	0.443	0.567			
0.001	0.0	0.0006	0.0004	0.0	0.0004	0.0	0.99994	0.972	0.892	0.944	0.944	1.110	0.742	
0.002	0.0	0.0014	0.0006	0.0	0.0006	0.0	0.99989	0.959	0.852	0.921	0.922	1.061	0.750	
0.005	0.0	0.0040	0.0011	0.0	0.0011	0.0	0.99973	0.934	0.784	0.881	0.883	1.000	0.725	
0.010	0.0	0.0085	0.0015	0.0	0.0015	0.0	0.99948	0.911	0.721	0.842	0.846	0.956	0.683	
0.020	0.0	0.0179	0.0021	0.0	0.0021	0.0	0.99901	0.884	0.650	0.795	0.801	0.914	0.627	
0.050	0.0	0.0469	0.0031	0.0	0.0031	0.0	0.99767	0.846	0.549	0.722	0.735	0.864	0.538	
0.100	0.0	0.0958	0.0042	0.0	0.0042	0.0	0.99553	0.817	0.473	0.662	0.683	0.828	0.466	
0.200	0.0	0.1945	0.0055	0.0	0.0055	0.0	0.99148	0.785	0.398	0.597	0.630	0.792	0.394	
0.400	0.0	0.3931	0.0070	0.0	0.0070	0.0	0.98390	0.746	0.325	0.531	0.580	0.751	0.323	
0.600	0.0	0.5922	0.0079	0.0	0.0079	0.0	0.97676	0.722	0.284	0.493	0.555	0.725	0.283	
0.800	0.0	0.7917	0.0085	0.0	0.0085	0.0	0.96983	0.706	0.258	0.468	0.542	0.709	0.257	
1.000	0.0	0.9913	0.0089	0.0	0.0089	0.0	0.96292	0.697	0.239	0.450	0.535	0.699	0.238	

Table 11. Osmotic and activity coefficients for sodium carbonate and bicarbonate solutions at 318 K. A solution with nominal composition given in the first two columns has an equilibrium composition given in columns 3 through 6, and an equilibrium vapor pressure of CO_2 given in column 7. See the text for definitions of stoichiometric properties. (X = CO_3 , Y = HCO_3 , and Z = OH .)

Nominal		Equilibrium Molalities				P_{CO_2}								
Molalities														
Na_2X	NaY	X^{2-}	Y^-	H_2X	Z^-	(bar)	a_w	ϕ	$\gamma_{\text{Na}_2\text{X}}$	γ_{NaY}	γ_{NaZ}	ϕ^{st}	$\gamma_{\text{Na}_2\text{X}}^{\text{st}}$	$\gamma_{\text{NaY}}^{\text{st}}$
0.0	0.001	0.0000	0.0010	0.0000	0.0	0.001	0.99996	0.988	0.927	0.963	0.963	0.990		0.952
0.0	0.002	0.0000	0.0020	0.0000	0.0	0.001	0.99993	0.983	0.900	0.949	0.949	0.984		0.938
0.0	0.005	0.0001	0.0049	0.0001	0.0	0.003	0.99982	0.974	0.850	0.922	0.923	0.974		0.911
0.0	0.010	0.0001	0.0098	0.0001	0.0	0.006	0.99965	0.964	0.800	0.894	0.896	0.964		0.883
0.0	0.020	0.0003	0.0195	0.0003	0.0	0.012	0.99931	0.952	0.739	0.859	0.862	0.952		0.848
0.0	0.050	0.0007	0.0487	0.0007	0.0	0.031	0.99832	0.931	0.644	0.798	0.806	0.931		0.788
0.0	0.100	0.0014	0.0972	0.0014	0.0	0.065	0.99672	0.913	0.566	0.743	0.757	0.913		0.733
0.0	0.200	0.0029	0.1942	0.0029	0.0	0.139	0.99358	0.894	0.487	0.683	0.705	0.894		0.673
0.0	0.400	0.0060	0.3880	0.0060	0.0	0.303	0.98745	0.876	0.412	0.620	0.656	0.876		0.610
0.0	0.600	0.0092	0.5816	0.0092	0.0	0.484	0.98140	0.868	0.371	0.584	0.631	0.868		0.575
0.0	0.800	0.0123	0.7751	0.0123	0.0	0.682	0.97538	0.865	0.344	0.561	0.616	0.865		0.552
0.0	1.000	0.0156	0.9686	0.0156	0.0	0.898	0.96936	0.864	0.324	0.544	0.608	0.864		0.535
0.001	0.001	0.0007	0.0013	0.0000	0.0003	0.000	0.99991	0.969	0.871	0.932	0.933			
0.002	0.002	0.0017	0.0023	0.0000	0.0003	0.000	0.99982	0.955	0.825	0.906	0.908			
0.005	0.005	0.0046	0.0054	0.0000	0.0004	0.000	0.99957	0.931	0.749	0.861	0.864			
0.010	0.010	0.0096	0.0104	0.0000	0.0004	0.000	0.99918	0.909	0.681	0.818	0.823			
0.020	0.020	0.0197	0.0204	0.0000	0.0004	0.000	0.99840	0.886	0.606	0.767	0.776			
0.050	0.050	0.0497	0.0503	0.0000	0.0003	0.001	0.99616	0.853	0.504	0.691	0.708			
0.100	0.100	0.0998	0.1002	0.0000	0.0003	0.001	0.99257	0.828	0.429	0.628	0.655			
0.200	0.200	0.1998	0.2001	0.0001	0.0002	0.003	0.98565	0.802	0.356	0.565	0.606			
0.400	0.400	0.3999	0.4000	0.0001	0.0002	0.006	0.97237	0.777	0.289	0.505	0.568			
0.600	0.600	0.6000	0.5998	0.0001	0.0001	0.011	0.95930	0.769	0.254	0.475	0.556			
0.800	0.800	0.8001	0.7997	0.0002	0.0001	0.017	0.94602	0.770	0.234	0.459	0.558			
1.000	1.000	1.0001	0.9996	0.0002	0.0001	0.023	0.93229	0.778	0.220	0.451	0.569			
0.001	0.0	0.0005	0.0005	0.0	0.0005	0.0	0.99994	0.974	0.892	0.944	0.944	1.138	0.704	
0.002	0.0	0.0012	0.0008	0.0	0.0008	0.0	0.99988	0.960	0.851	0.921	0.922	1.085	0.722	
0.005	0.0	0.0037	0.0013	0.0	0.0013	0.0	0.99973	0.936	0.782	0.880	0.882	1.017	0.706	
0.010	0.0	0.0081	0.0019	0.0	0.0019	0.0	0.99948	0.912	0.717	0.840	0.844	0.968	0.669	
0.020	0.0	0.0174	0.0026	0.0	0.0026	0.0	0.99900	0.884	0.645	0.792	0.799	0.922	0.615	
0.050	0.0	0.0461	0.0039	0.0	0.0039	0.0	0.99766	0.845	0.543	0.719	0.732	0.867	0.528	
0.100	0.0	0.0947	0.0053	0.0	0.0053	0.0	0.99553	0.815	0.466	0.658	0.678	0.829	0.458	
0.200	0.0	0.1931	0.0070	0.0	0.0070	0.0	0.99148	0.783	0.391	0.594	0.625	0.792	0.387	
0.400	0.0	0.3912	0.0089	0.0	0.0089	0.0	0.98388	0.746	0.320	0.529	0.576	0.752	0.317	
0.600	0.0	0.5901	0.0100	0.0	0.0100	0.0	0.97667	0.724	0.280	0.493	0.552	0.728	0.279	
0.800	0.0	0.7894	0.0107	0.0	0.0107	0.0	0.96961	0.711	0.255	0.469	0.540	0.714	0.254	
1.000	0.0	0.9890	0.0112	0.0	0.0112	0.0	0.96251	0.704	0.237	0.453	0.535	0.707	0.236	

Figure Captions

Figure 1. The effect of hydrolysis upon Na_2CO_3 heat capacity. The data are those of Perron, Desnoyers, and Millero [19]. The uppermost curve represents the authors' optimized values of Na_2CO_3 heat capacity; see the text for explanations of the remaining curves.

Figure 2. The effect upon Na_2CO_3 heat capacity of the addition of a small amount of NaOH. The data are those of Perron, Desnoyers, and Millero [19]: $\square$ Na_2CO_3 , $\odot$ $\text{Na}_2\text{CO}_3 + 0.005 \text{ mol kg}^{-1} \text{ NaOH}$, $\triangle$ $\text{Na}_2\text{CO}_3 + 0.010 \text{ mol kg}^{-1} \text{ NaOH}$. The lowest curve represents a hypothetical, nonhydrolysed solution of Na_2CO_3 .

Figure 1. The effect of hydrolysis upon Na_2CO_3 heat capacity. The data are those of Perron, Desnoyers, and Millero [19]. The upper-most curve represents the authors' optimized values of Na_2CO_3 heat capacity; see the text for explanations of the remaining curves.

Figure 2. The effect upon Na_2CO_3 heat capacity of the addition of a small amount of NaOH . The data are those of Perron, Desnoyers, and Millero [19]: $\square$ Na_2CO_3 , $\odot$ $\text{Na}_2\text{CO}_3 + 0.005 \text{ mol kg}^{-1} \text{NaOH}$, Δ $\text{Na}_2\text{CO}_3 + 0.010 \text{ mol kg}^{-1} \text{NaOH}$. The lowest curve represents a hypothetical, nonhydrolyzed solution of Na_2CO_3 .

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720