

# UCLA

## UCLA Previously Published Works

### Title

Isolated Hepatitis B Core Antibody is Associated With Advanced Hepatic Fibrosis in HIV/HCV Infection But Not in HIV Infection Alone

### Permalink

<https://escholarship.org/uc/item/0qs619sc>

### Journal

J AIDS Journal of Acquired Immune Deficiency Syndromes, 72(1)

### ISSN

1525-4135

### Authors

Bhattacharya, Debika  
Tseng, Chi-hong  
Tate, Janet P  
[et al.](#)

### Publication Date

2016-05-01

### DOI

10.1097/qai.0000000000000941

Peer reviewed

**Title:**

**Isolated Hepatitis B Core Antibody is Associated with Advanced Hepatic Fibrosis in HIV/HCV Infection but not in HIV infection alone**

**Authors and Affiliations:**

Debika Bhattacharya MD, MS<sup>1,2</sup>, Chi-hong Tseng PhD<sup>2</sup>, Janet P. Tate ScD<sup>3,4</sup>, Vincent Lo Re III MD, MSCE<sup>5,6</sup>, Cynthia L. Gibert MD<sup>7,8</sup>, Adeel A. Butt MD<sup>9,10</sup>, Sheldon T. Brown MD<sup>11,12</sup>, Joseph K. Lim MD<sup>3,4</sup>, Maria C. Rodriguez-Barradas MD<sup>13,14</sup>, David Rimland MD<sup>15,16</sup>, Erica Kaufman MD<sup>17</sup>, Amy C. Justice MD, PhD<sup>3,4</sup>, Matthew Bidwell Goetz MD<sup>1,2</sup>

1. VA Greater Los Angeles Healthcare System, Los Angeles, CA, USA, 2. David Geffen School of Medicine at UCLA, Los Angeles, CA, USA, 3. VA Connecticut Healthcare System, West Haven, CT, USA, 4. Yale University School of Medicine, New Haven, CT, USA, 5. Philadelphia VA Medical Center, Philadelphia, PA, USA, 6. Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA, 7. Washington DC VA Medical Center, Washington, DC, USA, 8. George Washington University Medical Center, Washington, DC, USA, 9. Hamad Healthcare Quality Institute and Hamad Medical Corporation, Doha, Qatar, 10. VA Pittsburgh Healthcare System, Pittsburgh, PA, USA, 11. James J. Peters VA Medical Center, Bronx, NY, USA, 12. Icahn School of Medicine at Mt. Sinai, New York, NY, USA, 13. Michael E. DeBakey VA Medical Center, Houston, TX, USA, 14. Baylor College of Medicine, Houston, TX, USA, 15. Atlanta VA Medical Center, Atlanta, GA, USA, 16. Emory University School of Medicine, Atlanta, GA, USA, 17. Saint Louis University School of Medicine, Saint Louis, MO, USA

**Keywords:** Hepatitis B Core Antibody, HIV, hepatitis C

**Running Head:** Isolated HBcAb and Hepatic Fibrosis

**40 Word Summary:** HIV+/HCV+ persons with isolated HBcAb have a higher prevalence of advanced fibrosis than persons who are non-immune to HBV, who have resolved HBV, or who are HbsAb+ only.

**Corresponding Author:**

Debika Bhattacharya MD MS  
Associate Clinical Professor of Medicine  
Division of Infectious Diseases and UCLA CARE Center  
David Geffen School of Medicine at the University of California, Los Angeles  
VA Greater Los Angeles Healthcare System  
11301 Wilshire Blvd  
111F  
Los Angeles, CA 90073

Phone: (310) 825-0137  
Fax: (310) 825-3632  
Email: [debika.bhattacharya@va.gov](mailto:debika.bhattacharya@va.gov)

**Alternate Author:**

Matthew Bidwell Goetz MD  
Chief, Division of Infectious Diseases  
VA Greater Los Angeles Healthcare System  
11301 Wilshire Blvd  
111F  
Los Angeles, CA 90073  
Phone: (310) 268-3763  
Fax: (310) 268-4928  
Email: [Matthew.Goetz@va.gov](mailto:Matthew.Goetz@va.gov)

**Potential Conflicts of Interest and Source of Funding:** The corresponding author, Debika Bhattacharya, has received payment by IAS-USA for development of educational presentations and research support from AbbVie, Bristol-Myers Squibb, and Merck for research outside the scope of this work. Janet Tate has received research grants from National Institutes of Health for the publication under consideration paid to her institution. Vincent Lo Re III has received research grant support from AstraZeneca outside of this submitted work paid to his institution. Cynthia Gibert has no conflicts of interest to disclose. Adeel Butt has received research grants/grants pending from Gilead Sciences and AbbVie outside of this submitted work paid to his institution. Sheldon Brown has no conflicts of interest to disclose. Joseph Lim has received payment from Bristol-Myers Squibb, Gilead Sciences, and Janssen for serving as a consultant outside of this submitted work. Maria Rodriguez-Barradas has received research grants from National Institutes of Health for the publication under consideration paid to

her institution; she also received support for travel to meetings for study or other purposes from National Institutes of Health and Veterans Affairs for the publication under consideration paid to her institution. David Rimland has received research grants from National Institutes of Health for the publication under consideration paid to his institution. Erica Kaufman has received payment from Gilead Sciences for serving on its Hepatitis C Advisory Board Panel outside of this submitted work. Amy Justice has received research grants from National Institutes of Health for the publication under consideration paid to her institution. Matthew Bidwell Goetz has received research grants from National Institute on Alcohol Abuse and Alcoholism for the publication under consideration paid to his institution.

This work was supported by the National Institute on Alcohol Abuse and Alcoholism (NIAAA) [U10 AA013566-completed, U24 AA020794, and U01 AA020790].

## **Introduction**

Isolated hepatitis B virus (HBV) core antibody (HBcAb+), defined as the presence of HBcAb in the absence of HBV surface antigen (HBsAg) and surface antibody (HBsAb), occurs in up to 34%, 50%, and 59% of HIV-HCV-coinfected, HCV monoinfected, and HIV monoinfected persons, respectively<sup>1-6</sup>. Isolated HBcAb can indicate occult HBV viremia (the presence of HBV viremia in the absence of HBsAg, resolved infection with low titers of HBsAb, a window period during acute infection, or a false positive result. However, the relationship between isolated HBcAb pattern and liver disease remains unclear with some, but not all, studies describing an association with liver disease<sup>3,9-12</sup>. Our objective was to characterize the prevalence of isolated HBcAb pattern among HIV and HIV/HCV-infected veterans in the Veterans' Aging Cohort Study (VACS) and to determine whether the isolated HBcAb pattern was associated with advanced fibrosis in HIV and HIV/HCV-coinfected Veterans.

## **Methods**

### **Study Design and Setting**

We conducted a cross-sectional study among HIV-infected individuals in the Veterans Aging Cohort Study Virtual Cohort (VACS-VC)<sup>13</sup>. Comprised of electronic medical record data from all HIV-infected veterans at Veterans Affairs (VA) medical facilities in the US, data include hospital and outpatient diagnoses, laboratory results, and pharmacy data<sup>14</sup>.

## Study Patients

The study sample included all HIV-infected individuals who enrolled in the VACS-VC between October 1996 and September 2010, had at least 180 days of follow-up, and who had all three HBcAb, HBsAg, and HBsAb serologies tested. We then searched for the first positive or negative HBcAb test and included only those who had subsequent HBsAb and HBsAg testing +/- 12 months from the first positive or negative HBcAb result. We further evaluated those for whom data variables to calculate FIB-4 and APRI scores were available within +/- 12 months of the earliest complete serologic data. The following five serologic patterns were assessed:

- 1) Isolated HBcAb: HBsAg negative, HBcAb positive, HBsAb negative
- 2) Resolved HBV infection: HBsAg negative, HBcAb positive, HBsAb positive
- 3) Non-immune HBV: HBsAg negative, HBcAb negative, HBsAb negative
- 4) Immunized HBV: HBsAg negative, HBcAb negative, HBsAb positive
- 5) Active HBV infection: HBsAg positive, HBcAb negative or positive, HBsAb negative

The main outcome was advanced hepatic fibrosis, defined by a FIB-4 >3.25 at the time of serologic evaluation (within one year of markers obtained). Secondary outcomes defining advanced hepatic fibrosis were assessed individually and included: 1) aspartate aminotransferase (AST)-to-platelet ratio index (APRI) >2.0, and 2) platelet count <140,000/ $\mu$ L. Both FIB-4 (based upon age, AST, alanine aminotransferase (ALT)<sup>15</sup> and APRI have been shown to validly identify patients

with advanced hepatic fibrosis/cirrhosis<sup>16</sup>. Similarly, platelet counts of <140,000/ $\mu$ l have been associated with advanced fibrosis/cirrhosis<sup>17</sup>.

HCV infection was defined as HCV Ab positive, HCV RNA positive, or an ICD-9 diagnosis of HCV. Age, race/ethnicity, CD4 count, log HIV viral load, diabetes, body mass index, alcohol-related disorders, drug use, smoking were collected from all patients at inclusion. Diabetes was defined by a random glucose level  $\geq$ 200 mg/dL or anti-diabetic medication use. Alcohol-related disorders and injection/non-injection drug use were defined by ICD-9 codes.

Logistic regression was used to evaluate associations between serological patterns of HBV and advanced hepatic fibrosis for patients with and without chronic HCV infection, adjusting for age, race/ethnicity, CD4 count, log HIV RNA, diabetes, body mass index, alcohol-related disorders, injection/non-injection drug use, and smoking. As substance use can be associated with immune dysfunction (ref: Friedman H, Newton C, Klein TW. Microbial infections, immunomodulation, and drugs of abuse. *Clinical microbiology reviews*. Apr 2003;16(2):209-219), which in turn can be associated with the isolated HBcAb pattern, interactions between serological patterns and alcohol-related disorders, injection/non-injection drug use, and smoking were also evaluated. All analyses were carried out using SAS software (Cary, NC).

## Results

Among 44,180 HIV infected veterans (14,568 with and 29,612 without HCV infection), 41,088 had at least 180 days of follow-up, 26,924 had all three serologies tested, and 23,970 veterans had all three serologies within one year of the first positive or negative HBcAb result. Among 23,970 HIV infected veterans, 9,327 with and 14,643 without HCV infection, 7,290 and 12,196, respectively, had complete data allowing for calculation of FIB-4 and APRI). Complete information on persons with missing hepatitis data included race, smoking status, and age. Those that were included were more likely to be white (37% vs 30%), older (mean age 47 vs 46), and less likely to be current smokers (59% vs. 64%), all  $p < 0.01$ .

In this cohort, 12.3% (1504/12196) and 37% (2707/7290) of HIV and HIV/HCV individuals, respectively, had the isolated HBcAb pattern (Table 1). Persons with isolated HBcAb were the same age or older and more likely to be nonwhite, or to have an alcohol related disorder, than were persons with resolved HBV, non-immune HBV, or HBsAb only. HIV-monoinfected persons with isolated HBcAb were more likely to have  $< 200$  CD4 cells/ $\mu$ L and a higher median level of HIV-1 RNA copies/mL. In addition, coinfecting persons who were non-immune to HBV and persons who were HBsAb+ only had a higher median HIV-1 RNA copies/mL than did persons with isolated HBcAb. In both groups there was not a statistically significant interaction noted between substance use and the isolated HBcAb pattern.



In both HIV monoinfected and HIV/HCV coinfecting groups, AST, FIB-4, and APRI scores were higher while platelets were lower in isolated HBcAb patients, when compared to those with resolved HBV, non-immune HBV, or HBsAb only serologic profiles. However, in multivariable analysis, when compared to HBsAg-negative groups, only HIV/HCV coinfecting patients with isolated HBcAb exhibited an association with advanced hepatic fibrosis. HIV+/HCV+ patients with isolated HBcAb had higher OR for advanced hepatic fibrosis by FIB-4 compared with those who had resolved HBV (OR, 1.34; 95% CI, 1.15-1.58), were non-immune (OR, 1.32; 95% CI, 1.09-1.60), or were HBsAb+ (OR, 1.65; 95% CI, 1.20-2.28). Similar results were observed when APRI and platelet counts <140,000/ $\mu$ L were evaluated. Not unexpectedly, advanced fibrosis was less in patients with isolated HBcAb than in those who were HBsAg positive. We performed a sensitivity analysis (not shown) that included HCV viremic patients which showed a similar association with HCV viremia, the isolated HBcAb profile, and advanced fibrosis, but this was not statistically significant, likely a reflection of the smaller sample size.

## **Discussion**

In this large cross-sectional study of isolated HBcAb patterns of both HIV-infected and HIV/HCV-coinfecting Veterans, we found that the isolated HBcAb pattern was associated with advanced hepatic fibrosis in HIV/HCV-coinfecting patients, but not in HIV infection without concomitant HCV infection. While most clearly seen

using a FIB-4 threshold of >3.25, similar results were seen with an APRI threshold of >2.0 or a platelet count of <140,000/ $\mu$ L.

The fact that the isolated HBcAb pattern was associated with advanced liver disease in HIV/HCV coinfecting, but not HIV monoinfected patients, suggests that there might be a multiplicative effect of HCV, HIV, and previous or ongoing low-grade HBV infection on the liver (as demonstrated by the isolated HBcAb pattern). Alternatively, both chronic HCV and the isolated HBcAb pattern may reflect a suboptimal immune response that facilitates HCV persistence, isolated HBcAb persistence, and ongoing liver damage, as manifested by an association with lower CD4 counts in a cohort of HIV infected women and in this study<sup>18</sup>. However, CD4 was incorporated into the logistic regression model for this study, suggesting that other components of the immune response may be involved. Although there were no interaction effects, an increased prevalence of substance use in those with the isolated HBcAb pattern, may be indicative of other defects in immune response, as has been seen with opiates, marijuana, and alcohol<sup>19</sup>. As this is a cross-sectional study, it was not possible to distinguish whether the advanced fibrosis reflected previous versus ongoing liver damage.

Strengths of the current analysis included a large sample size of patients with a full panel of HBV serologies and direct comparison of HCV infected and uninfected veterans with HIV infection. Limitations of the study include the cross-sectional nature of the analysis and missing data in those who had complete

serologies, which represented 19% of the cohort. This may have led to selection bias, especially in the HCV uninfected group where hepatitis B serologies were disproportionately not evaluated. However, as patients who did not have serologies were likely to be healthier than those who did, it is unlikely that this would have affected our results. Additionally, we did not assess HBV DNA in the serum, as only a fraction of the study population had serum HBV DNA measured. Moreover, FIB-4 and APRI might not be accurate in assessing intermediate degrees of fibrosis. Finally, our study was conducted in HIV-infected veterans who may have differing risk factors, including socioeconomic factors, for liver disease than other HIV infected cohorts potentially limiting generalizability to other cohorts.

In summary, we found that the isolated HBcAb pattern was associated with advanced hepatic fibrosis in HIV/HCV-coinfected, but not HIV-monoinfected, Veterans. These findings suggest that HIV- and HCV-infected persons with isolated HBcAb may be at increased risk of clinical liver complications and support the need for further studies in the longitudinal assessment of liver disease.

### **Funding**

This work was supported by the National Institute on Alcohol Abuse and Alcoholism (NIAAA) [U10 AA013566-completed, U24 AA020794, and U01 AA020790].

## **Acknowledgments**

This material is based upon work supported by the Department of Veterans Affairs, Veterans Health Administration, and Office of Research and Development. We must disclaim that the views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs.

*Potential conflicts of interest:* The corresponding author, Debika Bhattacharya, has received research support from AbbVie and Bristol-Myers Squibb and payment by IAS-USA for development of educational presentations. Janet Tate has received research grants from National Institutes of Health for the publication under consideration paid to her institution. Vincent Lo Re III has received research grants/pending grants from AstraZeneca outside of this submitted work paid to his institution. Cynthia Gibert has no conflicts of interest to disclose. Adeel Butt has received research grants/grants pending from Gilead Sciences and AbbVie outside of this submitted work paid to her institution. Sheldon Brown has no conflicts of interest to disclose. Joseph Lim has received payment from Bristol-Myers Squibb, Gilead Sciences, and Janssen for serving as a consultant outside of this submitted work. Maria Rodriguez-Barradas has received research grants from National Institutes of Health for the publication under consideration paid to her institution; she also received support for travel to meetings for study or other purposes from National Institutes of Health and Veterans Affairs for the

publication under consideration paid to her institution. David Rimland has received research grants from National Institutes of Health for the publication under consideration paid to his institution. Erica Kaufman has received payment from Gilead Sciences for serving on its Hepatitis C Advisory Board Panel outside of this submitted work. Amy Justice has received research grants from National Institutes of Health for the publication under consideration paid to her institution. Matthew Bidwell Goetz has received research grants from National Institute on Alcohol Abuse and Alcoholism for the publication under consideration paid to his institution.

## References

1. Fukuda R, Ishimura N, Niigaki M, et al. Serologically silent hepatitis B virus coinfection in patients with hepatitis C virus-associated chronic liver disease: clinical and virological significance. *Journal of medical virology*. Jul 1999;58(3):201-207.
2. Rodriguez-Torres M, Gonzalez-Garcia J, Brau N, et al. Occult hepatitis B virus infection in the setting of hepatitis C virus (HCV) and human immunodeficiency virus (HIV) co-infection: clinically relevant or a diagnostic problem? *Journal of medical virology*. Jun 2007;79(6):694-700.
3. Cacciola I, Pollicino T, Squadrito G, Cerenzia G, Orlando ME, Raimondo G. Occult hepatitis B virus infection in patients with chronic hepatitis C liver disease. *The New England journal of medicine*. Jul 1 1999;341(1):22-26.
4. Palacios R, Mata R, Hidalgo A, et al. Very low prevalence and no clinical significance of occult hepatitis B in a cohort of HIV-infected patients with isolated anti-HBc seropositivity: the BHOI study. *HIV clinical trials*. Sep-Oct 2008;9(5):337-340.
5. Lo Re V, 3rd, Wertheimer B, Localio AR, et al. Incidence of transaminitis among HIV-infected patients with occult hepatitis B. *Journal of clinical virology : the official publication of the Pan American Society for Clinical Virology*. Sep 2008;43(1):32-36.
6. Osborn MK, Guest JL, Rimland D. Hepatitis B virus and HIV coinfection: relationship of different serological patterns to survival and liver disease. *HIV medicine*. Jul 2007;8(5):271-279.
7. Torbenson M, Thomas DL. Occult hepatitis B. *Lancet Infect Dis*. Aug 2002;2(8):479-486.
8. Witt MD, Lewis RJ, Rieg G, Seaberg EC, Rinaldo CR, Thio CL. Predictors of the isolated hepatitis B core antibody pattern in HIV-infected and -uninfected men in the multicenter AIDS cohort study. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. Feb 2013;56(4):606-612.
9. Nirei K, Kaneko M, Moriyama M, Arakawa Y. The clinical features of chronic hepatitis C are not affected by the coexistence of hepatitis B virus DNA in patients negative for hepatitis B surface antigen. *Intervirolgy*. 2000;43(2):95-101.

10. Kazemi-Shirazi L, Petermann D, Muller C. Hepatitis B virus DNA in sera and liver tissue of HBsAg negative patients with chronic hepatitis C. *Journal of hepatology*. Nov 2000;33(5):785-790.
11. Filippini P, Coppola N, Pisapia R, et al. Impact of occult hepatitis B virus infection in HIV patients naive for antiretroviral therapy. *AIDS (London, England)*. Jun 12 2006;20(9):1253-1260.
12. Lo Re V, 3rd, Frank I, Gross R, et al. Prevalence, risk factors, and outcomes for occult hepatitis B virus infection among HIV-infected patients. *Journal of acquired immune deficiency syndromes (1999)*. Mar 1 2007;44(3):315-320.
13. Fultz SL, Skanderson M, Mole LA, et al. Development and verification of a "virtual" cohort using the National VA Health Information System. *Medical care*. Aug 2006;44(8 Suppl 2):S25-30.
14. Lo Re V, 3rd, Kallan MJ, Tate JP, et al. Hepatic decompensation in antiretroviral-treated patients co-infected with HIV and hepatitis C virus compared with hepatitis C virus-monoinfected patients: a cohort study. *Annals of internal medicine*. Mar 18 2014;160(6):369-379.
15. Sterling RK, Lissen E, Clumeck N, et al. Development of a simple noninvasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology (Baltimore, Md.)*. Jun 2006;43(6):1317-1325.
16. Wai CT, Greenson JK, Fontana RJ, et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology (Baltimore, Md.)*. Aug 2003;38(2):518-526.
17. Iacobellis A, Fusilli S, Mangia A, et al. Ultrasonographic and biochemical parameters in the non-invasive evaluation of liver fibrosis in hepatitis C virus chronic hepatitis. *Alimentary pharmacology & therapeutics*. Nov 1 2005;22(9):769-774.
18. Tsui JI, French AL, Seaberg EC, et al. Prevalence and long-term effects of occult hepatitis B virus infection in HIV-infected women. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. Sep 15 2007;45(6):736-740.
19. Friedman H, Newton C, Klein TW. Microbial infections, immunomodulation, and drugs of abuse. *Clinical microbiology reviews*. Apr 2003;16(2):209-219.

