

UCLA

UCLA Previously Published Works

Title

Group B streptococcus (GBS) osteomyelitis in a diabetic patient following pasteurella multocida infection

Permalink

<https://escholarship.org/uc/item/0qw9x4mf>

Journal

Journal of Clinical Images and Medical Case Reports, 4(4)

ISSN

2766-7820

Author

E, Oscar

Publication Date

2023-04-04

DOI

10.52768/2766-7820/2356

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Clinical Image*Open Access, Volume 4***Group B streptococcus (GBS) osteomyelitis in a diabetic patient following pasteurilla multocida infection****Oscar E*; Gallardo Huizar**

UCLA Olive View, Department of Medicine, Room 2B-182, 14445 Olive View Dr. Sylmar, CA 91342, USA.

***Corresponding Author: Oscar E**

UCLA Olive View, Department of Medicine, room 2B-182, 14445 Olive View Dr. Sylmar, CA 91342, USA.

Phone: +1-619-684-0268;

Email: ogallardohuizar@dhs.lacounty.gov

Abstract

Skin and Soft Tissue Infections (SSTI) resulting from cat bites most commonly occur due to Pasteurella infection. However, other pathogens such as B-Hemolytic Streptococci (BHS) have been identified and tend to be highly virulent in diabetic patients. Here we present a case of a 59 year-old man presenting with left wrist osteomyelitis following a cat bite.

Received: Jan 23, 2023

Accepted: Mar 28, 2023

Published: Apr 04, 2023

Archived: www.jcimcr.org

Copyright: © Oscar E (2023).

DOI: www.doi.org/10.52768/2766-7820/2356

Case description

A 59 year-old male with a history of uncontrolled diabetes was bitten by cat 6 weeks before presenting to an outside hospital. He was found to have Pasteurella multocida soft tissue infection and was discharged with oral antibiotics. However, he did not complete full course of outpatient therapy, and returned to the ED with increased left hand swelling and pain for three weeks.

On admission, worsening left hand swelling, pain, and erythema were noted. Pertinent laboratory data included elevated CRP (313 mg/L), leukocytosis (28 k/cumm), and sedimentation rate of 119 mm/hr. Clinical presentation was suggestive of osteomyelitis and he was treated with intravenous antibiotics and surgical wash-out. Cultures grew Streptococcus agalactiae and patient was transitioned to oral amoxicillin-clavulanate for 6-8 weeks, and scheduled for outpatient follow up.

Teaching points

Skin and Soft Tissue Infections (SSTI) resulting from cat bites tend to most commonly grow Pasteurella. Although uncom-

**Figure 1:**

Citation: Oscar E, Huizar G. Group B streptococcus (GBS) osteomyelitis in a diabetic patient following pasteurella multocida infection. *J Clin Images Med Case Rep.* 2023; 4(4): 2356.

mon, Group B streptococcus infections have been associated with cat bites [1] and in diabetics, soft tissue infections by this pathogen are highly virulent [2]. Although the initial presentation seemed to suggest a complicated infection by Pasteurella, it is important to remember that the most common pathogens in SSTIs continue to be Staphylococcus aureus and B-Hemolytic Streptococci (BHS) [3].

Conflicts of interest: None.

References

1. Ashrafian H, Griselli M, Rubens MB, Mullen MJ, Sethia B, et al. Pulmonary homograft endocarditis 19 years after a Ross procedure. *The Thoracic and Cardiovascular Surgeon.* 2007; 55: 55-56.
2. Collin SM, Shetty N, Lamagni T. Invasive Group B Streptococcus Infections in Adults, England, 2015-2016. *Emerg Infect Dis.* 2020; 26: 1174-1181.
3. Jeng A, Beheshti M, Li J, Nathan R. The role of beta-hemolytic streptococci in causing diffuse, nonculturable cellulitis: A prospective investigation. *Medicine (Baltimore).* 2010; 89: 217-226.