UCSF UC San Francisco Previously Published Works

Title

A Sinus Venosus Atrial Septal Defect Is Diagnosed by Echocardiography with an Unusual Bubble Study

Permalink https://escholarship.org/uc/item/0r04d7tg

Journal Echocardiography, 30(6)

ISSN 0742-2822

Authors

Dorosz, Jennifer L Fonseca, Brian M Kay, Joseph D <u>et al.</u>

Publication Date 2013-07-01

DOI

10.1111/echo.12191

Peer reviewed



HHS Public Access

Author manuscript *Echocardiography*. Author manuscript; available in PMC 2022 August 31.

Published in final edited form as:

Echocardiography. 2013 July ; 30(6): E182–E183. doi:10.1111/echo.12191.

A Sinus Venosus Atrial Septal Defect Is Diagnosed by Echocardiography with an Unusual Bubble Study

Jennifer L. Dorosz, M.D.^{*}, Brian M. Fonseca, M.D.[†], Joseph D. Kay, M.D.^{*,†}, Brian B. Graham, M.D.[‡], Ernesto E. Salcedo, M.D.^{*}, Robert A. Quaife, M.D.^{*}

*Division of Cardiology, School of Medicine, University of Colorado Denver, Aurora, Colorado

[†]Division of Pediatric Cardiology, School of Medicine, University of Colorado Denver, Aurora, Colorado

[‡]Division of Pulmonary - Critical Care Medicine, School of Medicine, University of Colorado Denver, Aurora, Colorado

Keywords

Eisenmeger's syndrome; sinus venosus atrial septal defect; contrast echocardiography

A 68-year-old man with a history of shortness of breath and hypoxia was undergoing workup for severe pulmonary hypertension. He was referred to pulmonary hypertension clinic for the first time. An echocardiography (iE33, Philips Medical Systems, Bothell, WA, USA) with agitated saline was performed. The calculated pulmonary artery pressure was 113 mmHg which was associated with right ventricular hypertrophy and dilatation. On contrast injection with agitated saline, the bubbles surprisingly filled the left heart rapidly before the right side (Fig. 1 and movie clips S1 and S2). Based on this bubble study, a diagnosis of Eisenmeger's syndrome from a sinus venosus atrial septal defect was made and later confirmed on cardiac magnetic resonance imaging (Avanto, Siemens Medical Solutions, Erlangen, Germany) (Fig. 2).

The fact that the bubbles first appear on the left side suggests that the shunt occurs above the level of the atria. Because of high right-sided pressures in this case, flow from the superior vena cava (SVC) preferentially travels through the defect to the right superior pulmonary vein and left atrium.

A residual left SVC with a direct connection to the left atrium via an unroofed coronary sinus was also considered as it would have also shown a similar filling pattern with agitated saline. This defect was excluded, however, because the agitated saline was serendipitously injected into the right hand.

Address for correspondence and reprint requests: Jennifer L. Dorosz, M.D., Division of Cardiology, University of Colorado Denver, 12631 East 17th Avenue, Mail Stop B130, Aurora, CO 80045. Fax: (303)-724-2094; Jennifer.Dorosz@ucdenver.edu. Supporting Information

Additional Supporting Information may be found in the online version of this article:

In this case, the shunt was not diagnosed until late in life. He had been treated for years with vasodilators for presumed primary pulmonary hypertension. Unfortunately, he died 3 months later from severe Eisenmeger's syndrome. In this case, the initial echocardiography with agitated saline was essential in making the diagnosis, but because of the large right-to-left shunt, further bubble studies should be avoided in patients with known Eisenmeger's syndrome.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Dorosz et al.




Figure 1.

Echocardiography with agitated saline injection in a patient with a sinus venosus atrial septal defect. **A.** shows a four-chamber view prior to the injection of agitated saline, highlighting the right ventricular enlargement and hypertrophy associated with severe pulmonary hypertension. **B.-D**. show the progress of the bubbles after the 2nd, 4th, and 6th beats, respectively. LA = left atrium; LV = left ventricle; RA = right atrium; RV = right ventricle.

Dorosz et al.




Figure 2.

The MRI of the sinus venosus atrial septal defect. LA = left atrium; RA = right atrium; SVC = superior vena cava.