

UC Berkeley

UC Berkeley Previously Published Works

Title

A reversible single-molecule switch based on activated antiaromaticity

Permalink

<https://escholarship.org/uc/item/0s94p7sw>

Journal

Science Advances, 3(10)

ISSN

2375-2548

Authors

Yin, Xiaodong
Zang, Yaping
Zhu, Liangliang
[et al.](#)

Publication Date

2017-10-06

DOI

10.1126/sciadv.aao2615

Peer reviewed

CHEMICAL PHYSICS

A reversible single-molecule switch based on activated antiaromaticity

Xiaodong Yin,^{1*} Yaping Zang,^{2*} Liangliang Zhu,^{3*} Jonathan Z. Low,¹ Zhen-Fei Liu,⁴ Jing Cui,² Jeffrey B. Neaton,^{4†} Latha Venkataraman,^{1,2†} Luis M. Campos^{1†}

Single-molecule electronic devices provide researchers with an unprecedented ability to relate novel physical phenomena to molecular chemical structures. Typically, conjugated aromatic molecular backbones are relied upon to create electronic devices, where the aromaticity of the building blocks is used to enhance conductivity. We capitalize on the classical physical organic chemistry concept of Hückel antiaromaticity by demonstrating a single-molecule switch that exhibits low conductance in the neutral state and, upon electrochemical oxidation, reversibly switches to an antiaromatic high-conducting structure. We form single-molecule devices using the scanning tunneling microscope–based break-junction technique and observe an on/off ratio of ~ 70 for a thiophenylidene derivative that switches to an antiaromatic state with 6-4-6- π electrons. Through supporting nuclear magnetic resonance measurements, we show that the doubly oxidized core has antiaromatic character and we use density functional theory calculations to rationalize the origin of the high-conductance state for the oxidized single-molecule junction. Together, our work demonstrates how the concept of antiaromaticity can be exploited to create single-molecule devices that are highly conducting.

INTRODUCTION

The ability to miniaturize macroscopic devices and electronic components at the molecular level provides unprecedented fundamental information on chemical structure and, importantly, bridges the gap toward single-molecule devices (1–4). Although the concept for a molecular rectifier was proposed in 1974 (5), recent breakthroughs, which stem from advanced nanofabrication and device characterization techniques, have led to the observation of thermoelectric behavior (6), the Kondo effect (7, 8), rectification (9), and switching (10–14). These advances have been due, in part, to the ability to use different aromatic molecular components as the functional elements of the devices. Surprisingly, structures that focus on Hückel antiaromaticity, a classical physical organic chemistry concept, have been overlooked. Although it has been shown that the aromatic character of a molecular component can control conductance, the concept of antiaromaticity has not been successfully implemented in molecular switches, because antiaromatic compounds are generally unstable (15).

Herein, we designed an electrochemically activated redox-active molecular wire that is based on the thieno derivative [8,8'-biindeno[2,1-*b*]thiophenylidene (BTP)] (16) of 9,9'-bifluorenylidene (Fig. 1) (17). These systems are known to be efficient electron acceptors, because upon reduction, they gain 6-6-6- π electron aromaticity. The important feature that we exploit is oxidation to the 6-4-6- π electron antiaromatic central rings, which does not involve making or breaking σ bonds, as in the case of most commonly studied switching mechanisms (10, 18, 19). Upon oxidation, the central ring of the BTP system takes on antiaromatic characteristics with 4 π electrons. Because antiaromatic systems are known to be unstable, the carbocycles flanking the central ring delocalize the charge and provide stability (20, 21). To form single-

molecule junctions to investigate the charge transport properties of the parent BTP core, we added thiochroman linkers (22), yielding the structure shown in Fig. 2A [thiochromatin derivative of BTP (TBTP)]. Synthetic and molecular characterization details are given in the Supplementary Materials.

RESULTS AND DISCUSSION

We first characterize the solution-based optical and electronic properties of the TBTP by cyclic voltammetry (CV) and spectroelectrochemistry. Figure 2B shows a CV plot where we see four reversible redox peaks, two corresponding to oxidation and two corresponding to reduction. We note that the two oxidation peaks at $E_{\text{ox}1} = 0.41$ V and $E_{\text{ox}2} = 0.56$ V (versus ferrocene [$\text{Fc}^{+/0}$]) are close in energy, whereas the reduction peaks are well separated. To characterize the changes in optical properties of TBTP under various applied electrochemical potentials, we measure ultraviolet–visible (UV-vis) absorption spectra of TBTP at different potentials. We start with the neutral molecule that exhibits an optical energy gap of 1.77 eV (fig. S1). Upon applying a potential of 0.9 V (versus ferrocene [$\text{Fc}^{+/0}$]), the molecule is doubly oxidized (+2 state), and the optical gap decreases to about 0.85 eV (Fig. 2C). The spectroelectrochemical changes are reversible, as is visible from the CV and additional UV-vis spectra (fig. S2), which shows the spectral characteristics of the transient radical cation. To probe changes to the molecular structure after oxidation, we chemically oxidize TBTP with $\text{AgSbF}_6/\text{I}_2$, and this shows further evidence of the +2 state by matching the UV-vis spectra to the spectroelectrochemistry data (the radical cation is not observed by chemical oxidation). TBTP can be chemically oxidized reversibly (fig. S3), and the dication exhibits remarkable stability at ambient conditions for several hours. Considering that the antiaromaticity of this stable system can be probed by proton nuclear magnetic resonance ($^1\text{H-NMR}$) (23), we collected the $^1\text{H-NMR}$ spectra for the neutral and chemically oxidized TBTP at room temperature. The NMR spectra show that the proton signal from the benzene ring of the BTP core exhibits a marked upshielding shift, indicating antiaromatic character (fig. S4). Although this upshielding shift is not seen for all protons on the core due to charge delocalization, this key result is

Copyright © 2017
The Authors, some
rights reserved;
exclusive licensee
American Association
for the Advancement
of Science. No claim to
original U.S. Government
Works. Distributed
under a Creative
Commons Attribution
NonCommercial
License 4.0 (CC BY-NC).

¹Department of Chemistry, Columbia University, New York, NY 10027, USA. ²Department of Applied Physics, Columbia University, New York, NY 10027, USA. ³State Key Laboratory of Molecular Engineering of Polymers, Department of Macromolecular Science, Fudan University, Shanghai 200433, China. ⁴Molecular Foundry, Lawrence Berkeley National Laboratory, and Department of Physics, University of California, Berkeley, Berkeley, CA 94720, USA.

*These authors contributed equally to this work.

†Corresponding author. Email: lv2117@columbia.edu (L.V.); jbnaton@lbl.gov (J.B.N.); lcampos@columbia.edu (L.M.C.)




Fig. 1. Model of the electrochemically activated BTP switch. BTP shows the neutral, low-conductance state. Upon oxidation, BTP gains antiaromatic character to the 6-4-6- π electron state that exhibits high conductance.

consistent with the NMR spectra predicted by density functional theory (DFT) calculations, as detailed in Materials and Methods. Thus, it can be concluded that the BTP core takes on antiaromatic character in the +2 state, consistent with the simple electron-counting picture presented above.

To test the hypothesis that antiaromaticity can lead to highly conductive molecular devices, we performed single-molecule measurements under an electrochemical potential using the scanning tunneling microscope break-junction (STM-BJ) technique (24, 25). The details of the method have been described before by Capozzi *et al.* (24). Briefly, we use a gold STM tip and substrate, which forms the working electrodes, whereas an additional Pt wire introduced into the fluid cell serves as the counter (or gate) electrode. Measurements are carried out in 0.1 mM TBTP solution in propylene carbonate (PC), a polar solvent, with 0.1 M tetrabutylammonium perchlorate as the supporting electrolyte, and potential calibrations are carried out by adding ferrocene to the solution and measuring its reversible oxidation potential. The tip is insulated with Apiezon wax to reduce the background ionic current (26). The conductance of TBTP is first measured without applying a gate voltage. Figure 2D shows the one-dimensional (1D) conductance histograms of TBTP at a tip bias of 0.09, 0.45, and -0.45 V relative to the substrate. Each conductance histogram reveals a clear narrow peak at a bias-dependent conductance value. The inset of Fig. 2D shows the peak conductance plotted against the tip bias (see fig. S5 for full data sets), where a modest conductance increase of a factor of 5 is seen within this bias range. Such a bias polarity-dependent conductance indicates that the highest occupied molecular orbital (HOMO) dominates charge transport (27).

The electrochemical switching of the single-molecule junction was accomplished by changing the voltage applied to the Pt gate electrode (relative to the substrate) and repeating the measurements while maintaining a low bias voltage of 25 mV on the tip (also relative to the substrate). Note that in this STM-BJ measurement, the voltage is applied to the gate electrode, relative to the substrate. Therefore, oxidation events are observed at negative gate voltages. This is corroborated by the in situ electrochemistry shown in fig. S6A. By contrast, in traditional CV measurements, the potential is applied to the working electrode, relative to the reference. Conductance histograms measured at different voltages applied to the gate are shown in Fig. 3A. At the positive gate voltage, the conductance peak shifts to lower values, consistent with transport being dominated by the HOMO. At negative gate voltages, a modest increase of the conductance is first observed until an applied gate voltage of about -1.4 V (+1 V measured potential relative to $\text{Fc}^{+/0}$). Beyond -1.4 V, the conductance peak shifts markedly by a factor of ~ 70 (Fig. 3A), and the histograms broaden—an aspect we will discuss further below. This sharp increase in conductance is visible in the inset of Fig. 3A, where we plot the measured conductance against the gate potential (see fig. S5B for full data sets). We show further that




Fig. 2. STM-BJ measurements and electrochemical characterizations of TBTP. (A) A schematic diagram of a single Au-TBTP-Au junction. (B) Cyclic voltammogram of TBTP in dichloromethane solution (1 mM) with Bu_4NPF_6 (0.1 M) as the supporting electrolyte. Scan rate used is 100 mV/s. (C) Steady-state linear absorption spectroscopy of the neutral TBTP (orange) and the dication (green) obtained at a potential of 0.9 V (versus ferrocene $\text{Fc}^{+/0}$). a.u., absorbance units. (D) Logarithm-binned 1D histograms (100 bins/decade) for TBTP in PC with tetrabutylammonium perchlorate at a tip bias of 0.09, 0.45, and -0.45 V (tip relative to the substrate). Inset: Peak conductance value versus applied tip bias.

the switching is reversible by changing the gate potential between negative (< -1.5 V) and positive (> 1 V) values, in an alternating fashion, after every 100 conductance traces. The conductance peak values determined from the histogram of these traces (fig. S6B) are plotted in Fig. 3B, yielding an average on/off ratio value of ~ 70 . Note that the redox-switching experiments were conducted at ambient conditions, without the rigorous exclusion of air and moisture. In addition, to show that the gate potential does change the oxidation state of the molecules in solution in our STM setup, we record the in situ linear sweep voltammetry (LSV), measuring the current through the STM tip as a function of the applied gate voltage. We do this measurement after adding ferrocene to the solution to calibrate the applied gate voltage and confirm oxidation states (fig. S6A). We see a broad oxidation peak at a gate voltage of -1.4 V, consistent with our finding that a negative gate voltage needs to be applied to measure the conductance of the oxidized TBTP.

To confirm that we are measuring the conductance of a TBTP molecule bridging the gap between two Au electrodes, we compare 2D conductance-displacement histograms compiled from these data in Fig. 3 (C and D) (28). Both figures show a molecular feature that extends by about 1 nm relative to the point where the gold point contact is ruptured. We have previously shown that the extent of the 2D histogram feature relates to the molecular backbone length (28), and thus, these data reveal that a stable molecular junction is formed at the high negative gate bias. The key result of this study is that changing the oxidation state of TBTP without making/breaking covalent bonds results




Fig. 3. STM-BJ results showing the reversible single-molecule switch. (A) Logarithm-binned 1D histograms for **TBTP** in PC with tetrabutylammonium perchlorate as the supporting electrolyte at a gate voltage (V_G) of +2, -0.5, -1.4, and -1.5 V. The gate voltage is applied relative to the substrate. Inset: Peak conductance value versus applied gate voltage. (B) Cyclic on/off conductance values determined by switching between the positive gate voltage and negative gate voltage chosen to be beyond the oxidation/reduction thresholds. Error bars correspond to the full width at half maximum of the conductance histogram peaks. (C and D) 2D conductance-displacement histogram for **TBTP** at off state ($V_G = 0$ V) (C) and on state ($V_G = -1.5$ V) (D). Note that the bias applied across the molecular junction is fixed at 25 mV (tip relative to the substrate) during these measurements.

in a large change in conductance, which can be attributed to the increased antiaromatic character of the oxidized species.

To provide further support for the observed increased conductance upon oxidation, we turn to DFT-based calculations of the single-molecule junction. We first construct the molecular junction containing the neutral **TBTP** (see Fig. 4A for the geometry) and perform conductance calculations with a DFT-based nonequilibrium Green's function (NEGF) approach, with the Transiesta package (see Materials and Methods for details) (29). Figure 4B shows the resulting transmission as a function of energy (orange trace). The conductance at zero bias (transmission at Fermi energy) is small and corresponds to a junction in its "off" state. We note that for a quantitative comparison between theory and experiment, self-energy corrections to the DFT mean-field eigensystem are required, as in, for example, the approximate DFT + Σ method (see Materials and Methods and the Supplementary Materials) (30). Here, we focus on the comparison between the neutral and the oxidized **TBTP** molecule. To model the molecule in a +2 oxidation state, eight fluorine atoms are added around the **BTP** core at approximately 1 Å above and below the central rings, such that the total Mulliken charge on the molecule reaches +2 in the gas phase (see Fig. 4A for the geometry). We then perform a DFT-NEGF calculation of the molecular junction with the fluorine atoms present. The resulting transmission is shown in Fig. 4B.




Fig. 4. DFT-based transmission calculations. (A) The molecular junction containing the neutral **TBTP** molecule (H, light pink; C, brown; and S, yellow) and eight fluorine atoms at about 1 Å above (dark blue) and below (light blue) the central rings to model the **TBTP** molecule in a +2 oxidized state. (B) Transmission as a function of energy for the neutral (orange) and the oxidized molecule (green). E_F is the Fermi level of the junction.

The transmission functions of both the neutral and oxidized species show stark differences. The oxidized **TBTP** leads to a partially occupied resonance and much higher transmission close to the Fermi energy. This increase corresponds to a conductance enhancement of a factor of ~60, consistent with the experimental results, where the average on/off ratio is ~70. Although the actual conductance of the oxidized species may be sensitive to the parameters of the calculation, the qualitative trend of increasing conductance after oxidation is correctly captured here. Because the resonance close to the Fermi level is quite narrow, small changes in the metal work function as a result of changes in junction geometry or ionic environment could result in large changes in conductance. This is consistent with the experimental data that shows a broader conductance peak for the +2 state of **TBTP**.

MATERIALS AND METHODS

Molecular characterizations

^1H -NMR and ^{13}C -NMR were recorded on a Bruker Avance III 500 (500 MHz) spectrometer in dichloromethane- d_2 [residual solvent peak at $\delta = 5.32$ parts per million (ppm) for ^1H NMR] and chloroform- d solution (residual solvent peak at $\delta = 77.16$ ppm for ^{13}C NMR). Mass spectra were obtained at the Columbia University Mass Spectrometry Facility using a Xevo G2-XS (Waters) equipped with a quadrupole time-of-flight detector with multiple inlet and ionization capabilities, including electrospray ionization, atmospheric pressure chemical ionization, and atmospheric solids analysis probe. The base peaks were usually obtained as $[\text{M}]^+$ or $[\text{M} + \text{H}]^+$ ions. UV-vis absorption data were acquired on a Varian Cary 5000 UV-Vis-NIR spectrophotometer. Cyclic voltammograms were recorded on a CHI 66 electrochemical workstation using Pt plate electrode as the working electrode, Pt wire as the counter electrode, and Ag/AgCl electrode as the reference electrode at room temperature.

STM-BJ measurements

Conductance measurements were performed using a custom STM that has been described in detail before (31). Conductance measurements were performed in dilute solutions (10 to 100 μM) in PC with tetrabutylammonium perchlorate as the supporting electrolyte. The insulated tips were created by driving a mechanically cut gold tip through Apiezon wax (26). 1D conductance histograms were constructed using

14. F. Chen, J. He, C. Nuckolls, T. Roberts, J. E. Klare, S. Lindsay, A molecular switch based on potential-induced changes of oxidation state. *Nano Lett.* **5**, 503–506 (2005).
15. W. Chen, H. Li, J. R. Widawsky, C. Appayee, L. Venkataraman, R. Breslow, Aromaticity decreases single-molecule junction conductance. *J. Am. Chem. Soc.* **136**, 918–920 (2014).
16. C.-Y. Chiu, H. Wang, F. G. Brunetti, F. Wudl, C. J. Hawker, Twisted but conjugated: Building blocks for low bandgap polymers. *Angew. Chem. Int. Ed.* **53**, 3996–4000 (2014).
17. F. G. Brunetti, X. Gong, M. Tong, A. J. Heeger, F. Wudl, Strain and Hückel aromaticity: Driving forces for a promising new generation of electron acceptors in organic electronics. *Angew. Chem. Int. Ed.* **49**, 532–536 (2010).
18. R. Breslow, Antiaromaticity. *Acc. Chem. Res.* **6**, 393–398 (1973).
19. C. Jia, A. Migliore, N. Xin, S. Huang, J. Wang, Q. Yang, S. Wang, H. Chen, D. Wang, B. Feng, Z. Liu, G. Zhang, D.-H. Qu, H. Tian, M. A. Ratner, H. Q. Xu, A. Nitzan, X. Guo, Covalently bonded single-molecule junctions with stable and reversible photoswitched conductivity. *Science* **352**, 1443–1445 (2016).
20. W. C. Lothrop, Biphenylene. *J. Am. Chem. Soc.* **63**, 1187–1191 (1941).
21. S. Schneebeli, M. Kamenetska, F. Foss, H. Vazquez, R. Skouta, M. Hybertsen, L. Venkataraman, R. Breslow, The electrical properties of biphenylenes. *Org. Lett.* **12**, 4114–4117 (2010).
22. Y. S. Park, J. R. Widawsky, M. Kamenetska, M. L. Steigerwald, M. S. Hybertsen, C. Nuckolls, L. Venkataraman, Frustrated rotations in single-molecule junctions. *J. Am. Chem. Soc.* **131**, 10820–10821 (2009).
23. M. D. Peeks, T. D. W. Claridge, H. L. Anderson, Aromatic and antiaromatic ring currents in a molecular nanoring. *Nature* **541**, 200–203 (2017).
24. B. Capozzi, Q. Chen, P. Darancet, M. Kotiuga, M. Buzzeo, J. B. Neaton, C. Nuckolls, L. Venkataraman, Tunable charge transport in single-molecule junctions via electrolytic gating. *Nano Lett.* **14**, 1400–1404 (2014).
25. B. Xu, N. J. Tao, Measurement of single-molecule resistance by repeated formation of molecular junctions. *Science* **301**, 1221–1223 (2003).
26. L. A. Nagahara, T. Thundat, S. M. Lindsay, Preparation and characterization of STM tips for electrochemical studies. *Rev. Sci. Instrum.* **60**, 3128–3130 (1989).
27. B. Capozzi, J. Z. Low, J. Xia, Z.-F. Liu, J. B. Neaton, L. M. Campos, L. Venkataraman, Mapping the transmission functions of single-molecule junctions. *Nano Lett.* **16**, 3949–3954 (2016).
28. M. Kamenetska, M. Koentopp, A. C. Whalley, Y. S. Park, M. L. Steigerwald, C. Nuckolls, M. S. Hybertsen, L. Venkataraman, Formation and evolution of single-molecule junctions. *Phys. Rev. Lett.* **102**, 126803 (2009).
29. M. Brandbyge, J.-L. Mozos, P. Ordejón, J. Taylor, K. Stokbro, Density-functional method for nonequilibrium electron transport. *Phys. Rev. B* **65**, 165401 (2002).
30. S. Y. Quek, L. Venkataraman, H. J. Choi, S. G. Louie, M. S. Hybertsen, J. B. Neaton, Amine–gold linked single-molecule circuits: Experiment and theory. *Nano Lett.* **7**, 3477–3482 (2007).
31. L. Venkataraman, J. E. Klare, C. Nuckolls, M. S. Hybertsen, M. L. Steigerwald, Dependence of single-molecule junction conductance on molecular conformation. *Nature* **442**, 904–907 (2006).
32. J. L. Malandra, N. S. Mills, D. E. Kadlecik, J. A. Lowery, Dications of tetrabenzofulvalenes. Paratropicity and σ donation in perpendicular antiaromatic systems. *J. Am. Chem. Soc.* **116**, 11622–11623 (1994).
33. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, D. J. Fox, *Gaussian 09, Revision D.01* (Gaussian Inc., 2009).
34. N. S. Mills, K. B. Llagostera, Summation of nucleus independent chemical shifts as a measure of aromaticity. *J. Org. Chem.* **72**, 9163–9169 (2007).
35. Z. Chen, C. S. Wannere, C. Corminboeuf, R. Puchta, P. v. R. Schleyer, Nucleus-independent chemical shifts (NICS) as an aromaticity criterion. *Chem. Rev.* **105**, 3842–3888 (2005).
36. J. P. Perdew, K. Burke, M. Ernzerhof, Generalized gradient approximation made simple. *Phys. Rev. Lett.* **77**, 3865–3868 (1996).
37. J. M. Soler, E. Artacho, J. D. Gale, A. García, J. Junquera, P. Ordejón, D. Sánchez-Portal, The SIESTA method for *ab initio* order-*N* materials simulation. *J. Phys. Condens. Matter* **14**, 2745–2779 (2002).
38. J. B. Neaton, M. S. Hybertsen, S. G. Louie, Renormalization of molecular electronic levels at metal-molecule interfaces. *Phys. Rev. Lett.* **97**, 216405 (2006).
39. S. Refaely-Abramson, S. Sharifzadeh, N. Govind, J. Autschbach, J. B. Neaton, R. Baer, L. Kronik, Quasiparticle spectra from a nonempirical optimally tuned range-separated hybrid density functional. *Phys. Rev. Lett.* **109**, 226405 (2012).
40. D. A. Egger, Z.-F. Liu, J. B. Neaton, L. Kronik, Reliable energy level alignment at physisorbed molecule–metal interfaces from density functional theory. *Nano Lett.* **15**, 2448–2455 (2015).

Acknowledgments

Funding: The experimental work was supported primarily by the NSF (award no. DMR-1507440). The computational work was supported by the Molecular Foundry and by the Materials Sciences and Engineering Division (Theory FWP), U.S. Department of Energy, Office of Basic Energy Sciences (contract no. DE-AC02-05CH11231). Portions of the computation work were performed at the National Energy Research Scientific Computing Center. J.Z.L. would like to thank the Agency for Science, Technology and Research (A*STAR) Graduate Academy in Singapore for a graduate fellowship. **Author contributions:** The project was planned by L.M.C. and L.V. **TBTP** was synthesized and characterized by L.Z. and X.Y. Electrochemistry and UV-vis spectroscopy measurements were conducted by X.Y. and J.Z.L. STM-BJ measurements were conducted by L.V., Y.Z., and J.C. The data analysis was performed by L.V., L.M.C., Y.Z., and X.Y. The calculations were carried out by Z.-F.L. and J.B.N. The manuscript was prepared by J.B.N., L.M.C., L.V., X.Y., Y.Z., and Z.-F.L. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors.

Submitted 1 July 2017

Accepted 5 October 2017

Published 27 October 2017

10.1126/sciadv.aao2615

Citation: X. Yin, Y. Zang, L. Zhu, J. Z. Low, Z.-F. Liu, J. Cui, J. B. Neaton, L. Venkataraman, L. M. Campos, A reversible single-molecule switch based on activated antiaromaticity. *Sci. Adv.* **3**, eao2615 (2017).