

# UC Riverside

## International Organization of Citrus Virologists Conference Proceedings (1957-2010)

### Title

Current Situation on Citrus Virus and Virus-like Diseases and their  
Vectors in Mexico

### Permalink

<https://escholarship.org/uc/item/0sr3q361>

### Journal

International Organization of Citrus Virologists Conference Proceedings  
(1957-2010), 16(16)

### ISSN

2313-5123

### Authors

Rocha-Peña, M. A.  
López Arroyo, J. I.  
Peña del Rio, M. A.  
et al.

### Publication Date

2005

### DOI

10.5070/C50sr3q361

Peer reviewed

## SURVEYS, CERTIFICATION AND OTHER TOPICS

### Current Situation on Citrus Virus and Virus-like Diseases and their Vectors in Mexico

M. A. Rocha-Peña<sup>1</sup>, J. I. López Arroyo<sup>2</sup>, M. A. Peña del Río<sup>2</sup>,  
and I. H. Almeida-León<sup>1</sup>

<sup>1</sup>INIFAP/UANL, Unidad de Investigación en Biología Celular y Molecular,  
Apartado Postal 128-F, Cd. Universitaria, San Nicolás de los Garza, NL 66450, Mexico;

<sup>2</sup>INIFAP, Campo Experimental General Terán, Apartado Postal 3, General Terán,  
NL 67400, Mexico

**ABSTRACT.** The available information on citrus virus and virus-like pathogens in Mexico is presented here. *Citrus exocortis viroid* (CEVd) is common in old plantations of sweet-fruited species grafted on the predominant sour orange rootstock. CEVd occurs commonly in mixed infections with either *Hop stunt viroid* variants or *Citrus viroid III* on Tahiti lime plantations from different locations. *Citrus tristeza virus* has been reported to occur in symptomless infections in 17 states of the country. Psorosis-like symptoms are common in some old groves, but occur at relatively low incidence throughout the country. Citrus blight has been identified in some particular groves grafted on rough lemon in the Yucatan Peninsula. No information is available on the occurrence of other viral pathogens such as vein enation virus and tatter leaf virus.

Citrus in Mexico is of major economic and social importance and is a permanent source of income. Currently the citrus industry consists of more than five hundred thousand hectares distributed in different ecological regions around the country. There are approximately 67,000 citrus growers and more than 138 packinghouses and 19 fruit processing plants nationwide. Annual production is 5.5 million tons of fruit, of which 75% is for fresh market, 15% for fruit processing and 10% for export.

**Citrus viroids.** Surveys conducted in selected groves in the northeastern states of Nuevo León and Tamaulipas showed a 50% incidence of *Citrus exocortis viroid* (CEVd), as determined by inoculation into citron indicator plants. Infections of CEVd were symptomless and hence overlooked by citrus growers, because of the predominance of sour orange rootstock in most citrus plantations (26, 33). When fruit production and tree

growth were evaluated in a grove of sweet orange on Troyer citrange rootstock affected by CEVd, the results showed that infected trees produced only one third of the fruit that was produced on the CEVd-free trees. Trunk diameter and canopy volume of infected trees also showed values reduced by close to one-third (33). The occurrence CEVd along with other citrus viroids is presumed to be widespread in old line cultivars of sweet-fruited citrus species (orange, grapefruit, mandarin and mandarin hybrids) around the country, mostly because of the predominance of sour orange as a rootstock (28). In Tahiti lime groves from several parts of the country, CEVd occurs commonly in mixed infections with *Citrus viroid II* (CVdII) (*Hop stunt viroid* variants), and *Citrus viroid III* (CVdIII) (1, 2). The typical symptoms of cachexia disease caused by CVdIIb or CVdIIc (25) occur commonly on trees grafted on alemow rootstock (2). These trees commonly show stunting and

reduced trunk diameter below the bud union, along with brownish lesions on the wood which are noticeable when a portion of bark is removed at the bud union (2). The presence of citrus viroids in either sweet citrus species or Tahiti lime samples collected from diverse locations indicates the continuing use of infected budwood sources for the establishment of new plantations.

**Tristeza.** *Citrus tristeza virus* (CTV) was reported in Mexico as early as 1982 in a foundation block in the northeastern state of Tamaulipas (36). CTV was found later in several other parts of the country (29, 37), and this subsequently led to the establishment of a nationwide campaign conducted by the federal government for a serological survey for CTV (31). Currently, the presence of symptomless CTV-infected trees has been reported in 17 of the citrus growing states of the country (32).

Relatively limited research has been conducted on the serological and molecular characterization of CTV isolates. Both mild and decline-inducing CTV isolates were reported in the state of Nuevo Leon (29, 34) as identified by tests using the strain specific monoclonal antibody MCA 13 (23). Further studies of these isolates using single stranded conformational polymorphism analysis (17) and molecular cloning and sequencing (11), revealed the presence of mild, decline-inducing and stem-pitting CTV isolates. Some additional information on the molecular characterization of some CTV isolates from Tamaulipas and Veracruz is also available (9, 13).

To assess the distribution and dissemination of CTV in the field, a survey conducted twice a year during 1996-1998 which looked for CTV in individual trees in three commercial groves in the state of Nuevo Leon, showed an increase in CTV occurrence of 0.27 and 0.18%, in the groves of Montemorelos and Linares, respectively. No CTV spread was

detected in the grove of Hualahuises (34). Even though some samples were reactive with the decline strain discriminating MCA-13 monoclonal antibody (23), no disease symptoms were observed under field conditions (29, 34). Similar studies conducted in the neighboring state of Tamaulipas during 2001-2002 showed some increase in the CTV incidence as measured from initial infected foci, showing the relative spread of CTV in the groves (13).

The aphid species that vector CTV and which are prevalent in most of the citrus groves are *Aphis gossypii* Glover, *A. spiraecola* Patch and *Toxoptera aurantii* Boyer de Fonscolombe (7, 18, 22, 24, 34). The presence of *Toxoptera citricida* (Kirkaldy) (the brown citrus aphid or BrCA), the most efficient CTV vector (27) was first reported in February of 2000 in the Peninsula of Yucatan (38), and in the following four years, this vector moved into the neighboring states of Campeche, Tabasco and Veracruz on the Gulf of Mexico (32). Upon discovery of the BrCA infestations in the Yucatan Peninsula in 2000, the federal government jointly with local Citrus Plant Protection Boards established a campaign for the permanent release of aphid predators. These predators included the coccinellid beetles *Harmonia axyridis* (Pallas), *Cycloneda sanguinea* L., and *Olla v-nigrum* (Hagen), and the chrysopid lacewings *Ceraeochrysa claveri* (Navás), and *C. cubana*, (Mulsant) (14, 16). The aphid predators have not prevented significant long distance moving of BrCA. In addition these insect predators, some field and laboratory experiments have been conducted with entomopathogenic fungi. *Beauveria bassiana* (Balsamo) Vuillemin, *Paecilomyces fumoroseus* (Wize) Brown & Smith and *Verticillium lecanii* (Zimmermann) Viégas, have shown variable infection rates in BrCA populations (3). The long distance movement of the BrCA has been mostly associated with incidental movement by people and/or trans-

portation vehicles, rather than natural spread of the insect. Therefore, the effectiveness of natural enemies has not been fully assessed.

**Citrus psorosis.** Symptoms of bark scaling typical of citrus psorosis are commonly found in some old groves in different regions of the country, particularly on old line cultivars of sweet-fruited species (19). Positive identification of *Citrus psorosis virus* has been documented in the northeastern state of Nuevo Leon (12).

**Disorders of unknown etiology.** Citrus blight has been identified in some groves on rough lemon rootstock in the Yucatan Peninsula (39). There are other virus-like disorders named “Amachamiento” in sweet orange in Veracruz and Tamaulipas (20), and “Lethal yellowing of Tahiti Lime” in the state of Veracruz (5). The plants affected by these two disorders show poor growth and reduced fruit set and production (5, 20). While “Amachamiento” is of unknown etiology (20), “Lethal yellowing of Tahiti Lime” has been associated with the presence of citrus viroids and phytoplasmas (35). However, proof of their role in causing the disease symptoms remains to be determined.

**Other insect vectors.** The presence of the Asiatic citrus psyllid (ACP) (*Diaphorina citri* Kawasaki), the vector of the Huanglongbing disease bacteria (6), was reported in the Peninsula of Yucatan in 2001, and its presence has been subsequently recorded in several other citrus growing states of the country (15). In addition, the presence of several mite species of the genus *Brevipalpus*, which are known vectors of Citrus leprosis virus, has been reported to occur in several citrus growing states (10). The presence of both the ACP and mites warrants attention because of the recent reports of Huanglongbing disease in Brazil (4) and Citrus leprosis virus in Central America (21, 30).

**Citrus certification program.** In 2000, the federal government established some specific regulatory actions and published federal guidelines and policies regarding the nationwide campaign against CTV and the brown citrus aphid, as well as the establishment and maintenance of the Citrus Budwood Certification Program. Currently there are ten certified foundation blocks that provide clean budwood of diverse citrus cultivars and rootstock seed nationwide (8).

## LITERATURE CITED

- Almeyda-León, I. H., M. M. Iracheta-Cárdenas, J. Jasso-Argumedo, S. A. Curti-Díaz, P. Ruíz-Beltrán, and M. A. Rocha-Peña  
2002. Reexamination of citrus viroids of Tahiti lime in Mexico. *Rev. Mex. Fitopatol.* 20: 152-160.
- Alvarado-Gómez, O. G., M. A. Rocha-Peña, S. Silva-Vara, J. P. Martínez-Soriano, R. F. Lee, R. Rivera Bustamante, and P. Ruíz-Beltrán  
2000. Citrus exocortis and citrus cachexia in commercial groves of Tahiti lime in México. In: *Proc. 14th Conf. IOCV*, 289-293. IOCV, Riverside, CA.
- Berlanga-Padilla, A.  
2003. Avances en el control microbiano del pulgón café de los cítricos en México. In: *Memorias Simposio Internacional de Limas Acidas*. M. A. Rocha-Peña and V. M. Medina-Urrutia (eds.), 215-225. Colima, Colima. México.
- Colleta-Filho, H. D., M. L. P. N. Targon, M. A. Takita, J. D. De Negri, A. M. Do Amaral, G. W. Müller, J. Pompeu, Jr., and M. A. Machado  
2005. Identification of *Candidatus Liberibacter asiaticus* associated with Huanglongbing symptoms in the State of São Paulo, Brazil. In: *Proc. 16th Conf. IOCV*, 429-431. IOCV, Riverside, CA.
- Curti-Díaz, S. A., X. Loredó-Salazar, U. Díaz-Zorrilla, J. A. Sandoval-Rincón, and J. Hernández-Hernández  
2000. Tecnología para producir limón Persa. Libro Técnico Num. 8. INIFAP. Campo Experimental Ixtacuaco. México. 145 p.

6. da Graça, J. V., and L. Korsten  
2004. Citrus huanglongbing: Review, Present Status, and Future Strategies. In: *Diseases of Fruits and Vegetables*. Vol. I (S.A.M.H.Naqvi ed.), 229-245. Kluwer Academic Publishers, Dordrecht.
7. Gaona-García, G., E. Ruíz-Cancino, and R. Peña-Martínez  
2000. Los pulgones (HOMOPTERA:APHIDIDAE) y sus enemigos naturales en la naranja, *Citrus sinensis* (L.) en la Zona Centro de Tamaulipas, México. *Acta Zool. Mex.* 81: 11-12.
8. Hernández-Baeza, J., and H. M. Sánchez-Anguiano  
2004. Avances del programa nacional de reconversión productiva de la cadena cítrica en México. In: *Taller Internacional de Citricultura*. INIFAP. Cd. Obregón, Sonora. México.
9. Herrera-Isidrón, L., J. C. Ochoa-Sánchez, M. Cruz, L. A. Villarreal-García, R. Rivera-Bustamante, and J. P. Martínez-Soriano  
2003. Caracterización molecular de aislamientos mexicanos del virus de la tristeza de los cítricos (CTV). *Proc. Pan Amer. Plant Dis. Conf., S. Padre Is. TX:* 21.
10. Hoffmann, A., and G. López-Campos  
2000. Biodiversidad de los ácaros en México. Ed. UNAM, CONABIO. México, D.F. 230 pp.
11. Iracheta-Cárdenas, M. M., I. H. Almeyda-León, K. L. Manjunath, B. Cevik, C. L. Niblett, R. F. Lee, and M. A. Rocha-Peña  
2002. Molecular cloning and expression of the p25 gene of two Mexican isolates of *Citrus tristeza virus*. *Rev. Mex. Fitopatol.* 20: 174-181.
12. Iracheta-Cárdenas, M. M., F. Orón-Castro, I. H. Almeyda-León, and M. A. Rocha-Peña  
2004. Ocurrencia de distribución y síntomas de psorosis en plantaciones comerciales de cítricos de Nuevo León, México. *Rev. Mex. Fitopatol.* 22: 299-307.
13. Loeza-Kuk, E.  
2003. Epidemiología y caracterización molecular de aislamientos mexicanos del citrus tristeza closterovirus. Tesis de Maestría en Ciencias. Colegio de Postgraduados, Montecillo, Texcoco, Estado de México. 85 p.
14. López-Arroyo, J. I.  
2002. Control biológico del pulgón café de los cítricos en México. Memorias Simposio Virus de la Tristeza de los Cítricos. XXV Congreso Nacional de la Sociedad Mexicana de Control Biológico. I. G. Romo-López, J. M. Avila-Salazar, L. J. Durón-Noriega, M. A. Rocha-Peña, and R. Báez-Sañudo (eds.), 33-43. Hermosillo, Sonora. México.
15. López-Arroyo, J. I., M. A. Peña del Río, M. A. Rocha-Peña, and J. Loera-Gallardo  
2005. Occurrence of the Asiatic citrus psyllid *Diaphorina citri* (Homoptera: Psyllidae) in Mexico. In: *Proc. 16th Conf. IOCV*, 508. IOCV, Riverside, CA.
16. Munguía-Rosales, R. A.  
2002. Experiencias en la liberación de agentes de control biológico del pulgón café de los cítricos en la Península de Yucatán. Control biológico del pulgón café *Toxoptera citricida*, vector del virus de la tristeza de los cítricos. H. C. Arredondo-Bernal, M. Medellín-Rosas, P. Pérez-Serrato, and J. P. Martínez-Soriano (eds.), 165-169. Dirección General de Sanidad Vegetal. SAGARPA. México.
17. Nava-Coronel, M., S. Silva-Vara, A. Morales-Loredo, V. J. Febres, C. L. Niblett, R. F. Lee, and M. A. Rocha-Peña  
1999. Caracterización serológica y molecular de algunos aislamientos del virus tristeza de los cítricos de Nuevo León, México. Resúmenes, XXVI Congreso Nacional de la Sociedad Mexicana de Fitopatología X Congreso Latinoamericano de Fitopatología. Guadalajara, Jalisco. México.
18. Orozco-Santos, M.  
1994. Principales especies de áfidos (Homoptera: Aphididae) en las áreas cítricas de México. Memorias, XXI Congreso Nacional de la Sociedad Mexicana de Fitopatología. Cuernavaca, Morelos. México.
19. Orozco-Santos, M.  
1995. Enfermedades presentes y potenciales de los cítricos en México. Universidad Autónoma Chapingo. México. 150 p.
20. Orozco-Santos, M. and R. González-Garza  
1985. Descripción del "amachamiento" de naranja Valencia (*Citrus sinensis* (L.) Osbeck) en el estado de Veracruz. *Rev. Mex. Fitopatol.* 3: 66-69.
21. Palmeri, M., I. Donis, A. L. Salazar, S. Blanco, M. Porres, R. H. Brlansky, A. S. Guerra-Moreno, K. L. Manjunath, and R. F. Lee  
2005. Leprosis in Guatemala. In: *Proc. 16th Conf. IOCV*, 510. IOCV, Riverside, CA.
22. Peña del Río, M. A.  
1987. Afidofauna asociada al cultivo de los cítricos en los estados de Tamaulipas, San Luis Potosí y Veracruz. Informe de Labores. CEGET-INIFAP-SARH. México (mimeo).
23. Permar, T. A., S. M. Garnsey, D. J. Gumpf, and R. F. Lee  
1990. A monoclonal antibody that discriminates strains of citrus tristeza virus. *Phytopathology* 80: 224-228.

24. Ramírez-Ch., J. L., and M. R. Guerrero  
1999. Monitoreo de *Toxoptera citricidus* Kirkaldy y otros áfidos vectores del VTC en huertos cítricos de Yucatán, México. Memorias, VIII Congreso Nacional Sociedad Mexicana de Ciencias Hortícolas, 185. Manzanillo, Colima, México.
25. Reanwarakorn, K., and J. S. Semancik  
1999. Correlation of hop stunt viroid variants to cachexia and xyloporosis diseases of citrus. *Phytopathology* 89: 568-574.
26. Robles-Serna, R.  
1985. Presencia de exocortis en los cítricos de Tamaulipas. Resúmenes XII Congreso Nacional de Fitopatología. Guanajuato, México.
27. Rocha-Peña, M. A., R. F. Lee, R. Lastra, C. L. Niblett, F. M. Ochoa-Corona, S. M. Garnsey, and R. K. Yokomi  
1995. Citrus tristeza virus and its aphid vector *Toxoptera citricida*: Threats to citrus production in the Caribbean and Central and North America. *Plant Dis.* 79: 437-445.
28. Rocha-Peña, M. A., J. P. Martínez-Soriano, and K. F. Byerly-Murphy  
1995. Viroides de cítricos: Amenaza oculta para la citricultura de México. *Rev. Mex. Fitopatol.* 13: 146-149.
29. Rocha-Peña, M. A., S. Silva-Vara, and J. P. Martínez-Soriano  
1996. Presencia del virus de la tristeza en plantaciones comerciales de cítricos del estado de Nuevo León, México. Memorias XXIII Congreso Nacional de Fitopatología. Guadalajara, Jalisco, México.
30. Saavedra de Domínguez, F., A. Bernal, C. C. Childers, and E. W. Kitajima  
2001. First report of *Citrus leprosis virus* in Panama. *Plant Dis.* 85: 228.
31. SAGAR  
1997. Plan de acción contra el virus tristeza de los cítricos y el pulgón café, *Toxoptera citricida*. Dirección General de Sanidad Vegetal. Comisión Nacional Agropecuaria. Secretaría de Agricultura, Ganadería y Desarrollo Rural. México (mimeo).
32. SAGARPA  
2004. Norma Oficial Mexicana NOM-031-FITO-2000, por la que se establece la campaña contra el virus tristeza de los cítricos. Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación. México.
33. Sánchez-Salas, J. A., R. Robles-Serna, and R. González-Garza  
1979. Ocurrencia de exocortis de los cítricos en el Estado de Nuevo León y su efecto en una combinación patrón-injerto susceptible. *Proc. Tropical Region Amer. Soc. Hort. Sci.* 23: 75-78.
34. Silva-Vara, S., M. A. Peña del Río, R. Peña-Martínez, N. Villegas-Jiménez, K. F. Byerly-Murphy, and M. A. Rocha-Peña  
2001. Distribución del virus de la tristeza en tres plantaciones comerciales de cítricos del estado de Nuevo León, México. *Agrociencia* 35: 441-450.
35. Uc-Vázquez, A., D. L. Ochoa-Martínez, E. Cárdenas-Soriano, G. Mora-Aguilera, and A. Villegas-Monte  
2001. Avances en el estudio etiológico del amarillamiento letal del limón Persa (*Citrus latifolia* Tan.) en Martínez de la Torre, Veracruz, México. Memorias XXVII Congreso Nacional de Fitopatología. Querétaro, Querétaro, México.
36. Vela-Mancilla, M. C.  
1984. Detección del virus de la tristeza de los cítricos en los bancos de germoplasma de cítricos en México mediante la técnica de inmunoadsorción ligada a enzima. Tesis de Químico Bacteriólogo Parasitólogo. Universidad Autónoma de Nuevo León. Monterrey, México.
37. Villarreal, L., L. Aguirre, and M. Cruz-M.  
1996. Detección del virus de la tristeza en México. Memorias XXIII Congreso Nacional de Fitopatología. Guadalajara, Jalisco, México.
38. Villarreal-G., L. A., A. M. Ramírez, and P. L. Robles  
2000. Campaña contra el virus tristeza de los cítricos en México. Memorias Séptima Reunión Anual del Consejo Nacional Consultivo Fitosanitario. Secretaría de Agricultura Ganadería y Desarrollo Rural. Comisión Nacional Agropecuaria. Puebla, Puebla, México.
39. Wutscher, H. K., A. Cervantes, and K. S. Derrick  
1997. Citrus blight found in México. *Plant Dis.* 81: 551.