

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

CARBON-10 AND MASS MEASUREMENTS FOR LIGHT NUCLEI

### Permalink

<https://escholarship.org/uc/item/0sz2p4r3>

### Authors

Brunnader, H.  
Hardy, J.C.  
Cerny, Joseph.

### Publication Date

1968-06-01

Cy. 2

# University of California Ernest O. Lawrence Radiation Laboratory

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

**CARBON-10 AND MASS MEASUREMENTS FOR LIGHT NUCLEI**

H. Brunnader, J. G. Hardy and Joseph Cerny

June 1968

**RECEIVED  
LAWRENCE  
RADIATION LABORATORY**

JUL 8 1968

**LIBRARY AND  
DOCUMENTS SECTION**

**Berkeley, California**

UCRL - 18272  
Cy. 2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

To be submitted to Phys. Rev.  
Section 53.1

UCRL-18272

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

CARBON-10 AND MASS MEASUREMENTS FOR LIGHT NUCLEI

H. Brunnader, J. C. Hardy and Joseph Cerny

June 1968

CARBON-10 AND MASS MEASUREMENTS FOR LIGHT NUCLEI \*

H. Brunnader, J. C. Hardy and Joseph Cerny

Lawrence Radiation Laboratory and  
Department of Chemistry  
University of California  
Berkeley, California

June 1968

ABSTRACT

The mass excess of  $^{10}\text{C}$  and the excitation energy of its first excited state have been measured using the  $^{10}\text{B}(^3\text{He}, t)^{10}\text{C}$  reaction. The results confirm one of three earlier measurements which gave the mass excess as  $(15.7025 \pm .0018)$  MeV, and an average of all relevant measurements for the energy of the first excited state yields  $(3.344 \pm .008)$  MeV. These values differ significantly from those previously accepted, and since they provided energy calibrations for measurements on a number of light nuclei, appropriately revised results are quoted for the mass excesses of  $^{21}\text{Mg}$  and  $^{37}\text{Ca}$ , as well as for the energies of the second excited state in  $^8\text{B}$  and  $T = 3/2$  states in  $^7\text{Be}$ ,  $^{21}\text{Na}$  and  $^{37}\text{K}$ .

## I. INTRODUCTION

Accurate values now exist for the excitation energies of the  $T = 2$  analogue states in  $^{20}\text{Ne}$  and  $^{24}\text{Mg}$ . These states have been observed as final states in  $(p,t)$  and  $(^3\text{He},n)$  reactions,<sup>1</sup> and as compound nuclear resonances appearing in proton induced reactions.<sup>2</sup> For both nuclei, the averages of values for these excitation energies compiled<sup>3</sup> from all available sources are accurate to better than  $\pm 5$  keV.

A recent investigation of analogue states throughout the  $(1d5/2)$  shell<sup>4</sup> included a remeasurement of the  $(p,t)$  and  $(p,^3\text{He})$  reactions on  $^{22}\text{Ne}$  and  $^{26}\text{Mg}$  targets in which carbon impurities had been added to provide an energy calibration. In particular, the ground and first excited states of  $^{10}\text{C}$ , produced in the reaction  $^{12}\text{C}(p,t)^{10}\text{C}$ , result in triton peaks whose energies bracket that of the  $T = 2$  states in  $^{20}\text{Ne}$  and  $^{24}\text{Mg}$ . Since the ground-state mass of  $^{10}\text{C}$  taken from the current mass table<sup>5</sup> was 15.658 MeV with a quoted error of  $\pm 13$  keV, it was therefore surprising to observe a discrepancy of  $\sim 45$  keV with the accepted energies of the  $T = 2$  states. This has led to the present re-evaluation of the mass of  $^{10}\text{C}$ , and the energy of its first excited state.

There have been three previous measurements<sup>6-8</sup> of the mass of  $^{10}\text{C}$ , and these are summarized in Table I. Only the first two results are included in the average value quoted in Ref. 5. Although the error bar on the third measurement is significantly smaller than the others, the fact that it differed from them by  $\sim 45$  keV, while using a similar experimental method, made it desirable to perform an independent measurement of a different type. For this purpose, the reaction  $^{10}\text{B}(^3\text{He},t)^{10}\text{C}$  was chosen.

## II. EXPERIMENTAL PROCEDURE

The experiment was carried out using the external beam of the Berkeley 88-inch spiral-ridge cyclotron. A diagram and general description of the experimental layout have been given elsewhere.<sup>9</sup> The beam was magnetically energy-analyzed by being deflected  $40^\circ$  and passed through a 1 mm slit. It was then focused to a spot  $2 \text{ mm} \times 2 \text{ mm}$  on a target at the center of the scattering chamber. The exact angle at which the beam intersected the target was measured using two luminous foils, one at the target position, and the other 70 cm downstream. The beam energy was inferred from measuring its range in aluminum.

Reaction products were detected in two independent counter telescopes on opposite sides of the beam. Each consisted of a  $150 \mu$  phosphorus-diffused silicon  $\Delta E$  transmission counter and a 3.0 mm lithium-drift silicon E counter operated in coincidence, followed by a  $500 \mu$  lithium-drift silicon E-reject counter operated in anticoincidence to eliminate long range particles. The signals from these telescopes were fed to Goulding-Landis particle identifiers,<sup>10</sup> producing output signals characteristic of the particle type, and the corresponding energy signals were routed accordingly into 1024-channel groups of a 4096-channel analyzer. In addition to the telescopes, two single 1 mm "monitor" counters were mounted at  $\theta_{\text{lab}} = +27.5^\circ$  and  $-18.0^\circ$ . The angular resolution of the telescopes and monitors was about  $0.25^\circ$ .

Figure 1 shows the energy spectrum of tritons observed from the ( $^3\text{He}, t$ ) reaction on a  $280 \mu\text{g}/\text{cm}^2$  boron target, 92% enriched in  $^{10}\text{B}$ . The  $^3\text{He}$  beam energy was measured to be  $(30.3 \pm 0.2) \text{ MeV}$ , and the laboratory angle, after correction for a measured "beam angle" of  $-0.4^\circ$ , was  $+22.7^\circ$ . It can be seen from the figure that the energy of the triton peak corresponding to the ground state of

$^{10}\text{C}$  is about 200 keV greater than that of the peak corresponding to the first excited state of  $^{11}\text{C}$  ( $E_x = 1.995 \pm .003$  MeV).<sup>11</sup> Similarly, the first excited state of  $^{10}\text{C}$  appears close to the second and third excited states of  $^{11}\text{C}$  ( $E_x = 4.305 \pm .006$  MeV and  $4.794 \pm .006$  MeV).<sup>11</sup> Kinematic calculations for peaks in this region show a change in the energy of  $^{10}\text{C}$  peaks relative to those from  $^{11}\text{C}$  of about 10 keV per degree, and thus, if the Q-values for the production of states in  $^{10}\text{C}$  are to be determined by comparison with known states in  $^{11}\text{C}$ , a precise knowledge of the laboratory angle must be maintained. Consequently, during this and all other runs, elastically scattered  $^3\text{He}$  particles were recorded in the fixed monitor counters. Prior measurements of the ( $^3\text{He}, ^3\text{He}$ ) cross section had shown variations of  $\sim 60\%$  and  $\sim 13\%$  in absolute magnitude per degree change in scattering angle at  $\theta_{\text{lab}} = 18.0^\circ$  and  $27.5^\circ$  respectively. For this reason, an integration of the elastically scattered particles, monitored frequently throughout a run, permitted a continuous sensitive determination (to about  $0.1^\circ$ ) of variations in the beam angle. No beam-angle excursions were detected during any of the experiments.

As a further precaution, in two runs spectra were recorded with counter telescope 1 set at the same nominal angle on opposite sides of the beam, while telescope 2 remained fixed at  $\theta_{\text{lab}} = -60.1^\circ$  to detect any significant beam energy change. It will be shown in the following section that the peak energies extracted from these runs were consistent with the independent determinations of the laboratory angle. Additional spectra using the enriched boron target were also taken at  $\theta_{\text{lab}} = +18.3^\circ$  and  $-55.2^\circ$ , and for comparison, a  $500 \mu\text{g}/\text{cm}^2$  natural boron target was bombarded at the same energy. A spectrum from the latter target, taken at  $\theta_{\text{lab}} = 22.7^\circ$ , is shown in Fig. 2.



## III. RESULTS AND DISCUSSION

The data were analyzed using the computer program LORNA.<sup>12</sup> This program establishes an energy scale by finding a least squares fit to peaks whose Q-values are known, after correcting all incoming and outgoing particles for kinematic effects and absorber losses. The states taken as known were the ground and first three excited states of  $^{11}\text{C}$ . Although the energy scale was determined by these states at all observed angles, the unknown states were evaluated independently at each angle, and the results for the  $^{10}\text{C}$  ground and first excited states are listed in Table II. The weighting for each individual measurement used in calculating the average was determined from the counting statistics.

A comparison of our final result in Table II with the measurements listed in Table I shows the former to be consistent with the value obtained in Ref. 8 and inconsistent with the other two. Thus we confirm the mass excess of  $^{10}\text{C}$  to be  $(15.7025 \pm .0018)$  MeV.

There have been three previous measurements<sup>13-15</sup> relating to the first excited state of  $^{10}\text{C}$ : two have determined its mass excess, and the other, its excitation energy. Using the correct ground state mass excess, good agreement is obtained among these measurements and the value derived in Table II. A summary of all experimental data on this state is given in Table III, together with the averaged result; this corresponds to an excitation energy of  $(3.344 \pm .008)$  MeV.

An earlier investigation<sup>13</sup> of the reaction  $^{10}\text{B}(^3\text{He}, t)^{10}\text{C}$  observed states in  $^{10}\text{C}$  at  $(5.28 \pm .06)$  and  $(6.58 \pm .06)$  MeV, and made a tentative identification of states at 5.03 and 5.60 MeV. An examination of all angles taken in this experiment confirms the existence of states at 5.28 and 6.58 MeV; however,

we see no indication of the other two states. In fact, it should be noted that at a bombarding energy of 30 MeV and  $\theta_{\text{lab}} = 30^\circ$  (the experimental conditions in Ref. 13) the peaks observed in the spectrum of Mangelson et al. would correspond exactly to states in  $^{11}\text{C}$  at 6.906 and 7.509 MeV. Similar peaks are observed in our spectrum of Fig. 1 and by comparison with Fig. 2 it is evident that they can be adequately accounted for as being entirely due to these states in  $^{11}\text{C}$ . Thus, the tentative assignment of states at 5.03 and 5.60 MeV in  $^{10}\text{C}$  should be dropped.

The earlier values for the masses of the ground and first excited states of  $^{10}\text{C}$  were used in determining the energies of analogue states and certain mass excesses for a number of light nuclei. Thus, the results of these measurements must be changed to take account of the revised  $^{10}\text{C}$  masses. Table 4 lists references,<sup>19,16</sup> previous values, and the revised excitation energies for those earlier measurements which were sufficiently precise to warrant adjustment. Not all values have been changed by as much as 45 keV since in most cases  $^{10}\text{C}$  was not the only source of calibration. Not listed in the table are the  $T = 2$  states in  $^{20}\text{Ne}$  and  $^{24}\text{Mg}$  whose measurement - as discussed earlier - first indicated that a discrepancy existed; using  $^{10}\text{C}$  for calibration the derived excitation energies of these states are within 10 keV of the accepted values, providing final confirmation of the present results.

One of us (JCH) gratefully acknowledges a fellowship from the Miller Institute for Basic Research in Science.

FOOTNOTE AND REFERENCES

\*Work performed under the auspices of the U.S. Atomic Energy Commission.

1. G. T. Garvey, J. Cerny, and R. H. Pehl, Phys. Rev. Letters 12, 726 (1964);  
J. Cerny, R. H. Pehl, and G. T. Garvey, Phys. Letters 12, 234 (1964);  
E. Adelberger and A. B. McDonald, Phys. Letters 24B, 270 (1967).
2. H. M. Kuan, D. W. Heikkinen, K. A. Snover, F. Riess, and S. S. Hanna,  
Phys. Letters 25B, 217 (1967);  
R. Bloch, R. E. Pixley, and P. Truöbl, Phys. Letters 25B, 215 (1967);  
F. Riess, W. J. O'Connell, D. W. Heikkinen, H. M. Kuan, and S. S. Hanna,  
Phys. Rev. Letters 19, 367 (1967).
3. J. Cerny, Ann. Rev. Nucl. Sci., to be published.
4. J. C. Hardy, H. Brunnader, and J. Cerny, Bull. Am. Phys. Soc. 13, 561 (1968);  
H. Brunnader, J. C. Hardy, J. Cerny and J. Jänecke, to be published.
5. J. H. E. Mattauch, W. Thiele, and A. H. Wapstra, Nucl. Phys. 67, 1 (1965);  
C. C. Maples, G. W. Goth, and J. Cerny, Nuclear Data 2A, 429 (1966).
6. S. Takayanagi, N. H. Gale, J. B. Garg, and J. M. Calvert, Nucl. Phys. 28,  
494 (1961).
7. F. J. Bartis, Phys. Rev. 132, 1763 (1963).
8. J. M. Freeman, J. G. Jenkins, and G. Murray, Phys. Letters 22, 177 (1966);
9. G. W. Butler, J. Cerny, S. W. Cosper, and R. L. McGrath, Phys. Rev. 166,  
1096 (1968).
10. F. S. Goulding, D. A. Landis, J. Cerny and R. H. Pehl, Nucl. Instr. Methods,  
31, 1 (1964).
11. F. Ajzenberg-Selove and T. Lauritzen, Energy Levels of Light Nuclei (VII),  
to be published.

12. LORNA is a program written by C. C. Maples, whom we thank for making it available.
13. N. Mangelson, F. Ajzenberg-Selove, M. Reed, and C. C. Lu, Nucl. Phys. 88, 137 (1966).
14. L. G. Earwaker, J. G. Jenkins, and E. W. Titterton, Nucl. Phys. 42, 521 (1963).
15. R. E. Segel, P. P. Singh, S. S. Hanna, and M. A. Grace, Phys. Rev. 145, 736 (1966).
16. R. L. McGrath, J. Cerny, and E. Norbeck, Phys. Rev. Letters 19, 1442 (1967).

Table I. Previous measurements of the mass of  $^{10}\text{C}$ .

Experiment	Ref.	Mass excess (MeV)
(p,n) threshold	6	15.663 $\pm$ .025
$\beta^+$ end point	7	15.656 $\pm$ .015
(p,n) threshold	8	15.7025 $\pm$ .0018

Table II. Summary of results from the  $^{10}\text{B}(^3\text{He},t)^{10}\text{C}$  reaction measuring the ground and first excited states of  $^{10}\text{C}$ .

Run No.	Telescope 1			Telescope 2		
	$\theta_{\text{lab}}$	Mass excess for:		$\theta_{\text{lab}}$	Mass excess for:	
		ground state (MeV)	1 <sup>st</sup> excited (MeV)		ground state (MeV)	1 <sup>st</sup> excited (MeV)
1	+22.7	15.695	19.041	-60.1	15.708	19.052
2	-21.9	15.702	19.045	-60.1	15.704	19.048
3	+22.7	15.701	19.036	-60.1	15.703	19.054
4	+18.3	15.692	19.039	-55.2	15.700	19.036
Weighted overall averages:		$^{10}\text{C}$ ground state		(+15.700 $\pm$ .010) MeV		
		$^{10}\text{C}$ 1 <sup>st</sup> excited state		(+19.042 $\pm$ .015) MeV		

Table III. Summary of experimental data on the first excited state of  $^{10}\text{C}$ .

Experiment	Ref.	Mass excess (MeV)
$^{10}\text{B}(^3\text{He},\text{t})^{10}\text{C}$	13	$19.028 \pm .030$
(p,n) threshold	14	$19.039 \pm .020$
$\gamma$ decay	15	$(19.053 \pm .010)^\dagger$
$^{10}\text{B}(^3\text{He},\text{t})^{10}\text{C}$	this work	$19.042 \pm .015$
Average		$19.047 \pm .008$

$^\dagger$  This measurement was of the excitation energy rather than the mass excess; it has been converted using the best value for the ground-state mass excess of  $^{10}\text{C}$  (15.703 MeV).


Table IV. Original and revised experimental values which depend for calibration upon the ground and first excited states of  $^{10}\text{C}$ .

Reaction	Ref.	Measured	Previous value (MeV)	New value (MeV)
$^{24}\text{Mg}(^3\text{He}, ^6\text{He})^{21}\text{Mg}$ g.s.	9	Mass excess	$+10.62 \pm 0.12$	$+10.66 \pm 0.12$
$^{40}\text{Ca}(^3\text{He}, ^6\text{He})^{37}\text{Ca}$ g.s.	9	Mass excess	$-13.24 \pm 0.05$	$-13.23 \pm 0.05$
$^9\text{Be}(p,\text{t})^7\text{Be}^*$ ( $T=3/2$ )	16	excitation	$10.97 \pm 0.04$	$11.01 \pm 0.04$
$^{10}\text{B}(p,\text{t})^8\text{B}^*$ (2nd excited)	16	excitation	$2.33 \pm 0.04$	$2.34 \pm 0.04$
$^{23}\text{Na}(p,\text{t})^{21}\text{Na}^*$ ( $T=3/2$ )	9	excitation	$8.92 \pm 0.03$	$8.97 \pm 0.03$
$^{39}\text{K}(p,\text{t})^{37}\text{K}^*$ ( $T=3/2$ )	9	excitation	$5.035 \pm 0.025$	$5.046 \pm 0.025$

FIGURE CAPTIONS

Fig. 1. A triton energy spectrum obtained at  $\theta_{\text{lab}} = 22.7^\circ$ , from a boron target 92% enriched in  $^{10}\text{B}$ .

Fig. 2. A triton energy spectrum obtained from a natural boron target at  $\theta_{\text{lab}} = 22.7^\circ$ .



XBL684-2298

Fig. 1






Fig. 2

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

