UC San Diego

UC San Diego Previously Published Works

Title

Honey robbing: could human changes to the environment transform a rare foraging tactic into a maladaptive behavior?

Permalink

<https://escholarship.org/uc/item/0v38c46p>

Authors

Rittschof, Clare C

Nieh, James C

Publication Date

2021-06-01

DOI

10.1016/j.cois.2021.02.005

Peer reviewed

Journal Pre-proof

Honey robbing: could human alterations to the environment change a rare foraging tactic into a maladaptive behavior?

Clare C Rittschof, James C Nieh

PII: S2214-5745(21)00013-4

DOI: <https://doi.org/10.1016/j.cois.2021.02.005>

Reference: COIS 784

To appear in: *Current Opinion in Insect Science*

Please cite this article as: Rittschof CC, Nieh JC, Honey robbing: could human alterations to the environment change a rare foraging tactic into a maladaptive behavior?, *Current Opinion in Insect Science* (2021), doi: <https://doi.org/10.1016/j.cois.2021.02.005>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Honey robbing: could human alterations to the environment change a rare foraging tactic into a maladaptive behavior?

Clare C. RITTSCHOF^{a*} & James C. NIEH^b

a. University of Kentucky, Department of Entomology
S-225 Agriculture Science Center North
Lexington, KY, 40546
*corresponding author: clare.rittschhof@uky.edu

b. University of California, San Diego, Division of Biological Sciences
9500 Gilman Drive, MC 0116
La Jolla, CA 92093-0116
jnieh@ucsd.edu

Highlights

- Human changes to the landscape may promote an extreme honey bee foraging tactic, honey robbing.
- Honey robbing is a complex behavioral syndrome with substantial health risks.
- Robbing is most common in the managed species *Apis mellifera*, perhaps because of human practices.
- Hybrid optimal foraging and game theory models may be useful for understanding robbing.
- The ecological and physiological factors underlying robbing require better elucidation.

Abstract

Human environmental modifications have outpaced honey bees' ability to evolve adaptive regulation of foraging tactics, possibly including a tactic associated with extreme food shortage, honey robbing. Honey robbing is a high risk, high reward, and understudied honey bee tactic whereby workers attack and often kill neighboring colonies to steal honey. Humans have exacerbated the conditions that provoke such robbing and its consequences. We describe robbing as an individual- and colony-level behavioral syndrome, implicating worker bees specialized for foraging, food processing, and defense. We discuss how colony signaling mechanisms could regulate this syndrome and then explore the ecological underpinnings of robbing—highlighting its unusual prevalence in the commonly managed *Apis mellifera* and outlining the conditions that provoke robbing. We advocate for studies that identify the cues that modulate this robbing syndrome. Additionally, studies that apply behavioral ecology modeling approaches to generate testable predictions about robbing could clarify basic bee biology and have practical implications for colony management.

Keywords: commercial beekeeping; optimal foraging theory; energetic constraints; dance language; aggression

Human agricultural practices impact honey bee foraging tactics like robbing

Human agricultural practices have diverse and complex implications for honey bee behavior, health, and beekeeping costs. Colony management, presently a requirement for successful pollination of many cash crops worldwide, is itself a source of honey bee stress. For example, during crop bloom in intensive agricultural landscapes, honey bees are densely packed into small areas, leading to nutritional stress from competition and lack of floral diversity [1]. The world-wide transport of *A. mellifera* has introduced novel pathogens and invasive parasites, most notably the *Varroa* mite, which has decimated colony survivorship and productivity in most regions where it has taken hold. In addition to beekeeping practices, agricultural land use generally has large effects on bee nutrition. Crop bloom results in boom-and-bust periods of resource availability, and honey bee colony health strongly tracks these feast-famine cycles [2]. In the U.S.A., a general period of nectar dearth arises in the mid-late summer, in part from the loss of native flowering prairie lands that naturally bloom at this time [3]. The replacement of natural floral communities with crop and grazing lands can increase competition for food among bee species, or alternatively, decrease competition by eliminating natural refuges and wild bees altogether [4]. Changing climate and weather patterns may further limit nectar abundance [5], harming bee populations [6]. Honey bees therefore face diverse stressors in human agricultural environments, including the complex and significant effects of floral resource limitation.

Honey bees forage for kilometers across the landscape, and they have sophisticated adaptations that allow them to take advantage of sparse, ephemeral floral resources [7]. During extreme resource scarcity, they can deploy a comparably extreme foraging tactic known as honey robbing. Robbing colonies invade neighboring colonies and collect honey stores, sometimes numbering in the kilograms. Because honey is processed, concentrated nectar collected from millions of different flowers, bee colonies provide a large, diverse, and valuable food resource. However, robbing is relatively rare under natural conditions because it is a high-risk tactic in which foragers attack, fight, and kill members of the “victim” colony [8]. This places unique demands on the robbing colony, which must mount an offense, but also collect and store an unusually large quantity of food. Moreover, victim colonies can be infected with transmissible diseases and parasites, weakening them and reducing their defensive behavior, which simultaneously makes them more attractive and more costly robbing targets [9-13]. Because robbing can result in widespread colony mortality, it is understudied, and it remains unclear whether or not colonies adaptively deploy this foraging tactic under the extreme conditions presented by agricultural practices. For example, foraging adaptations like nestmate recruitment using the waggle dance may actually work against colony productivity in extreme conditions of resource scarcity [14]. Thus, there is evidence for maladaptive expression of colony level foraging tactics in highly modified environments, and honey robbing, which integrates nutritional stress with other disease and mortality costs, may also be maladaptive.

Here, we provide an overview of robbing in honey bees, focusing on *Apis mellifera*, the commonly managed western honey bee. We describe honey robbing as a complex behavioral syndrome at individual and colony levels, and we discuss how well-established honey bee communication mechanisms may operate in a robbing context. We explore the scope of this foraging tactic in other *Apis* species. Finally, we discuss the ecological contexts that promote robbing, and consider how humans have altered robbing risks and

benefits. Robbing may be a sophisticated colony-level foraging tactic that becomes dysregulated under human-altered conditions. As a result, robbing may be triggered in maladaptive scenarios, ultimately increasing colony losses.

Honey robbing as a behavioral syndrome

In social insects, behavioral syndromes (suites of correlated behavioral traits), occur at both individual and colony levels. This behavioral organization is particularly complex in honey bees because of their large colony sizes (20,000 to 60,000 worker bees). Colony foraging regulation is sophisticated: tactics change in response to colony demography, size, and nutritional deficits [15, 16], the quality, quantity, and proximity of available resources [17, 18], and abiotic factors including temperature [19]. To adjust food collection, colonies can deploy different ratios of nectar and pollen foraging specialists [20], and individuals can adjust their foraging trip frequency, duration, and distance, as well as their use of social information about resource quality and location (the famous dance language) [21], which is likely also involved in robbing.

At the individual level, little is known about traits that characterize robbing foragers. Foragers are divided into individuals that specialize on resin, nectar, pollen, or water collection. The latter three specializations are correlated with sensitivity to sugar concentration in a logical manner (nectar, pollen, and water foragers have respectively low, medium, and high sensitivities, [22]). This logic implies that robbing bees could have unusually low sensitivity and thus high sugar response thresholds. Robbers are unlikely to be a distinct foraging specialization given the seasonal and opportunistic nature of the tactic. However, there is some evidence they possess correlated traits. For example, even when kept safely in the lab, robbers have decreased longevity compared to non-robbing foragers, a pattern that lessens the potential costs to the riskier tactic [10]. There is also some evidence that robbing foragers, even without engaging in fights, are unusually aggressive [23], a potentially adaptive trait since aggression may be required to overwhelm victim colony defenses.

At the colony level, robbing is characterized by a conspicuous increase in foraging activity, requiring a concomitant increase in food storing abilities. Nectar foragers do not directly store their nectar load, but instead transfer it to receiver bees. The length of time a forager waits for a receiver affects the speed at which that forager leaves for another trip. During robbing, the rapid increase in foraging trip rate [23] likely requires rapid, increased recruitment of receiver bees [24], and possibly a change in the feedback mechanism that regulates foraging activity (discussed below). The robbing colony's defensive specialists also have been implicated in the syndrome [23]. Guard bees are specialists who defend against invading robber honey bees. Guards smell odors on incoming foragers and chase away non-nestmates [25]. Unsurprisingly, experiencing robber intrusion results in increased victim colony guard numbers and higher rates of non-nestmate rejection [26, 27]. However, guards from robbing colonies that have not experienced such intrusions can also elevate their defensiveness, even towards their own foragers [23]. The causes and consequences of this shift are largely unknown, although it suggests a correlation between robbing activity and invasion risk (Fig 1).

Colony signaling mechanisms and the regulation of robbing

Honey bees use social information and sophisticated communication mechanisms to modulate foraging behavior. Foragers perform waggle dances to signal the presence of good food resources [21], likely including a nest that can be robbed. In fact, the classic artificial sugar feeders used by researchers to decode the characteristics of the waggle dance more closely simulate a nest than a floral resource because flowers do not offer concentrated nectar *ad libitum*. Artificial feeders can provide very large quantities of high-concentration sugar solution that may simulate nest robbing [28]. They also elicit guarding and patrolling (robbing behaviors), and attract hundreds of foragers to a single location [29]. Managed honey bee colonies are routinely fed during times of food dearth with artificial feeders containing sugar solutions, a human manipulation that may increase robbing, particularly given the lack of other food sources.

Using rich “robbable” feeders placed outside the nest, researchers have identified two signals, tremble dancing and stop signaling, that regulate food collection and storage processes relevant to robbing. During a massive increase in nectar inflow (which could occur during robbing), there are insufficient receiver bees to collect food from incoming foragers. Forager wait time increases, and in response, they shift from waggle dancing to tremble dancing [30]. Tremble dancing may stimulate bees to engage in food receiving because the number of receivers increases after tremble dancing rises. This regulation is elegant since foragers cease recruiting new foragers to a food resource when the colony is unable to process it. New receiver recruitment can happen rapidly [31], but it is unknown whether this typical mechanism is sufficient to accommodate the influx of honey during robbing.

Robbing could trigger an additional signaling mechanism associated with resource competition. When foragers overwhelm an artificial feeder, they experience overcrowding [32] and fight with foragers from competing colonies [28]. In this situation, foragers return to the nest and produce brief vibratory pulses called stop signals, directed at other waggle dancers recruiting for the same overwhelmed resource. These signals cause foragers to stop dancing [28]. Unlike the passive process by which tremble dancing supplants waggle dancing, the stop signal provides negative feedback to counter the positive recruitment effects of waggle dancing. The result is more rapidly reduced recruitment and a faster foraging behavioral shift at the colony level.

Different signals may allow colonies to weigh robbing risks and benefits during its incipient stages and abandon situations that are overly risky. For example, foragers that are not attacked at an overwhelmed feeder continue to waggle dance and recruit, while individuals who are attacked at a rich feeder (as could occur during the initial stages of robbing when a victim colony defends itself) produce stop signals and tremble dance [28]. This case suggests a broader role for the tremble dance as a catalyst for reallocating colony labor [33]. Agent-based modeling of robbing suggests that stop signaling allows a colony to more efficiently stop robbing when the robbed colony can mount an effective defense [34]. Thus, stop signals can theoretically inform a conservative robbing decision-making process. Such a process predicts that colonies preferentially attack weak colonies (which is supported by empirical evidence) and quickly call off attacks when strong colonies mount an effective defense [34], as supported by evidence of abandoned robbing attempts [8].

Robbing in different honey bee species.

The existence of the defensive guard specialist whose effort is directly modulated by conspecific robber intrusion suggests that robbing occurs outside of the managed colony context (i.e., between non-managed *Apis mellifera* colonies). However, among the different *Apis* species, the most common observations of robbing come from apiaries, illustrating a potential impact of human manipulation. Colony management conditions may simultaneously increase passive worker drift and provoke non-nestmate aggression when resources become scarce, or when a robbing target presents itself. In *A. mellifera*, for example, managed colonies rob both within and among apiaries [12], and based on patterns of disease transmission, evidence of density-dependent robbing exists at the landscape, not just apiary, scale [35].

All *Apis* species create nests with highly concentrated resources that are attractive to interspecifics [36]. *Apis florea* [37], an open-nesting species, can rob *A. mellifera* colonies, and, surprisingly, given the generally smaller sizes of *A. florea* colonies and workers, can overcome *A. mellifera* [38]. *Apis cerana* (a cavity nesting species) is robbed by *A. mellifera* and vice versa, with *A. mellifera* possessing the competitive advantage [39]. Similarly, the open nesting *A. dorsata* [37] can rob *A. cerana* [39], and *A. cerana* and *A. koshevnikovi* (both cavity nesters) have been observed landing on *A. dorsata* nests to rob them [11].

Although robbing between species is broadly supported, it is less clear how common intraspecific robbing is for feral and wild honey bees. Compared to *A. mellifera*, there is little evidence of robbing, and surprisingly little intraspecific aggression in *A. florea*, *A. andreniformis* (an open nester), *A. dorsata*, and *A. cerana* [40]. However, *A. cerana* robbing and fighting was observed in managed colonies during honey harvesting [40], just as with *A. mellifera*. Robbing observations may be more frequent in apiaries because they are closely monitored, but the association with honey harvesting suggests that increased colony densities and honey availability due to management practices could increase robbing.

Apis dorsata nest in clusters that could facilitate robbing. However, genetic analyses found only about 1.4% worker drift (defined as passive movement of workers to a non-natal colony) between colonies [41], suggesting adaptations to prevent drift, non-nestmate intrusion, or both. *A. dorsata* are relatively docile towards conspecifics, but they do become more defensive towards non-nestmates during the migratory season when a new nest has settled in their vicinity. Actual robbing is still rarely observed [42]. Interestingly, managed *A. mellifera* colonies kept at a similar density show a drift rate that is 35-fold higher. *Apis mellifera* colonies naturally occur at much lower densities than the density of colonies found in most apiaries, which may have weakened natural selection against worker drift [43].

In addition to passive drift, robbing is a key mechanism for mite and virus transfer among colonies [12]. The prevalence of both stressors are correlated with landscape colony density, suggesting interactions among colonies from widely spaced apiaries. Forfert et al. [35] found a positive correlation between density and pathogen prevalence, comparing a minimum density of 0.32 colonies per km² to a maximum density of 1.4 colonies per km². Honey bee colonies are often managed at densities well above this maximum. In general, species variation in robbing could result from differences in the

quantity of stored resources, nesting habits, dissimilarities in the risks and benefits of intraspecific aggression, management practices, or a combination of these factors.

Ecological conditions that provoke honey robbing

Floral resource limitation, specifically nectar scarcity, seems to be the main ecological condition that promotes robbing and can be exacerbated by a lack of natural floral resources in landscapes altered by humans. Nectar scarcity can be assessed by measuring visitation to sugar feeders placed in the vicinity of honey bee colonies. As floral resources dwindle, bees visit and collect food from feeders with progressively lower sugar concentrations [27, 44, 45]. Although there is substantial landscape-level variation in floral resource abundance, in most moist temperate areas, nectar scarcity shows a predictable seasonal pattern that tracks robbing activity. Floral resources decline substantially in late summer [46]. Simultaneously, other factors compound the effects of food shortage, including increased competition from other bee species at their peak population abundance [47], heightened pressure to accumulate sufficient honey for the winter [48, 49], and declining temperatures, which make nectar processing more energetically costly [50]. Further studies across a range of climate regions are needed to determine how robbing risk is predicted by any one of these seasonal changes. For example, in parts of Florida dominated by citrus, robbing risk increases after the end of the citrus bloom, even though this decline in floral resources occurs in early spring, removing the other seasonal pressures associated with the approach of winter [27].

Conclusions

Robbing is a sophisticated foraging tactic that carries great risks to honey bee colonies in human altered landscapes. Improved understanding of the mechanisms underlying robbing are needed to understand its impacts and regulation. For example, the cues driving the simultaneous increase in foraging activity and aggression during robbing are unclear [23]. Guards could be confused by changes in nestmate odors that result from bees entering another nest. Guards may also respond to the heightened aggression displayed by robbing foragers, even though they are nestmates. Minor intrusion attempts by neighboring colonies could also cause these changes [26], and eventually escalate to robbing. If so, the use of migratory colonies for pollination could exacerbate robbing when colonies are set out closely spaced and bees begin to orient themselves in a new location. Other costs like pathogen and parasite transmission may be higher in high density conditions, and the impact of robbing in these contexts remains to be considered [51, 52]. For beekeepers, the costs associated with robbing-related mortality and disease risk are compensated, to a degree, by the financial gains from pollination services. The state of this fragile balance could substantially impact the future of commercial beekeeping [53, 54].

If colonies use floral resource abundance or other seasonal cues to anticipate robbing threats, they may proactively shift their defensiveness over the season. Food scarcity and defensive behaviors are correlated and peak in the fall when robbing is common, even without evidence of an acute robbing event [44]. Interestingly, brain titers of aggression-related biogenic amines, serotonin and dopamine [55, 56], peak during

August and September [57] when robbing peaks, but it not known if this variation arises from abiotic cues or robbing experience. Simultaneous expression of two high energy behaviors, aggression and foraging activity is difficult to sustain, suggesting robbing has unknown colony level energetic consequences that may manifest over time. For individual robbers, it is unclear how foragers accommodate an increase in aggression, as these two phenotypes are typically mutually exclusive, including at the level of neural regulation [58, 59]. Future work should address how the colony level features of the robbing syndrome are regulated by existing signals, and how robbing related behaviors are regulated at the individual level.

Determining the physiological and ecological mechanisms that modulate robbing (Fig. 1) should improve our understanding of robbing and its consequences. For example, the cost of fighting over rich food sources is not necessarily, as might be expected, a mortality cost, but rather a cost of lost time. In the case of two stingless bee species, time spent fighting for an unwinnable resource takes away from time spent foraging for other resources, and thus mechanisms have evolved that allow species to avoid time-wasting combat [60]. A similar decision may be made by robbing honey bees. Is it better to rob or to forage for floral resources? It may also be illuminating to integrate optimal foraging theory with game theory, an approach commonly used in studies of facultative kleptoparasitism (selective stealing of food resources [61]). Such modeling suggests that food stealing is affected by multiple factors, including fighting, but it is particularly influenced by population density (in our case, the number of colonies in each area, [35]), a factor that could be adjusted during agricultural pollination. In addition, decision analysis modeling could be used to better understand factors, such as the time costs and the probability of winning a rich resource, that a robbing or aggressive colony weighs [60]. In general, such modeling would be useful for understanding how collectives adapt to dynamic ecological conditions. Robbing therefore provides a case-study in the precariousness of social coordination in the face of accumulating stressors, particularly those exacerbated by human actions. In addition to its direct impacts on colony health, human intervention may result in maladaptive foraging tactics that lead to colony losses.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture Hatch Program (accession number 1012993, C.C.R.), and the Foundation for Food and Agriculture Research Pollinator Health Fund (549049, C.C.R.).

Annotated references

Grume, G.J., S. Biedenbender, and C.C. Rittschof, *Honey robbing causes coordinated changes in foraging and nest defense in the honey bee, *Apis mellifera**. Animal Behaviour, 2020. **in press**.

*Empirically describes the simultaneous increase in aggression and foraging that characterizes a robbing event, provides empirical methodology to induce controlled robbing.

Peck, D.T. and T.D. Seeley, *Mite bombs or robber lures? The roles of drifting and robbing in Varroa destructor transmission from collapsing honey bee colonies to their neighbors*. PLoS One, 2019. **14**(6): p. e0218392.

*Provides evidence that robbing is a key means to transmit invasive *Varroa* mites and disease from dead and dying colonies.

Garbuzov, M., N.J. Balfour, K. Shackleton, H. Al Toufalia, L. Scandian, and F.L.W. Ratnieks, *Multiple methods of assessing nectar foraging conditions indicate peak foraging difficulty in late season*. Insect Conservation and Diversity, 2020.

*Shows correlated seasonal changes in visitation to low-value sugar feeders and defensive effort. Either metric could predict late summer declines in floral resources.

Garay, J., R. Cressman, F. Xu, M. Broom, V. Csiszar, and T.F. Mori, *When optimal foragers meet in a game theoretical conflict: a model of kleptoparasitism*. Journal of Theoretical Biology, 2020. **502**.

*Presents an approach to combine optimal foraging theory and evolutionary game theory in the study of food stealing behavior.

Dolezal, A.G., A.L. St Clair, G. Zhang, A.L. Toth, and M.E. O'Neal, *Native habitat mitigates feast-famine conditions faced by honey bees in an agricultural landscape*. Proc Natl Acad Sci U S A, 2019. **116**(50): p. 25147-25155.

*Shows that relocation to natural habitats during the late summer improves colony productivity and health.

Nieh, J.C., *A negative feedback signal that is triggered by peril curbs honey bee recruitment*. Current Biology, 2010. **20**: p. 310-315.

*Competition at food resource can elicit robbing-like behaviors and leads to stop signaling, which can halt recruitment in the losing colony for the contested resource.

l'Anson Price, R., N. Dulex, N. Vial, C. Vincent, and C. Gruter, *Honeybees forage more successfully without the "dance language" in challenging environments*. Science Advances, 2019. **5**: p. 1-9.

*Disrupting the waggle dance improved colony growth because foragers saved time by relying on private information.

References

[1] Smart, M., J. Pettis, N. Rice, Z. Browning, and M. Spivak, *Linking Measures of Colony and Individual Honey Bee Health to Survival among Apiaries Exposed to Varying Agricultural Land Use*. PLoS One, 2016. **11**(3): p. e0152685.

[2] Dolezal, A.G., A.L. St Clair, G. Zhang, A.L. Toth, and M.E. O'Neal, *Native habitat mitigates feast-famine conditions faced by honey bees in an agricultural landscape*. Proc Natl Acad Sci U S A, 2019. **116**(50): p. 25147-25155.

- [3] Carr-Markell, M.K., C.M. Demler, M.J. Couvillon, R. Schurch, and M. Spivak, *Do honey bee (Apis mellifera) foragers recruit their nestmates to native forbs in reconstructed prairie habitats?* PLoS One, 2020. **15**(2): p. e0228169.
- [4] Wojcik, V.A., L.A. Morandin, L. Davies Adams, and K.E. Rourke, *Floral Resource Competition Between Honey Bees and Wild Bees: Is There Clear Evidence and Can We Guide Management and Conservation?* Environ Entomol, 2018. **47**(4): p. 822-833.
- [5] Fernie, A.R., L. Perez de Souza, M. Borghi, and T. Yoshida, *Flowers and climate change: a metabolic perspective.* New Phytologist, 2019. **224**: p. 1425-1441.
- [6] Flores, J.M., S. Gil-Lebrero, V. Gamiz, M.I. Rodriguez, M.A. Ortiz, and F.J. Quiles, *Effect of the climate change on honey bee colonies in a temperate Mediterranean zone assessed through remote hive weight monitoring system in conjunction with exhaustive colonies assessment.* Sci Total Environ, 2019. **653**: p. 1111-1119.
- [7] Nurnberger, F., A. Keller, S. Hartel, and I. Steffan-Dewenter, *Honey bee waggle dance communication increases diversity of pollen diets in intensively managed agricultural landscapes.* Mol Ecol, 2019. **28**(15): p. 3602-3611.
- [8] Free, J.B., *The behaviour of robber honeybees.* Behaviour, 1954. **7**(2/3): p. 233-240.
- [9] Geffre, A.C., T. Gernat, G.P. Harwood, B.M. Jones, D. Morselli Gysi, A.R. Hamilton, B.C. Bonning, A.L. Toth, G.E. Robinson, and A.G. Dolezal, *Honey bee virus causes context-dependent changes in host social behavior.* Proc Natl Acad Sci U S A, 2020. **117**(19): p. 10406-10413.
- [10] Kuszewska, K. and M. Woyciechowski, *Risky robbing is a job for short-lived and infected worker honeybees.* Apidologie, 2014. **45**(5): p. 537-544.
- [11] Koeniger, G., N. Koeniger, D.L. Anderson, C. Lekprayoon, and S. Tingek, *Mites from debris and sealed brood cells of Apis dorsata colonies in Sabah (Borneo) Malaysia, including a new haplotype of Varroa jacobsoni.* Apidologie, 2002. **33**: p. 15-24.
- [12] Peck, D.T. and T.D. Seeley, *Mite bombs or robber lures? The roles of drifting and robbing in Varroa destructor transmission from collapsing honey bee colonies to their neighbors.* PLoS One, 2019. **14**(6): p. e0218392.
- [13] Lindström, A., S. Korpela, and I. Fries, *Horizontal transmission of Paenibacillus larvaesporos between honey bee (Apis mellifera) colonies through robbing.* Apidologie, 2008. **39**(5): p. 515-522.
- [14] l'Anson Price, R., N. Dulex, N. Vial, C. Vincent, and C. Gruter, *Honeybees forage more successfully without the "dance language" in challenging environments.* Science Advances, 2019. **5**: p. 1-9.
- [15] Schmickl, T. and I. Karsai, *Integral feedback control is at the core of task allocation and resilience of insect societies.* Proc Natl Acad Sci U S A, 2018. **115**(52): p. 13180-13185.
- [16] Beekman, M., D.J. Sumpter, N. Seraphides, and F.L. Ratnieks, *Comparing foraging behaviour of small and large honey bee colonies by decoding waggle dances made by foragers.* Functional Ecology, 2004. **18**: p. 829-835.
- [17] Van Nest, B.N., A.E. Wagner, C.N. Hobbs, and D. Moore, *Dance floor clustering: food-anticipatory behavior in persistent and reticent honey bee foragers.* Behavioral Ecology and Sociobiology, 2016. **70**(11): p. 1961-1973.
- [18] Seeley, T.D., *Honey bee colonies are group-level adaptive units.* American Naturalist, 1997. **150**(S1): p. s22-s41.

- [19] Stabentheiner, A. and H. Kovac, *Honeybee economics: optimisation of foraging in a variable world*. Sci Rep, 2016. **6**: p. 28339.
- [20] Hendriksma, H.P., A.L. Toth, and S. Shafir, *Individual and Colony Level Foraging Decisions of Bumble Bees and Honey Bees in Relation to Balancing of Nutrient Needs*. Frontiers in Ecology and Evolution, 2019. **7**.
- [21] Frisch, v., K., *The dance language and orientation of bees*. Second ed. 1967, Cambridge, Massachusetts: Belknap Press.
- [22] Pankiw, T. and R.E. Page, *The effect of genotype, age, sex, and caste on response thresholds to sucrose and foraging behavior of honey bees (Apis mellifera L.)*. J Comparative Physiology A, 1999. **185**: p. 2017-213.
- [23] Grume, G.J., S. Biedenbender, and C.C. Rittschof, *Honey robbing causes coordinated changes in foraging and nest defense in the honey bee, Apis mellifera*. Animal Behaviour, 2020. **in press**.
- [24] Seeley, T.D., *Social foraging in honey bees: how nectar foragers assess their colony's nutritional status*. Behavioral Ecology and Sociobiology, 1989. **24**: p. 181-199.
- [25] Breed, M.D., M.F. Garry, A.N. Pearce, B.E. Hibbard, L.B. Bjostad, and R.E. Page, *The role of wax comb in honey bee nestmate recognition*. Animal Behaviour, 1995. **50**: p. 489-496.
- [26] Couvillon, M.J., E.J.H. Robinson, B. Atkinson, L. Child, K.R. Dent, and F.L.W. Ratnieks, *En garde: rapid shifts in honeybee, Apis mellifera, guarding behaviour are triggered by onslaught of conspecific intruders*. Animal Behaviour, 2008. **76**(5): p. 1653-1658.
- [27] Downs, S.G. and F.L. Ratnieks, *Adaptive shifts in honey bee (Apis mellifera L.) guarding behavior support predictions of the acceptance threshold model*. Behavioral Ecology, 2000. **11**(3): p. 326-333.
- [28] Nieh, J.C., *A negative feedback signal that is triggered by peril curbs honey bee recruitment*. Current Biology, 2010. **20**: p. 310-315.
- [29] Roubik, D.W., *Ecology and natural history of tropical bees*. 1989, New York: Cambridge University Press.
- [30] Seeley, T.D., *The tremble dance of the honey bee: message and meanings*. Behavioral Ecology and Sociobiology, 1992. **31**: p. 375-383.
- [31] Seeley, T.D., S. Kuhnholz, and A. Weidenmuller, *The honey bee's tremble dance stimulates additional bees to function as nectar receivers*. Behavioral Ecology and Sociobiology, 1996. **39**: p. 419-427.
- [32] Lau, C.W. and J.C. Nieh, *Honey bee stop-signaling production: temporal distribution and effect of feeder crowding*. Apidologie, 2010. **41**: p. 87-95.
- [33] Lam, C., Y. Li., T. Landgraf, and J.C. Nieh, *Dancing attraction: followers of honey bee tremble and waggle dances exhibit similar behaviors*. Biology Open, 2017. **6**: p. 810-817.
- [34] Johnson, B.R. and J.C. Nieh, *Modeling the Adaptive Role of Negative Signaling in Honey Bee Intraspecific Competition*. J Insect Behav, 2010. **23**(6): p. 459-471.
- [35] Forfert, N., M.E. Natsopoulou, R.J. Paxton, and R.F.A. Moritz, *Viral prevalence increases with regional colony abundance in honey bee drones (Apis mellifera L.)*. Infect Genet Evol, 2016. **44**: p. 549-554.

- [36] Michener, C.D., *Comparative social behavior of bees*. Annual Review of Entomology, 1969. **14**: p. 299-342.
- [37] Seeley, T.D., R.H. Seeley, and P. Akkratanakul, *Colony defense strategies of the honeybees in Thailand*. Ecological Monographs, 1982. **52**: p. 43-63.
- [38] Chahal, B.S., H.S. Brar, G.S. Gatoria, and H.S. Jhaji, *Aggressive behaviour of Apis florea towards Apis mellifera in hive robbing and in foraging*. Journal of Apiculture Research, 2015. **134-138**.
- [39] Atwal, A.S. and G.S. Dhaliwal, *Robbing between Apis indica F. and Apis mellifera L.* American Bee Journal, 1969. **109**(12).
- [40] Breed, M.D., X.-B. Deng, and R. Buchwald, *Comparative nestmate recognition in Asian honey bees, Apis florea, Apis andreniformis, Apis dorsata, and Apis cerana*. Apidologie, 2007. **38**: p. 411-418.
- [41] Paar, J.R., B.P. Oldroyd, E. Huettinger, and G. Kastberger, *Drifting workers in nest aggregations of the giant honeybee Apis dorsata*. Apidologie, 2002. **33**: p. 553-561.
- [42] Weihmann, F., D. Waddoup, T. Hotzl, and G. Kastberger, *Intraspecific aggression in giant honey bees (Apis dorsata)*. Insects, 2014. **5**: p. 689-704.
- [43] Seeley, T.D. and M.L. Smith, *Crowding honeybee colonies in apiaries can increase their vulnerability to the deadly ectoparasite Varroa destructor*. Apidologie, 2015. **46**(6): p. 716-727.
- [44] Garbuzov, M., N.J. Balfour, K. Shackleton, H. Al Toufailia, L. Scandian, and F.L.W. Ratnieks, *Multiple methods of assessing nectar foraging conditions indicate peak foraging difficulty in late season*. Insect Conservation and Diversity, 2020.
- [45] Scheiner, R., M. Barnert, and J. Erber, *Variation in water and sucrose responsiveness during the foraging season affects proboscis extension learning in honey bees*. Apidologie, 2003. **34**(1): p. 67-72.
- [46] Corby-Harris, V., J.H. Bowsher, M. Carr-Markell, M.J. Carroll, M. Centrella, S.C. Cook, M. Couvillon, G. DeGrandi-Hoffman, A. Dolezal, J.C. Jones, et al., *Emerging Themes from the ESA Symposium Entitled "Pollinator Nutrition: Lessons from Bees at Individual to Landscape Levels"*. Bee World, 2018. **96**(1): p. 3-9.
- [47] Wignall, V.R., I. Campbell Harry, N.L. Davies, S.D. Kenny, J.K. McMinn, and F.L.W. Ratnieks, *Seasonal variation in exploitative competition between honeybees and bumblebees*. Oecologia, 2020. **192**(2): p. 351-361.
- [48] Danner, N., A. Keller, S. Hartel, and I. Steffan-Dewenter, *Honey bee foraging ecology: Season but not landscape diversity shapes the amount and diversity of collected pollen*. PLoS One, 2017. **12**(8): p. e0183716.
- [49] Requier, F., J.-F. Odoux, T. Tamic, N. Moreau, M. Henry, A. Decourtye, and V. Bretagnolle, *Honey bee diet in intensive farmland habitats reveals an unexpectedly high flower richness and a major role of weeds*. Ecological Applications, 2015. **25**(4): p. 881-890.
- [50] Lau, P., V. Bryant, J.D. Ellis, Z.Y. Huang, J. Sullivan, D.R. Schmehl, A.R. Cabrera, and J. Rangel, *Seasonal variation of pollen collected by honey bees (Apis mellifera) in developed areas across four regions in the United States*. PLoS One, 2019. **14**(6): p. e0217294.

- [51] Bartlett, L.J., C. Rozins, B.J. Brosi, K.S. Delaplane, J.C. de Roode, A. White, L. Wilfert, and M. Boots, *Industrial bees: The impact of apicultural intensification on local disease prevalence*. J Appl Ecol, 2019. **56**(9): p. 2195-2205.
- [52] Tehel, A., M.J. Brown, and R.J. Paxton, *Impact of managed honey bee viruses on wild bees*. Curr Opin Virol, 2016. **19**: p. 16-22.
- [53] Potts, S.G., S.P.M. Roberts, R. Dean, G. Marris, M.A. Brown, R. Jones, P. Neumann, and J. Settele, *Declines of managed honey bees and beekeepers in Europe*. Journal of Apicultural Research, 2015. **49**(1): p. 15-22.
- [54] Degrandi-Hoffman, G., H. Graham, F. Ahumada, M. Smart, and N. Ziolkowski, *The Economics of Honey Bee (Hymenoptera: Apidae) Management and Overwintering Strategies for Colonies Used to Pollinate Almonds*. J Econ Entomol, 2019. **112**(6): p. 2524-2533.
- [55] Nouvian, M., S. Mandal, C. Jamme, C. Claudianos, P. d'Ettorre, J. Reinhard, A.B. Barron, and M. Giurfa, *Cooperative defence operates by social modulation of biogenic amine levels in the honey bee brain*. Proceedings of the Royal Society B, 2018. **285**(1871).
- [56] Nouvian, M., N. Deisig, J. Reinhard, and M. Giurfa, *Seasonality, alarm pheromone and serotonin: insights on the neurobiology of honeybee defence from winter bees*. Biol Lett, 2018. **14**(8).
- [57] Harris, J.W. and J. Woodring, *Effects of stress, age, season, and source colony on levels of octopamine, dopamine, and serotonin in the honey bee (Apis mellifera L.) brain*. Journal of Insect Physiology, 1992. **38**(1): p. 29-35.
- [58] Nouvian, M., L. Hotier, C. Claudianos, M. Giurfa, and J. Reinhard, *Appetitive floral odours prevent aggression in honeybees*. Nat Commun, 2015. **6**: p. 10247.
- [59] Rittschof, C.C., H.J. Vekaria, J.H. Palmer, and P.G. Sullivan, *Biogenic amines and activity levels alter the neural energetic response to aggressive social cues in the honey bee Apis mellifera*. Journal of Neuroscience Research, 2019. **00**: p. 1-13.
- [60] Lichtenberg, E.M., *Eavesdropping selects for conspicuous signals*. Current Biology, 2014. **24**: p. R598-R599.
- [61] Garay, J., R. Cressman, F. Xu, M. Broom, V. Csiszar, and T.F. Mori, *When optimal foragers meet in a game theoretical conflict: a model of kleptoparasitism*. Journal of Theoretical Biology, 2020. **502**.
- [62] Harrison, J.M., *Caste-specific changes in honeybee flight capacity*. Physiological Zoology, 1986. **59**(2): p. 175-187.
- [63] Southwick, E.E. and R.F.A. Moritz, *Metabolic response to alarm pheromone in honey bees*. Journal of Insect Physiology, 1985. **31**(5): p. 389-392.
- [64] Rivera-Marchand, B., T. Giray, and E. Guzmán-Novoa, *The cost of defense in social insects: insights from the honey bee*. Entomologia Experimentalis et Applicata, 2008. **129**(1): p. 1-10.
- [65] Suarez, R.K., J.R.B. Lighton, B. Joos, S.P. Roberts, and J.F. Harrison, *Energy metabolism, enzymatic flux capacities, and metabolic flux rates in flying honeybees*. Proceedings of the National Academy of Sciences, 1996. **93**: p. 12616-12620.
- [66] Rittschof, C.C., H.J. Vekaria, J.H. Palmer, and P.G. Sullivan, *Brain mitochondrial bioenergetics change with rapid and prolonged shifts in aggression in the honey bee, Apis mellifera*. J Exp Biol, 2018. **221**(Pt 8).

Figure 1. Mechanisms that co-regulate aggression and foraging in a robbing context. Robbing colonies display a behavioral syndrome characterized by simultaneous increases in foraging activity and aggression [23, 27, 44]. Because both foraging and aggression are high-energy behaviors [62, 63], this co-regulation is surprising [64], and likely the result of interacting processes at the individual, colony, and landscape levels that are also under human influence. At the landscape level, floral resources alter competition among colonies [27, 44], while colony density can increase aggressive interactions due to random processes like drift or increased availability of robbing targets. At the colony level, the decision to rob requires both increased aggression to fight bees from the victim colony and increased foraging activity to exploit the high-value food resource [23]. Risks associated with robbing cause colonies to target “winnable” small and often diseased colonies [10, 12, 13]. The lost bees and time due to robbing pose additional costs that may modulate both foraging and aggression. Such costs may only appear over a long-term timeframe. At the individual level, studies document seasonal fluctuations in biogenic amines, including a peak in brain serotonin and dopamine levels during the robbing season [57]. These biogenic amines modulate honey bee aggression [55]. Seasonal shifts could reflect increased aggressive interactions with neighboring colonies, or a correlated response to floral resource scarcity that prepares colonies for defense. Bioenergetics are important for both foraging and aggression in honey bees, due in part to the high metabolic costs of flight [65]. However, brain metabolic dynamics, which are modulated by biogenic amines, distinguish foraging and aggressive specialists [59, 66]. It is unclear how this neural mechanism is altered in the context of robbing.