

# Lawrence Berkeley National Laboratory

## Recent Work

**Title**

DYNAMICAL CALCULATIONS OF THE DIVISION OF IDEALIZED NUCLEI

**Permalink**

<https://escholarship.org/uc/item/0vs8r5k9>

**Author**

Nix, James Rayford.

**Publication Date**

1967-06-23

cg 2

# University of California Ernest O. Lawrence Radiation Laboratory

DYNAMICAL CALCULATIONS OF THE DIVISION OF IDEALIZED NUCLEI

James Rayford Nix

June 23, 1967

RECEIVED  
ERNEST O. LAWRENCE  
RADIATION LABORATORY  
AUG 2 1967  
LIBRARY AND  
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

Berkeley, California

UCRL-17540  
cg 2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

For International Conference on  
Nuclear Structure, September 7-13,  
1967, Tokyo, Japan

UCRL-17540  
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

DYNAMICAL CALCULATIONS OF THE DIVISION OF IDEALIZED NUCLEI

James Rayford Nix

June 23, 1967

Dynamical Calculations of the Division of Idealized Nuclei\*  
 James Rayford Nix  
 Lawrence Radiation Laboratory  
 University of California  
 Berkeley, California

We study the dynamics of the division of nuclei idealized as incompressible uniformly charged liquid drops with constant surface tension and nonviscous irrotational hydrodynamical flow. The shape of the nuclear surface is specified with six degrees of freedom, in terms of three smoothly joined portions of quadratic surfaces of revolution (e.g. two spheroids connected by a hyperboloidal neck). The potential energy is calculated in a standard way, whereas the kinetic energy is calculated by the approximate method of Werner and Wheeler.<sup>1-3</sup> The resulting classical equations of motion are solved numerically.

Figure 1 illustrates for three values of the fissility parameter  $x$  the sequences of shapes from the saddle point to scission, for initial conditions at the saddle point corresponding to 1 MeV of kinetic energy in the fission mode and 1 MeV in the mass-asymmetry mode. The parameter  $x$  is defined as  $\frac{1}{2}E_C(0)/E_S(0)$ , where  $E_S(0)$  and  $E_C(0)$  denote respectively the surface and Coulomb energies of the original spherical drop (with a sharp surface); the nuclei that correspond approximately to these values of  $x$  are indicated in parentheses.<sup>4</sup> We observe that: (1) The time from saddle to scission increases substantially with increasing  $x$ . (2) For small values of  $x$  the motion consists primarily of a constriction of the drop's neck, whereas for large  $x$  the drop also elongates substantially. (3) The mass-asymmetric component of motion is not amplified into a large mass asymmetry during the descent as proposed by Hill<sup>5</sup> and by Hill and Wheeler,<sup>6</sup> but instead represents stable oscillations about a symmetrical division. This means that within the limitation of a parameterization that permits only binary division, the liquid-drop model does not suggest an explanation of the observed mass asymmetry in fission.

In Fig. 2 we compare as a function of  $x$  the calculated (solid curve) and experimental most-probable total translational kinetic energies of fission fragments at infinity. (According to classical statistical mechanics, the most probable division occurs when the system starts from rest at the saddle point.) The dashed curve represents approximately the translational kinetic energy acquired by the fragments from their Coulomb repulsion after scission. The energy difference between the solid and dashed curves gives the translational kinetic energy acquired by the fragments prior to scission. The increase in this quantity as  $x$  increases above  $\sim 0.7$  reflects the increased distance from saddle to scission and the greater pre-scission elongation at larger values of  $x$ .<sup>7</sup> The experimental data are most-probable (open symbols) and average (solid symbols) fragment kinetic energies from a variety of sources but include only cases in which the mass distribution is symmetrical. The calculations (with no adjustable parameters<sup>4</sup>) reproduce both the correct order of magnitude and the correct trend with  $x$  of the kinetic energies, but a systematic difference of about 5% is evident.

The widths of the distributions in mass, translational kinetic energy, and excitation energy are now being calculated as functions of  $x$ .




Fig. 1. Illustrations of the descent from saddle to scission.




Fig. 2. Calculated and experimental fission-fragment kinetic energies.

Footnotes and references:

\* This work was performed under the auspices of the U.S. Atomic Energy Commission.

1. F. G. Werner and J. A. Wheeler, unpublished manuscript.
2. J. R. Nix, UCRL-11338; J. R. Nix and W. J. Swiatecki, Nucl. Phys. **71** (1965) 1.
3. I. Kelson, Phys. Rev. **136** (1964) B1667.
4. In converting from natural liquid-drop units to conventional units, and vice versa, we use the latest semiempirical mass-formula constants of W. D. Myers and W. J. Swiatecki, in *Proceedings of the Lysekil Symposium, 1966* (to be published in Arkiv Fysik).
5. D. L. Hill, Ph. D. dissertation, Princeton University, April 1951.
6. D. L. Hill and J. A. Wheeler, Phys. Rev. **89** (1953) 1102.
7. The calculated translational kinetic energies at scission for the fission of  $^{236}\text{U}$  and  $^{252}\text{Cf}$  essentially agree with the values deduced from experimental data on long-range alpha particles emitted during fission. See I. Halpern (preprint) and Y. Boneh, Z. Fraenkel and I. Nebenzahl, Phys. Rev. **156** (1967) 1305.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

