UCLA

UCLA Previously Published Works

Title

Dissociated Prostate Regeneration under the Renal Capsule.

Permalink

<https://escholarship.org/uc/item/0w71h28s>

Journal

Cold Spring Harbor protocols, 2015(11)

ISSN

1940-3402

Authors

Zong, Yang
Goldstein, Andrew S
Witte, Owen N

Publication Date

2015-11-01

DOI

10.1101/pdb.prot078063

Peer reviewed

Dissociated Prostate Regeneration under the Renal Capsule

Yang Zong,¹ Andrew S. Goldstein,^{2,3,5,6,7} and Owen N. Witte^{1,2,4,5,6}

¹Howard Hughes Medical Institute, University of California, Los Angeles, California 90095; ²Department of Molecular and Medical Pharmacology, University of California, Los Angeles, California 90095; ³Department of Urology, University of California, Los Angeles, California 90095; ⁴Department of Microbiology, Immunology, and Molecular Genetics, University of California, Los Angeles, California 90095; ⁵Jonsson Comprehensive Cancer Center, David Geffen School of Medicine, University of California, Los Angeles, California 90095; ⁶Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California, Los Angeles, California 90095

Tissue recombination models are useful for studying cancer initiation, progression, and metastasis. They also provide an in vivo environment in which to investigate the functional role of stem cells in tissue repair. In this protocol, we describe in detail the dissociated prostate regeneration assay. Dissociated adult murine prostate cells are combined with embryonic urogenital sinus mesenchymal cells and implanted under the renal capsule. Morphological tissue structures with appropriate epithelial–stroma interactions are reconstituted in the grafts.

MATERIALS

It is essential that you consult the appropriate Material Safety Data Sheets and your institution's Environmental Health and Safety Office for proper handling of equipment and hazardous material used in this protocol.

RECIPES: Please see the end of this protocol for recipes indicated by <R>. Additional recipes can be found online at <http://cshprotocols.cshlp.org/site/recipes>.

Reagents

BFS growth medium <R>
 Collagenase type I (1900 units/mL in RPMI medium)
 DNase I
 Dulbecco's Modified Eagle Medium (DMEM)
 Ethanol
 Fetal bovine serum (FBS)
 Ketamine/xylazine
 Mice (C57/BL6, male)
 Mice (severe combined immunodeficient [SCID], male)
 Neutralized collagen (freshly prepared, ice-cold) <R>
 Povidone-iodine solution
 Testosterone pellets (12.5 mg/pellet, 90-d release) (Innovative Research of America)

⁷Correspondence: andrew.goldstein@ucla.edu

Trypan blue
 Trypsin (0.05%)
 Urogenital sinus mesenchymal cells (see Protocol: **Preparation of Urogenital Sinus Mesenchymal Cells for Prostate Tissue Recombination Models** [Zong et al. 2014])

Equipment

Cell-culture plates (six well)
 Cell strainer (40 μ m)
 Centrifuge (benchtop)
 Dissecting macroscope
 Dissecting tools (scissors and fine forceps)
 Glass pipette (fire-polished)
 Hemocytometer
 Incubator
 Razor blade
 Shaker
 Shaver
 Surgical supplies (sutures and metal clips)
 Syringes with 18-, 20-, and 22-gauge needles
 Tissue-culture dishes
 Tubes (1.5-mL microcentrifuge)
 Tubes (50-mL conical)
 Warming plate
 Vortex

METHOD

Dissociation of Mouse Prostate Tissues

1. Harvest and dissect the prostate tissues from male C57/BL6 mice as previously described in detail (Lukacs et al. 2010).
2. Place the prostate tissue into a culture dish and mince thoroughly using a razor blade. Transfer the minced prostate fragments into a 50-mL conical tube by flushing the dish with DMEM + 10% FBS, and add more fresh DMEM + 10% FBS to bring the final volume to 27 mL.
3. Add 3 mL of collagenase solution (1900 units/mL in RPMI medium) to the tissue and incubate the tube on a shaker at 37°C for 2 h.
These conditions are suitable for the digestion of prostates from as many as five adult mice. To dissociate more prostate tissue, simply split the tissue to ensure efficient digestion.
4. Centrifuge the prostate chunks at 550g for 5 min at 25°C. Discard the supernatant and add 2 mL of 0.05% trypsin to the pellet, mix briefly by vortex, and incubate at 37°C for 5 min.
5. Add 3 mL of DMEM + 10% FBS containing 500 units of DNase I to quench the trypsin and break up the genomic DNA of the dead cells. Serially pass the suspension through 18-gauge, 20-gauge, and 22-gauge syringes by pipetting up and down five times per syringe.
6. Filter the cells through a 40- μ m cell strainer into a 50-mL conical tube, and then wash the cell strainer with 40 mL of DMEM + 10% FBS.
7. Centrifuge the tube at 550 g for 5 min at 25°C to collect the cell pellet. Resuspend the cells in 1 mL of DMEM + 10% FBS. Count the live cells using a hemocytometer after trypan blue staining.

Preparation of Prostate Cell Recombinants in Collagen Plugs

8. Aliquot $1.5\text{--}2 \times 10^5$ dissociated prostate cells into a 1.5-mL microcentrifuge tube and add 2×10^5 UGSM cells. Centrifuge the combined cells at 550 g for 3 min at room temperature and remove the supernatant completely.
9. Resuspend the cell pellet thoroughly in 15 μL of ice-cold neutralized collagen and place the cell/collagen slurry into one well of a six-well cell-culture plate.
Caution should be taken to avoid introducing air bubbles when resuspending the cells in collagen.
10. Incubate the plate at 37°C for 20 min to allow the collagen plug(s) to solidify. Add 3 mL of BFS medium to cover the plugs and incubate at 37°C until the engrafting surgery.

Surgery can be performed the same day or after overnight incubation. Occasionally, the collagen plugs will lift off the plate and become condensed after incubation; this is unlikely to affect the regeneration.

Engraftment of Prostate Cell Recombinants into the Subrenal Capsules

11. Anesthetize the SCID mice via intraperitoneal injection of ketamine/xylazine (100 mg/kg and 10 mg/kg body weight, respectively).
12. Prepare the surgical area by shaving and sterilization with povidone-iodine and ethanol.
Ensure that the mice are fully anesthetized, and increase the dose of ketamine/xylazine if necessary.
13. With a mouse lying on its side on a warming plate, make a skin incision ($\sim 6\text{--}8$ mm) on the middle flank above the kidney region. Cut through the peritoneal wall and expose the kidney. Exteriorize the kidney out of the abdominal cavity by applying gentle pressure on the abdomen using the thumb and forefinger.

See Troubleshooting.

14. Gently pinch and lift the thin fibrous membrane of the kidney with fine forceps, and then puncture it and create a pocket between the lifted membrane and the renal parenchyma by blunt dissection. Transfer the collagen plug(s) from the six-well plate onto the cut edge of the renal capsule, and insert the grafts into the pocket using a sterile fire-polished glass pipette. Take care not to damage the parenchyma, which can cause severe bleeding.

Two to three prostate cell recombinants can be placed under the renal capsule of one kidney.

15. Gently push the kidney back into the abdominal cavity while dilating the incision on the peritoneal wall. Suture the peritoneal wall and insert a testosterone pellet subcutaneously into the dorsal neck and shoulder region. Close the skin incision using metal clips, which should be removed 7 d after surgery.

See Troubleshooting.

TROUBLESHOOTING

Problem (Step 13): The kidney cannot be slipped out of the abdominal cavity using gentle pressure on the abdomen.

Solution: Alternatively, grab the fat tissue surrounding the tip of the kidney with fine forceps, and pull it and the kidney out.

Problem (Step 15): Prostate cell recombinants fall out of the subrenal capsules.

Solution: Before pushing the kidney back into the abdominal cavity, confirm that the grafts are inserted into the subrenal capsules and completely covered with the fibrous membrane of the kidney.

RECIPES

BFS Growth Medium

Reagent	Final concentration
Dulbecco's Modified Eagle Medium (DMEM)	1×
Fetal bovine serum (FBS)	5% (v/v)
Nu-Serum IV	5% (v/v)
Glutamine	2 mM
Penicillin	100 units/mL
Streptomycin	100 µg/mL
Insulin	5 µg/mL

Store medium at 4°C for up to 1 mo.

Neutralized Collagen

Reagent	Volume
Collagen I, rat tail	500 µL
NaOH (1 N)	11.5 µL
PBS (10×	56.8 µL

The final pH value should be 7.0. Store the solution on ice for up to 1 h.

REFERENCES

Lukacs RU, Goldstein AS, Lawson DA, Cheng D, Witte ON. 2010. Isolation, cultivation and characterization of adult murine prostate stem cells. *Nat Protoc* 5: 702–713.

Zong Y, Goldstein AS, Witte ON. 2014. Preparation of urogenital sinus mesenchymal cells for prostate tissue recombination models. *Cold Spring Harb Protoc* doi: 10.1101/pdb.prot078055.