

Lawrence Berkeley National Laboratory

Recent Work

Title

THERMODYNAMIC PROPERTIES OF POLAR FLUIDS FROM A PERTURBED-DIPOLAR-HARD-SPHERE EQUATION OF STATE

Permalink

<https://escholarship.org/uc/item/0wh5w4qb>

Authors

Bryan, P.F.

Prausnitz, J.M.

Publication Date

1986-12-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

FEB 20 1987

Submitted to Fluid Phase Equilibria

THERMODYNAMIC PROPERTIES OF POLAR FLUIDS FROM A
PERTURBED-DIPOLAR-HARD-SPHERE EQUATION OF STATE

P.F. Bryan and J.M. Prausnitz

December 1986

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.*



LBL-22693
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THERMODYNAMIC PROPERTIES OF POLAR FLUIDS FROM A PERTURBED-DIPOLAR-HARD-SPHERE EQUATION OF STATE

P.F. Bryan and J.M. Prausnitz

*Materials and Molecular Research Division, Lawrence Berkeley Laboratory and
Chemical Engineering Department, University of California, Berkeley 94720*

ABSTRACT

Based on theoretical results for a system of hard spheres with dipoles, a new equation of state is applied to the correlation of thermodynamic properties for four fluids: argon, ammonia, water, and acetonitrile. The reference system has the same dependence on density as that given by the Carnahan-Starling equation, but the coefficients are now functions of temperature through the reduced dipole moment. These coefficients are chosen to match the Padé approximant developed by Stell, Rushbrooke and Hoyer for the Helmholtz energy of dipolar hard spheres. The reference system proposed here shows a phase transition for reduced dipole moments greater than 1.9. A simple, empirical perturbation term is added to the reference system to account for induction and dispersion forces. For polar fluids, the equation gives results significantly better than those obtained from conventional cubic equations of state, when using the same limited experimental data for determining equation-of-state parameters.

*This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

INTRODUCTION

While much attention has been given to the equation of state for nonpolar fluids, relatively little work has been reported on the equation of state for those fluids whose molecules possess an appreciable dipole moment. For practical applications to systems containing polar fluids, it has become common practice to use empirical extensions of methods originally intended for nonpolar fluids (e.g. Soave(1980), Lüdecke and Prausnitz(1985), Guo, et al (1985a, 1985b), Ikonomou and Donahue (1986), Vimalchand and Donahue (1985), and Huron et al(1978)) or else to use "chemical" equations of state such as those of Grenzheuser and Gmehling(1986, 1981), Gmehling et al (1979), Wenzel et al (1982), Kulasinska, et al (1983), and others.

Recently, however, there has been much interest in the use of perturbation theory to describe properties of polar fluids. Perturbation theories have been reported for dipolar hard spheres (Stell, et al (1974), (1973), and (1972)) and multipolar Lennard-Jones fluids (Annath et al (1974), Twu and Gubbins (1975), Flytzani-Stephanopoulos, et al (1975), and Gray, et al(1978)). These theories have led to some successful applications for representing the properties of real fluids (Gubbins and Twu (1978), Twu, et al (1978), and (1976), Winkelmann (1983), (1981), and (1979), and Chung, et al (1984)). All of these perturbation theories use a reference system which exhibits only repulsive forces; typical reference systems are assemblies of hard spheres or other rigid convex bodies. However, for highly polar fluids, these reference systems can lead to difficulties in convergence of the perturbation series. Among others, Chung et al (1984) state the need for a more sophisticated reference system to surmount this obstacle.

In this work, we use a reference system containing orientation-averaged attractive (dipole-dipole) forces in addition to repulsive forces; this reference system is combined with a semi-empirical perturbation term to represent thermodynamic properties of real polar fluids.

EQUATION OF STATE FOR POLAR FLUIDS

We use an equation of state of the van der Waals form where compressibility factor z is the sum of two contributions, one from the reference system and one from a perturbation,

$$z = \frac{Pv}{RT} = z^{\text{ref}} + z^{\text{pert}} \quad (1)$$

It has been customary to use for the reference system an assembly of convex bodies (usually spheres) which exhibit only repulsive forces at contact. Attractive forces, at somewhat larger distances, are taken into account in the perturbation term.

In a departure from previous practice, we use here a reference system that contains hard spheres with ideal dipole moments at their centers. This reference system therefore includes not only the usual forces of repulsion but, in addition, dipole-dipole forces which, on average, are attractive. The perturbation term takes into account attractive forces arising from induction and dispersion.

For the compressibility factor of the reference system, we write a generalized Carnahan-Starling expression:

$$z^{\text{ref}} = \frac{1 + f^{(1)}\eta + f^{(2)}\eta^2 + f^{(3)}\eta^3}{(1 - \eta)^3} \quad (2)$$

where the reduced density is:

$$\eta = bp/4 = \pi\rho\sigma^3/6 \quad (3)$$

and the coefficients depend on reduced dipole moment $\tilde{\mu}$:

$$f^{(1)} = 1 - 0.61357\tilde{\mu} + 3.06030\tilde{\mu}^2 - 4.63519\tilde{\mu}^3 + 0.86956\tilde{\mu}^4 \quad (4)$$

$$f^{(2)} = 1 + 2.79108\tilde{\mu} - 12.6917\tilde{\mu}^2 + 15.8026\tilde{\mu}^3 - 2.77528\tilde{\mu}^4 \quad (5)$$

$$f^{(3)} = 1 + 2.98994\tilde{\mu} - 13.1452\tilde{\mu}^2 + 14.5664\tilde{\mu}^3 - 2.50544\tilde{\mu}^4 \quad (6)$$

Here ρ is density and σ is the hard-sphere diameter. In the original Carnahan-Starling equation, all coefficients $f^{(1)} = f^{(2)} = f^{(3)} = 1$. In this work, however, these coefficients are functions of temperature through the reduced dipole moment $\tilde{\mu} = \mu/(\sigma^3 kT)^{1/2} = (2\pi\mu^2/3bRT)^{1/2}$. When μ is in Debye, b in cm^3/mole , and T in Kelvins, $\tilde{\mu} = 95.59 \mu/(bT)^{1/2}$.

Dimensionless dipole moment $\tilde{\mu}$ depends on dipole moment, temperature, and molecular size. For the fluids examined here, at their normal boiling points, the values of $\tilde{\mu}$ are: argon (0.00), ammonia (1.33), water (1.41), and acetonitrile (1.42); at their critical temperatures, they are: argon (0.00), ammonia (1.02), water (1.07), and acetonitrile (1.14).

The coefficients in equation (2) were determined by fitting to a complex equation of state for hard spheres with point dipoles at their centers. That equation was developed by Stell, Rushbrooke, and Hoyer (1973) who devised a Padé approximant to the Helmholtz energy of a system of hard spheres using the perturbation theory of Stell, Rasaiah, and Narang (1974, 1972). Their Padé approximant ("Stell's equation"), discussed in Appendix I, gives the Helmholtz energy of a system of dipolar hard spheres as a function of reduced density η and reduced dipole moment $\tilde{\mu}$; it is valid for values of $\tilde{\mu}$ to at least 2.00 and perhaps slightly higher.

The Helmholtz energy obtained from Stell's equation is in good agreement with Monte-Carlo calculations for dipolar hard spheres. The third and fifth columns of Table 1 show "experimental" (Monte-Carlo) Helmholtz energies and those calculated from Stell's equation. Monte-Carlo calculations are taken from Patey and Valleau (1974).

Table 1. Effect of Reduced Dipole Moment on the Helmholtz Energy of a System of Dipolar Hard Spheres. Results from Monte-Carlo Simulation, from Dipole-Dipole Carnahan-Starling (DDCS), and from Stell's (SRH) Equations

$-(A - A^{HS})/RT$				

η	$\tilde{\mu}$	Monte-Carlo	DDCS	SRH

0.474	0.24	0.0037	0.0042	0.0034
"	0.47	0.052	0.053	0.051
"	0.71	0.22	0.24	0.23
"	0.95	0.58	0.61	0.62
"	1.19	1.2	1.2	1.3
"	1.42	2.1	2.1	2.2
"	1.66	3.3	3.4	3.5
0.437	0.24	0.0035	0.0039	0.0030
"	0.71	0.21	0.22	0.20
"	1.19	1.1	1.1	1.2
"	1.66	3.2	3.2	3.2

A^{HS} is the Helmholtz energy at the same η with $\tilde{\mu} = 0$

As discussed in the Appendix I, Stell's equation for the Helmholtz energy gives an equation of state that is mathematically complex. To simplify the reference system, the Carnahan-Starling (1969) form is retained here as shown by equation (2). Functions $f^{[1]}$, $f^{[2]}$, and $f^{[3]}$ (Equations (4) - (6)) reduce to the Carnahan-Starling values when $\tilde{\mu} = 0$. These functions are determined by fitting to "data" generated from Stell's equation. The compressibility factors calculated from equations (2), (4), (5), and (6) are in good agreement with those calculated from Stell's equation, as shown in Figure 1.

Although not immediately apparent by visual inspection, Figure 1 shows that when $\tilde{\mu} \geq 1.90$, dipolar hard spheres exhibit a phase transition. This behavior was also noted by Stell, Rushbrooke, and Hoyer (1973), who made representative VLE calculations but did not report the critical dipole moment (or, equivalently, critical temperature) for dipolar hard spheres. When Stell's equation is used, the critical reduced dipole moment is 1.91, again showing excellent agreement with equations (2), (4), (5), and (6). Figure 2 shows the phase transition more clearly in an expanded view of a portion of Figure 1: an isotherm at $\tilde{\mu} = 2.5$ is included to show the behavior at high reduced dipole moments (low temperatures).

Table 1 compares our results for the Helmholtz energies of dipolar hard spheres (equations (2), (4), (5), and (6)) to those obtained from Stell's equation and by Monte-Carlo simulation.

The dipole-dipole Carnahan-Starling (DDCS) equation provides a reference system that represents the behavior of dipolar hard spheres. For the perturbation term we use a semi-empirical form, suggested by Wong and Prausnitz (1985), similar to that of Redlich and Kwong (1949). The proposed equation of state is:

$$z = \frac{1 + f[1]\eta + f[2]\eta^2 - f[3]\eta^3}{(1 - \eta)^3} - \frac{a}{RT(v + 0.2b)} \quad (7)$$

where the second term on the RHS is a form intermediate between the well-known attractive forms proposed by van der Waals and by Redlich and Kwong.

At the critical temperature T_c , parameters a and b in Equation (7) can be determined from the usual criteria; at the critical point,

$$\left(\frac{\partial P}{\partial v}\right)_{\tilde{\mu}_c, T_c} = 0 \quad (8)$$

$$\left(\frac{\partial^2 P}{\partial v^2}\right)_{\tilde{\mu}_c, T_c} = 0 \quad (9)$$

where $\tilde{\mu}_c = \mu / (\sigma^3 k T_c)^{1/2}$.

Equations (8) and (9) give values of (b_c/v_c) and $(a_c/RT_c v_c)$. Since accurate experimental values for v_c are often not available, parameters a_c and b_c are here determined by:

$$a_c = \left(\frac{a_c}{RT_c v_c}\right) z_c^{\text{theo}} \left(\frac{R^2 T_c^2}{P_c}\right) \quad (10)$$

and:

$$b_c = \left(\frac{b_c}{v_c}\right) z_c^{\text{theo}} \left(\frac{RT_c}{P_c}\right) \quad (11)$$

where z_c^{theo} is the compressibility factor calculated from equation (7) at the critical point; that is, the point where equations (8) and (9) are satisfied. When applied to argon, equation (11) gives a value for b_c which is about 10% smaller than that which gives the most accurate calculated values for the critical compressibility and the liquid molar volumes. Since these properties are extremely sensitive to b_c , the direct use of equation (11) leads to poor results. Therefore, the value of b_c was adjusted upward by 10%, providing an improvement for all calculated properties for all fluids examined, especially liquid molar volumes. The calculation of properties from Equation (10) is further improved if b_c is fixed so as to match a single experimental value of the liquid molar volume. The procedure for matching b_c to experimental liquid molar volumes is explained in detail by Panagiotopoulos and Kumar (1985), who demonstrate the technique for three popular two-parameter equations of state. Appendix II describes application of this technique to Equation (7).

Reduced values of b_c (adjusted) and a_c can be found from:

$$\tilde{b}_c = b_c P_c / RT_c = 0.178 + 1.124 \mu / (RT_c^2 / P_c)^{1/2} \quad (12)$$

$$\tilde{a}_c = a_c P_c / R^2 T_c^2 = 0.5352 + 0.1288 \mu_c - 0.1783 \mu_c^2 \quad (13)$$

(For dimensional consistency, equation (12) must have μ in Debye, T_c in Kelvins, P_c in bars, and $R = 83.14 \text{ cm}^3\text{-bar/mole-K}$)

For our engineering-oriented purposes, parameter b , however determined, is independent of temperature to a good approximation. Parameter a , on the other hand, increases markedly with falling temperature. This dependence must be given accurately, especially for the calculation of vapor pressures, where an error of a few percent in a can lead to an error of 100% in the calculated vapor pressure. Soave (1972) showed that, for

the Redlich-Kwong equation of state, a plot of $(a/a_c)^{1/2}$ vs. $(T/T_c)^{1/2}$ is linear for non-polar fluids at temperatures below the critical. An examination of the properties of argon with Equation (7) confirms this linearity, but the plot of $(a/a_c)^{1/2}$ vs. $(T/T_c)^{1/2}$ gives one line for $(T/T_c) \leq 0.8$ and another for $(T/T_c) \geq 1.0$, with nonlinear behavior in the intermediate region $0.8 \leq (T/T_c) \leq 1.0$. Therefore, we propose that the dependence of a on T be given by:

$$(a/a_c)^{1/2} = (0.41 + 0.075(T/T_c)^{1/2})(1-\Phi) + (\alpha + \beta(T/T_c)^{1/2})(1+\Phi) \quad (14)$$

where the weighting function Φ is defined by:

$$\Phi = \tanh [10((T_c/T) - 1)] \quad (15)$$

Function Φ approaches +1 as T/T_c approaches zero, and approaches -1 as T/T_c approaches infinity. The transition between these two values occurs almost entirely in a small temperature range near $T/T_c = 0.9$.

The first expression on the RHS of Equation (14) represents the high-temperature ($T/T_c > 1.0$) behavior of a . The constants shown are suitable for all of the substances examined here; these values should be used unless a large body of data is available at high temperatures over a wide range of densities.

The second expression on the RHS of equation (14) represents the low-temperature ($T/T_c < 0.8$) behavior of a . Constants α and β can be determined in many ways but, for engineering purposes, the most convenient is to use data for the vapor pressure and enthalpy of vaporization at a single specified temperature, most commonly the normal boiling point. The appendix describes the procedure for fixing α and β from such data.

Function Φ , defined by Equation (15), blends the high- and low-temperature functions for a smoothly together in the range of temperatures just below the critical temperature.

RESULTS

Figure 3 shows calculated and experimental results for ammonia; the data were taken from the compilation of Haar and Gallagher (1978). Calculations are based exclusively on experimental data for critical temperature, critical pressure, dipole moment, normal boiling point, and enthalpy of vaporization at the normal boiling point. The calculations are in good agreement with experiment, although there is some discrepancy in the near-critical region, as is common with equations of state of the Van der Waals form. The calculated boundaries of the two-phase region are also in good agreement with experiment, again with the exception of the critical region.

Figure 4 compares calculated and experimental vapor pressures for ammonia (Haar and Gallagher (1978)), water (Keenan, et al (1978)), argon (Angus and Armstrong (1971)), and acetonitrile (Francesconi, et al (1975)). Required input data are the same as those indicated above. Agreement is excellent; about the same as would be obtained with the original Soave-Redlich-Kwong (SRK) equation for argon (and other non-polar fluids), and somewhat better than that from the SRK equation for polar fluids. With more experimental data as input, an even better correlation for a (and hence vapor pressure) can be obtained, as suggested by Soave (1980).

Figure 5 shows that calculated saturated-liquid molar volumes are also in good agreement with experimental data for ammonia (Haar and Gallagher (1978)), water (Keenan, et al (1978)), and argon (Angus and Armstrong (1971)), even though no volumetric data were used to determine the parameters. The calculated liquid volumes for acetonitrile are less accurate, for reasons that are not clear. Calculations for other highly polar fluids give much better results, suggesting that the polarity of acetonitrile alone does not fully explain the relatively poor performance of Equation (7) in predicting its saturated-liquid molar volume. Better values of b , and hence of liquid volume, can be obtained if a single liquid-volume datum is available to fix parameter b .

Figure 6 compares calculated second virial coefficients with experimental results taken from the compilation of Dymond and Smith (1980). Agreement is excellent for argon, good for ammonia and water, and fair for acetonitrile. The results are surprisingly good, in view of the well-known difficulty of **simultaneously** correlating high-density properties (vapor pressures and liquid volumes) and low-density properties (second virial coefficients) with a simple equation of state. Appendix III discusses a fundamental problem concerning calculation of the second virial coefficient from Equation (7).

COMPARISON WITH CONVENTIONAL EQUATIONS

To determine the effect of dipolar forces in the reference system, calculations were performed for polar fluids with and without reduced dipole moment $\tilde{\mu}$. For polar fluids, the change in reference system has almost no effect on the accuracy of the calculated vapor pressures, because parameter a is determined from vapor-pressure data. However, there is significant improvement in calculated saturated-liquid molar volumes and in second virial coefficients when the dipolar reference system is used. Table 2 compares the accuracy of Equation (2) with and without a dipole moment in the reference system.

The results obtained here suggest that, when using a semi-empirical perturbed hard-sphere equation, it is worthwhile to include the effect of reduced dipole moment on the reference term. For polar fluids, significant improvement is obtained with little additional computational effort and without adding any additional adjustable parameters.

ACKNOWLEDGEMENT

This work was supported by the Director, Office of Energy Research, Basic Energy Science, Chemical Sciences Division of the US Department of Energy under Contract DE-AC03-76SF00098. Additional support was provided by the Gas Research Institute.

Table 2. Influence of Dipole Moment on Calculated Liquid Volumes and on
 Second Virial Coefficients for Ammonia and for Water
 (constants are determined from critical and vapor-pressure data)

Property	Temperature (K)	Average Absolute Deviation Between DDCS Equation and Experimental Data	
		$\mu = \mu_{\text{exptl}}$	$\mu = 0$
		Percent	
$v_{\text{sat liq}}(\text{NH}_3)$	197 - 400	2.2	7.1
$v_{\text{sat liq}}(\text{H}_2\text{O})$	273 - 590	1.9	8.4
		cm^3/mole	
$B(\text{NH}_3)$	273 - 523	32	86
$B(\text{H}_2\text{O})$	323 - 723	97	203

Literature Cited

- Anath, M.S., Gubbins, K.E., and Gray, C.G., 1974. Perturbation Theory for Equilibrium Properties of Molecular Fluids. *Mol. Phys.*, **28**: 1005 - 1030.
- Angus, S. and Armstrong, B. (eds.), 1971. *International Thermodynamic Tables of the Fluid State: Argon*. Butterworths, London, England.
- Carnahan, N.F. and Starling, K.E., 1969. Equation of State for Nonattracting Rigid Spheres. *J. Chem. Phys.*, **51**: 635 - 636.
- Chung, T.H., Khan, M.M., Lee, L.L., and Starling, K.E., 1984. A New Equation of State for Polar and Nonpolar Fluids. *Fluid Phase Equilibria*, **17**: 351 - 372.
- Deiters, U., 1979, Doctoral Dissertation, University of Bochum.
- Dymond, J.H. and Smith, E.B., 1980. *The Virial Coefficients of Pure Gases and Mixtures*. Clarendon Press, Oxford, England.
- Flytzani-Stephanopoulos, M., Gubbins, K.E., and Gray, C.G., 1975. Thermodynamics of Mixtures of Nonspherical Molecules II. Strong Polar, Quadrupolar, and Overlap Forces. *Mol. Phys.*, **30**: 1649 - 1676.
- Francesconi, A.Z., Franck, E.U., and Lentz, H., 1975. Die PVT-Daten des Acetonitrils bis 450° C und 2500 bar. *Ber. Bunsen. Ges.*, **79**: 897 - 901.
- Gray, C.G., Gubbins, K.E., and Twu, C.H., 1978. Perturbation Theory for Molecular Fluids: Third-order Term in the Pople Expansion. *J. Chem. Phys.*, **69**: 182 - 193.
- Grenzheuser, F. and Gmehling, J., 1986. An Equation of State for the Description of Phase-Equilibrium and Caloric Quantities on the Basis of the "Chemical Theory." *Fluid Phase Equilibria*, **25**: 1 - 29.

- Grenzheuser, F. and Gmehling, J., 1981. Calculation of Phases Equilibria Containing Strongly Polar or Associating Compounds by an Equation of State. Seventh Int. Congr. Chem. Eng., Chem. Equip. Design. Autom. (CHISA), Prague.
- Gubbins, K.E. and C.H. Twu, 1978. Thermodynamics of Polyatomic Fluid Mixtures I. Theory. Chem. Eng. Sci., 33: 863 - 878.
- Guo, T.M., Kim, H., Lin, H.M., and Chao, K.C., 1985a. Ind. Eng. Chem. Proc. Des. Dev., 24: 764 - 768.
- Guo, T.M., Lin, H.M., and Chao, K.C., 1985b. Ind. Eng. Chem. Proc. Des. Dev., 24: 768 - 773.
- Haar, L. and Gallagher, J.S., 1978. Thermodynamic Properties for Ammonia. J. Phys. Chem. Ref. Data, 7: 635.
- Huron, M.-J., Dufour, G.-N., and Vidal, J., 1978. Vapor-Liquid-Equilibrium and Critical-Loci-Curve Calculations with the Soave Equation for Hydrocarbon Systems with Carbon Dioxide and Hydrogen Sulfide. Fluid Phase Equilibria, 1: 247 - 265.
- Ikonomou, G.D. and Donahue, M.D., 1986. Thermodynamics of Hydrogen-Bonded Molecules: The Associated Perturbed-Anisotropic-Chain Theory. AIChE J., 32: 1716 - 1725.
- Keenan, J.H., Keyes, F.G., Hill, P.G., and Moore, J.G., 1978. Steam Tables. Wiley, New York, NY, USA.
- Keesom, W.H., 1912. Comm. Phys. Lab. Leiden., Suppl 24b, Section 6.
- Kulasinska, G., Moorwood, R.A.S., and Wenzel, H., 1983. Calculation of Vapor-Liquid and Liquid-Liquid Equilibria by an Equation of State. Fluid Phase Equilibria, 13: 121 - 132.

- Ludecke, B. and Prausnitz, J.M., 1985. Phase Equilibria for Strongly Non-ideal Mixtures from an Equation of State with Density-Dependent Mixing Rules. *Fluid Phase Equilibria*, **22**: 1 - 19.
- Oellrich, L.R., Knapp, H., and Prausnitz, J.M., 1978. A Simple Perturbed-Hard-Sphere Equation of State Applicable to Subcritical and Supercritical Temperatures. *Fluid Phase Equilibria*, **2**: 163 - 171.
- Panagiotopoulos, A.Z. and Kumar, S.K., 1985. A Generalized Technique to Obtain Pure-Component Parameters for Two-Parameter Equations of State. *Fluid Phase Equilibria*, **22**: 77 - 88.
- Patey, G.N. and Valleau, J.P., 1974. Dipolar Hard Spheres: A Monte-Carlo Study. *J. Chem. Phys.*, **61**: 534 - 540.
- Redlich, O. and Kwong, J.N.S., 1949. On the Thermodynamics of Solutions V. *Chem. Rev.*, **44**: 233 - 244.
- Soave, G., 1980. Rigorous and Simplified Procedures for Determining the Pure-Component Parameters in the Redlich-Kwong-Soave Equation of State. *Chem. Eng. Sci.*, **35**: 1725 - 1729.
- Soave, G., 1972. Equilibrium Constants from a Modified Redlich-Kwong Equation of State. *Chem. Eng. Sci.*, **27**: 1197 - 1203.
- Stell, G., Rasaiah, J.C., and Narang, H., 1974. Thermodynamic Perturbation Theory for Simple Polar Fluids, II. *Mol. Phys.*, **27**: 1393 - 1414.
- Stell, G., Rushbrooke, G.S., and Hoye, J.S., 1973. Theory of Polar Fluids I. Dipolar Hard Spheres. *Mol. Phys.*, **26**: 1199 - 1215.
- Stell, G., Rasaiah, J.C., and Narang, H., 1972. Thermodynamic Perturbation Theory for Simple Polar Fluids, I. *Mol. Phys.*, **23**: 393 - 406.

- Twu, C.H. and Gubbins, K.E., 1978. Thermodynamics of Polyatomic Fluid Mixtures II. Polar, Quadrupolar, and Octopolar Molecules. *Mol. Phys.*, **33**: 879 - 887.
- Twu, C.H., Gubbins, K.E., and Gray, C.G., 1976. Thermodynamics of Mixtures of Nonspherical Molecules III. Fluid Phase Equilibria and Critical Loci. *J. Chem. Phys.*, **64**: 5186 - 5197.
- Twu, C.H., Gubbins, K.E., and Gray, C.G., 1975. Excess Thermodynamic Properties for Liquid Mixtures of Nonspherical Molecules. *Mol. Phys.*, **29**: 713 - 729.
- Vera, J.H. and Prausnitz, J.M., 1971. A Semi-Theoretical Van der Waals Equation for Calculation of Vapor-Liquid Equilibria at High Pressure. *Can. J. Chem.*, **49**: 2037 - 2043.
- Vimalchand, P. and Donahue, M.D., 1985. Thermodynamics of Quadrupolar Molecules: The Perturbed-Anisotropic Chain Theory. *Ind. Eng. Chem.*, **24**: 246.
- Wenzel, H., Moorwood, R.A.S., and Baumgaertner, M., 1982. Calculation of Vapor-Liquid Equilibria of Associated Systems by an Equation of State. *Fluid Phase Equilibria*, **9**: 225 - 266.
- Winkelmann, J., 1983. Perturbation Theory of Dipolar Hard Spheres: The Vapour-Liquid Equilibria of Strongly Polar Mixtures. *Fluid Phase Equilibria*, **11**: 207 - 224.
- Winkelmann, J., 1981. Perturbation Theory of Dipolar Hard Spheres: The Vapour-Liquid Equilibria of Strongly Polar Mixtures. *Fluid Phase Equilibria*, **7**: 207 - 217.
- Winkelmann, J., 1979. The Vapour Pressure of Dipolar-Hard-Sphere Systems: A Study Using the Thermodynamic Perturbation Theory. *SINET (Ethiop. J. Sci.)*, **2**: 23 - 35.
- Wong, J.O. and Prausnitz, J.M., 1985. Comments Concerning a Simple Equation of State of the van der Waals Form. *Chem. Eng. Commun.*, **37**: 41 - 53.

APPENDIX I

DIPOLAR-HARD-SPHERE EQUATION OF STATE OF STELL, ET AL

For Helmholtz energy A, the Padé approximant of Stell, et al (1973) is:

$$\frac{A - A^{HS}}{RT} = \frac{-4/3 \tilde{\mu}^4 \eta I_2(\eta)}{1 + 5/6 \tilde{\mu}^2 I_3(\eta)/I_2(\eta)} \quad (I-1)$$

where A^{HS} is the Helmholtz energy of a system of hard spheres (no dipoles) at temperature T and reduced density η . Here:

$$I_2(\eta) = \frac{1 - 0.6910 \eta - 1.1690 \eta^2 + 0.7510 \eta^3}{(1 - \eta)^3} \quad (I-2)$$

$$I_3(\eta) = \frac{1 + 1.1913 \eta - 0.4252 \eta^2}{1 - 1.1279 \eta - 0.7317 \eta^2} \quad (I-3)$$

and the reduced dipole moment $\tilde{\mu}$ is defined by $\tilde{\mu} = \mu/(\sigma^3 kT)^{1/2} = (2\pi\mu^2/3bRT)^{1/2}$. The equation of state is obtained from:

$$P = - \left(\frac{\partial A}{\partial V} \right)_T \quad (I-4)$$

APPENDIX II

DETERMINATION OF EQUATION-OF-STATE PARAMETERS

In dimensionless form, the Perturbed DDCS equation (Equation (7)) is:

$$\tilde{P} = \frac{Pb}{4RT} - f(\eta, \tilde{\mu}) - \frac{\zeta a}{bRT} \frac{\eta}{1.25 + \eta} \quad (\text{II-1})$$

If pressure, temperature, and dipole moment are known, the only remaining variables are parameters a and b and reduced density η . To fit an experimental vapor-pressure datum, we use two criteria, equality of pressure and equality of chemical potentials in the two phases; both criteria must be satisfied. For the perturbed DDCS equation, the vapor-liquid-equilibrium (VLE) criterion may be conveniently stated as:

$$\frac{\zeta a}{bRT} = F(\eta^{\text{sat liq}}) - F(\eta^{\text{sat vap}}) \quad (\text{II-2})$$

where function F is defined by:

$$F(\eta) = \frac{\ln \eta + (1-f^{(3)}) \ln(1-\eta) + \frac{(3 + f^{(1)} - 3f^{(3)}) + \eta(-4f^{(3)} + 2f^{(2)} - 2)}{2(1-\eta)}}{\ln(1.25 + \eta)} \quad (\text{II-3})$$

If we know the saturated-liquid molar volume corresponding to the vapor-pressure datum, we can determine the unique pair of parameters a and b that simultaneously satisfies the VLE criterion and gives the proper value of η for the liquid phase by the following procedure:

1. Make initial estimates of a and b.
2. Compute $\tilde{\mu}$.
3. Compute $\eta^{\text{sat liq}}$ and $\eta^{\text{sat vap}}$ from (II-1).
4. Compute new value of a from (II-2) and (II-3).
5. New value of a = old value of a ?
 - NO : Replace old value of a with new value and go to step 3.
 - YES : Go to step 6.
6. $\eta^{\text{sat liq}}$ (calculated) = $\eta^{\text{sat liq}}$ (experimental) ?
 - NO : Adjust value of b and go to step 2.
 - YES : Exit. The values of a and b are consistent with the experimental values for p^{sat} and $v^{\text{sat liq}}$.

The algorithm described above is only an example; for converging the "inner loop" (steps 3 - 5), there may be more efficient methods than the resubstitution scheme described, and there are many formulae for adjusting b in the "outer loop" (steps 2 - 6). The procedure indicated above is tedious but, with sufficient repetition, it is possible to develop tables and/or equations relating the dimensionless groups $Pb/4RT$, $5a/bRT$, $\tilde{\mu}$, and $\eta^{\text{sat liq}}$ (and, if desired, $\eta^{\text{sat vap}}$) to one another.

This repetition has been carried out by Panagiotopoulos and Kumar (1985) for three popular two-parameter equations of state. The Perturbed DDCS equation, however, has one more independent dimensionless group in addition to those in popular equations of the Van der Waals or Redlich-Kwong type.

Instead of fitting simultaneously a and b, it is more convenient to fix b at the critical value and to develop a simple method for determining the temperature dependence of a. When b is known, there are a number of simple ways to determine constants α and β in equation (14), and hence to fix the temperature dependence of a in the sub-critical region:

(1) The original approach of Soave (1972) can be applied to the DDCS equation if the acentric factor is known.

(2) If the vapor pressure is known at two different temperatures, Equations (II-1) - (II-3) allow calculation of a at each temperature. Constants α and β are then obtained by straightforward algebra.


(3) If the vapor pressure and the enthalpy of vaporization are known at a given temperature (such as the normal boiling point), the value of a can be fixed at that point from the vapor-pressure datum, and from the derivative (da/dT) which is related to the enthalpy of vaporization. The latter relation is a complex one which requires solution by numerical methods. When such data are used, it is much simpler to use the enthalpy of vaporization with the Clausius-Clapeyron equation to estimate vapor pressures at temperatures a few degrees above and below the temperature of interest. These estimated vapor pressures are used to calculate a at each point; da/dT is calculated from these two values by finite-difference approximation. In either case, α and β are easily calculated when a and its derivative with respect to temperature are known.

APPENDIX III

CALCULATION OF SECOND VIRIAL COEFFICIENTS

The relatively large errors in the calculated second virial coefficients follow from a fundamental problem. Neither Stell's equation nor equation (2) give the correct second virial coefficient for a system of dipolar hard spheres, as determined by Keesom (1912). It is possible to relate the function $f^{(1)}$ to this theoretical value by: $f^{(1)} = B^{\text{theo}} - 3$, where B^{theo} is the reduced theoretical second virial coefficient ($B^{\text{theo}} = 3B/2\pi\sigma^3$), which is equal to 4 for non-polar hard spheres and decreases with increasing dipole moment. If $f^{(1)}$ is determined in this way, agreement between experimental second virial coefficients and those calculated from the resulting equation is excellent, even for acetonitrile. However, since the virial series converges slowly or not at all for liquid-like densities, use of the theoretical second virial coefficient leads to serious problems in the calculation of high-density fluid properties. In this work, it was considered more important to represent accurately the high-density behavior. Therefore, Keesom's result was not used in the development of the DDCS equation.

Figure 1



Isotherms for a Dipolar-Hard-Sphere Fluid.

Comparison of Dipole-Dipole Carnahan-Starling Equation to Stell's Equation

NOTATION

- a = equation-of-state constant reflecting attractive forces
- b = equation-of-state constant reflecting repulsive forces
- B = second virial coefficient
- k = Boltzmann constant
- P = pressure
- R = gas constant
- T = absolute temperature
- v = molar volume
- z = compressibility factor

Greek Letters


- α, β = constants reflecting temperature dependence of parameter a
- ρ = molar density
- Φ = function providing smooth transition for temperature dependence of parameter a between super- and sub-critical temperatures
- η = $bp/4$ = dimensionless density
- σ = hard-sphere diameter
- μ = dipole moment

Superscripts

- ref = reference part of equation of state
- pert = perturbation part of equation of state
- theo = theoretically calculated
- expt = experimentally determined
- HS = hard sphere
- = dimensionless


Subscript

- c = critical




Isotherms for a Dipolar-Hard-Sphere Fluid
Calculated by Modified Carnahan-Starling
Equation. A Phase Transition occurs when
the Reduced Dipole Moment is at Least 1.90.

Figure 3




Calculated and Observed Ammonia Isotherms

Figure 4




Calculated and Observed Vapor Pressures



Calculated and Observed Saturated-Liquid
Molar Volumes

Figure 6



Calculated and Observed Second Virial Coefficients

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*