

Lawrence Berkeley National Laboratory

LBL Publications

Title

Interferometric measurements of many-body topological invariants using mobile impurities.

Permalink

<https://escholarship.org/uc/item/0wm996r1>

Authors

Abanin, D

Fleischhauer, M

Demler, E

et al.

Publication Date

2016-06-17

DOI

10.1038/ncomms11994

Peer reviewed

ARTICLE

Received 23 Dec 2015 | Accepted 16 May 2016 | Published 17 Jun 2016

DOI: 10.1038/ncomms11994

OPEN

Interferometric measurements of many-body topological invariants using mobile impurities

F. Grusdt^{1,2,3}, N.Y. Yao⁴, D. Abanin⁵, M. Fleischhauer¹ & E. Demler³

Topological quantum phases cannot be characterized by Ginzburg-Landau type order parameters, and are instead described by non-local topological invariants. Experimental platforms capable of realizing such exotic states now include synthetic many-body systems such as ultracold atoms or photons. Unique tools available in these systems enable a new characterization of strongly correlated many-body states. Here we propose a general scheme for detecting topological order using interferometric measurements of elementary excitations. The key ingredient is the use of mobile impurities that bind to quasiparticles of a host many-body system. Specifically, we show how fractional charges can be probed in the bulk of fractional quantum Hall systems. We demonstrate that combining Ramsey interference with Bloch oscillations can be used to measure Chern numbers characterizing the dispersion of individual quasiparticles, which gives a direct probe of their fractional charges. Possible extensions of our method to other many-body systems, such as spin liquids, are conceivable.

¹Department of Physics and Research Center OPTIMAS, University of Kaiserslautern, Kaiserslautern 67663, Germany. ²Graduate School Materials Science in Mainz, Gottlieb-Daimler-Strasse 47, 67663 Kaiserslautern, Germany. ³Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA. ⁴Department of Physics, University of California, Berkeley, California 94720, USA. ⁵Department of Theoretical Physics, University of Geneva, 24 quai Ernest-Ansermet, 1207 Geneva, Switzerland. Correspondence and requests for materials should be addressed to F.G. (email: fgrusdt@physics.harvard.edu).

Many-body systems with spontaneous symmetry breaking can be described by Ginzburg–Landau theories, formulated in terms of local order parameters. This powerful approach provides a universal description of systems with very different microscopic Hamiltonians but with similar type of symmetry breaking, such as superfluids and ferromagnets. The integer and fractional quantum Hall effects^{1–3} in contrast are examples of quantum phases of matter, for which no local order parameters exist. Instead, these systems are described by non-local topological invariants⁴. The fractional charges of elementary excitations^{2,3}, the many-body Chern number \mathcal{C} (ref. 5) and, in the case of quantum spin liquids, fractional quantum Hall systems and fractional Chern insulators^{6–13}, the groundstate degeneracy on a torus¹⁴, constitute important examples of topological order parameters.

Probing—and, in some cases, even defining—the non-local order parameters of topological systems with strong correlations represents a major experimental and theoretical challenge. Many indicators used in the theoretical description of such systems, such as the entanglement entropy^{15,16} and spectrum¹⁷, are difficult to probe directly in current experiments, although first steps in this direction have been undertaken^{18–21}. Previously, it has been shown that edge excitations can be used to detect topological orders by measuring their fractional charges^{4,22–24} and statistics⁴.

Motivated by the coherent control of mesoscopic quantum systems achievable in recent experiments, we explore new ways of detecting topological order. Our method is ideally suited to systems of ultracold atoms, which recently emerged as a new promising platform for realizing and probing various topological states of matter. The ability to perform interferometric measurements in such systems is one of their key technical advantages in comparison with other experimental set-ups. Cold atoms provide a versatile toolbox, allowing to engineer not only single-particle properties of Hamiltonians, such as the shape of optical lattice potentials, but also the interactions between particles^{25–28}. Recently, the Chern number has been measured in transport experiments²⁹ and the celebrated Haldane model has been realized³⁰ in systems of weakly interacting ultracold atoms. An experimental realization of the fractional quantum Hall effect in such systems^{6,7,31–36} should be within reach with the currently available tools. In addition, direct and fully coherent control over individual atoms has been demonstrated in experiments with ultracold quantum gases, see, for example, refs 37,38.

Here we present a concept of using impurity atoms as coherent probes of the topological invariants of strongly correlated many-body systems of host atoms. Our approach allows one to measure topological order parameters directly in the bulk of the system, without the need of relying on the bulk-edge correspondence. The main idea is to map out the topology characterizing the effective bandstructure of elementary quasi-particle (qp) excitations. As will be pointed out, it is intimately related to the topological order of the groundstate. In particular, we show how the Chern numbers of the effective qp bandstructures can be measured by combining Bloch oscillations with a Ramsey interferometric sequence. We point out that they are directly related to the corresponding (fractional) charges for arbitrary Abelian quantum Hall states, and show how the Chern number of the many-body groundstate can be derived. Our scheme extends earlier ideas^{39–42}, which have been developed to measure topological invariants of essentially non-interacting particles (ultracold atoms in particular), to the realm of strongly correlated quantum many-body systems. More generally, our interferometric method paves the way for a detailed investigation of qp properties, including, possibly, their braiding statistics.

Results

Interferometric detection of many-body Chern numbers. The key idea of our approach is to measure the Chern number of the effective qp bandstructure using a generalization of the interferometric technique developed for non-interacting systems in refs 39,40. First, let us briefly summarize the main idea of the interferometric protocol for a weakly interacting Bose–Einstein condensate loaded in a two-dimensional (2D) Bloch band in a system with an effective magnetic field. Because of their experimental relevance for ultracold atoms we will discuss lattice systems. The first ingredient to measure the corresponding Chern number \mathcal{C} is a direct detection of the geometric Zak phase $\varphi_{\text{Zak}}(k_y)$ ³⁹ for a given value of the quasimomentum k_y .

To this end, the condensate is moved to (k_x, k_y) , where a $\pi/2$ Ramsey pulse is used to prepare a superposition of two internal states $\sigma = \uparrow, \downarrow$. The initial wavefunction thus reads $(|\uparrow\rangle + |\downarrow\rangle) \otimes |k_x, k_y\rangle/\sqrt{2}$. Next, sufficiently weak opposite forces $\mathbf{F}(\downarrow) = -\mathbf{F}(\uparrow)$ are applied such that the two components undergo Bloch oscillations (Fig. 1b). When the force is applied along the x axis for a time $\Delta t = G_x/2F$, where G_x denotes the width of the magnetic Brillouin zone (BZ) containing an integer number of magnetic flux quanta, the two components pick up a relative geometric phase $\varphi_{\text{Zak}}(k_y)$ (ref. 43) and the wavefunction reads $(|\uparrow\rangle + e^{i\varphi_{\text{Zak}}(k_y)}|\downarrow\rangle) \otimes |G_x/2, k_y\rangle/\sqrt{2}$. By recombining the two states using a second $\pi/2$ Ramsey pulse, the Zak phase can be read out. This measurement can be repeated for different values of k_y (ref. 40), and the winding of the Zak phase across the




Figure 1 | Topological polarons. We propose a scheme for the measurement of many-body topological invariants of interacting states with topological order. It can be applied to measure the Chern numbers of Abelian quantum Hall states (for example, of Laughlin type) and of their topological excitations, as illustrated in **a**. An elementary excitation (here a quasihole) is coupled to a mobile two-component impurity. When the impurity is tightly bound to the quasihole, a topological polaron is formed. It can be labelled by its quasimomentum q , and its two internal degrees of freedom \uparrow, \downarrow and can be used to perform interferometry. The spectrum $\epsilon(q_x)$ of the many-body system is depicted in **b** for a generic one-dimensional case. The topological invariant of the qp bandstructure can be measured using tools developed for non-interacting systems by a combination of Bloch oscillations and Ramsey interferometry.

magnetic BZ in the k_y direction (size G_y) gives the Chern number⁴⁴,

$$C = \frac{1}{2\pi} \int_0^{G_y} dk_y \partial_{k_y} \varphi_{\text{Zak}}(k_y). \quad (1)$$

For more details, including the discussion of dynamical phases and gauge dependence of the Zak phase, the reader is referred to refs 39–42.

Now we extend the interferometric protocol to strongly correlated systems. The key idea is to apply the exact same sequence as described above to a single qp excitation in the host system. To obtain fully coherent control of the qp, it is coupled to a mobile impurity. The resulting composite object (the impurity bound to the topological excitation) will be called a topological polaron (TP), which is at the heart of our scheme. (We call this bound state a TP because, as will be shown, it inherits the topological properties of the qp excitation.) We assume that the impurity has an internal degree of freedom σ , the only role of which is to enable Ramsey interferometry as described above. By introducing two pseudospin components $\sigma = \uparrow, \downarrow$ a measurement of their relative phase is sufficient, that is, one of the pseudospin states effectively acts as a reference for the second. By applying opposite forces $\sigma^z F e_x$ directly to the two impurity components for the same time $\Delta t = G_x/2F$ as in the non-interacting case (where G_x is defined for host particles), a topological invariant can be measured, which will be identified as the Chern number C_{TP} of the effective TP bandstructure (for more details see Methods section). Note that now $|k_x, k_y\rangle$ stands for the full many-body wavefunction describing the TP at quasimomentum (k_x, k_y) , and in this sense true many-body Zak phases $\varphi_{\text{Zak}}(k_y)$ are measured.

Now our main results are summarized. We will establish that the Chern number of the TP bandstructure is directly related to the fractional charges of the qps. We will use Chern–Simons effective field theory, valid at low energies, for the description of Abelian fractional quantum Hall states and their excitations. In addition, we introduce a numerically exact technique for calculating Zak phases of TPs for small lattice systems relevant for experiments. As a concrete example, Laughlin-type fractional Chern insulator states at filling fraction $\nu = 1/m$ (defined as a ratio of particle density to flux density) will be considered. For that case, we show that the TP Chern number, discussed in the protocol above, is given by the inverse of the fractional qp charge $e^* = e/m$ (e is the charge of particles in the host many-body system),

$$C_{\text{TP}} = \frac{e}{e^*} = m. \quad (2)$$

Therefore, a measurement of the TP Chern number directly yields the fractional qp charge.

The result in equation (2), moreover, indicates a direct relation to the many-body Chern number $C_0 = 1/m$ (ref. 5) of the incompressible $\nu = 1/m$ Laughlin groundstate, $C_0 = 1/C_{\text{TP}}$. (By a fractional Chern number $C = q/p$, with q, p integers, we refer to a situation where p degenerate states on a torus share a total Chern number q , giving rise to a fractionally quantized Hall conductivity $\sigma_{xy} = Ce^2/h$ (ref. 5). q is an integer topological invariant characterizing the vector bundle defined by the set of degenerate states on a torus.) Similar relations are true for any Abelian quantum Hall state, where e/e^* needs to be expressed in terms of the K -matrix. In that case there exist $J = 1, \dots, n$ different qp types, with fractional charges e_j^* and qp Chern numbers $C_{\text{TP}}^{(j)} = e/e_j^*$. The Chern number of the incompressible fractional quantum Hall state is given by $C = \sum_{j=1}^n 1/C_{\text{TP}}^{(j)}$ (see Methods section).

TPs and qp Chern numbers. Our discussion of TPs will be organized as follows. In this section we focus on the physical picture of TPs in fractional quantum Hall systems. We discuss the origin of equation (2), assuming that an impurity particle is strongly bound to a qp. Therefore, the impurity current j_μ^σ (where $\mu = x, y, t$ and $\sigma = \uparrow, \downarrow$) is equivalent to the qp current, which is conserved because of a global $U(1)$ symmetry. Detailed discussion of the binding mechanism between the impurity and a qp will be presented in the next subsection.

As in the theoretical proposal of ref. 40 and the experimental demonstration of ref. 39, we assume that the two-component TP can be controlled by the force \mathbf{F} acting on the impurity. (In experiments a spin-dependent force \mathbf{F} could be realized by a magnetic field gradient.) We model this by coupling the TP current j_μ^σ to an external field that acts differently on the two internal states of the impurity but does not affect host particles. We describe this field by the potential $B_\mu(\sigma)$ so that the external force acting on the impurity is given by $F_i(\sigma) = -q(\partial_i B_i + \partial_i B_0)$, where q denotes the charge of the impurity associated with the field B_μ . We emphasize that the impurity is not affected directly by the gauge field creating a fractional quantum Hall state for host particles. However, by binding a qp of the surrounding fractional quantum Hall state it acquires topologically non-trivial dynamics. This can be understood as arising from the particle–vortex duality of the fractional quantum Hall effect, which implies that the moving qp bound to the impurity sees surrounding host particles as a source of effective flux.

To avoid complications arising because of an underlying lattice, we first employ an effective field theory to capture the universal topological properties of our system (in particular, Chern numbers). It is valid when all relevant length scales arising from gauge fields and interactions are large compared with the lattice constant. Using this Chern–Simons description of the fractional quantum Hall effect one can show (see Methods section for details) that the effective Lagrangian describing the TP is given by (throughout the paper we set $\hbar = 1$)

$$\mathcal{L}_{\text{TP}} = \sum_{\sigma=\uparrow,\downarrow} (qB_\mu(\sigma) + e^* A_\mu) j_\mu^\sigma. \quad (3)$$

Here A_μ denotes the external gauge field corresponding to the homogeneous magnetic field b_z seen by the host particles in the many-body quantum Hall system, $\partial_x A_y - \partial_y A_x = b_z$. In the case of ultracold atoms, b_z corresponds to an artificial gauge field⁴⁵. The second term in the Lagrangian (3) describes the emergent coupling of the TP to the gauge field A_μ , albeit with a fractional A_μ charge $e^* = e/m$. Hence, according to equation (3) the TP sees an effective magnetic field $b_z^* = b_z e^*/e$. Note that we consider a situation where the host system has a global $U(1)$ symmetry, giving rise to conserved charges (or particle numbers in the context of ultracold atoms).

We emphasize again that the emergent coupling of impurity particles to the gauge field A_μ arises only through binding of a qp excitation. For Laughlin states, a qp acquires an Aharonov–Bohm phase 2π when going around a single host particle. Hence, the effective flux density seen by the qp is equal to the density of host particles, which is $1/m$ times the magnetic flux density of A_μ .

Now the Chern number C_{TP} of the TP can easily be calculated. To this end we note that, in a homogeneous magnetic field b_0 , the Berry curvature \mathcal{F} is constant, $\mathcal{F} = 1/(eb_0)$. Similar to the manner in which we described the interferometric protocol above, we defined the TP Chern number by integrating the Berry curvature over the entire magnetic BZ of the host many-body system. Its size is $G_x \times$

$G_y = 2\pi e b_z$, where b_z is the magnetic field seen by host particles. Because the Berry curvature seen by the TP is $\mathcal{F}_{\text{TP}} = 1/(e b_z^*)$ we obtain

$$C_{\text{TP}} = \frac{1}{2\pi} \int_{\text{BZ}} d^2k \mathcal{F}_{\text{TP}} = \frac{b_z}{b_z^*} = \frac{e}{e^*}, \quad (4)$$

as claimed in equation (2).

Note that because of gauge invariance $C_{\text{TP}} \in \mathbb{Z}$ has to be an integer^{46,47}, as long as the TP groundstate is non-degenerate for all k (see also Methods section). Therefore, when the qp charge is not the inverse of an integer, the band of TP groundstates needs to have degeneracies. When $e^*/e = p/q$, where p and q are relative prime, we expect p degenerate bands sharing a total Chern number of $C_{\text{TP}}^{\text{tot}} = q$. We find numerical signatures for this degeneracy, and a specific example is discussed in the Supplementary Note 1.

Alternatively, the qp charge e^*/e of the TP could be obtained directly from an interferometric measurement of the Aharonov–Bohm phase $\Phi_{\text{AB}} = 2\pi \frac{e^*}{e} A b_z / \Phi_0$ in the magnetic field b_z . Here A denotes the area encircled by the TP and Φ_0 is the magnetic flux quantum. Because obtaining control over the impurity momentum is technically less challenging than achieving full spatial resolution, the interferometric measurement of the Chern number should be easier to implement.

Microscopic description of TPs. Now we provide a microscopic description of TPs in correlated many-body systems interacting with a single impurity particle. To this end, we investigate concrete models of interacting fermions on a lattice, whose groundstates are integer and fractional Chern insulators^{6–12}. We develop an exact numerical method to solve this problem and compare our results to an approximate strong coupling theory of TPs, which we also introduce below. Like before, we concentrate on states in the Laughlin universality class, with a single qp type. Although the field theory employed so far assumed a continuous system in a homogeneous magnetic field, it should describe correctly the universal topological properties of the lattice systems, which will be investigated below.

We consider a topologically ordered phase in a many-body system (referred to as the host system), described by the groundstate of a Hamiltonian $\hat{\mathcal{H}}_0$. For probing qp excitations, we introduce a mobile impurity with two internal states, described by $\hat{\mathcal{H}}_1 = \hat{\mathcal{H}}_1^0 - \hat{\sigma}^z \mathbf{F} \cdot \hat{\mathbf{r}}$. Here $\hat{\mathcal{H}}_1^0$ denotes the kinetic energy and $\hat{\mathbf{r}}$ the position operator of the impurity. The two internal states \uparrow and \downarrow of the impurity experience opposite forces $\pm \mathbf{F}$, as described by the Pauli matrix $\hat{\sigma}^z$. To bind qps to the impurity, a local interaction $\hat{\mathcal{H}}_{\text{int}}$ with the many-body system is introduced. Its concrete form can differ from model to model; however, for simplicity we will assume throughout that it is independent of the internal state of the impurity. Thus, our system is described by the Hamiltonian $\hat{\mathcal{H}} = \hat{\mathcal{H}}_0 + \hat{\mathcal{H}}_1 + \hat{\mathcal{H}}_{\text{int}}$. Local interactions between particles with multiple internal states are routinely realized in systems of ultracold atoms.

In equilibrium, that is, for $\mathbf{F} = 0$, the groundstate $|\psi_{\text{TP}}(\mathbf{q}, \sigma)\rangle$ describes a TP and can be labelled by its quasimomentum \mathbf{q} and its internal state $\sigma = \uparrow, \downarrow$. The external force \mathbf{F} couples to the quasimomentum \mathbf{q} of the TP. This drives Bloch oscillations where the quasimomentum changes according to $\frac{d}{dt} \mathbf{q} = \sigma^z \mathbf{F}$, where $\sigma^z = 1$ ($\sigma^z = -1$) for $\sigma = \uparrow$ ($\sigma = \downarrow$), respectively). By applying the scheme described in ref. 39 to the states $|\psi_{\text{TP}}(\mathbf{q}, \sigma)\rangle$ the geometric Zak phases characterizing the TP bandstructure can be measured (Fig. 1). As discussed in the Methods section,

the corresponding Chern number is obtained by integrating the Berry curvature seen by the TP over the magnetic BZ of the host many-body system.

Strong coupling theory. Before turning our attention to concrete models, we introduce the strong coupling theory of TPs, which is inspired by Landau’s and Pekar’s treatment of the polaron problem in polar crystals^{48,49}. There are several important energy scales for describing TPs. First, qp excitations are characterized by the bandwidth J_{qp} of their effective dispersion and by the energy required for their creation, which is set by the bulk excitation gap Δ_0 . Second, the impurity is characterized by the effective hopping J_I and the coupling strength to the host particles V . Our description of TPs requires the following hierarchy of scales. The qp gap Δ_0 , which is larger than J_{qp} , should be larger than J_I and V , that is, $\Delta_0 \gg J_I, V$. The impurity–host particle interaction strength V should be chosen such that the impurity binds precisely one qp. We also need J_{qp} to be smaller than J_I , that is, $J_{\text{qp}} \ll J_I$ so that the total momentum of the TP, which is effectively shared by the impurity and the qp, resides predominantly in the qp. Another way of understanding this requirement is that the impurity should be fast compared with the qp and thus follow its dynamics adiabatically.

Next, we introduce a frame where the total momentum of the TP is conserved explicitly. To this end, we restrict ourselves to a single qp and approximate $\hat{\mathcal{H}}_0 \approx \sum_{\mathbf{k}} \epsilon_{\text{qp}}(\mathbf{k}) |\psi_{\text{qp}}(\mathbf{k})\rangle \langle \psi_{\text{qp}}(\mathbf{k})|$, with $\epsilon_{\text{qp}}(\mathbf{k})$ being the effective dispersion of qps and $|\psi_{\text{qp}}(\mathbf{k})\rangle$ the qp state with momentum \mathbf{k} . We apply the unitary transformation $\hat{U}_{\text{LLP}} = e^{i\hat{\mathbf{p}} \cdot \hat{\mathbf{R}}_{\text{qp}}}$ introduced by Lee, Low and Pines (LLP)⁵⁰, where $\hat{\mathbf{p}}$ is the impurity momentum operator and $\hat{\mathbf{R}}_{\text{qp}}$ is qp position operator conjugate to its momentum operator $\hat{\mathbf{P}}_{\text{qp}} = \sum_{\mathbf{k}} \mathbf{k} |\psi_{\text{qp}}(\mathbf{k})\rangle \langle \psi_{\text{qp}}(\mathbf{k})|$. The transformed Hamiltonian $\hat{U}_{\text{LLP}}^\dagger \hat{\mathcal{H}} \hat{U}_{\text{LLP}}$ reads

$$\hat{\mathcal{H}}_{\text{SC}} = \hat{\mathcal{H}}_1^0 + \hat{\mathcal{H}}_{\text{int}}(\hat{\mathbf{r}}, \hat{\mathbf{R}}_{\text{qp}} = 0) - \sigma^z \mathbf{F} \cdot (\hat{\mathbf{r}} + \hat{\mathbf{R}}_{\text{qp}}) + \sum_{\mathbf{k}} \epsilon_{\text{qp}}(\mathbf{k} - \hat{\mathbf{p}}) |\psi_{\text{qp}}(\mathbf{k})\rangle \langle \psi_{\text{qp}}(\mathbf{k})|, \quad (5)$$

as will be explained now.

The kinetic part of the impurity Hamiltonian commutes with the impurity momentum, $[\hat{\mathbf{p}}, \hat{\mathcal{H}}_1^0] = 0$, and remains unchanged. $\hat{\mathcal{H}}_{\text{int}}(\hat{\mathbf{r}}, \hat{\mathbf{R}}_{\text{qp}} = 0)$ denotes the interaction Hamiltonian for a qp localized in the origin of the new polaron frame, that is, $\hat{\mathbf{R}}_{\text{qp}} = 0$. Here we assumed that (within the single qp approximation) $\hat{\mathcal{H}}_{\text{int}}(\hat{\mathbf{r}}, \hat{\mathbf{R}}_{\text{qp}}) = \hat{\mathcal{H}}_{\text{int}}(\hat{\mathbf{r}} - \hat{\mathbf{R}}_{\text{qp}})$ depends only on the relative distance between impurity and qp, and we used that $\hat{U}_{\text{LLP}}^\dagger \hat{\mathbf{r}} \hat{U}_{\text{LLP}} = \hat{\mathbf{r}} + \hat{\mathbf{R}}_{\text{qp}}$. Finally, because $[\hat{\mathbf{R}}_{\text{qp}}, \hat{\mathbf{P}}_{\text{qp}}] = i$ the qp momentum is shifted by an amount $\hat{\mathbf{p}}$ in the last line, $\mathbf{k} \rightarrow \mathbf{k} - \hat{\mathbf{p}}$.

Under the strong coupling conditions outlined above, we can make a product ansatz for the TP wavefunction, $|\psi_{\text{TP}}(\mathbf{q})\rangle = |\psi_{\text{qp}}(\mathbf{q})\rangle \otimes |\phi_I\rangle$, where the impurity follows the qp adiabatically. In the polaron frame the impurity sees a quasistatic potential created by the qp and its wavefunction $|\phi_I\rangle$ is determined by the strong coupling impurity Hamiltonian

$$\hat{\mathcal{H}}_{\text{I,SC}} = \hat{\mathcal{H}}_1^0 - \sigma^z \mathbf{F} \cdot \hat{\mathbf{r}} + \hat{\mathcal{H}}_{\text{int}}(\hat{\mathbf{r}}, \mathbf{R}_{\text{qp}} = 0). \quad (6)$$

This leads to a modification of the effective qp dispersion in equation (5), which we approximate by $\langle \phi_I | \epsilon_{\text{qp}}(\mathbf{k} - \hat{\mathbf{p}}) | \phi_I \rangle$. We proceed by eliminating the last term in the first line of equation (5) by a time-dependent gauge transformation $\hat{U}_{\text{qp}}(t) = e^{i\hat{\mathbf{R}}_{\text{qp}} \cdot \hat{\sigma}^z \mathbf{F} t}$. In the resulting effective Hamiltonian the

force \mathbf{F} couples to the conserved momentum \mathbf{k} and the qp Hamiltonian in the strong coupling theory thus reads

$$\hat{\mathcal{H}}_{\text{qp,SC}} = \sum_{\mathbf{k}} \langle \phi_1 | \epsilon_{\text{qp}}(\mathbf{k} - \hat{\sigma}^z \mathbf{F}t - \hat{\mathbf{p}}) | \phi_1 \rangle | \psi_{\text{qp}}(\mathbf{k}) \rangle \langle \psi_{\text{qp}}(\mathbf{k}) |. \quad (7)$$

Next, we discuss how the topological invariant measured by the TP relates to the Chern number of the effective qp bandstructure using the strong coupling wavefunction. During the protocol described above (see also Fig. 1) the force $\hat{\sigma}^z \mathbf{F}$ is applied and the TP wavefunction $|\psi_{\text{TP}}(\mathbf{q}, t)\rangle = e^{i\varphi} |\psi_{\text{qp}}(\mathbf{q} - \hat{\sigma}^z \mathbf{F}t)\rangle \otimes |\phi_1\rangle$ follows its groundstate adiabatically. Thereby, it picks up a phase φ containing a geometric part of $2\pi\nu_{\text{TP}}$, which is measured by the Ramsey sequence (dynamical phases are discussed in refs 39,40). Owing to the product form of the strong coupling wavefunction we find two contributions, $\nu_{\text{TP}} = \nu_{\text{qp}} + \nu_1$.

The first contribution is picked up by the qp wavefunction, $2\pi\nu_{\text{qp}} = \oint d\mathbf{k} \cdot \langle \psi_{\text{qp}}(\mathbf{k}) | i\nabla_{\mathbf{k}} | \psi_{\text{qp}}(\mathbf{k}) \rangle$. When the path in momentum space described by the TP in the interferometer encloses the (magnetic) BZ, the qp invariant is related to the TP Chern number defined above,

$$\nu_{\text{qp}} = \mathcal{C}_{\text{TP}}. \quad (8)$$

The second contribution ν_1 is picked up by the impurity part of the wavefunction. In the adiabatic limit of small \mathbf{F} it is $2\pi\nu_1 = -\sigma^z \oint dt \mathbf{F} \cdot \langle \phi_1 | \hat{\mathbf{r}} | \phi_1 \rangle$. This term corresponds to a geometric phase because it does not vanish in the limit when $\mathbf{F} \rightarrow 0$ and needs to be considered in general. It measures the displacement of the impurity wavefunction relative to the qp located in the origin of the polaron frame (recall that $\mathbf{R}_{\text{qp}} = \mathbf{0}$ in the interaction Hamiltonian) and can become relevant in lattice systems. However, when a closed loop $\oint dt \mathbf{F} = 0$ is considered as in the interferometric sequence we discuss, the impurity invariant vanishes, $\nu_1 = 0$.

Tps in Chern insulators. Now we turn our attention to a concrete model of interacting particles on a lattice, described by the Hofstadter–Hubbard Hamiltonian

$$\hat{\mathcal{H}}_0 = -t \sum_{m,n} [e^{-i2\pi\alpha n} \hat{c}_{m+1,n}^\dagger \hat{c}_{m,n} + \hat{c}_{m,n+1}^\dagger \hat{c}_{m,n} + \text{h.c.}] + U \sum_{\langle(m,n),(m',n')\rangle} \hat{c}_{m,n}^\dagger \hat{c}_{m,n} \hat{c}_{m',n'}^\dagger \hat{c}_{m',n'}. \quad (9)$$

The first term is the celebrated Hofstadter model⁵¹, and it describes free particles hopping between the sites (m, n) of a square lattice in a magnetic field (using Landau gauge), where α

denotes the magnetic flux density per plaquette (in units of the flux quantum) and t is the hopping amplitude. The second term describes nearest neighbour interactions of strength U between the particles. Here we consider fermions for concreteness, $\{\hat{c}_{m,n}, \hat{c}_{m',n'}^\dagger\} = \delta_{m,m'} \delta_{n,n'}$, but a similar Hofstadter–Hubbard model has also been discussed for bosons with contact interactions^{6,7}. For sufficiently small values of α , the groundstates of equation (9) show the *integer and fractional quantum Hall effect* depending on the filling fraction ν .

To study TPs we consider the Hofstadter–Hubbard model equation (9) at filling $\nu = 1/m$ on a torus. We choose the number of flux quanta in the host many-body system $N_\phi = Nm + 1$ such that the groundstate of $\hat{\mathcal{H}}_0$ contains one quasihole (qh) excitation. Next, we add a single impurity, described by $\hat{b}_{m,n}^\dagger$, hopping between the sites (located at positions $\mathbf{r}_{m,n}$) of the same 2D lattice,

$$\hat{\mathcal{H}}_1 = -J \sum_{m,n} [\hat{b}_{m+1,n}^\dagger \hat{b}_{m,n} + \hat{b}_{m,n+1}^\dagger \hat{b}_{m,n} + \text{h.c.}] - \mathbf{F} \cdot \sum_{m,n} \mathbf{r}_{m,n} \hat{b}_{m,n}^\dagger \hat{b}_{m,n}. \quad (10)$$

To bind the impurity to the qh, its interaction with the surrounding fermions is modelled by a repulsive contact potential, $\hat{\mathcal{H}}_{\text{int}} = V \sum_{m,n} \hat{c}_{m,n}^\dagger \hat{c}_{m,n} \hat{b}_{m,n}^\dagger \hat{b}_{m,n}$. In the following we consider an impurity with only a single internal state for simplicity.

The model proposed above (equations (9) and (10)) can be implemented with ultracold atoms. In refs 52,53 the Hofstadter Hamiltonian was realized for bosons, and interactions can be introduced by using deep optical lattices. The impurity could be realized by adding a second atomic species, with different metastable internal states.

In the Methods section we present a formalism for calculating the full many-body TP wavefunction $|\psi_{\text{TP}}(\mathbf{q})\rangle$ exactly for a given total momentum \mathbf{q} , based on the LLP polaron transformation. Here we use this approach to obtain both the dispersion relation and the Chern number of TPs.

In Fig. 2a our results are shown for an integer Chern insulator, with and without fermion–fermion interactions U . We consider the case $\alpha = 1/4$ and use Landau gauge where the size of the magnetic unit cell is $(a_x = 4a) \times a$, with a denoting the lattice constant. We calculate the Chern number of the TP from the winding of the many-body Zak phase, $\mathcal{C} = \frac{1}{2\pi} \int_{-\pi/a}^{\pi/a} dk_y \partial_{k_y} \varphi_{\text{Zak}}(k_y)$ ⁴⁴, see Methods section for details. For comparison, the result of a simple strong coupling analysis is shown, where the impurity is bound to a free hole. Although the




Figure 2 | Many-body Chern numbers. A single mobile impurity can be coupled to an elementary hole excitation of an integer (a) or fractional (b) Chern insulator, and form a TP. The winding of the many-body Zak phase $\varphi_{\text{Zak}}(k_y)$ across the BZ defines the many-body Chern number of the TP. In a we compare predictions for integer Chern insulators without ($U = 0$) and with interfermion interactions ($U \neq 0$). Parameters are $J = t/2$, $V = 2t$, $\alpha = 1/4$ and we simulated 4×4 sites with $N = 3$ fermions. In b, a $\nu = 1/3$ fractional Chern insulator is considered, for parameters $J = t/2$, $V = 2t$, $\alpha = 1/4$, $U = 10t$. We simulated 4×7 sites filled with $N = 2$ fermions, corresponding to $3N + 1 = 7$ flux quanta as required for having one quasihole excitation.

distribution of Berry curvature differs, the strong coupling theory predicts correctly the TP Chern number, $\mathcal{C}=1$.

In Fig. 2b we repeat our calculations for a fractional Chern insulator at $\nu=1/3$, corresponding to a $1/3$ Laughlin-type state. (We checked numerically that the incompressible groundstate of $\hat{\mathcal{H}}_0$ has a fractional Chern number $\mathcal{C}_0=1/3$ and the expected threefold groundstate degeneracy on a torus.) The Chern number of the TP, which is determined as the winding of the Zak phase of the TP in Fig. 2b, has the value $\mathcal{C}=3$. This is expected from the field-theoretical arguments given in the beginning (equation (2)).

An important quantity for estimating the robustness of our protocol is the energy gap of the TP state. To get some analytical understanding, we first consider the continuum limit $\alpha \rightarrow 0$ when the system enters the quantum Hall regime. Here the impurity (mass M_I) interacts with fermions (mass M_F) in a filling- ν Laughlin state through a local contact interaction $g_{IF}\delta^{(2)}(\mathbf{r})$. Within the strong coupling theory introduced above, we find a finite TP-binding energy given by

$$\Delta_{\text{TP}} = \sqrt{\nu \frac{g_{IF}}{2\pi\ell_B^2} \frac{M_F}{M_I} \omega_c}, \quad (11)$$

where $\ell_B=1/\sqrt{eb_0}$ is the magnetic length and $\omega_c=1/(M_F\ell_B^2)$ the cyclotron frequency.

In Fig. 3 we calculate the TP spectra numerically for small lattice systems and find low-energy states where the impurity is bound to the qh forming the TP. We observe that the TP gap Δ_{TP} shows large finite-size effects. The TP-binding energy oscillates at a frequency given by the system size, but at the discrete momenta allowed by the finite quantization volume—which are indicated in the figures by circles—we find the largest values of the TP gap. In the case of integer Chern insulators, this TP gap constitutes a sizable fraction of the cyclotron gap $\Delta_{\text{TP}}\omega_c \approx 4\pi\alpha t$. Note that ω_c provides an upper bound for the TP gap, $\Delta_{\text{TP}} < \omega_c$, because at this energy unbound excitons can be created in the bulk. In the case of fractional Chern insulators in contrast, the TP gap is only a small fraction of the cyclotron energy, but so is the bulk gap Δ_{LN} of the fractional Chern insulator, $\Delta_{\text{LN}} < \omega_c$ (ref. 7). In this case Δ_{LN} similarly provides an upper bound for the TP gap as the cyclotron energy in the integer Chern insulator case, $\Delta_{\text{TP}} < \Delta_{\text{LN}}$.

To form a TP for mapping out the topological order of the host many-body system, we wish to realize the strong coupling limit described above. Thus, for impurity dynamics to be fast compared with qp tunnelling, a weak qp dispersion is desirable. This can be achieved in lattice systems by including long-range hoppings for the host particles, see ref. 54. We simulated TPs in systems with long-range tunnellings of this kind (see also

Supplementary Note 1) and verified that the TP Chern number $\mathcal{C}=3$ is a robust feature of the groundstates.

Discussion

By coupling a mobile impurity to the topological qp excitations of an incompressible many-body groundstate of host particles, a TP can be formed. Using internal degrees of freedom of the impurity, fully coherent control can be gained over individual qp excitations of the host many-body system. We demonstrated that TPs can be used to measure the topological invariants characterizing such qp excitations. In particular, we developed an interferometric measurement scheme for qp Chern numbers and showed that these are directly related to the fractional charges of topological excitations in two dimensions. To this end, we generalized schemes developed earlier for non-interacting systems and showed by explicit calculations that our scheme can be applied to integer- and fractional quantum Hall systems and Chern insulators.

In systems of ultracold atoms, impurities can be realized, for example, by introducing another atomic species and their internal degree of freedom corresponds to different hyperfine states. To neglect the effects of interactions between TPs, the density of impurity atoms needs to be well below one atom per cyclotron orbit. We expect that, in order to realize TPs in an experiment, a main challenge will be their preparation. Because TPs can carry fractional charges as in the case of fractional Chern insulators, their preparation requires non-local operations in general. Therefore, we suggest to build impurity atoms into the state as defects already before preparing the incompressible many-body state of host atoms. One concrete approach would be to start from this system and cool it down into its groundstate. If the densities of majority and impurity atoms are suitably chosen, this leads to the formation of TPs. Alternative approaches have also been discussed in the literature^{55–58}.

The method presented in this article can provide a powerful experimental tool for detecting topological order in interacting many-body systems. It is ideally suited for cold atom experiments, which offer a rich toolbox with precise and fully coherent control over individual atoms. The concept of TPs as experimental probes for topological order can be generalized to other systems, however, including, for example, qp excitations on topological superconductors, or systems with symmetry-protected topological orders. Another interesting direction would be to probe the (non-Abelian) braiding statistics of anyons by coupling them to impurities and forming TPs. In this case, one can envision interferometric sequences designed to probe braiding statistics, which work in real-space rather than momentum space, as considered in this work.




Figure 3 | TP spectrum. The energies of the lowest TP eigenstates are shown for $k_y=0$ as a function of the lattice moment k_x , calculated in the polaron frame (see Methods section). In **a** an integer Chern insulator is considered for $U=10t$, $J=t/2$, $V=2t$, $\alpha=1/4$ and we simulated 4×4 sites with $N=3$ fermions. In **b** the case of a fractional Chern insulator is shown, for parameters $J=t/2$, $V=2t$, $\alpha=1/4$, $U=10t$. We simulated 4×7 sites filled with $N=2$ fermions. Circles mark the positions of the discrete momenta defined by the finite quantization volume.

We expect that our idea—which is to witness topological order in a groundstate by probing the topology characterizing the effective bandstructure of its qp excitations—may also be of broader theoretical interest. It provides a direct route how the concepts developed for non-interacting systems can be generalized to correlated many-body systems: we can define a set of topological invariants for a many-body state by considering single-particle topological invariants of all qp excitations. These may include Chern numbers, \mathbb{Z}_2 invariants or quantized Zak phases. The approach may be useful, for example, in the study of fractional topological insulators⁵⁹ or quantum spin liquid states¹⁴. In the case of chiral quantum spin liquids the analogy to our work is particularly close. On a mean-field level the chiral quantum spin liquid can be described by an integer Chern insulator formed by spinons with a unity Chern number (for example, ref. 60). Therefore, it would be interesting to study the binding of a spinon to a mobile impurity and apply the interferometric method introduced in our paper to detect the spinon Chern number.

Methods

Chern number of TPs. In the main text we generalized the interferometric protocol developed for the measurement of Chern numbers of non-interacting particles^{39,40} to a single qp excitation in a strongly correlated many-body system. The coupling to an impurity particle was necessary for adapting the interferometric protocol. In this way a topological invariant of the TP was defined which, as we will now argue, is the Chern number of the TP.

Consider a qp state $|\psi_{\text{TP}}(k_x, k_y)\rangle$, which is characterized by its quasimomentum (k_x, k_y) . Let us assume that the state is non-degenerate for all quasimomenta. The corresponding Chern number C_{TP} is then defined by the quantized Hall response to an external force \mathbf{F} . In the case of TPs the external force \mathbf{F} acts directly on the impurity and couples to the quasimomentum according to $\frac{d}{dt}\mathbf{k}=\mathbf{F}$. Using the Kubo formula, Thouless *et al.*⁴⁶ have shown that the Chern number can be defined as an integral of the TP Berry curvature \mathcal{F}_{TP} over the (magnetic) BZ,

$$C_{\text{TP}} = \frac{1}{2\pi} \int_{\text{BZ}} d^2k \mathcal{F}_{\text{TP}}(\mathbf{k}). \quad (12)$$

The Berry curvature $\mathcal{F}_{\text{TP}}=\nabla_{\mathbf{k}} \times \langle u_{\mathbf{k}} | i \nabla_{\mathbf{k}} | u_{\mathbf{k}} \rangle$ is defined through the Bloch wavefunction $|u_{\mathbf{k}}\rangle$ constructed from $|\psi_{\text{TP}}(\mathbf{k})\rangle$. The magnetic BZ is defined by the periodicity of the qp Hamiltonian, including gauge-dependent vector potentials⁶¹. The periodicity of the TP wavefunction $|\psi_{\text{TP}}(\mathbf{k}+\mathbf{G})\rangle=|\psi_{\text{TP}}(\mathbf{k})\rangle$, where \mathbf{G} is a reciprocal lattice vector, guarantees the integer quantization of the Chern number^{46,47}.

The magnetic BZ of the TP is determined by the microscopic details of the model for both the impurity as well as the host many-body system. Let us assume that the impurity either lives in the continuum (such that effectively the impurity lattice constant $a \rightarrow 0$ vanishes) or that the unit cell of the impurity lattice fits into the magnetic unit cell of the host many-body system an integer number of times in a commensurate way. In either case, the magnetic BZ of the TP is then equal to the magnetic BZ of the host many-body system. This explains why, in our interferometric protocol, the TP Chern number needs to be defined as the winding of the Zak phase of the TP $\varphi_{\text{Zak}}(k_y)$ over the magnetic BZ of the host many-body system, $k_y \rightarrow k_y + G_y$.

Field theory of TPs. Here we discuss a field theory description of TPs, allowing us to derive the topological invariants characterizing their effective bandstructure. We consider a more general situation than discussed in the main text, and allow quantum Hall states with an arbitrary Abelian topological order. The generalization of our interferometric protocol to this case is discussed in the end.

Our starting point is a field-theoretical description of the topologically ordered host many-body system. We consider an incompressible groundstate with an Abelian topological order, described by a Chern–Simons theory of level n . Such theories are believed to classify all Abelian topological orders and are relevant, for example, for the hierarchical description of the fractional quantum Hall effect⁶² or multilayer fractional quantum Hall systems. They are characterized by the symmetric integer n -by- n matrix \underline{K} and the charge vector \underline{t} . The Lagrangian is⁴

$$\mathcal{L} = \frac{1}{4\pi} \underline{a}_\mu^T \underline{K} \partial_\nu \underline{a}_\nu e^{i\nu\lambda} - \frac{e}{2\pi} \underline{A}_\mu \underline{t}^T \partial_\nu \underline{a}_\nu e^{i\nu\lambda} + \sum_{I=1}^n a_{I\mu} \ell_I j_{I\mu} + \text{kin. energy}. \quad (13)$$

Here $\underline{a}_\mu = a_{1\dots n, \mu}$ are the auxiliary compact $U(1)$ gauge fields from which the conserved current J_μ of the many-body system can be derived, $J_\mu = \frac{e}{2\pi} \sum_I \partial_\nu a_{I\nu} e^{i\nu\lambda}$. As usual, $\mu, \nu, \dots = t, x, y$ denote temporal and spatial coordinates.

The first two terms of the Lagrangian equation (13) describe the response of the many-body system to the external $U(1)$ gauge field A_μ . Here e denotes the A_μ charge of the indistinguishable host particles constituting the many-body system.

From the Euler–Lagrange equations the quantized Hall response is obtained, $J_\mu = C \frac{e}{2\pi} \epsilon^{\mu\nu\lambda} \partial_\nu A_\lambda$, where the many-body Chern number is given by

$$C = \sum_{I,J=1}^n t_I (K^{-1})_{IJ}. \quad (14)$$

The third term in equation (13) describes the conserved currents $j_{I\mu}$ of the $I=1\dots n$ different qps (in the bulk of the system). The integers ℓ_I denote the number of qps that are bound together, in particular, $\ell_I = +1$ ($\ell_I = -1$) for elementary qp (qh) excitations. Edge terms and kinetic energy corrections will be ignored in the Lagrangian equation (13) in the following. The qps carry fractional charges $e_I^*/e = \sum_{I=1}^n t_I (K^{-1})_{IJ}$ (ref. 4). Thus, we note that the Chern number of the incompressible groundstate is given by the sum of the fractional charges of elementary topological excitations, $C = \sum_I e_I^*/e$.

When the impurity (in a given internal state) binds ℓ_I qps of type I , their currents can be directly related to the TP current j_μ by $j_{I\mu} = \ell_I j_\mu$. This is demonstrated by a microscopic calculation in the main text. Using this expression and integrating out the auxiliary $U(1)$ gauge fields $a_{I\mu}$ in the Lagrangian equation (13) (ref. 63), we obtain

$$\mathcal{L}_{\text{eff}} = - \frac{e^2}{4\pi} \underline{t}^T \underline{K}^{-1} \underline{t} e^{i\nu\lambda} A_\mu \partial_\nu A_\lambda + \pi \underline{t}^T \underline{K}^{-1} \underline{\ell}_\mu \frac{e^{i\nu\lambda} \partial_\nu}{\partial^2} j_\lambda + (qB_\mu + e A_\mu \underline{t}^T \underline{K}^{-1} \underline{\ell}) j_\mu, \quad (15)$$

where $\partial^2 = \partial_t \partial_t$. We also included a coupling $qB_\mu j_\mu$ of the impurity to an additional external field B_μ . The first term in equation (15) is a Chern–Simons term for the external gauge field A_μ . The second term describes the braiding statistics of the TP, which coincides with the expected qp statistics, see, for example, ref. 64. The statistical phase $e^{i\theta}$ picked up when interchanging two TPs adiabatically is given by $\theta = \pi \underline{\ell}^T \underline{K}^{-1} \underline{\ell}$ (ref. 4). The last term, most important to our discussion, corresponds to an effective gauge field $(A_\mu e_{\text{TP}}^* + B_\mu q)/e$ seen by the TP, where the A_μ charge of the TP is given by $e_{\text{TP}}^* = e \underline{t}^T \underline{K}^{-1} \underline{\ell}$.

By applying the interferometric protocol introduced in the main text to different flavours $J=1, \dots, n$ of TPs, all the Chern numbers (or, equivalently, all the fractional charges)

$$C_{\text{TP}}^{(J)} = \frac{1}{C_J} = \frac{e}{e_J^*} \quad (16)$$

of elementary topological excitations can be measured. To this end a single qp (or qh) of flavour J is bound to the impurity. The last equation is then derived as equation (4) in the main text. When the TP Chern numbers of all qps are known, the Chern number of the incompressible groundstate can be derived from equation (14),

$$C = \sum_{J=1}^n \frac{(\pm 1)}{C_{\text{TP}}^{(J)}}. \quad (17)$$

Here (-1) needs to be inserted if the elementary qh excitation of flavour J is used, with $\ell_I = -\delta_{IJ}$ and $(+1)$ for elementary qps with $\ell_I = \delta_{IJ}$.

In the main text we discuss the case of $\nu = 1/m$ Laughlin states where $n = t = 1$ and $K = m$. In this case there exists one qh type, with fractional charge $e^* = -e/m$. According to equation (16) the Chern number of the TP consisting of an impurity bound to a hole is given by $C_{\text{TP}} = -m$. We confirm this by a microscopic calculation for a fractional Chern insulator in Fig. 3. Then, equation (17) predicts a fractional Chern number $C = 1/m$ of the incompressible Laughlin state, in agreement with the established result by Niu *et al.*⁵

Exact polaron transformation. To calculate the topological invariants characterizing TPs exactly, we need its full wavefunction $|\psi_{\text{TP}}(\mathbf{q})\rangle$ for any given value of the total TP quasimomentum \mathbf{q} . Here we consider lattice systems for which we develop a method allowing to calculate $|\psi_{\text{TP}}(\mathbf{q})\rangle$ using exact numerical methods. Our approach is based on the LLP unitary transformation⁵⁰ introduced in the context of conventional polaron physics, which makes the conservation of the polaron momentum explicit. The effect of the external force \mathbf{F} acting on the impurity is also discussed in this framework.

Our starting point is the impurity-centred LLP transformation

$$\hat{U}_{\text{LLP}}(t) = \exp[i\hat{\mathbf{R}}_1 \cdot (\hat{\mathbf{p}}_c + \mathbf{F}t)]. \quad (18)$$

To define the impurity position operator $\hat{\mathbf{R}}_1$ and the fermion momentum operator $\hat{\mathbf{p}}_c$, a gauge choice is made. We introduce the magnetic unit cell of size $a_x \times a_y$, for the fermion Hamiltonian \hat{T}_0 , see equation (9), which contains an integer number of flux quanta. Using the Landau gauge as in equation (9) and assuming $\alpha = r/s$ with r, s integers, we have $a_x = a$ and $a_y = sa$. Next, we label sites within the unit cell by an integer μ and define the impurity position operator

$$\hat{\mathbf{R}}_1 = \sum_{\mathbf{j}, \mu} \frac{(\hat{j}_x a_x, \hat{j}_y a_y)^T}{=_{\mathbf{j}}} \hat{b}_{\mathbf{j}, \mu}^\dagger \hat{b}_{\mathbf{j}, \mu}. \quad (19)$$

Here the integers j_{xy} label unit cells, and (\mathbf{j}, μ) is merely an alternative way of parametrizing the site indices (m, n) that were used previously in the definition of the model. Hence, we see that $\hat{\mathbf{R}}_1$ represents only the position of the unit cell, but not the

positions of individual sites *within* one cell. Similarly, the fermion momentum operator is

$$\hat{\mathbf{p}}_c = \sum_{\mathbf{k}, \mu} \mathbf{k} \hat{c}_{\mathbf{k}, \mu}^\dagger \hat{c}_{\mathbf{k}, \mu}, \quad (20)$$

where we introduced operators in momentum space

$$\hat{c}_{\mathbf{k}, \mu} := \sqrt{\frac{a_x a_y}{L_x L_y}} \sum_{\mathbf{j}} e^{i\mathbf{k} \cdot (\mathbf{j}_x a_x \mathbf{e}_x + \mathbf{j}_y a_y \mathbf{e}_y)} \hat{c}_{\mathbf{j}, \mu}.$$
 The wavevector \mathbf{k} takes quantized values

$\mathbf{k} = 2\pi(i_x/L_x, i_y/L_y)^T$ for integers $i_x = 1, \dots, L_x/a_x$ and $i_y = 1, \dots, L_y/a_y$, and with $L_{x,y}$ denoting system size in x and y directions. Note that, although the impurity lattice has a smaller period of a , we have chosen the larger magnetic unit cell of the fermion model in equation (19). This is necessary to distinguish between inequivalent sites μ within one magnetic unit cell for both the fermions and the impurity.

We proceed by applying the LLP transformation defined above to the Hamiltonian $\hat{\mathcal{H}} = \hat{\mathcal{H}}_0 + \hat{\mathcal{H}}_1 + \hat{\mathcal{H}}_{\text{int}}$ (equations (9) and (10)). The new effective Hamiltonian in the polaron frame reads $\hat{\mathcal{H}}(t) = \hat{U}_{\text{LLP}}^\dagger(t) \hat{\mathcal{H}} \hat{U}_{\text{LLP}}(t) - i \hat{U}_{\text{LLP}}^\dagger(t) \partial_t \hat{U}_{\text{LLP}}(t)$. First we note that the many-body (fermion) Hamiltonian $\hat{\mathcal{H}}_0$ trivially commutes with the LLP transformation because of its translational invariance by multiples of one magnetic unit cell. The potential term $\mathbf{F} \cdot \sum_{m,n} \mathbf{r}_{m,n} \hat{b}_{m,n}^\dagger \hat{b}_{m,n}$ in equation (10) also commutes with the LLP transformation, as can easily be checked.

To transform the kinetic energy of the impurity, we introduce the single-particle band Hamiltonian $h_{\mu, \mu'}^i(\mathbf{k})$ defined in the magnetic unit cell of the fermions. This allows us to write $\hat{\mathcal{H}}_1 = \sum_{\mathbf{k}, \mu, \mu'} \hat{b}_{\mathbf{k}, \mu}^\dagger \hat{b}_{\mathbf{k}, \mu'} h_{\mu, \mu'}^i(\mathbf{k})$ in the absence of the force \mathbf{F} . The momentum operators $\hat{b}_{\mathbf{k}, \mu}$ are defined as in the case of fermions $\hat{c}_{\mathbf{k}, \mu}$ discussed above. The impurity position operator $\hat{\mathbf{R}}_1$ is the infinitesimal generator of displacements in quasimomentum space. Therefore, it holds

$$\hat{U}_{\text{LLP}}^\dagger(t) \hat{b}_{\mathbf{k}, \mu} \hat{U}_{\text{LLP}}(t) = \hat{b}_{\mathbf{k} + \hat{\mathbf{p}}_c + \mathbf{F}t, \mu} \quad (21)$$

as can also be checked directly from the definition of the impurity momentum operators $\hat{b}_{\mathbf{k}, \mu}$. Hence, after the LLP transformation we obtain the effective impurity Hamiltonian

$$\begin{aligned} \hat{\mathcal{H}}_1(t) &= \hat{U}_{\text{LLP}}^\dagger(t) \hat{\mathcal{H}}_1 \hat{U}_{\text{LLP}}(t) - i \hat{U}_{\text{LLP}}^\dagger(t) \partial_t \hat{U}_{\text{LLP}}(t) \\ &= \sum_{\mathbf{k}, \mu, \mu'} h_{\mu, \mu'}^i(\mathbf{k} - \hat{\mathbf{p}}_c - \mathbf{F}t) \hat{b}_{\mathbf{k}, \mu}^\dagger \hat{b}_{\mathbf{k}, \mu'}. \end{aligned} \quad (22)$$

Finally, we apply the LLP transformation to the impurity-fermion interaction $\hat{\mathcal{H}}_{\text{int}}$. To keep the discussion general, we consider a density-density interaction of the form

$$\hat{\mathcal{H}}_{\text{int}} = \sum_{\mathbf{j}, \mathbf{l}, \mu, \nu} V_{\mu, \nu}(\mathbf{r}_\mathbf{j} - \mathbf{r}_\mathbf{l}) \hat{b}_{\mathbf{l}, \mu}^\dagger \hat{b}_{\mathbf{j}, \nu} \hat{c}_{\mathbf{j}, \nu}^\dagger \hat{c}_{\mathbf{l}, \nu}, \quad (23)$$

where $V_{\mu, \nu}(\mathbf{r})$ denotes an arbitrary potential. Because the LLP transformation displaces the many-body system (recall that $\hat{\mathbf{p}}_c$ is the infinitesimal generator of fermion translations), it holds

$$\hat{U}_{\text{LLP}}^\dagger(t) \hat{c}_{\mathbf{j}, \mu} \hat{U}_{\text{LLP}}(t) = \hat{c}_{\mathbf{j} - \hat{\mathbf{R}}_1, \mu}. \quad (24)$$

Using this relation and restricting ourselves to the subspace of one impurity, we obtain

$$\hat{U}_{\text{LLP}}^\dagger(t) \hat{\mathcal{H}}_{\text{int}} \hat{U}_{\text{LLP}}(t) = \sum_{\mathbf{i}, \mu} \hat{b}_{\mathbf{i}, \mu}^\dagger \hat{b}_{\mathbf{i}, \mu} \sum_{\mathbf{j}, \nu} V_{\mu, \nu}(\mathbf{r}_\mathbf{j}) \hat{c}_{\mathbf{j}, \nu}^\dagger \hat{c}_{\mathbf{j}, \nu}. \quad (25)$$

This corresponds to a static potential for the many-body fermion system, centred around the central unit cell (where $\mathbf{r}_\mathbf{j} = 0$). For the local interaction assumed in the main text it holds $V_{\mu, \nu}(\mathbf{r}) = V \delta_{\mathbf{r}, 0} \delta_{\mu, \nu}$.

For a single impurity, the resulting Hamiltonian in the polaron frame (written in second quantization for notational convenience) is

$$\hat{\mathcal{H}}(t) = \sum_{\mathbf{k}, \mu, \mu'} \hat{b}_{\mathbf{k}, \mu}^\dagger \hat{b}_{\mathbf{k}, \mu'} \otimes [h_{\mu, \mu'}^i(\mathbf{k} - \hat{\mathbf{p}}_c - \mathbf{F}t) + \delta_{\mu, \mu'} \sum_{\mathbf{j}, \nu} V_{\mu, \nu}(\mathbf{r}_\mathbf{j}) \hat{c}_{\mathbf{j}, \nu}^\dagger \hat{c}_{\mathbf{j}, \nu}] + \hat{\mathcal{H}}_0. \quad (26)$$

Here we used the relation $\sum_{\mathbf{i}} \hat{b}_{\mathbf{i}, \nu}^\dagger \hat{b}_{\mathbf{i}, \nu} = \sum_{\mathbf{k}} \hat{b}_{\mathbf{k}, \nu}^\dagger \hat{b}_{\mathbf{k}, \nu}$ to make the conservation of the TP momentum apparent, $[\hat{\mathcal{H}}, \sum_{\mathbf{k}} \hat{c}_{\mathbf{k}, \mu}^\dagger \hat{c}_{\mathbf{k}, \mu}] = 0$. Equation (26) demonstrates that the force \mathbf{F} directly couples to the TP momentum. Using exact diagonalization techniques we solved the TP band Hamiltonian equation (26) for different values of \mathbf{k} (Figs 2 and 3). This allows us to extract the TP Chern number \mathcal{C} for systems with a small number of fermions.

Fractional TP Chern numbers. Now we discuss TPs with qp charge $e^*/e = p/q$, which is not the inverse of an integer, $p \neq 1$, assuming p, q are relative prime. In this case, the topological field theory of the TP presented above predicts a Chern number

$$\mathcal{C}_{\text{TP}} = \frac{q}{p}. \quad (27)$$

Because the TP Chern number needs to be an integer because of gauge invariance, the groundstate manifold has to be at least p -fold degenerate. In that case all states share a total Chern number $\mathcal{C}_{\text{TP}}^{\text{tot}} = q$.

To understand the origin of this degeneracy we investigate the $\nu = 1/3$ Laughlin state. To this end we bind $p = 2$ qhs to the impurity, which realizes the simplest non-trivial fraction $e^*/e = -2/3$ because each of the qhs has charge $e^*/e = -1/3$. Every qh sees the original host particles as a source of magnetic flux. In a mean-field theory this leads to a homogeneous effective magnetic field $b_z^* = -\frac{2}{3} b_z$ seen by the impurity.

Our reasoning so far relies on the effective low-energy topological field theory, which does not take into account lattice effects. Let us now treat the lattice as a perturbation, which is justified when the magnetic flux per plaquette $\alpha \ll 1$ is sufficiently small. The lattice defines the magnetic unit cell of the TP, the size of which, $2\pi\ell_B^2$, is equal to that of the original magnetic unit cell of the host particles. The size of the magnetic unit cell in the effective field b_z^* , on the other hand, is larger by a factor of $3/2$, $2\pi\ell_B^{*2} = 3\pi\ell_B^2$ (Supplementary Fig. 1). To construct groundstates in the full lattice model from the groundstate in the effective continuum model, we choose the smallest commensurable unit cell, with a size $6\pi\ell_B^2$.

In reciprocal space, we thus arrive at a description of the system in a reduced zone with size $1/3$ of the original magnetic BZ. Because it has half the size of the effective magnetic BZ, it contains two degenerate groundstates. (As a result of the lattice, small gaps may open in some places in the BZ.) We can thus understand the twofold degeneracy as a consequence of the incommensurability of the original magnetic unit cell and the effective magnetic unit cell seen by the impurity.

We used our numerics to verify the existence of a twofold degenerate TP band in small systems (Supplementary Fig. 2a). The corresponding Berry curvature is compatible with a total Chern number $\mathcal{C}_{\text{TP}}^{\text{tot}} = 3$ shared by the two degenerate states (Supplementary Fig. 2b). Details of our calculations are discussed in Supplementary Note 1.

Data availability. The data that support the findings of this study are available from the corresponding author upon request.

References

- Von Klitzing, K., Dorda, G. & Pepper, M. New method for high-accuracy determination of the fine-structure constant based on quantized hall resistance. *Phys. Rev. Lett.* **45**, 494–497 (1980).
- Tsui, D. C., Stormer, H. L. & Gossard, A. C. Two-dimensional magnetotransport in the extreme quantum limit. *Phys. Rev. Lett.* **48**, 1559–1562 (1982).
- Laughlin, R. B. Anomalous quantum hall-effect - an incompressible quantum fluid with fractionally charged excitations. *Phys. Rev. Lett.* **50**, 1395–1398 (1983).
- Wen, X. G. Topological orders and edge excitations in fractional quantum hall states. *Adv. Phys.* **44**, 405–473 (1995).
- Niu, Q., Thouless, D. J. & Wu, Y. S. Quantized hall conductance as a topological invariant. *Phys. Rev. B* **31**, 3372–3377 (1985).
- Sorensen, A. S., Demler, E. & Lukin, M. D. Fractional quantum hall states of atoms in optical lattices. *Phys. Rev. Lett.* **94**, 086803 (2005).
- Hafezi, M., Sorensen, A. S., Demler, E. & Lukin, M. D. Fractional quantum hall effect in optical lattices. *Phys. Rev. A* **76**, 023613 (2007).
- Neupert, T., Santos, L., Chamon, C. & Mudry, C. Fractional quantum hall states at zero magnetic field. *Phys. Rev. Lett.* **106**, 236804 (2011).
- Tang, E., Mei, J. & Wen, X. G. High-temperature fractional quantum hall states. *Phys. Rev. Lett.* **106**, 236802 (2011).
- Regnault, N. & Bernevig, B. A. Fractional chern insulator. *Phys. Rev. X* **1**, 021014 (2011).
- Sun, K., Gu, Z., Katsura, H. & Das Sarma, S. Nearly flatbands with nontrivial topology. *Phys. Rev. Lett.* **106**, 236803 (2011).
- Sheng, D. N., Gu, Z., Sun, K. & Sheng, L. Fractional quantum hall effect in the absence of landau levels. *Nat. Commun.* **2**, 389 (2011).
- Parameśwaran, S. A., Roy, R. & Sondhi, S. L. Fractional quantum hall physics in topological flat bands. *C. R. Phys.* **14**, 816–839 (2013).
- Kitaev, A. Anyons in an exactly solved model and beyond. *Ann. Phys.* **321**, 2–111 (2006).
- Kitaev, A. & Preskill, J. Topological entanglement entropy. *Phys. Rev. Lett.* **96**, 110404 (2006).
- Levin, M. & Wen, X. G. Detecting topological order in a ground state wave function. *Phys. Rev. Lett.* **96**, 110405 (2006).
- Li, H. & Haldane, F. D. M. Entanglement spectrum as a generalization of entanglement entropy: identification of topological order in non-abelian fractional quantum hall effect states. *Phys. Rev. Lett.* **101**, 010504 (2008).
- Abanin, D. A. & Demler, E. Measuring entanglement entropy of a generic many-body system with a quantum switch. *Phys. Rev. Lett.* **109**, 020504 (2012).
- Daley, A. J., Pichler, H., Schachenmayer, J. & Zoller, P. Measuring entanglement growth in quench dynamics of bosons in an optical lattice. *Phys. Rev. Lett.* **109**, 020505 (2012).
- Mazza, L., Rossini, D., Fazio, R. & Endres, M. Detecting two-site spin-entanglement in many-body systems with local particle-number fluctuations. *New J. Phys.* **17**, 013015 (2015).

21. Islam, R. *et al.* Measuring entanglement entropy in a quantum many-body system. *Nature* **528**, 77–83 (2015).
22. Goldman, V. J. & Su, B. Resonant tunneling in the quantum hall regime: measurement of fractional charge. *Science* **267**, 1010–1012 (1995).
23. de Picciotto, R. *et al.* Direct observation of a fractional charge. *Nature* **389**, 162–164 (1997).
24. Saminadayar, L., Glattli, D. C., Jin, Y. & Etienne, B. Observation of the $e/3$ fractionally charged Laughlin quasiparticle. *Phys. Rev. Lett.* **79**, 2526–2529 (1997).
25. Bloch, I., Dalibard, J. & Zwierger, W. Many-body physics with ultracold gases. *Rev. Mod. Phys.* **80**, 885–964 (2008).
26. Lahaye, T., Menotti, C., Santos, L., Lewenstein, M. & Pfau, T. The physics of dipolar bosonic quantum gases. *Rep. Progr. Phys.* **72**, 126401 (2009).
27. Saffman, M., Walker, T. G. & Molmer, K. Quantum information with rydberg atoms. *Rev. Mod. Phys.* **82**, 2313–2363 (2010).
28. Chin, C., Grimm, R., Julienne, P. & Tiesinga, E. Feshbach resonances in ultracold gases. *Rev. Mod. Phys.* **82**, 1225–1286 (2010).
29. Aidelsburger, M. *et al.* Measuring the chern number of hofstadter bands with ultracold bosonic atoms. *Nat. Phys.* **11**, 162–166 (2015).
30. Jotzu, G. *et al.* Experimental realization of the topological haldane model with ultracold fermions. *Nature* **515**, 237–240 (2014).
31. Wilkin, N. K., Gunn, J. M. F. & Smith, R. A. Do attractive bosons condense? *Phys. Rev. Lett.* **80**, 2265–2268 (1998).
32. Regnault, N. & Jolicoeur, T. Quantum hall fractions in rotating bose-einstein condensates. *Phys. Rev. Lett.* **91**, 030402 (2003).
33. Baranov, M. A., Fehrmann, H. & Lewenstein, M. Wigner crystallization in rapidly rotating 2d dipolar fermi gases. *Phys. Rev. Lett.* **100**, 200402 (2008).
34. Grusdt, F. & Fleischhauer, M. Fractional quantum hall physics with ultracold rydberg gases in artificial gauge fields. *Phys. Rev. A* **87**, 043628 (2013).
35. Yao, N. Y. *et al.* Realizing fractional chern insulators in dipolar spin systems. *Phys. Rev. Lett.* **110**, 185302 (2013).
36. Schine, N., Ryou, A., Gromov, A., Sommer, A. & Simon, J. Synthetic landau levels for photons. Preprint at <http://arxiv.org/abs/arXiv:1511.07381> (2015).
37. Frese, D. *et al.* 'Single atoms in an optical dipole trap: towards a deterministic source of cold atoms. *Phys. Rev. Lett.* **85**, 3777–3780 (2000).
38. Karski, M. *et al.* Quantum walk in position space with single optically trapped atoms. *Science* **325**, 174–177 (2009).
39. Atala, M. *et al.* Direct measurement of the zak phase in topological bloch bands. *Nat. Phys.* **9**, 795–800 (2013).
40. Abanin, D. A., Kitagawa, T., Bloch, I. & Demler, E. Interferometric approach to measuring band topology in 2d optical lattices. *Phys. Rev. Lett.* **110**, 165304 (2013).
41. Grusdt, F., Abanin, D. & Demler, E. Measuring Z2 topological invariants in optical lattices using interferometry. *Phys. Rev. A* **89**, 043621 (2014).
42. Duca, L. *et al.* An aharonov-bohm interferometer for determining bloch band topology. *Science* **347** no. 6219, 288–292 (2014).
43. Zak, J. Berrys phase for energy-bands in solids. *Phys. Rev. Lett.* **62**, 2747–2750 (1989).
44. Xiao, D., Chang, M. & Niu, Q. Berry phase effects on electronic properties. *Rev. Mod. Phys.* **82**, 1959–2007 (2010).
45. Dalibard, J., Gerbier, F., Juzeliunas, G. & Oehberg, P. Colloquium: artificial gauge potentials for neutral atoms. *Rev. Mod. Phys.* **83**, 1523 (2011).
46. Thouless, D. J., Kohmoto, M., Nightingale, M. P. & denNijs, M. Quantized hall conductance in a two-dimensional periodic potential. *Phys. Rev. Lett.* **49**, 405–408 (1982).
47. Kohmoto, M. Topological invariant and the quantization of the hall conductance. *Ann. Phys.* **160**, 343–354 (1985).
48. Pekar, S. I. *Zh. Eksp. Teor. Fiz.* **16**, 341 (1946).
49. Landau, L. D. & Pekar, S. I. Effective mass of a po-laron. *Zh. Eksp. Teor. Fiz.* **18**, 419 (1948).
50. Lee, T. D., Low, F. E. & Pines, D. The motion of slow electrons in a polar crystal. *Phys. Rev.* **90**, 297–302 (1953).
51. Hofstadter, D. R. Energy-levels and wave-functions of bloch electrons in rational and irrational magnetic-fields. *Phys. Rev. B* **14**, 2239–2249 (1976).
52. Aidelsburger, M. *et al.* Realization of the hofstadter hamiltonian with ultracold atoms in optical lattices. *Phys. Rev. Lett.* **111**, 185301 (2013).
53. Miyake, H., Siviloglou, G. A., Kennedy, C. J., Burton, W. C. & Ketterle, W. Realizing the harper hamiltonian with laser-assisted tunneling in optical lattices. *Phys. Rev. Lett.* **111**, 185302 (2013).
54. Kapit, E. & Mueller, E. Exact parent hamiltonian for the quantum hall states in a lattice. *Phys. Rev. Lett.* **105**, 215303 (2010).
55. Paredes, B., Fedichev, P., Cirac, J. I. & Zoller, P. 1/2-anyons in small atomic bose-einstein condensates. *Phys. Rev. Lett.* **87**, 010402 (2001).
56. Grusdt, F., Letscher, F., Hafezi, M. & Fleischhauer, M. Topological growing of laughlin states in synthetic gauge fields. *Phys. Rev. Lett.* **113**, 155301 (2014).
57. Zhang, Y., Sreejith, G. J., Gemelke, N. D. & Jain, J. K. Fractional angular momentum in cold-atom systems. *Phys. Rev. Lett.* **113**, 160404 (2014).
58. Zhang, Y., Sreejith, G. J. & Jain, J. K. Creating and manipulating non-abelian anyons in cold atom systems using auxiliary bosons. *Phys. Rev. B* **92**, 075116 (2015).
59. Levin, M. & Stern, A. Fractional topological insulators. *Phys. Rev. Lett.* **103**, 196803 (2009).
60. Wen, X. G. *Quantum Field Theory of Many-Body Systems* (Oxford University Press, 2004).
61. Zak, J. Magnetic translation group. *Phys. Rev. A* **134**, 1602 (1964).
62. Halperin, B. I. Statistics of quasiparticles and the hierarchy of fractional quantized hall states. *Phys. Rev. Lett.* **52**, 1583–1586 (1984).
63. Moore, J. *An Introduction to Topological Phases of Electrons* (Oxford University Press, 2014).
64. Altland, A. & Simons, B. *Condensed Matter Field Theory* (Cambridge University Press, 2010).

Acknowledgements

We acknowledge useful discussions with M. Knap, M. Hafezi, F. Letscher, W. Phillips, J. Sau, J. Taylor, I. Bloch, E. Mueller, M. Greiner, F. Gerbier, J. Dalibard and C. Weitenberg. F.G. acknowledges support by the Excellence Initiative (DFG/GSC 266), the Marion Köser Stiftung and the Moore foundation. E.D. and F.G. acknowledge support from Harvard-MIT CUA, NSF Grant No. DMR-1308435, AFOSR Quantum Simulation MURI, the ARO-MURI on Atomtronics and the ARO-MURI Quism programme. E.D. acknowledges support from the Humboldt foundation, Dr Max Rössler, the Walter Haefner Foundation and the ETH Foundation. D.A. acknowledges support by the Alfred Sloan Foundation. N.Y.Y. acknowledges support from the Miller Institute for Basic Research in Science.

Author contributions

All authors contributed substantially to the writing of the manuscript. F.G. performed the calculations. F.G. and E.D. conceived the method. N.Y.Y., D.A. and M.F. contributed to clarifying the initial idea of the method.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npng.nature.com/reprintsandpermissions/>

How to cite this article: Grusdt, F. *et al.* Interferometric measurements of many-body topological invariants using mobile impurities. *Nat. Commun.* **7**:11994 doi: 10.1038/ncomms11994 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>