

# UC Irvine

## UC Irvine Previously Published Works

### Title

Mitochondrial Zn<sup>2+</sup> Accumulation: A Potential Trigger of Hippocampal Ischemic Injury.

### Permalink

<https://escholarship.org/uc/item/0ww471f7>

### Journal

The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry, 25(2)

### ISSN

1073-8584

### Authors

Ji, Sung G  
Medvedeva, Yuliya V  
Wang, Hwai-Lee  
et al.

### Publication Date

2019-04-01

### DOI

10.1177/1073858418772548

Peer reviewed

# The Neuroscientist

## Mitochondrial Zn<sup>2+</sup> accumulation: A potential trigger of hippocampal ischemic injury

Journal:	<i>Neuroscientist</i>
Manuscript ID	NRO-18-RE-0014.R1
Manuscript Type:	Review
Date Submitted by the Author:	n/a
Complete List of Authors:	Ji, Sung; University of California Irvine School of Medicine, Anatomy & Neurobiology Medvedeva, Yuliya; University of California Irvine School of Medicine, Neurology Wang, Hwai; University of California Irvine School of Medicine, Neurology; China Medical University, Graduate Institute of Clinical Medical Science Yin, Hong; University of California Irvine School of Medicine, Neurology Weiss, John; University of California Irvine School of Medicine, Neurology and Anatomy & Neurobiology
Keywords:	Calcium, Cell death, Excitotoxicity, Ischemia, Mitochondria, Reactive oxygen species, ROS, zinc

SCHOLARONE™  
Manuscripts

NEW

1 Review article

2  
3 Address correspondence to:

4  
5 John H. Weiss, MD, Ph.D.

6  
7 2101 Gillespie Building

8  
9 Department of Neurology, University of California, Irvine

10  
11 Irvine, CA 92697-4292

12  
13 Tel: (949) 824-6774

14  
15 Fax: (949) 824-1668

16  
17 E-mail: [jweiss@uci.edu](mailto:jweiss@uci.edu)

18  
19  
20  
21  
22  
23  
24  
25 **Mitochondrial Zn<sup>2+</sup> accumulation: A**  
26  
27 **potential trigger of hippocampal ischemic**  
28  
29 **injury**  
30  
31  
32  
33  
34  
35  
36  
37

38 Zinc, mitochondria and ischemic hippocampal injury.  
39  
40  
41  
42

43 Sung G. Ji<sup>1</sup>, Yuliya V. Medvedeva<sup>2</sup>, Hwai-Lee Wang<sup>2,3</sup>, Hong Z. Yin<sup>2</sup>  
44  
45 and John H. Weiss, MD, PhD<sup>1,2</sup>  
46  
47  
48  
49

50 From the <sup>1</sup>Department of Anatomy & Neurobiology, and the <sup>2</sup>Department of Neurology,

51  
52 University of California, Irvine and the <sup>3</sup>Graduate Institute of Clinical Medical Science, China

53  
54 Medical University, Taichung, Taiwan  
55  
56  
57  
58  
59  
60

## Abstract

Ischemic stroke is a major cause of death and disabilities worldwide, and it has been long hoped that improved understanding of relevant injury mechanisms would yield targeted neuroprotective therapies. While  $\text{Ca}^{2+}$  overload during ischemia-induced glutamate excitotoxicity has been identified as a major contributor, failures of glutamate targeted therapies to achieve desired clinical efficacy have dampened early hopes for the development of new treatments. However, additional studies examining possible contributions of  $\text{Zn}^{2+}$ , a highly prevalent cation in the brain, have provided new insights that may help to rekindle the enthusiasm. In this review, we discuss both old and new findings yielding clues as to sources of the  $\text{Zn}^{2+}$  that accumulates in many forebrain neurons after ischemia, and mechanisms through which it mediates injury. Specifically, we highlight the growing evidence of important  $\text{Zn}^{2+}$  effects on mitochondria in promoting neuronal injury. A key focus has been to examine  $\text{Zn}^{2+}$  contributions to the degeneration of highly susceptible hippocampal pyramidal neurons. Recent studies provide evidence of differences in sources of  $\text{Zn}^{2+}$  and its interactions with mitochondria in CA1 vs CA3 neurons that may pertain to their differential vulnerabilities in disease. We propose that  $\text{Zn}^{2+}$ -induced mitochondrial dysfunction is a critical and potentially targetable early event in the ischemic neuronal injury cascade, providing opportunities for the development of novel neuroprotective strategies to be delivered after transient ischemia.

## Key words

Calcium, Cell death, Excitotoxicity, Ischemia, Mitochondria, Reactive oxygen species, ROS, Zinc

## Ischemic stroke: the role of $\text{Ca}^{2+}$

Ischemic stroke is a leading cause of disability and death worldwide, reflecting the extreme sensitivity of brain to even brief (several minutes) disruption of blood flow. Despite extensive efforts to understand the basis of this unique vulnerability with the aim of developing neuroprotective interventions, attempts to date have failed, with the maintenance and prompt restoration of perfusion being the only presently available therapeutic approach.

Considerable evidence implicates a role for “excitotoxicity” (neuronal damage triggered by excessive release of the excitatory neurotransmitter glutamate) occurring in conditions including ischemia, prolonged seizures and trauma. Excitotoxic mechanisms have been extensively investigated, and a critical early finding was that brief strong activation of highly  $\text{Ca}^{2+}$  permeable NMDA type glutamate receptors (**NMDAR**) results in delayed  $\text{Ca}^{2+}$  dependent neurodegeneration (Choi 1987; Choi and others 1988). After the brief exposure, intracellular  $\text{Ca}^{2+}$  levels recover for a period of time before undergoing a sharp and sustained rise (termed “ **$\text{Ca}^{2+}$  deregulation**”) that is strongly correlated with cell death (Randall and Thayer 1992).

It is also apparent that oxidative mechanisms contribute to the neuronal injury, induced after production of the reactive oxygen species (**ROS**; including superoxide and nitric oxide) (Lafon-Cazal and others 1993; Sattler and others 1999).

Mitochondria have been implicated as important targets of  $\text{Ca}^{2+}$  effects.  $\text{Ca}^{2+}$  enters mitochondria through a specific channel (the mitochondrial  $\text{Ca}^{2+}$  uniporter, **MCU**), and under normal circumstances, physiological mitochondrial  $\text{Ca}^{2+}$  rises help to regulate mitochondrial metabolic function by matching ATP production to need (Nicholls and Budd 2000).

1  
2  
3  
4 Mitochondria are also important buffers of large cytosolic  $\text{Ca}^{2+}$  loads (Wang and Thayer 1996;  
5  
6 White and Reynolds 1997). However, with excess accumulation,  $\text{Ca}^{2+}$  can disrupt mitochondrial  
7  
8 function, with effects including increased superoxide production (Dugan and others 1995;  
9  
10 Reynolds and Hastings 1995), and opening of a large conductance inner membrane channel (the  
11  
12 mitochondrial permeability transition pore; **mPTP**), that can lead to mitochondrial swelling and  
13  
14 the release of cytochrome C and other pro-apoptotic peptides (Nicholls and Budd 2000). Recent  
15  
16 studies have also demonstrated the importance of another distinct mechanism of excitotoxic  
17  
18 superoxide generation, via  $\text{Ca}^{2+}$  dependent activation of the superoxide-generating cytosolic  
19  
20 enzyme NADPH oxidase (**NOX**) (Brennan and others 2009; Clausen and others 2013), and it is  
21  
22 likely that depending upon conditions both sources can contribute.  
23  
24  
25  
26  
27

28  
29 However, despite considerable early hope and some promising results in animals, use of  
30  
31 NMDAR antagonists (to prevent  $\text{Ca}^{2+}$  mediated injury and deregulation) have yielded little  
32  
33 benefit in human studies (Hoyte and others 2004; Ikonomidou and Turski 2002), necessitating a  
34  
35 further search for new targets yielding better efficacy.  
36  
37

## 38 $\text{Zn}^{2+}$ : a distinct ionic contributor to brain injury

39  
40  
41  
42

43  $\text{Zn}^{2+}$  is a critical and highly prevalent cation in all tissues. It is particularly prevalent in  
44  
45 brain, which has an overall  $\text{Zn}^{2+}$  content estimated to be 100-200  $\mu\text{M}$ , and is especially high in  
46  
47 certain limbic and forebrain regions including hippocampus, amygdala and cortex (Frederickson  
48  
49 1989). Despite the high total  $\text{Zn}^{2+}$ , virtually all of it is bound or sequestered; while precise  
50  
51 measurements are difficult (as it can bind numerous ligands with a wide range of affinities), it is  
52  
53 agreed that free intracellular  $\text{Zn}^{2+}$  levels are subnanomolar (Colvin and others 2010; Maret  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 2015). Reflecting its importance in all tissues, there are two families of transporters (with >20  
5  
6 variants identified to date) dedicated to movement of  $Zn^{2+}$  between compartments, with the Zrt-,  
7  
8 Irt-like protein (**ZIP**) family moving  $Zn^{2+}$  into cytosol, and the  $Zn^{2+}$  transporter (**ZnT**) family  
9  
10 moving  $Zn^{2+}$  from cytosol out of the cell or into subcellular compartments (Kambe and others  
11  
12 2014). In neurons, most (~ 90%) of the  $Zn^{2+}$  is bound to or associated with proteins, and it is an  
13  
14 integral component of numerous enzymes, transcription factors and structural proteins  
15  
16 (Frederickson 1989).  
17  
18  
19

## 20 21 **Synaptic $Zn^{2+}$ : a modulator of neurotransmission and contributor to** 22 23 **injury** 24 25 26 27

28  
29 A distinct and critical pool of brain  $Zn^{2+}$  is that which is sequestered within presynaptic  
30  
31 vesicles of some excitatory neurons. This pool of free or loosely bound  $Zn^{2+}$  is visualized by  
32  
33 histochemical procedures like Timm's silver sulfide staining or labeling with  $Zn^{2+}$  sensitive  
34  
35 fluorescent dyes, and is often referred to as chelatable or "histochemically reactive"  $Zn^{2+}$   
36  
37 (Frederickson 1989; Frederickson and others 1992). This  $Zn^{2+}$  has a distinctive distribution,  
38  
39 generally corresponding with areas of greatest total  $Zn^{2+}$ ; high levels are found in hippocampus  
40  
41 (particularly the dentate granule cells and their "mossy fiber" projections, accounting for the  
42  
43 distinctive appearance of hippocampus after Timm's staining; see **Fig. 1A**), as well as in cortex  
44  
45 and amygdala. In these neurons, the  $Zn^{2+}$  appears to be loaded into vesicles at millimolar  
46  
47 concentrations by the vesicular  $Zn^{2+}$  transporter, **ZnT3** (Cole and others 1999). It is further  
48  
49 evident that this  $Zn^{2+}$  is co-released with glutamate upon stimulation (Assaf and Chung 1984;  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60 Howell and others 1984; Sloviter 1985), and peak levels at synapses may reach into the 100  $\mu M$

1  
2  
3  
4 range with strong activation (Ueno and others 2002; Vogt and others 2000), constituting about a  
5  
6 10,000-fold increase over physiologic resting level of extracellular  $Zn^{2+}$  (Frederickson and others  
7  
8 2006).  
9

10  
11  
12 **[insert Figure 1 and Figure 1 caption]**  
13

14  
15 The identification of populations of forebrain excitatory neurons containing substantial  
16  
17 quantities of presynaptic vesicular  $Zn^{2+}$  begs understanding of the actions and effects of  
18  
19 synaptically released  $Zn^{2+}$ . While much is not known,  $Zn^{2+}$  has complex effects on extracellular  
20  
21 receptors, antagonizing NMDAR currents via both voltage dependent and voltage independent  
22  
23 mechanisms; electrophysiological studies have demonstrated  $Zn^{2+}$  release from mossy fibers to  
24  
25 provide tonic inhibition of NMDAR on CA3 pyramidal neurons (Vogt and others 2000). In  
26  
27 addition,  $Zn^{2+}$  has effects on GABA and glycinergic receptors, as well as on a  $Zn^{2+}$  sensing G-  
28  
29 protein linked metabotropic receptor, and synaptic  $Zn^{2+}$  likely has roles in forms of synaptic  
30  
31 plasticity (Sensi and others 2011).  
32  
33  
34  
35

36  
37 Observations that ischemia, prolonged seizures and brain trauma resulted in loss of  
38  
39 chelatable  $Zn^{2+}$  labeling in presynaptic pools (most evident in the mossy fibers) (see **Fig. 1B**),  
40  
41 and its appearance in somata of injured neurons led to the suggestion that synaptic  $Zn^{2+}$  release  
42  
43 and its translocation through channels into postsynaptic neurons contributed to their degeneration  
44  
45 in these conditions (Frederickson and others 1989; Suh and others 2000; Tonder and others  
46  
47 1990). Indeed, this idea was markedly strengthened by observations that application of an  
48  
49 extracellular  $Zn^{2+}$  chelator decreased both the postsynaptic  $Zn^{2+}$  accumulation and subsequent  
50  
51 neurodegeneration (Calderone and others 2004; Koh and others 1996; Yin and others 2002).  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4 Paralleling observations of neuronal  $Zn^{2+}$  accumulation after seizures or ischemia *in vivo*,  
5  
6 studies in neuronal culture models documented the potent toxic effects of  $Zn^{2+}$  and sought to  
7  
8 examine its mechanisms. One early aim was to identify the routes through which synaptically  
9  
10 released  $Zn^{2+}$  can enter post-synaptic neurons to trigger injury. These studies found  $Zn^{2+}$  to  
11  
12 permeate three distinct channels through which  $Ca^{2+}$  also permeates: (1) NMDAR (Koh and  
13  
14 Choi 1994); (2) L-type voltage gated  $Ca^{2+}$  channels (VGCC) (Freund and Reddig 1994;  
15  
16 Kerchner and others 2000; Weiss and others 1993); and (3) atypical  $Ca^{2+}$  permeable AMPA type  
17  
18 glutamate receptors (“Ca-AMPA”); whereas most AMPA receptors are  $Ca^{2+}$  impermeable,  
19  
20 these lack the GluA2 subunit in their tetrameric structure, and are only present in substantial  
21  
22 numbers on small subpopulations of neurons. We found these Ca-AMPA to be highly  $Zn^{2+}$   
23  
24 permeable (Jia and others 2002; Yin and Weiss 1995). However, direct comparison of these  
25  
26 routes indicated substantial differences in their  $Zn^{2+}$  permeabilities, and corresponding  
27  
28 differences in the potency with which  $Zn^{2+}$  entry through each of them triggers injury.  
29  
30 Consistent with its effective antagonism of NMDAR currents, very little  $Zn^{2+}$  permeates  
31  
32 NMDARs. Ubiquitously expressed VGCC showed an intermediate permeability, and the  
33  
34 selectively expressed Ca-AMPA had the greatest  $Zn^{2+}$  permeability (Sensi and others 1999).  
35  
36  
37  
38  
39  
40  
41

42 While brief moderate  $Zn^{2+}$  exposures to depolarized neurons resulted in sufficient  $Zn^{2+}$   
43  
44 entry through VGCC to trigger extensive degeneration over the subsequent day (Weiss and  
45  
46 others 1993), several considerations led us to believe that entry through Ca-AMPA might be of  
47  
48 particular importance. First, despite their selective expression (in contrast to the VGCC, they are  
49  
50 only present in large numbers on ~13% of neurons in cortical cultures, and preferentially found  
51  
52 in dendrites of some pyramidal neurons) (Lerma and others 1994; Ogoshi and Weiss 2003; Sensi  
53  
54 and others 1999; Yin and others 1994; Yin and others 1999), they permit substantially greater  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 rates of  $Zn^{2+}$  entry, and, when present, are concentrated at postsynaptic membranes where the  
5  
6 highest levels of extracellular  $Zn^{2+}$  are likely achieved. Furthermore, early  $Zn^{2+}$  accumulation has  
7  
8 been found to trigger a delayed increase in numbers of Ca-AMPA in many forebrain neurons 2-  
9  
10 3 days after transient ischemia (due to decreased expression of GluA2), a factor that likely  
11  
12 contributes to delayed neurodegeneration (Calderone and others 2004; Gorter and others 1997).  
13  
14 Indeed, supporting the significance of this route, a Ca-AMPA antagonist attenuated  $Zn^{2+}$   
15  
16 accumulation and injury both in a slice model of acute ischemia (Yin and others 2002), and when  
17  
18 delivered late after transient global ischemia *in vivo* (Noh and others 2005). However, this does  
19  
20 not mean VGCC are unimportant. Although VGCC are not concentrated specifically at  
21  
22 synapses, entry through this route would likely occur under pathologic conditions in which  
23  
24 extracellular  $Zn^{2+}$  accumulation is accompanied by widespread neuronal depolarization. Also,  
25  
26 VGCC activity increases with age (Thibault and Landfield 1996), possibly increasing the  
27  
28 contribution of this route in aging populations most at risk of brain ischemia.  
29  
30  
31  
32  
33  
34

35  
36 The generation of ZnT3 knockout mice, which are entirely lacking in chelatable  
37  
38 presynaptic  $Zn^{2+}$  (Cole and others 1999), provided a valuable tool to test the presumption that  
39  
40 presynaptic  $Zn^{2+}$  release and its translocation into postsynaptic neurons accounted for the  
41  
42 injurious postsynaptic  $Zn^{2+}$  accumulation. Consistent with this idea, when ZnT3 knockouts were  
43  
44 tested in a prolonged kainate seizure model, the knockouts showed modestly decreased  $Zn^{2+}$   
45  
46 accumulation and injury in CA3 pyramidal neurons (which are innervated by the very densely  
47  
48  $Zn^{2+}$  containing mossy fibers). Surprisingly, however,  $Zn^{2+}$  accumulation and injury were  
49  
50 markedly *increased* in CA1 pyramidal neurons of the knockouts, indicating an additional source  
51  
52 of  $Zn^{2+}$  that did not depend upon synaptic release and translocation (Lee and others 2000).  
53  
54  
55  
56  
57  
58  
59  
60

## Zn<sup>2+</sup> binding proteins: buffers of Zn<sup>2+</sup> loads or sources of non-synaptic Zn<sup>2+</sup> accumulation (or both)?

Metallothioneins (MT, I-IV), are cysteine rich peptides with multiple Zn<sup>2+</sup> binding sites that play critical roles in buffering Zn<sup>2+</sup> within cells (MT-III being the predominant neuronal isoform), making them likely candidate sources for the non-synaptic neuronal Zn<sup>2+</sup> accumulation (Maret 1995). Zn<sup>2+</sup> binding to MT's is highly sensitive to environmental conditions, with metabolic aberrations associated with pathological conditions (specifically oxidative stress and acidosis) destabilizing binding, resulting in release of free Zn<sup>2+</sup> into cytosol (Jiang and others 2000; Maret 1995). A seminal observation that simple application of a disulfide oxidant to cultured neurons was capable of causing cytosolic Zn<sup>2+</sup> rises that could trigger delayed neurodegeneration provided the first *proof of principle* that simple mobilization of Zn<sup>2+</sup> from intracellular buffers could result in neurodegeneration (Aizenman and others 2000). A subsequent study overexpressing MT-III found that depending upon conditions it could have divergent effects, either buffering excess Zn<sup>2+</sup> that enters the cell (and thereby diminishing its toxic effects), or providing a source of injurious Zn<sup>2+</sup> mobilization, under conditions of oxidative stress (Malaiyandi and others 2004).

Indeed, use of MT-III knockout mice (as well as double MT-III / ZnT3 knockouts) helped clarify the respective contributions of synaptic vs MT-III bound Zn<sup>2+</sup> in the kainate seizure model. In contrast to the increased Zn<sup>2+</sup> accumulation seen in ZnT3 knockouts in CA1 neurons, Zn<sup>2+</sup> accumulation and injury in MT-III knockouts were decreased in CA1, consistent with a dominant contribution of mobilization from MT-III. Conversely, these were increased in

1  
2  
3  
4 CA3 of MT-III knockouts, consistent with synaptic “translocation” predominating, with MT-III  
5  
6 in CA3 serving a protective role by helping to buffer  $Zn^{2+}$  entering the neurons (Lee and others  
7  
8 2003).  
9

10  
11  
12 Might these differences in sources of injurious  $Zn^{2+}$  accumulation be a factor contributing  
13  
14 to their differential disease susceptibilities, with CA3 neurons preferentially degenerating after  
15  
16 recurrent limbic seizures (associated with repetitive firing of the  $Zn^{2+}$  rich mossy fibers) and  
17  
18 CA1 neurons undergoing delayed degeneration after transient ischemia (Ben-Ari and others  
19  
20 1980; Sugawara and others 1999)?  
21  
22  
23

## 24 25 Discrimination of $Ca^{2+}$ and $Zn^{2+}$ reveals distinct contributions 26 27 28

29  
30 Despite the emerging evidence for contributions of  $Zn^{2+}$ , there is still much evidence for  
31  
32 important  $Ca^{2+}$  contributions in excitotoxicity associated conditions, and it is probable that both  
33  
34 ions contribute. However, early attempts to discriminate their contributions were confounded by  
35  
36 the fact that until relatively recently, there were no available  $Zn^{2+}$ -selective indicators.  
37  
38 Furthermore, it became apparent that some effects that had been attributed to  $Ca^{2+}$  might actually  
39  
40 be partly  $Zn^{2+}$  mediated, since available  $Ca^{2+}$  indicators bound and responded to  $Zn^{2+}$  with higher  
41  
42 affinity than  $Ca^{2+}$  (Cheng and Reynolds 1998), and fluorescence increases detected by a “ $Ca^{2+}$   
43  
44 indicator” in a slice model of ischemia (that would previously have been assumed to reflect  $Ca^{2+}$   
45  
46 rises) were found to be substantially diminished by selective  $Zn^{2+}$  chelation (Stork and Li 2006).  
47  
48 The development of  $Zn^{2+}$  selective indicators provided a breakthrough in attempts to study  $Zn^{2+}$ -  
49  
50 specific effects and discriminate them from those of  $Ca^{2+}$ . Furthermore, using a high affinity  
51  
52  $Zn^{2+}$  indicator in combination with a low affinity  $Ca^{2+}$  indicator, it became possible to  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 simultaneously track changes in both ions (Devinney and others 2005). We used this approach to  
5  
6 simultaneously track changes in both  $Zn^{2+}$  and  $Ca^{2+}$  in single pyramidal neurons in hippocampal  
7  
8 slices subjected to oxygen glucose deprivation (**OGD**) (see **Fig. 2A**). Interestingly, we found that  
9  
10 cytosolic  $Zn^{2+}$  rises both preceded and contributed to the onset of terminal  $Ca^{2+}$  deregulation  
11  
12 events, which still occurred but were significantly delayed by the presence of a  $Zn^{2+}$  chelator (see  
13  
14 **Fig. 2B**) (Medvedeva and others 2009). This provided new evidence that  $Zn^{2+}$  accumulation  
15  
16 might be an early event in the ischemic injury cascade, the appropriate targeting of which might  
17  
18 provide therapeutic benefit. As discussed further below, clues from this and other early studies  
19  
20 suggested that mitochondria might be an important target for these early  $Zn^{2+}$  effects (see **Fig.**  
21  
22 **2C**).  
23  
24  
25  
26  
27

28 **[insert Figure 2 and Figure 2 caption]**  
29  
30

## 31 Mitochondria: a critical target of $Zn^{2+}$

32  
33  
34  
35

36 Paralleling studies of  $Ca^{2+}$ , studies over several decades have highlighted ways in which  
37  
38  $Zn^{2+}$  impacts mitochondrial function. Below, we review the evolution of these data, leading up to  
39  
40 our proposition that mitochondrial  $Zn^{2+}$  accumulation may be an important early step in the  
41  
42 ischemic injury cascade of many neurons. Specifically, as it occurs upstream from terminal  $Ca^{2+}$   
43  
44 deregulation, its targeting may provide benefits distinct from those provided by attenuation of  
45  
46  $Ca^{2+}$  entry (via NMDAR blockade).  
47  
48  
49  
50

## 51 Isolated mitochondria: Evidence of potent $Zn^{2+}$ effects

52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 A number of studies dating back over 50 years have found that  $Zn^{2+}$  can enter  
5  
6 mitochondria, inducing effects including swelling, and inhibition of respiration with high  
7  
8 potency (Brierley 1967; Skulachev and others 1967). Over the subsequent decades, with  
9  
10 growing awareness that  $Zn^{2+}$  is a pathophysiologically important ion that contributes to neuronal  
11  
12 injury, there has been an increasing interest in determining how  $Zn^{2+}$  impacts mitochondria.  $Zn^{2+}$   
13  
14 was found to enter mitochondria specifically through the MCU (Saris and Niva 1994), and to  
15  
16 trigger opening of the mPTP (Wudarczyk and others 1999). Other studies found potent  
17  
18 (submicromolar)  $Zn^{2+}$  inhibition of the bc1 complex of the electron transport chain and of the  
19  
20 TCA cycle  $\alpha$ -ketoglutarate dehydrogenase enzyme complex (Brown and others 2000; Link and  
21  
22 von Jagow 1995). Highlighting the complexity of  $Zn^{2+}$  effects on mitochondria, we found low  
23  
24 (submicromolar) exposures to induce loss of mitochondrial membrane potential ( $\Delta\Psi_{mito}$ ),  
25  
26 decreased ROS production and increased  $O_2$  consumption (consistent with uncoupling of the  
27  
28 electron transport from ATP synthesis), while slightly higher levels increased ROS generation  
29  
30 and decreased  $O_2$  consumption (consistent with inhibition of electron transport) (Sensi and others  
31  
32 2003). A subsequent study reported  $Zn^{2+}$ , after entry through the MCU, to induce irreversible  
33  
34 inhibition of major thiol oxidoreductase enzymes involved in energy production and antioxidant  
35  
36 defense, an effect that appeared to be linked to mPTP opening (Gazaryan and others 2007).  
37  
38  
39  
40  
41  
42  
43

44 Using isolated brain mitochondria, we found  $Zn^{2+}$  (10-100 nM) to potently induce  
45  
46 swelling, that appeared to depend upon  $Zn^{2+}$  entry through the MCU and opening of the mPTP  
47  
48 (Jiang and others 2001). We further found that although  $Zn^{2+}$  triggered mitochondrial swelling  
49  
50 with far greater potency than  $Ca^{2+}$ , the effects of these ions were synergistic, with greater  
51  
52 swelling when  $Ca^{2+}$  was also present (Jiang and others 2001). Indeed, a number of other studies  
53  
54 have also suggested that the presence of  $Ca^{2+}$  may critically modulate effects of  $Zn^{2+}$  on isolated  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 mitochondria. Specifically,  $\text{Ca}^{2+}$  was found to markedly enhance  $\text{Zn}^{2+}$  entry through the MCU  
5  
6 (Saris and Niva 1994), and  $\text{Zn}^{2+}$  triggered mPTP opening of de-energized (but not energized)  
7  
8 mitochondria was found to be  $\text{Ca}^{2+}$  dependent (Wudarczyk and others 1999). Interestingly, a  
9  
10 relatively recent study exposed purified and substrate attached mitochondria using buffers pre-  
11  
12 treated to ensure complete elimination of  $\text{Ca}^{2+}$ , and found  $\text{Zn}^{2+}$  to have weak depolarizing effects  
13  
14 with no evidence of its entry into mitochondria (Devinney and others 2009). Of possible  
15  
16 relevance, the MCU and associated regulatory peptides were recently identified and two  
17  
18 regulatory peptides (MICU1 and 2), appear to sense  $\text{Ca}^{2+}$ , inhibiting MCU opening when  $\text{Ca}^{2+}$  is  
19  
20 near resting levels (<100-200 nM) and promoting opening when  $\text{Ca}^{2+}$  is elevated, thus conferring  
21  
22 a sigmoid shaped  $\text{Ca}^{2+}$  level / conductance relationship to the channel (De Stefani and others  
23  
24 2015; Kamer and Mootha 2015; Marchi and Pinton 2014). Indeed,  $\text{Ca}^{2+}$  dependence of MCU  
25  
26 opening to permit  $\text{Zn}^{2+}$  entry could help to explain apparent synergism between  $\text{Ca}^{2+}$  and  $\text{Zn}^{2+}$   
27  
28 effects on mitochondria.  
29  
30  
31  
32  
33  
34

35 Thus, it is apparent that  $\text{Zn}^{2+}$  effects on mitochondria are complex and a better definition  
36  
37 of its mechanisms and how its entry is regulated by the MCU are rich areas for further  
38  
39 investigation. Yet, the potency of its effects, taken together with the high levels of  $\text{Zn}^{2+}$  present  
40  
41 in neurons, highlight the strong potential for  $\text{Zn}^{2+}$  to contribute to mitochondrial dysfunction in  
42  
43 disease.  
44  
45  
46  
47

48 **Cell culture studies: Neuronal  $\text{Zn}^{2+}$  entry results in mitochondrial**  
49  
50  
51 **accumulation and dysfunction contributing to cell death**  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Culture studies permit investigation of  $Zn^{2+}$  effects in the neuronal environment, bringing  
5  
6 us a step closer to understanding possible effects in diseases like ischemia. Above, we  
7  
8 introduced studies examining routes through which synaptic  $Zn^{2+}$  could enter neurons, and  
9  
10 reported evidence for particularly rapid entry through selectively expressed Ca-AMPA, with  
11  
12 slower entry through VGCC. We subsequently examined effects of this  $Zn^{2+}$  entry, and found  
13  
14 brief Ca-AMPA activation, in the presence of 100-300  $\mu M Zn^{2+}$ , to induce rapid loss of  $\Delta\Psi_{mito}$   
15  
16 and ROS generation that persisted for at least an hour after the exposure, consistent with the  
17  
18 potent neurotoxicity of these exposures. Identical kainate exposures with physiological (1.8 mM)  
19  
20  $Ca^{2+}$ , but no  $Zn^{2+}$ , triggered smaller and transient episodes of ROS generation. However, if  $Zn^{2+}$   
21  
22 and  $Ca^{2+}$  were both present during the exposure, the ROS production was significantly greater  
23  
24 than with  $Zn^{2+}$  alone, again indicating synergistic effects of these ions (Sensi and others 1999;  
25  
26 Sensi and others 2000).

27  
28  
29  
30  
31  
32  
33 In other studies, we induced smaller  $Zn^{2+}$  loads, via similar brief  $Zn^{2+}$  exposures under  
34  
35 depolarizing conditions, to trigger entry through VGCC (rather than Ca-AMPA). Although  
36  
37 still causing considerable delayed neurotoxicity (Weiss and others 1993), these exposures did not  
38  
39 cause the acute ROS generation and loss of  $\Delta\Psi_{mito}$  seen with rapid entry through Ca-AMPA  
40  
41 (Sensi and others 1999; Sensi and others 2000). This, along with similar findings by others, have  
42  
43 led to questions as to the likelihood that mitochondria constitute important targets of  $Zn^{2+}$  effects  
44  
45 in disease (Pivovarova and others 2014). However, despite the absence of rapid ROS production,  
46  
47 these brief episodes of  $Zn^{2+}$  entry through VGCC had distinct and long lasting effects on  
48  
49 mitochondria, with low (50-100  $\mu M$ ) exposures resulting in  $Zn^{2+}$  accumulation within  
50  
51 mitochondria persisting for at least 2 hours after the exposure along with partial loss of  $\Delta\Psi_{mito}$   
52  
53 (Sensi and others 2002); similar brief exposures with 300  $\mu M Zn^{2+}$  (and 1.8 mM  $Ca^{2+}$ ) triggered  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4 mitochondrial swelling, and delayed release of apoptotic mediators (cytochrome C and apoptosis  
5 inducing factor) (Jiang and others 2001), possibly consistent with more slowly evolving cell  
6  
7 death occurring after these exposures.  
8  
9

10  
11  
12 Notably, cytosolic  $Zn^{2+}$  accumulation results not only from entry of extracellular  $Zn^{2+}$ ,  
13  
14 but also upon mobilization from cytosolic pools like MT-III, and studies of the effects of strong  
15  
16 cytosolic  $Zn^{2+}$  mobilization alone also have found it to induce effects on mitochondria,  
17  
18 contributing to loss of  $\Delta\Psi_{mito}$  and delayed degeneration (Bossy-Wetzel and others 2004; Sensi  
19  
20 and others 2003). In addition, recent studies have highlighted possible contributions of such  
21  
22  $Zn^{2+}$  mobilization and consequent mitochondrial dysfunction to the  $Ca^{2+}$  dependent excitotoxic  
23  
24 injury cascade (Granzotto and Sensi 2015). In pathologic conditions like ischemia or seizures,  
25  
26 where synaptic  $Zn^{2+}$  release and mobilization from cytosolic buffers both occur, it is likely that  
27  
28 both sources contribute to mitochondrial dysfunction. Indeed, in cell culture studies we find  
29  
30 evidence for synergistic impact upon mitochondria, with even brief and quite low levels of  $Zn^{2+}$   
31  
32 entry through VGCC (which alone had little or no acute effect on mitochondria), when combined  
33  
34 with disrupted buffering (using DTDP, that also by itself had little or no effect), resulting in  
35  
36 dramatic potentiation of acute mitochondrial ROS generation and loss of  $\Delta\Psi_{mito}$ , long lasting  
37  
38 inhibition of mitochondrial respiration, and cell death (Clausen and others 2013; Ji and Weiss  
39  
40 2018). Furthermore, although the presence of physiological  $Ca^{2+}$  during the brief  $Zn^{2+}$  exposure  
41  
42 attenuated cytosolic  $Zn^{2+}$  loading (due to competition with  $Zn^{2+}$  for entry through VGCC), the  
43  
44 effects on mitochondrial function and cell death were markedly enhanced, further highlighting  
45  
46 the synergistic effects of these two ions. Indeed, the strong correlation between effects of  
47  
48 disrupted buffering and presence of  $Ca^{2+}$  on mitochondrial function with those on consequent  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 cell death provide further support to the hypothesis that mitochondrial disruption contributes  
5  
6 directly to  $Zn^{2+}$  triggered neurotoxicity (Ji and Weiss 2018).  
7  
8  
9

10 Thus, these findings not only indicate the potency with which  $Zn^{2+}$  accumulation in  
11  
12 neurons can cause mitochondrial dysfunction, they further support the contention that during *in*  
13  
14 *vivo* ischemia, even low level  $Zn^{2+}$  entry from the extracellular space, when combined with  
15  
16 impaired intracellular  $Zn^{2+}$  buffering and mobilization from intracellular pools, has potential to  
17  
18 powerfully disrupt mitochondrial function and contribute to subsequent neuronal injury.  
19  
20  
21

## 22 Slice and *in vivo* studies support contributions of mitochondrial $Zn^{2+}$ to 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Although the studies discussed above demonstrate that exogenously applied  $Zn^{2+}$  can  
impact mitochondria and contribute to neuronal injury, this does not indicate that endogenous  
 $Zn^{2+}$  actually does so in ischemia. However, recent studies in more pathophysiologically  
relevant ischemia models provide compelling evidence that mitochondria are indeed important  
targets of endogenous  $Zn^{2+}$  effects. Specifically, in one study, addition of extracellular  $Zn^{2+}$   
chelators shortly after a transient episode of ischemia reduced the subsequent mitochondrial  
release of pro-apoptotic peptides (Calderone and others 2004). In another *in vivo* study,  $Zn^{2+}$   
was found to accumulate in mitochondria within 1 hour after transient ischemia, contributing to  
the opening of large, multi-conductance outer membrane channels (Bonanni and others 2006).  
However, whereas these studies demonstrate that  $Zn^{2+}$  contributes to mitochondrial dysfunction  
after *in vivo* ischemia, they do not address therapeutically crucial questions including the source  
and time course of the  $Zn^{2+}$  accumulation, and potential avenues for beneficial interventions.

1  
2  
3  
4 To examine these issues, we have undertaken studies using hippocampal slice OGD  
5  
6 models, a paradigm that models aspects of *in vivo* ischemia while permitting precise control of  
7  
8 the microenvironment and detailed measurement of cellular responses. Our early studies in this  
9  
10 model (see **Fig. 2A, B**) found cytosolic  $Zn^{2+}$  rises to precede and contribute to the onset of  
11  
12 delayed  $Ca^{2+}$  deregulation and cell death during prolonged, lethal OGD (Medvedeva and others  
13  
14 2009), with evidence for early  $Zn^{2+}$  entry into mitochondria. Subsequent studies provided strong  
15  
16 evidence that  $Zn^{2+}$  entry specifically through the MCU is a critical early step, triggering  
17  
18 mitochondrial dysfunction (including ROS production) that contributes to the occurrence of  
19  
20 acute  $Ca^{2+}$  deregulation and degeneration of CA1 neurons (**Fig. 2C**) (Medvedeva and Weiss  
21  
22 2014).  
23  
24  
25  
26  
27

28 In further studies using this slice OGD model, we have compared the contributions and  
29  
30 sources of  $Zn^{2+}$  between CA1 and CA3 neurons (Medvedeva and others 2017). First, we found  
31  
32 that neuronal  $Zn^{2+}$  accumulation contributes to a similar extent in both subdomains, with early  
33  
34  $Zn^{2+}$  rises preceding  $Ca^{2+}$  deregulation, and  $Zn^{2+}$  chelation similarly delaying the onset of the  
35  
36 terminal  $Ca^{2+}$  deregulation in both regions. However, our studies using ZnT3 and MT-III  
37  
38 knockout mice implicated distinct differences in the sources of the  $Zn^{2+}$  underlying acute OGD  
39  
40 induced injury. Paralleling the differences previously noted after prolonged *in vivo* seizures (Lee  
41  
42 and others 2000; Lee and others 2003), synaptic  $Zn^{2+}$  release and its translocation largely  
43  
44 through Ca-AMPA dominated in CA3, and  $Zn^{2+}$  mobilization from MT-III dominated in CA1  
45  
46 (see **Fig. 3A**).  
47  
48  
49  
50  
51

52 **[insert Figure 3 and Figure 3 caption]**  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Because most opportunities for intervention are after reperfusion, we examined events  
5  
6 occurring after sublethal episodes of OGD (which better model transient *in vivo* ischemia) and  
7  
8 found evidence for substantial difference between CA1 and CA3 mitochondria in their handling  
9  
10 of the  $Zn^{2+}$  loads. In these studies we terminated the OGD after the early  $Zn^{2+}$  rises had occurred  
11  
12 but shortly before the time of the terminal  $Ca^{2+}$  deregulation, and in both regions, the cytosolic  
13  
14  $Zn^{2+}$  rises gradually recovered, in part due to uptake into mitochondria via the MCU. However,  
15  
16 at 1 hr after OGD, there was still considerable  $Zn^{2+}$  retained within CA1 mitochondria, whereas  
17  
18 in CA3 mitochondrial  $Zn^{2+}$  loads recovered far more rapidly (generally within 20 min) (see **Fig.**  
19  
20 **3B**) (Medvedeva and others 2017). In light of the differential susceptibilities of CA1 vs CA3  
21  
22 neurons in disease, with CA1 neurons undergoing prominent delayed degeneration after transient  
23  
24 ischemia, associated with mitochondrial swelling and release of cytochrome C (Sugawara and  
25  
26 others 1999), might the persistent  $Zn^{2+}$  accumulation within CA1 mitochondria be a trigger of  
27  
28 events leading to the delayed degeneration of these neurons? Further elucidation of  
29  
30 mitochondrial  $Zn^{2+}$  interactions during and after ischemia in hippocampus as well as in other  
31  
32  $Zn^{2+}$  rich areas of brain (including cortex) may reveal new therapeutic approaches and time  
33  
34 windows for their delivery that may yield improved outcomes.  
35  
36  
37  
38  
39  
40  
41

## 42 Neurodegeneration: The culmination of cascades of injury-promoting 43 events 44 45 46 47 48

49 Cell death is multi-step process, occurring when a sequence of events leads to a state  
50  
51 from which the cell cannot recover. As discussed above, ROS production has been strongly  
52  
53 implicated as a trigger of the neurodegeneration occurring after excitotoxic  $Ca^{2+}$  loading, and  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 downstream events have been identified, including activation of Poly (ADP-ribose) polymerase  
5  
6 (PARP), a nuclear enzyme involved in DNA repair which becomes activated in response to ROS  
7  
8 induced DNA damage. PARP utilizes  $\text{NAD}^+$  as substrate, with strong activity leading to  $\text{NAD}^+$   
9  
10 depletion, glycolytic and mitochondrial inhibition, and release of the apoptotic mediator,  
11  
12 Apoptosis Inducing Factor (AIF) (Kauppinen and Swanson 2007). Paralleling these studies of  
13  
14  $\text{Ca}^{2+}$  excitotoxicity, moderate  $\text{Zn}^{2+}$  exposures have also been found to cause ROS production (in  
15  
16 part due to delayed induction of NOX and neuronal nitric oxide synthase) (Kim and Koh 2002;  
17  
18 Noh and Koh 2000), resulting in PARP activation, that contributes to the evolving injury (Kim  
19  
20 and Koh 2002).  
21  
22  
23  
24  
25

26 A number of pathways have also been described in which early  $\text{Zn}^{2+}$  signals can trigger  
27  
28 more delayed neurodegeneration. Studies of the delayed neurodegeneration caused by strong  
29  
30 intracellular  $\text{Zn}^{2+}$  mobilization (Aizenman and others 2000) have implicated a distinct pathway,  
31  
32 in which activation of p38 MAP kinase results in membrane insertion of Kv2.1  $\text{K}^+$  channels,  
33  
34 resulting in  $\text{K}^+$  efflux from neurons and consequent apoptosis (McLaughlin and others 2001).  
35  
36  
37  
38

39 Notably, these mechanisms contributing to delayed degeneration in response to early  
40  
41  $\text{Zn}^{2+}$  signals represent later steps in cascades, the inciting steps of which are not always apparent.  
42  
43 However, in light of our findings that mitochondrial accumulation of endogenous  $\text{Zn}^{2+}$  under  
44  
45 ischemic conditions triggers rapid mitochondrial ROS production (Medvedeva and Weiss 2014),  
46  
47 *perhaps mitochondrial ROS constitutes a critical upstream trigger of some of these downstream ,*  
48  
49 *neurodegeneration pathways.* Indeed, rapid  $\text{Zn}^{2+}$  triggered mitochondrial ROS could mediate  
50  
51 DNA damage that underlies PARP activation, and has been implicated in the activation of p38  
52  
53 MAP kinase occurring upstream from the insertion of Kv2.1  $\text{K}^+$  channels (Bossy-Wetzel and  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 others 2004), raising the possibility that early targeting of mitochondrial  $Zn^{2+}$  may have both  
5  
6 immediate and delayed therapeutic benefits.  
7  
8  
9

## 10 Therapeutic potential of targeting mitochondrial $Zn^{2+}$ : possible 11 12 13 14 future directions 15 16 17

18 In summary, studies at multiple levels of complexity—ranging from isolated  
19  
20 mitochondria and dissociated neurons, to hippocampal slice and *in vivo* models of ischemia—  
21  
22 indicate that  $Zn^{2+}$  is likely to contribute to mitochondrial dysfunction, ROS generation, and  
23  
24 neurodegeneration in ischemia (and may well do so in prolonged seizures and brain trauma as  
25  
26 well). Furthermore, emerging evidence supports the notion that the  $Zn^{2+}$  entry into mitochondria  
27  
28 is an early event in the ischemic injury cascade (especially in hippocampal CA1), which, as it  
29  
30 occurs upstream from onset of terminal  $Ca^{2+}$  deregulation, may not be adequately targeted by  
31  
32 simply slowing neuronal  $Ca^{2+}$  entry (as via NMDAR blockade). We suggest that:  $Zn^{2+}$   
33  
34 *accumulation in neuronal mitochondria is a targetable early event in the cell death cascade of*  
35  
36 *CA1 and other populations of forebrain neurons; this idea merits further investigation and*  
37  
38 *examination for therapeutic utility.*  
39  
40  
41  
42  
43

44 With strong and prolonged ischemia, mitochondrial  $Zn^{2+}$  loading may result in rapid  
45  
46 irreversible mitochondrial disruption and cell death (Medvedeva and others 2009; Medvedeva  
47  
48 and Weiss 2014) (see **Fig. 2**). However, with milder or transient ischemia, mitochondrial  $Zn^{2+}$   
49  
50 loading may contribute to the activation of downstream cell death pathways. Optimal  
51  
52 interventions might well vary depending on the stage at which they are delivered. We believe  
53  
54 that the targeting of specific events in the injury cascade has potential to yield benefit (see **Fig. 4**):  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 **[insert Figure 4 and Figure 4 caption]**  
5  
6  
7  
8  
9

10 **(1) Early mitochondrial Zn<sup>2+</sup> accumulation:** At the early stages, Zn<sup>2+</sup> chelators or MCU  
11 blockers might provide benefit by lessening early mitochondrial Zn<sup>2+</sup> accumulation. Indeed,  
12 delayed Zn<sup>2+</sup> chelation and MCU blockade have each shown beneficial effects in recent *in vitro*  
13 studies (Ji and Weiss 2018; Medvedeva and others 2017; Slepchenko and others 2017). Of note,  
14 these interventions could also act to promote injurious Ca<sup>2+</sup> loading, possibly complicating  
15 efforts to use them for therapeutic benefit *in vivo*. Specifically, while diminishing mitochondrial  
16 Zn<sup>2+</sup> accumulation, Zn<sup>2+</sup> chelation attenuates physiological antagonism of NMDAR by synaptic  
17 Zn<sup>2+</sup>, thereby increasing neuroexcitation (Cole and others 2000; Dominguez and others 2003;  
18 Vogt and others 2000) and MCU blockade during acute stages of ischemia could diminish  
19 mitochondrial buffering of cytosolic Ca<sup>2+</sup> loads (Velasco and Tapia 2000), both effects that could  
20 exacerbate early injurious cytosolic Ca<sup>2+</sup> loading and hasten Ca<sup>2+</sup> deregulation. For this reason,  
21 in acute stages of ischemia, these agents could show greatest benefit when combined with  
22 maneuvers (such as NMDAR blockade) to abrogate rapid Ca<sup>2+</sup> loading (Medvedeva and Weiss  
23 2014).  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 **(2) Mitochondrial ROS generation:** Antioxidants may provide benefit at slightly later  
44 stages, in two ways: **a.** By diminishing oxidative Zn<sup>2+</sup> mobilization from buffers (thereby  
45 helping to prevent delayed oxidative feedforward amplification of Zn<sup>2+</sup> triggered mitochondrial  
46 disruption), and **b.** By decreasing oxidative tissue damage and activation of oxidant triggered  
47 downstream pathways (including PARP and p38 MAP kinase).  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 **(3) Opening of the mPTP:** Mitochondrial  $Zn^{2+}$  loading may also act upstream to more delayed  
5 apoptotic forms of injury, with  $Zn^{2+}$  triggered mPTP opening (occurring up to several hours after  
6 the  $Zn^{2+}$  load) resulting in mitochondrial disruption and release of apoptotic mediators like  
7 (cytochrome C and AIF), effects against which mPTP blockers (like cyclosporine A) might  
8 provide benefit.  
9  
10  
11  
12  
13  
14  
15

16 **(4) Downstream injury pathways:** As noted above,  $Zn^{2+}$  signals have been found to contribute  
17 to delayed insertion of new ion channels that promote delayed neurodegeneration. Targeting of  
18 these channels (specifically Kv2.1 channels and Ca-AMPA) may yield benefit from hours to  
19 several days after the episode (Aizenman and others 2000; McLaughlin and others 2001; Noh  
20 and others 2005; Yeh and others 2017).  
21  
22  
23  
24  
25  
26  
27  
28

29 In summary, accumulating evidence supports the notion that early mitochondrial  $Zn^{2+}$   
30 accumulation after ischemia contributes to mitochondrial dysfunction and may well be a critical  
31 triggering event for a number of neurodegenerative cascades. The targeting of these  $Zn^{2+}$   
32 triggered events in the post ischemic period has been largely unexplored, yet has potential to  
33 yield substantial benefit, and merits further study.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Figure caption

**Figure 1. Synaptic  $Zn^{2+}$  is released after ischemia.** The mossy fiber pathway (MF) from dentate granule (DG) cells to CA3 pyramidal neurons contains high levels of vesicular  $Zn^{2+}$ , accounting for the dark labeling of this pathway upon Timm's silver sulfide staining (A). Note the loss of synaptic  $Zn^{2+}$  labeling after ischemia (B), resulting from release of this synaptic  $Zn^{2+}$ .

**Figure 2.  $Zn^{2+}$  rise precedes and contributes to lethal  $Ca^{2+}$  deregulation during prolonged oxygen glucose deprivation.** A single CA1 pyramidal neuron in an acute murine hippocampal slice was co-loaded via a patch pipette with the low affinity  $Ca^{2+}$  indicator Fura FF ( $K_d \sim 5.5 \mu M$ ) and the high affinity  $Zn^{2+}$  indicator FluoZin-3 ( $K_d \sim 15 nM$ ), prior to subjecting the slice to prolonged oxygen glucose deprivation (OGD), via perfusion. **A).  $Zn^{2+}$  and  $Ca^{2+}$  responses in a single CA1 hippocampal pyramidal neuron. LEFT:** Pseudocolor Images. Numbers indicate the duration of the OGD exposure (min). Note the early  $Zn^{2+}$  rise (FluoZin-3 fluorescence; 9.4 min), followed after several min by the sharp  $Ca^{2+}$  deregulation event (Fura FF fluorescence; 13.7 min). **RIGHT:** Traces show the time course of the  $Zn^{2+}$  and  $Ca^{2+}$  rises in the same neuron. Responses in this neuron are representative of published findings (Medvedeva and others 2009). **B).  $Zn^{2+}$  contributes to delayed  $Ca^{2+}$  deregulation.** To validate the role of  $Zn^{2+}$  in neuronal injury, hippocampal slices were exposed to OGD alone (control) or in the presence of the  $Zn^{2+}$  chelator N,N,N',N'-tetrakis(2-pyridylmethyl)ethane-1,2-diamine (TPEN;  $40 \mu M$ ). Note that TPEN significantly delayed the onset of the terminal  $Ca^{2+}$  deregulation. Traces show mean  $\pm$  SEM of  $n = 9$ ; from (Medvedeva and others 2017). **C). Schematic of events during lethal OGD.** Numbers refer to events occurring at time points indicated on the traces illustrated

1  
2  
3  
4 in **A. (1)  $Zn^{2+}$  influx into mitochondria:**  $Zn^{2+}$  and  $Ca^{2+}$  enter postsynaptic neurons through  
5 glutamate activated channels.  $Zn^{2+}$  is also mobilized from intracellular buffers (largely MT-III)  
6 as a result of ischemia-associated oxidative stress and acidosis. The cytosolic  $Zn^{2+}$  enters and  
7 accumulates in the mitochondria (via the MCU), contributing to early mitochondrial dysfunction  
8 (including ROS generation and loss of  $\Delta\Psi_{mito}$ ), prior to the sharp cytosolic  $Zn^{2+}$  rise. **(2)**  
9  
10 **Mitochondrial  $Zn^{2+}$  released to cytosol:** After a threshold level of  $Zn^{2+}$  (and  $Ca^{2+}$ ) has entered  
11 the mitochondria, they undergo a rapid depolarization (loss of  $\Delta\Psi_{mito}$ ), and the  $Zn^{2+}$  and  $Ca^{2+}$   
12 sequestered within them are released back into the cytosol. At this point, oxidative stress and  
13 acidosis prevent  $Zn^{2+}$  buffering by MT-III, and the cytosolic  $Zn^{2+}$  rises sharply. **(3)  $Ca^{2+}$**   
14  
15 **deregulation and cell death:** Severe disruption of mitochondrial function and strong ROS  
16 production results in loss of ATP, membrane damage, cellular depolarization, and inability to  
17 clear or sequester the large  $Ca^{2+}$  loads. The sharp cytosolic  $Ca^{2+}$  rises also contribute to  
18 activation of catabolic enzymes, further accelerating cellular disruption and death. Diagram  
19 modified from (Medvedeva and others 2017).  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 **Figure 3. Differential vulnerability of CA1 vs CA3: Dependence on  $Zn^{2+}$  sources and**  
41 **persistence of mitochondrial  $Zn^{2+}$  accumulation. A). Distinct sources of  $Zn^{2+}$  contribute to**  
42 **injury in CA1 vs CA3 pyramidal neurons.** Early cytosolic  $Zn^{2+}$  accumulation contributes to  
43 acute OGD induced injury in both CA1 and CA3 pyramidal neurons. However, in CA1, the  $Zn^{2+}$   
44 largely derives from mobilization from MT-III (**left**), whereas in CA3,  $Zn^{2+}$  translocation  
45 through Ca-AMPA predominates (**right**) (Medvedeva and others 2017). **B).  $Zn^{2+}$  enters**  
46 **mitochondria during OGD in both CA1 and CA3, but after sublethal OGD, persists in**  
47 **mitochondria for prolonged periods only in CA1.** CA1 and CA3 pyramidal neurons were co-  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 loaded with cytosolic  $\text{Ca}^{2+}$  and  $\text{Zn}^{2+}$  indicators, then exposed to either prolonged (lasting until  
5  
6  $\text{Ca}^{2+}$  deregulation; **TOP**) or sublethal (lasting until cytosolic  $\text{Zn}^{2+}$  rise; **MIDDLE AND**  
7  
8 **BOTTOM**) OGD. After sublethal OGD, Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone  
9  
10 (**FCCP**, which induces loss of  $\Delta\Psi_{\text{mito}}$ , releasing mitochondrial  $\text{Zn}^{2+}$  into the cytosol; 2  $\mu\text{M}$ ) and  
11  
12 the MCU blocker Ruthenium Red (**RR**; 10  $\mu\text{M}$ ) were added as indicated. Mitochondrial  
13  
14 diagrams illustrate the anticipated degree of  $\text{Zn}^{2+}$  accumulation (represented by black dots) at  
15  
16 time points indicated by red arrows. Traces and pseudocolor images are reprinted from  
17  
18 (Medvedeva and others 2017). **TOP:** OGD induces rapid mitochondrial  $\text{Zn}^{2+}$  influx in both CA1  
19  
20 and CA3. During OGD, rapid mitochondrial  $\text{Zn}^{2+}$  influx occurs early in both CA1 (**left**) and  
21  
22 CA3 (**right**) pyramidal neurons, contributing to the loss of  $\Delta\Psi_{\text{mito}}$ , release of mitochondrial  $\text{Zn}^{2+}$   
23  
24 into cytosol, and  $\text{Ca}^{2+}$  deregulation. Traces show mean  $\pm$  SEM response of  $n \geq 8$  neurons.  
25  
26 **MIDDLE:**  $\text{Zn}^{2+}$  persists in CA1 mitochondria but is rapidly cleared from CA3 mitochondria  
27  
28 after transient OGD. After sublethal OGD, cytosolic  $\text{Zn}^{2+}$  rises gradually recover in both CA1  
29  
30 and CA3 neurons. To examine the persistence of mitochondrial  $\text{Zn}^{2+}$  accumulation, FCCP was  
31  
32 added as indicated  $\sim 1$  hr after OGD, to depolarize the mitochondria, releasing sequestered  $\text{Zn}^{2+}$ .  
33  
34 Note the strong response to FCCP in CA1 (**left**), indicative of prolonged mitochondrial  $\text{Zn}^{2+}$   
35  
36 sequestration. In contrast, the lack of late FCCP response in CA3 neurons is indicative of the  
37  
38 rapidity with which CA3 mitochondria clear  $\text{Zn}^{2+}$  loads after ischemia (**right**). Traces and  
39  
40 pseudocolor images show responses from representative neurons. **BOTTOM:** Delayed  
41  
42 mitochondrial  $\text{Zn}^{2+}$  uptake depends upon entry through the MCU. Note that application of the  
43  
44 MCU blocker, RR, to CA1 neurons shortly after OGD, while cytosolic  $\text{Zn}^{2+}$  was still elevated,  
45  
46 blocked mitochondrial  $\text{Zn}^{2+}$  uptake, and prevented the protracted mitochondrial  $\text{Zn}^{2+}$   
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 accumulation (as indicated by the lack of FCCP response). Traces show responses of  
5  
6 representative neurons.  
7  
8  
9  
10  
11

12 **Figure 4.  $Zn^{2+}$ -induced mitochondrial dysfunction is a critical and targetable early**  
13 **contributor to ischemic neuronal injury.** During ischemia,  $Zn^{2+}$  accumulation in neurons  
14 reflects contributions from two primary sources:  $Zn^{2+}$  released from presynaptic vesicles that  
15 enters postsynaptic neurons (through Ca-AMPA and VGCC), and  $Zn^{2+}$  released from MT-III  
16 (due to oxidative stress and acidosis) (1). This  $Zn^{2+}$  rapidly enters mitochondria through the  
17 MCU (2). An early consequence of mitochondrial  $Zn^{2+}$  accumulation is acute ROS generation,  
18 which can further disrupt cytosolic  $Zn^{2+}$  buffering, resulting in more mitochondrial  $Zn^{2+}$  entry  
19 and consequent dysfunction, thereby initiating a feedforward “ $Zn^{2+}$ -ROS” cycle. (3). In  
20 addition,  $Zn^{2+}$  can induce delayed activation of NOX, producing more ROS, and possibly further  
21 amplifying this  $Zn^{2+}$ -ROS cycle (4). This protracted  $Zn^{2+}$  influx into mitochondria triggers  
22 mPTP opening, leading to mitochondrial depolarization, swelling, and cytochrome C release (5).  
23 These  $Zn^{2+}$  effects on mitochondria (ROS generation and mPTP opening) can activate major  
24 downstream events, including direct oxidative damage to proteins and DNA (that can lead to  
25 PARP activation), activation of the apoptotic pathway via Caspase 3, and activation of p38 MAP  
26 kinase, promoting the delayed insertion of Kv2.1  $K^+$  channels (6). Furthermore, cytosolic  $Zn^{2+}$ ,  
27 acting through incompletely defined mechanisms, can cause delayed insertion of Ca-AMPA,  
28 further promoting delayed neurodegeneration (7). As these steps are temporally discrete, optimal  
29 therapeutic strategies will likely target a combination of them at different time points, as  
30 highlighted in timeline.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Funding statement

Supported by NIH grants NS065219 and NS096987 (JHW), and grants from the American Heart Association 17GRNT33410181 (JHW) and 16PRE29560003 (SGJ). The authors declare no competing financial interests.

## References

- Aizenman E, Stout AK, Hartnett KA, Dineley KE, McLaughlin B, Reynolds IJ. 2000. Induction of neuronal apoptosis by thiol oxidation: putative role of intracellular zinc release. *J Neurochem* 75: 1878-88.
- Assaf SY, Chung SH. 1984. Release of endogenous Zn<sup>2+</sup> from brain tissue during activity. *Nature* 308: 734-6.
- Ben-Ari Y, Tremblay E, Ottersen OP, Meldrum BS. 1980. The role of epileptic activity in hippocampal and "remote" cerebral lesions induced by kainic acid. *Brain Res* 191: 79-97.
- Bonanni L, Chachar M, Jover-Mengual T, Li H, Jones A, Yokota H, and others. 2006. Zinc-dependent multi-conductance channel activity in mitochondria isolated from ischemic brain. *J Neurosci* 26: 6851-62.
- Bossy-Wetzel E, Talantova MV, Lee WD, Scholzke MN, Harrop A, Mathews E, and others. 2004. Crosstalk between Nitric Oxide and Zinc Pathways to Neuronal Cell Death Involving Mitochondrial Dysfunction and p38-Activated K(+) Channels. *Neuron* 41: 351-65.

- 1  
2  
3  
4 Brennan AM, Suh SW, Won SJ, Narasimhan P, Kauppinen TM, Lee H, and others. 2009.  
5  
6 NADPH oxidase is the primary source of superoxide induced by NMDA receptor activation.  
7  
8 Nat Neurosci 12: 857-63.  
9  
10  
11  
12 Brierley GP. 1967. Ion transport by heart mitochondria. VII. Activation of the energy-linked  
13  
14 accumulation of  $Mg^{++}$  by  $Zn^{++}$  and other cations. J Biol Chem 242: 1115-22.  
15  
16  
17 Brown AM, Kristal BS, Effron MS, Shestopalov AI, Ullucci PA, Sheu KF, and others. 2000.  
18  
19  $Zn^{2+}$  inhibits alpha-ketoglutarate-stimulated mitochondrial respiration and the isolated  
20  
21 alpha-ketoglutarate dehydrogenase complex. J Biol Chem 275: 13441-7.  
22  
23  
24  
25 Calderone A, Jover T, Mashiko T, Noh KM, Tanaka H, Bennett MV, and others. 2004. Late  
26  
27 calcium EDTA rescues hippocampal CA1 neurons from global ischemia-induced death. J  
28  
29 Neurosci 24: 9903-13.  
30  
31  
32  
33 Cheng C, Reynolds IJ. 1998. Calcium-sensitive fluorescent dyes can report increases in  
34  
35 intracellular free zinc concentration in cultured forebrain neurons. J Neurochem 71: 2401-  
36  
37 10.  
38  
39  
40  
41 Choi DW. 1987. Ionic dependence of glutamate neurotoxicity. J Neurosci 7: 369-79.  
42  
43  
44 Choi DW, Koh JY, Peters S. 1988. Pharmacology of glutamate neurotoxicity in cortical cell  
45  
46 culture: attenuation by NMDA antagonists. J Neurosci 8: 185-96.  
47  
48  
49 Clausen A, McClanahan T, Ji SG, Weiss JH. 2013. Mechanisms of Rapid Reactive Oxygen  
50  
51 Species Generation in response to Cytosolic  $Ca^{2+}$  or  $Zn^{2+}$  Loads in Cortical Neurons. Plos  
52  
53 One 8: e83347.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 Cole TB, Robbins CA, Wenzel HJ, Schwartzkroin PA, Palmiter RD. 2000. Seizures and  
5  
6 neuronal damage in mice lacking vesicular zinc. *Epilepsy Res* 39: 153-69.  
7  
8  
9  
10 Cole TB, Wenzel HJ, Kafer KE, Schwartzkroin PA, Palmiter RD. 1999. Elimination of zinc from  
11  
12 synaptic vesicles in the intact mouse brain by disruption of the ZnT3 gene. *Proc Natl Acad*  
13  
14 *Sci U S A* 96: 1716-21.  
15  
16  
17 Colvin RA, Holmes WR, Fontaine CP, Maret W. 2010. Cytosolic zinc buffering and muffling:  
18  
19 their role in intracellular zinc homeostasis. *Metallomics* 2: 306-17.  
20  
21  
22  
23 De Stefani D, Patron M, Rizzuto R. 2015. Structure and function of the mitochondrial calcium  
24  
25 uniporter complex. *Biochim Biophys Acta* 1853: 2006-11.  
26  
27  
28 Devinney MJ, 2nd, Reynolds IJ, Dineley KE. 2005. Simultaneous detection of intracellular free  
29  
30 calcium and zinc using fura-2FF and FluoZin-3. *Cell Calcium* 37: 225-32.  
31  
32  
33  
34 Devinney MJ, Malaiyandi LM, Vergun O, DeFranco DB, Hastings TG, Dineley KE. 2009. A  
35  
36 comparison of Zn<sup>2+</sup>- and Ca<sup>2+</sup>-triggered depolarization of liver mitochondria reveals no  
37  
38 evidence of Zn<sup>2+</sup>-induced permeability transition. *Cell Calcium* 45: 447-55.  
39  
40  
41  
42 Dominguez MI, Blasco-Ibanez JM, Crespo C, Marques-Mari AI, Martinez-Guijarro FJ. 2003.  
43  
44 Zinc chelation during non-lesioning overexcitation results in neuronal death in the mouse  
45  
46 hippocampus. *Neuroscience* 116: 791-806.  
47  
48  
49  
50 Dugan LL, Sensi SL, Canzoniero LM, Handran SD, Rothman SM, Lin TS, and others. 1995.  
51  
52 Mitochondrial production of reactive oxygen species in cortical neurons following exposure  
53  
54 to N-methyl-D-aspartate. *J Neurosci* 15: 6377-88.  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Frederickson CJ. 1989. Neurobiology of zinc and zinc-containing neurons. *Int Rev Neurobiol* 31:  
5  
6 145-238.

7  
8  
9  
10 Frederickson CJ, Giblin LJ, Krezel A, McAdoo DJ, Mueller RN, Zeng Y, and others. 2006.  
11  
12 Concentrations of extracellular free zinc (pZn)<sub>e</sub> in the central nervous system during simple  
13  
14 anesthetization, ischemia and reperfusion. *Exp Neurol* 198: 285-93.

15  
16  
17 Frederickson CJ, Hernandez MD, McGinty JF. 1989. Translocation of zinc may contribute to  
18  
19 seizure-induced death of neurons. *Brain Res* 480: 317-21.

20  
21  
22 Frederickson CJ, Rampy BA, Reamy-Rampy S, Howell GA. 1992. Distribution of  
23  
24 histochemically reactive zinc in the forebrain of the rat. *J Chem Neuroanat* 5: 521-30.

25  
26  
27 Freund WD, Reddig S. 1994. AMPA/Zn(2+)-induced neurotoxicity in rat primary cortical  
28  
29 cultures: involvement of L-type calcium channels. *Brain Res* 654: 257-64.

30  
31  
32  
33 Gazaryan IG, Krasinskaya IP, Kristal BS, Brown AM. 2007. Zinc irreversibly damages major  
34  
35 enzymes of energy production and antioxidant defense prior to mitochondrial permeability  
36  
37 transition. *J Biol Chem* 282: 24373-80.

38  
39  
40  
41 Gorter JA, Petrozzino JJ, Aronica EM, Rosenbaum DM, Opitz T, Bennett MV, and others. 1997.  
42  
43 Global ischemia induces downregulation of Glur2 mRNA and increases AMPA receptor-  
44  
45 mediated Ca<sup>2+</sup> influx in hippocampal CA1 neurons of gerbil. *J Neurosci* 17: 6179-88.

46  
47  
48  
49 Granzotto A, Sensi SL. 2015. Intracellular zinc is a critical intermediate in the excitotoxic  
50  
51 cascade. *Neurobiol Dis* 81: 25-37.



1  
2  
3  
4 Howell GA, Welch MG, Frederickson CJ. 1984. Stimulation-induced uptake and release of zinc  
5  
6 in hippocampal slices. *Nature* 308: 736-8.  
7

8  
9  
10 Hoyte L, Barber PA, Buchan AM, Hill MD. 2004. The rise and fall of NMDA antagonists for  
11  
12 ischemic stroke. *Curr Mol Med* 4: 131-6.  
13

14  
15 Ikonomidou C, Turski L. 2002. Why did NMDA receptor antagonists fail clinical trials for stroke  
16  
17 and traumatic brain injury? *Lancet Neurol* 1: 383-6.  
18

19  
20 Ji SG, Weiss JH. 2018. Zn(2+)-induced disruption of neuronal mitochondrial function:  
21  
22 Synergism with Ca(2+), critical dependence upon cytosolic Zn(2+) buffering, and  
23  
24 contributions to neuronal injury. *Exp Neurol* 302: 181-95.  
25  
26

27  
28 Jia Y, Jeng JM, Sensi SL, Weiss JH. 2002. Zn<sup>2+</sup> currents are mediated by calcium-permeable  
29  
30 AMPA/kainate channels in cultured murine hippocampal neurones. *J Physiol* 543: 35-48.  
31

32  
33 Jiang D, Sullivan PG, Sensi SL, Steward O, Weiss JH. 2001. Zn(2+) induces permeability  
34  
35 transition pore opening and release of pro-apoptotic peptides from neuronal mitochondria. *J*  
36  
37 *Biol Chem* 276: 47524-9.  
38  
39

40  
41 Jiang LJ, Vasak M, Vallee BL, Maret W. 2000. Zinc transfer potentials of the alpha - and beta-  
42  
43 clusters of metallothionein are affected by domain interactions in the whole molecule. *Proc*  
44  
45 *Natl Acad Sci U S A* 97: 2503-8.  
46  
47

48  
49 Kambe T, Hashimoto A, Fujimoto S. 2014. Current understanding of ZIP and ZnT zinc  
50  
51 transporters in human health and diseases. *Cell Mol Life Sci* 71: 3281-95.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Kamer KJ, Mootha VK. 2015. The molecular era of the mitochondrial calcium uniporter. *Nat*  
5  
6 *Rev Mol Cell Biol* 16: 545-53.  
7  
8

9  
10 Kauppinen TM, Swanson RA. 2007. The role of poly(ADP-ribose) polymerase-1 in CNS  
11  
12 disease. *Neuroscience* 145: 1267-72.  
13  
14

15 Kerchner GA, Canzoniero LM, Yu SP, Ling C, Choi DW. 2000. Zn<sup>2+</sup> current is mediated by  
16  
17 voltage-gated Ca<sup>2+</sup> channels and enhanced by extracellular acidity in mouse cortical  
18  
19 neurones. *J Physiol* 528 Pt 1: 39-52.  
20  
21

22  
23 Kim YH, Koh JY. 2002. The role of NADPH oxidase and neuronal nitric oxide synthase in zinc-  
24  
25 induced poly(ADP-ribose) polymerase activation and cell death in cortical culture. *Exp*  
26  
27 *Neurol* 177: 407-18.  
28  
29

30  
31 Koh JY, Choi DW. 1994. Zinc toxicity on cultured cortical neurons: involvement of N-methyl-  
32  
33 D-aspartate receptors. *Neuroscience* 60: 1049-57.  
34  
35

36 Koh JY, Suh SW, Gwag BJ, He YY, Hsu CY, Choi DW. 1996. The role of zinc in selective  
37  
38 neuronal death after transient global cerebral ischemia. *Science* 272: 1013-6.  
39  
40

41 Lafon-Cazal M, Pietri S, Culcasi M, Bockaert J. 1993. NMDA-dependent superoxide production  
42  
43 and neurotoxicity. *Nature* 364: 535-7.  
44  
45

46  
47 Lee JY, Cole TB, Palmiter RD, Koh JY. 2000. Accumulation of zinc in degenerating  
48  
49 hippocampal neurons of ZnT3-null mice after seizures: evidence against synaptic vesicle  
50  
51 origin. *J Neurosci* 20: RC79.  
52  
53

- 1  
2  
3  
4 Lee JY, Kim JH, Palmiter RD, Koh JY. 2003. Zinc released from metallothionein-iii may  
5  
6 contribute to hippocampal CA1 and thalamic neuronal death following acute brain injury.  
7  
8 Exp Neurol 184: 337-47.  
9  
10  
11  
12 Lerma J, Morales M, Ibarz JM, Somohano F. 1994. Rectification properties and Ca<sup>2+</sup>  
13  
14 permeability of glutamate receptor channels in hippocampal cells. Eur J Neurosci 6: 1080-8.  
15  
16  
17 Link TA, von Jagow G. 1995. Zinc ions inhibit the QP center of bovine heart mitochondrial bc1  
18  
19 complex by blocking a protonatable group. J Biol Chem 270: 25001-6.  
20  
21  
22  
23 Malaiyandi LM, Dineley KE, Reynolds IJ. 2004. Divergent consequences arise from  
24  
25 metallothionein overexpression in astrocytes: zinc buffering and oxidant-induced zinc  
26  
27 release. Glia 45: 346-53.  
28  
29  
30  
31 Marchi S, Pinton P. 2014. The mitochondrial calcium uniporter complex: molecular components,  
32  
33 structure and physiopathological implications. J Physiol 592: 829-39.  
34  
35  
36  
37 Maret W. 1995. Metallothionein/disulfide interactions, oxidative stress, and the mobilization of  
38  
39 cellular zinc. Neurochem Int 27: 111-7.  
40  
41  
42 Maret W. 2015. Analyzing free zinc(II) ion concentrations in cell biology with fluorescent  
43  
44 chelating molecules. Metallomics 7: 202-11.  
45  
46  
47 McLaughlin B, Pal S, Tran MP, Parsons AA, Barone FC, Erhardt JA, and others. 2001. p38  
48  
49 activation is required upstream of potassium current enhancement and caspase cleavage in  
50  
51 thiol oxidant-induced neuronal apoptosis. J Neurosci 21: 3303-11.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 Medvedeva YV, Ji SG, Yin HZ, Weiss JH. 2017. Differential Vulnerability of CA1 versus CA3  
5  
6 Pyramidal Neurons After Ischemia: Possible Relationship to Sources of Zn<sup>2+</sup> Accumulation  
7  
8 and Its Entry into and Prolonged Effects on Mitochondria. *J Neurosci* 37: 726-37.  
9  
10  
11  
12 Medvedeva YV, Lin B, Shuttleworth CW, Weiss JH. 2009. Intracellular Zn<sup>2+</sup> accumulation  
13  
14 contributes to synaptic failure, mitochondrial depolarization, and cell death in an acute slice  
15  
16 oxygen-glucose deprivation model of ischemia. *J Neurosci* 29: 1105-14.  
17  
18  
19  
20 Medvedeva YV, Weiss JH. 2014. Intramitochondrial Zn(2+) accumulation via the Ca(2+)  
21  
22 uniporter contributes to acute ischemic neurodegeneration. *Neurobiol Dis* 68: 137-44.  
23  
24  
25  
26 Nicholls DG, Budd SL. 2000. Mitochondria and neuronal survival. *Physiol Rev* 80: 315-60.  
27  
28  
29  
30 Noh KM, Koh JY. 2000. Induction and activation by zinc of NADPH oxidase in cultured cortical  
31  
32 neurons and astrocytes. *J Neurosci* 20: RC111.  
33  
34  
35  
36 Noh KM, Yokota H, Mashiko T, Castillo PE, Zukin RS, Bennett MV. 2005. Blockade of  
37  
38 calcium-permeable AMPA receptors protects hippocampal neurons against global ischemia-  
39  
40 induced death. *Proc Natl Acad Sci U S A* 102: 12230-5.  
41  
42  
43  
44 Ogoshi F, Weiss JH. 2003. Heterogeneity of Ca<sup>2+</sup>-permeable AMPA/kainate channel expression  
45  
46 in hippocampal pyramidal neurons: Fluorescence imaging and immunocytochemical  
47  
48 assessment. *J Neurosci* 23: 10521-30  
49  
50  
51  
52 Pivovarova NB, Stanika RI, Kazanina G, Villanueva I, Andrews SB. 2014. The interactive roles  
53  
54 of zinc and calcium in mitochondrial dysfunction and neurodegeneration. *J Neurochem* 128:  
55  
56 592-602.  
57  
58  
59  
60

- 1  
2  
3  
4 Randall RD, Thayer SA. 1992. Glutamate-induced calcium transient triggers delayed calcium  
5  
6 overload and neurotoxicity in rat hippocampal neurons. *J Neurosci* 12: 1882-95.  
7  
8  
9  
10 Reynolds IJ, Hastings TG. 1995. Glutamate induces the production of reactive oxygen species in  
11  
12 cultured forebrain neurons following NMDA receptor activation. *J Neurosci* 15: 3318-27.  
13  
14  
15 Saris NE, Niva K. 1994. Is Zn<sup>2+</sup> transported by the mitochondrial calcium uniporter? *FEBS Lett*  
16  
17 356: 195-8.  
18  
19  
20 Sattler R, Xiong Z, Lu WY, Hafner M, MacDonald JF, Tymianski M. 1999. Specific coupling of  
21  
22 NMDA receptor activation to nitric oxide neurotoxicity by PSD-95 protein. *Science* 284:  
23  
24 1845-8.  
25  
26  
27  
28 Sensi SL, Paoletti P, Koh JY, Aizenman E, Bush AI, Hershfinkel M. 2011. The neurophysiology  
29  
30 and pathology of brain zinc. *J Neurosci* 31: 16076-85.  
31  
32  
33  
34 Sensi SL, Ton-That D, Sullivan PG, Jonas EA, Gee KR, Kaczmarek LK, and others. 2003.  
35  
36 Modulation of mitochondrial function by endogenous Zn<sup>2+</sup> pools. *Proc Natl Acad Sci U S*  
37  
38 *A* 100: 6157-62.  
39  
40  
41 Sensi SL, Ton-That D, Weiss JH. 2002. Mitochondrial sequestration and Ca(2+)-dependent  
42  
43 release of cytosolic Zn(2+) loads in cortical neurons. *Neurobiol Dis* 10: 100-8.  
44  
45  
46  
47 Sensi SL, Yin HZ, Carriedo SG, Rao SS, Weiss JH. 1999. Preferential Zn<sup>2+</sup> influx through  
48  
49 Ca<sup>2+</sup>-permeable AMPA/kainate channels triggers prolonged mitochondrial superoxide  
50  
51 production. *Proc Natl Acad Sci U S A* 96: 2414-9.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 Sensi SL, Yin HZ, Weiss JH. 2000. AMPA/kainate receptor-triggered Zn<sup>2+</sup> entry into cortical  
5  
6 neurons induces mitochondrial Zn<sup>2+</sup> uptake and persistent mitochondrial dysfunction. *Eur J*  
7  
8 *Neurosci* 12: 3813-8.  
9  
10  
11  
12 Skulachev VP, Chistyakov VV, Jasaitis AA, Smirnova EG. 1967. Inhibition of the respiratory  
13  
14 chain by zinc ions. *Biochem Biophys Res Commun* 26: 1-6.  
15  
16  
17 Slepchenko KG, Lu Q, Li YV. 2017. Cross talk between increased intracellular zinc (Zn<sup>2+</sup>) and  
18  
19 accumulation of reactive oxygen species in chemical ischemia. *Am J Physiol Cell Physiol*  
20  
21 313: C448-C59.  
22  
23  
24  
25 Sloviter RS. 1985. A selective loss of hippocampal mossy fiber Timm stain accompanies granule  
26  
27 cell seizure activity induced by perforant path stimulation. *Brain Res* 330: 150-3.  
28  
29  
30  
31 Stork CJ, Li YV. 2006. Intracellular zinc elevation measured with a "calcium-specific" indicator  
32  
33 during ischemia and reperfusion in rat hippocampus: a question on calcium overload. *J*  
34  
35 *Neurosci* 26: 10430-7.  
36  
37  
38  
39 Sugawara T, Fujimura M, Morita-Fujimura Y, Kawase M, Chan PH. 1999. Mitochondrial release  
40  
41 of cytochrome c corresponds to the selective vulnerability of hippocampal CA1 neurons in  
42  
43 rats after transient global cerebral ischemia. *J Neurosci* 19: RC39.  
44  
45  
46  
47 Suh SW, Chen JW, Motamedi M, Bell B, Listiak K, Pons NF, and others. 2000. Evidence that  
48  
49 synaptically-released zinc contributes to neuronal injury after traumatic brain injury. *Brain*  
50  
51 *Res* 852: 268-73.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Thibault O, Landfield PW. 1996. Increase in single L-type calcium channels in hippocampal  
5  
6 neurons during aging. *Science* 272: 1017-20.  
7  
8

9  
10 Tonder N, Johansen FF, Frederickson CJ, Zimmer J, Diemer NH. 1990. Possible role of zinc in  
11  
12 the selective degeneration of dentate hilar neurons after cerebral ischemia in the adult rat.  
13  
14 *Neurosci Lett* 109: 247-52.  
15  
16

17  
18 Ueno S, Tsukamoto M, Hirano T, Kikuchi K, Yamada MK, Nishiyama N, and others. 2002.  
19  
20 Mossy fiber Zn<sup>2+</sup> spillover modulates heterosynaptic N-methyl-D-aspartate receptor  
21  
22 activity in hippocampal CA3 circuits. *J Cell Biol* 158: 215-20.  
23  
24

25  
26 Velasco I, Tapia R. 2000. Alterations of intracellular calcium homeostasis and mitochondrial  
27  
28 function are involved in ruthenium red neurotoxicity in primary cortical cultures. *J Neurosci*  
29  
30 *Res* 60: 543-51.  
31  
32

33  
34 Vogt K, Mellor J, Tong G, Nicoll R. 2000. The actions of synaptically released zinc at  
35  
36 hippocampal mossy fiber synapses. *Neuron* 26: 187-96.  
37  
38


39  
40 Wang GJ, Thayer SA. 1996. Sequestration of glutamate-induced Ca<sup>2+</sup> loads by mitochondria in  
41  
42 cultured rat hippocampal neurons. *J Neurophysiol* 76: 1611-21.  
43  
44

45  
46 Weiss JH, Hartley DM, Koh JY, Choi DW. 1993. AMPA receptor activation potentiates zinc  
47  
48 neurotoxicity. *Neuron* 10: 43-9.  
49  
50

51  
52 White RJ, Reynolds IJ. 1997. Mitochondria accumulate Ca<sup>2+</sup> following intense glutamate  
53  
54 stimulation of cultured rat forebrain neurones. *J Physiol* 498 ( Pt 1): 31-47.  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 Wudarczyk J, Debska G, Lenartowicz E. 1999. Zinc as an inducer of the membrane permeability  
5 transition in rat liver mitochondria. *Arch Biochem Biophys* 363: 1-8.  
6  
7  
8  
9  
10 Yeh CY, Bulas AM, Moutal A, Saloman JL, Hartnett KA, Anderson CT, and others. 2017.  
11 Targeting a Potassium Channel/Syntaxin Interaction Ameliorates Cell Death in Ischemic  
12 Stroke. *J Neurosci* 37: 5648-58.  
13  
14  
15  
16  
17 Yin H, Turetsky D, Choi DW, Weiss JH. 1994. Cortical neurones with Ca<sup>2+</sup> permeable  
18 AMPA/kainate channels display distinct receptor immunoreactivity and are GABAergic.  
19 *Neurobiol Dis* 1: 43-9.  
20  
21  
22  
23  
24  
25 Yin HZ, Sensi SL, Carriedo SG, Weiss JH. 1999. Dendritic localization of Ca(2+)-permeable  
26 AMPA/kainate channels in hippocampal pyramidal neurons. *J Comp Neurol* 409: 250-60.  
27  
28  
29  
30  
31 Yin HZ, Sensi SL, Ogoshi F, Weiss JH. 2002. Blockade of Ca<sup>2+</sup>-permeable AMPA/kainate  
32 channels decreases oxygen-glucose deprivation-induced Zn<sup>2+</sup> accumulation and neuronal  
33 loss in hippocampal pyramidal neurons. *J Neurosci* 22: 1273-9.  
34  
35  
36  
37  
38  
39 Yin HZ, Weiss JH. 1995. Zn(2+) permeates Ca(2+) permeable AMPA/kainate channels and  
40 triggers selective neural injury. *Neuroreport* 6: 2553-6.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





45 Figure 1. Synaptic  $Zn^{2+}$  is released after ischemia. The mossy fiber pathway (MF) from dentate granule  
46 (DG) cells to CA3 pyramidal neurons contains high levels of vesicular  $Zn^{2+}$ , accounting for the dark labeling  
47 of this pathway upon Timm's silver sulfide staining (A). Note the loss of synaptic  $Zn^{2+}$  labeling after  
48 ischemia (B), resulting from release of this synaptic  $Zn^{2+}$ .

49 161x196mm (300 x 300 DPI)




Figure 2.  $Zn^{2+}$  rise precedes and contributes to lethal  $Ca^{2+}$  deregulation during prolonged oxygen glucose deprivation. A single CA1 pyramidal neuron in an acute murine hippocampal slice was co-loaded via a patch pipette with the low affinity  $Ca^{2+}$  indicator Fura FF ( $K_d \sim 5.5 \mu M$ ) and the high affinity  $Zn^{2+}$  indicator FluoZin-3 ( $K_d \sim 15 nM$ ), prior to subjecting the slice to prolonged oxygen glucose deprivation (OGD), via perfusion. A).  $Zn^{2+}$  and  $Ca^{2+}$  responses in a single CA1 hippocampal pyramidal neuron.

LEFT: Pseudocolor Images. Numbers indicate the duration of the OGD exposure (min). Note the early  $Zn^{2+}$  rise (FluoZin-3 fluorescence; 9.4 min), followed after several min by the sharp  $Ca^{2+}$  deregulation event (Fura FF fluorescence; 13.7 min). RIGHT: Traces show the time course of the  $Zn^{2+}$  and  $Ca^{2+}$  rises in the same neuron. Responses in this neuron are representative of published findings (Medvedeva and others 2009). B).  $Zn^{2+}$  contributes to delayed  $Ca^{2+}$  deregulation. To validate the role of  $Zn^{2+}$  in neuronal injury, hippocampal slices were exposed to OGD alone (control) or in the presence of the  $Zn^{2+}$  chelator N,N,N',N'-tetrakis(2-pyridylmethyl)ethane-1,2-diamine (TPEN; 40  $\mu M$ ). Note that TPEN significantly delayed the onset of the terminal  $Ca^{2+}$  deregulation. Traces show mean  $\pm$  SEM of  $n = 9$ ; from (Medvedeva and others 2017). C). Schematic of events during lethal OGD. Numbers refer to events occurring at time points indicated on the traces illustrated in A. (1)  $Zn^{2+}$  influx into mitochondria:  $Zn^{2+}$  and  $Ca^{2+}$  enter postsynaptic neurons through glutamate activated channels.  $Zn^{2+}$  is also mobilized from intracellular buffers (largely MT-III) as a result of ischemia-associated oxidative stress and acidosis. The cytosolic  $Zn^{2+}$  enters and accumulates in the mitochondria (via the MCU), contributing to early mitochondrial dysfunction (including ROS generation and loss of  $\Delta\Psi_{mito}$ ), prior to the sharp cytosolic  $Zn^{2+}$  rise. (2) Mitochondrial  $Zn^{2+}$  released to cytosol: After a threshold level of  $Zn^{2+}$  (and  $Ca^{2+}$ ) has entered the mitochondria, they undergo a rapid depolarization (loss of  $\Delta\Psi_{mito}$ ), and the  $Zn^{2+}$  and  $Ca^{2+}$  sequestered within them are released back into the cytosol. At this point, oxidative stress and acidosis prevent  $Zn^{2+}$  buffering by MT-III, and the cytosolic  $Zn^{2+}$  rises sharply. (3)  $Ca^{2+}$  deregulation and cell death: Severe disruption of mitochondrial function and strong ROS production results in loss of ATP, membrane damage, cellular depolarization, and inability to clear or sequester the large  $Ca^{2+}$  loads. The sharp cytosolic  $Ca^{2+}$  rises also contribute to activation of catabolic enzymes, further accelerating cellular disruption and death. Diagram modified from (Medvedeva and others 2017).

309x190mm (300 x 300 DPI)




Figure 3. Differential vulnerability of CA1 vs CA3: Dependence on  $Zn^{2+}$  sources and persistence of mitochondrial  $Zn^{2+}$  accumulation. A). Distinct sources of  $Zn^{2+}$  contribute to injury in CA1 vs CA3 pyramidal neurons. Early cytosolic  $Zn^{2+}$  accumulation contributes to acute OGD induced injury in both CA1 and CA3 pyramidal neurons. However, in CA1, the  $Zn^{2+}$  largely derives from mobilization from MT-III (left), whereas in CA3,  $Zn^{2+}$  translocation through Ca-AMPA receptor predominates (right) (Medvedeva and others 2017). B).  $Zn^{2+}$  enters mitochondria for prolonged periods only in CA1. CA1 and CA3 pyramidal neurons were co-loaded with cytosolic  $Ca^{2+}$  and  $Zn^{2+}$  indicators, then exposed to either prolonged (lasting until  $Ca^{2+}$  deregulation; TOP) or sublethal (lasting until cytosolic  $Zn^{2+}$  rise; MIDDLE AND BOTTOM) OGD. After sublethal OGD, Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone (FCCP, which induces loss of  $\Delta\Psi_{mito}$ , releasing mitochondrial  $Zn^{2+}$  into the cytosol; 2  $\mu M$ ) and the MCU blocker Ruthenium Red (RR; 10  $\mu M$ ) were added as indicated. Mitochondrial diagrams illustrate the anticipated degree of  $Zn^{2+}$  accumulation (represented by black dots) at time points indicated by red arrows. Traces and pseudocolor images are reprinted from

1  
2  
3 (Medvedeva and others 2017). TOP: OGD induces rapid mitochondrial Zn<sup>2+</sup> influx in both CA1 and  
4 CA3. During OGD, rapid mitochondrial Zn<sup>2+</sup> influx occurs early in both CA1 (left) and CA3 (right) pyramidal  
5 neurons, contributing to the loss of  $\Delta\Psi_{\text{mito}}$ , release of mitochondrial Zn<sup>2+</sup> into cytosol, and Ca<sup>2+</sup>  
6 deregulation. Traces show mean  $\pm$  SEM response of  $n \geq 8$  neurons. MIDDLE: Zn<sup>2+</sup> persists in CA1  
7 mitochondria but is rapidly cleared from CA3 mitochondria after transient OGD. After sublethal OGD,  
8 cytosolic Zn<sup>2+</sup> rises gradually recover in both CA1 and CA3 neurons. To examine the persistence of  
9 mitochondrial Zn<sup>2+</sup> accumulation, FCCP was added as indicated  $\sim 1$  hr after OGD, to depolarize the  
10 mitochondria, releasing sequestered Zn<sup>2+</sup>. Note the strong response to FCCP in CA1 (left), indicative of  
11 prolonged mitochondrial Zn<sup>2+</sup> sequestration. In contrast, the lack of late FCCP response in CA3 neurons is  
12 indicative of the rapidity with which CA3 mitochondria clear Zn<sup>2+</sup> loads after ischemia (right). Traces and  
13 pseudocolor images show responses from representative neurons. BOTTOM: Delayed mitochondrial Zn<sup>2+</sup>  
14 uptake depends upon entry through the MCU. Note that application of the MCU blocker, RR, to CA1 neurons  
15 shortly after OGD, while cytosolic Zn<sup>2+</sup> was still elevated, blocked mitochondrial Zn<sup>2+</sup> uptake, and  
16 prevented the protracted mitochondrial Zn<sup>2+</sup> accumulation (as indicated by the lack of FCCP  
17 response). Traces show responses of representative neurons.

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

148x338mm (300 x 300 DPI)




Figure 4. Zn<sup>2+</sup>-induced mitochondrial dysfunction is a critical and targetable early contributor to ischemic neuronal injury. During ischemia, Zn<sup>2+</sup> accumulation in neurons reflects contributions from two primary sources: Zn<sup>2+</sup> released from presynaptic vesicles that enters postsynaptic neurons (through Ca-AMPA and VGCC), and Zn<sup>2+</sup> released from MT-III (due to oxidative stress and acidosis) (1). This Zn<sup>2+</sup> rapidly enters mitochondria through the MCU (2). An early consequence of mitochondrial Zn<sup>2+</sup> accumulation is acute ROS generation, which can further disrupt cytosolic Zn<sup>2+</sup> buffering, resulting in more mitochondrial Zn<sup>2+</sup> entry and consequent dysfunction, thereby initiating a feedforward "Zn<sup>2+</sup>-ROS" cycle. (3). In addition, Zn<sup>2+</sup> can induce delayed activation of NOX, producing more ROS, and possibly further amplifying this Zn<sup>2+</sup>-ROS cycle (4). This protracted Zn<sup>2+</sup> influx into mitochondria triggers mPTP opening, leading to mitochondrial depolarization, swelling, and cytochrome C release (5). These Zn<sup>2+</sup> effects on mitochondria (ROS generation and mPTP opening) can activate major downstream events, including direct oxidative damage to proteins and DNA (that can lead to PARP activation), activation of the apoptotic pathway via Caspase 3, and activation of p38 MAP kinase, promoting the delayed insertion of Kv2.1 K<sup>+</sup> channels (6). Furthermore,

1  
2  
3 cytosolic Zn<sup>2+</sup>, acting through incompletely defined mechanisms, can cause delayed insertion of Ca-AMPA,  
4 further promoting delayed neurodegeneration (7). As these steps are temporally discrete, optimal  
5 therapeutic strategies will likely target a combination of them at different time points, as highlighted in  
6 timeline.

7 190x338mm (300 x 300 DPI)

8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review