

UC Irvine

UC Irvine Previously Published Works

Title

On the domain of analyticity and small scales for the solutions of the damped-driven 2D Navier-Stokes equations

Permalink

<https://escholarship.org/uc/item/0xp23266>

Journal

Dynamics of Partial Differential Equations, 4(2)

ISSN

1548-159X

Authors

Ilyin, Alexi A.
Titi, Edriss A.

Publication Date

2007-06-01

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

ON THE DOMAIN OF ANALYTICITY AND SMALL SCALES FOR THE SOLUTIONS OF THE DAMPED-DRIVEN 2D NAVIER–STOKES EQUATIONS

ALEXEI A. ILYIN¹ AND EDRISS S. TITI²

Abstract. We obtain a logarithmically sharp estimate for the space-analyticity radius of the solutions of the damped-driven 2D Navier–Stokes equations with periodic boundary conditions and relate this to the small scales in this system. This system is inspired by the Stommel–Charney barotropic ocean circulation model.

Key words: Analyticity, Gevrey regularity, Navier–Stokes equations, dissipative length scales, Stommel–Charney model.

AMS subject classification: 35B41, 35Q30, 37L30.

1. INTRODUCTION

It was shown in [15] (see also [3], [12]) that the solutions of the 2D Navier–Stokes equations with periodic boundary conditions belong to the Gevrey class of analytic functions (if the forcing term does). Using the Gevrey regularity approach the following estimate for the spatial analyticity radius for the solutions that lie on the global attractor (or are near it) was obtained

$$l_a \geq \frac{c|\Omega|^{1/2}}{G^2 \log G}, \quad (1.1)$$

where $G = \|f\|_{L_2}|\Omega|/\nu^2$ is the Grashof number and $|\Omega| = L^2/\gamma$ is the area of the periodic domain $\Omega = [0, L/\gamma] \times [0, L]$, $\gamma \leq 1$.

Therefore, the Fourier coefficients \hat{u}_k are exponentially small for $|k| \gg L/l_a$, and l_a naturally forms a lower bound for the small dissipative length scale for the system (see, for instance, [11]).

There are other ways of estimating the dissipative small length scale for the Navier–Stokes system, for instance, in terms of the dimension of the global attractor [1], [6], [7], [12], [39]. The Hausdorff and fractal dimensions of the global attractor satisfy the following estimate [8] (see also [6], [39]):

$$\dim_F \mathcal{A} \leq c_1 G^{2/3} (\log(1 + G))^{1/3}, \quad c_1 = c_1(\gamma)$$

which has been shown in [33] (following ideas of [1]) to be logarithmically sharp.

If we accept the point of view that the small length scale can be defined as follows (see [7], [12], [36], [39])

$$l_f \sim \left(\frac{|\Omega|}{\dim_F \mathcal{A}} \right)^{1/2}, \quad (1.2)$$

DATE: JANUARY 18, 2007.

¹Keldysh Institute of Applied Mathematics, Russian Academy of Sciences, Miusskaya Sq. 4, 125047 Moscow, Russia, E-mail: ilyin@keldysh.ru

²Department of Mathematics and Department of Mechanical and Aerospace Engineering, University of California, Irvine, California 92697, USA, E-mail: etiti@math.uci.edu. Also: Department of Computer Science and Applied Mathematics, Weizmann Institute of Science, P.O. Box 26, Rehovot, 76100, Israel, E-mail: edriss.titi@weizmann.ac.il

then up to logarithmic correction we have

$$l_f \sim \frac{|\Omega|^{1/2}}{G^{1/3}}. \quad (1.3)$$

This heuristic estimate for the small length scale is probably the best one can hope for since it matches, up to logarithmic term, the physically asserted estimates for the enstrophy dissipation length scale [30]. We also observe that the estimate (1.3) is extensive, that is, independent of the size of the spatial domain provided that its shape is fixed.

Another rigorous definition of the small length scale can be given in terms of the number of determining modes, nodes, or volume elements (see [12], [14], [16], [28] and the references therein). It was shown that if N is sufficiently large and N equal squares of size l_{dn} tile the periodic spatial domain, then any collection of points (one in each square) are determining for the long time dynamics of the 2D Navier–Stokes system. The best to date estimate for N was obtained in [28]:

$$N \leq c_2 G,$$

where $c_2 = c_2(\gamma)$ depends only on the aspect ratio $\gamma \leq 1$. (An explicit estimate for c_2 was obtained in [26]: $c_2(\gamma) = (68/(\gamma\pi))^{1/2}$.)

Therefore the small length scale defined in terms of the lattice of determining nodes satisfies

$$l_{dn} \geq c_2(\gamma)^{-1/2} \frac{|\Omega|^{1/2}}{G^{1/2}}. \quad (1.4)$$

We observe that this estimate is not extensive, that is, l_{dn} scales like $\lambda^{-1/2}$ if Ω is replaced by $\lambda\Omega$, $\lambda > 0$.

We point out here that for the 2D Navier–Stokes system with analytic forcing the results of [17], [18] provide the existence of a finite number N of instantaneously determining nodes comparable with the fractal dimension of the attractor. These nodes, however, can be chosen arbitrarily (up to a subset of Ω^N with $2N$ -dimensional Lebesgue measure zero) and therefore do not naturally define a regular lattice of determining nodes.

The best to date estimate for the analyticity radius of the solutions of the Navier–Stokes equations with analytic forcing term f was obtained in [31]:

$$l_a \geq c_3(\gamma) \frac{|\Omega|^{1/2}}{G^{1/2}(1 + \log G)^{1/4}}. \quad (1.5)$$

Relating the radius of analyticity to the dissipative small length scale (see also [23] in this regard) we note that up to a logarithmic correction the estimate (1.5) coincides with (1.4), but both are worse than (1.2), where the latter coincides, as we have already pointed out, with the physically asserted estimate of [30].

In this paper we focus on the 2D space periodic Navier–Stokes system with damping

$$\begin{aligned} \partial_t u + \sum_{i=1}^2 u^i \partial_i u &= -\mu u + \nu \Delta u - \nabla p + f, \\ \operatorname{div} u &= 0. \end{aligned} \quad (1.6)$$

By adding the Coriolis forcing term to (1.6) one obtains the well-known Stommel–Charney barotropic model of ocean circulation [4], [10], [35], [37]. Here the damping μu represents the Rayleigh friction term and f is the wind stress. For an analytical study of this system see, for instance [5], [22], [24], [41], and the references therein. In a follow up work we will be studying the effect of adding rotation (Coriolis parameter) on the size of small scales and the complexity of the dynamics of (1.6). Therefore, we will focus in this work on the system (1.6). We also point out that in this geophysical context the viscosity plays a much

smaller role in the mechanism of dissipating energy than the Rayleigh friction. That is why in this work the friction coefficient $\mu > 0$ will be fixed and we consider the system at the limit when $\nu \rightarrow 0^+$.

Sharp estimates (as $\nu \rightarrow 0$) for the Hausdorff and the fractal dimensions of the global attractor of the system (1.6) were first obtained in the case of the square-shaped domain in [25] ($\gamma = 1$). Then the case of an elongated domain was studied in [27] ($\gamma \rightarrow 0$), where it was shown that

$$\dim_F \mathcal{A} \leq c_4 D, \quad D = \frac{\|\operatorname{rot} f\|_\infty |\Omega|}{\mu\nu}, \quad (1.7)$$

where c_4 is an absolute constant ($c_4 \leq 12$). This estimate is sharp as both $\nu \rightarrow 0$ and $\gamma \rightarrow 0$. Therefore the small length scale defined as in (1.2) is of the order of

$$l_f \sim \left(\frac{|\Omega|}{\dim_F \mathcal{A}} \right)^{1/2} \sim \left(\frac{\mu\nu}{\|\operatorname{rot} f\|_\infty} \right)^{1/2} \sim \frac{|\Omega|^{1/2}}{D^{1/2}}. \quad (1.8)$$

This heuristic estimate is, in fact, a rigorous bound for the small length scale expressed in terms of the number of determining modes and nodes [26]:

$$l_{dn} = c_5 \left(\frac{|\Omega|}{D} \right)^{1/2} = c_5 \left(\frac{\mu\nu}{\|\operatorname{rot} f\|_\infty} \right)^{1/2}, \quad c_5 = 68^{1/4}. \quad (1.9)$$

This means that any lattice of points in Ω at a typical distance $l \leq l_{dn}$ is determining.

The main result of this paper is in showing that the analyticity radius l_a of the solutions of the damped-driven Navier–Stokes system (1.6) lying on the global attractor is bounded from below and satisfies the estimate:

$$l_a \geq \frac{c|\Omega|^{1/2}}{D^{1/2}(1 + \log D)^{1/2}}, \quad (1.10)$$

which up to a logarithmic correction agrees both with the smallest scale estimate (1.8) and the rigorously defined typical distance between the determining nodes (1.9).

It is worth mentioning that this point of view of relating the radius of analyticity of solutions on the Navier–Stokes equations to small scales in turbulence was also presented in [23].

This paper is organized as follows. In section 2 we employ the Gevrey–Hilbert space technique of [15] to derive a lower bound for the radius of analyticity of the order

$$\frac{c|\Omega|^{1/2}}{D^2 \log D}. \quad (1.11)$$

This bound considerably improves, for a fixed $\mu > 0$, the lower bound (1.1) for the classical Navier–Stokes system as $\nu \rightarrow 0^+$ (see also Remark 2.1). Let us remark that as an alternative to the Gevrey regularity technique for estimating small scales one can apply the ladder estimates approach presented in [9] to obtain estimates for the small scales in (1.6) (see also [19]).

In section 3 the estimate (1.10) is proved for the system (1.6) following [31].

2. GEVREY REGULARITY OF THE DAMPED NAVIER–STOKES SYSTEM

As usual (see, for instance, [1],[6],[32],[38]), we write (1.6) as an evolution equation in the Hilbert space H which is the closed subspace of solenoidal vectors in $(L_2(\Omega))^2$ with zero average over the torus $\Omega = [0, L/\gamma] \times [0, L]$:

$$\partial_t u + B(u, u) + \nu Au + \mu u = f, \quad u(0) = u_0. \quad (2.1)$$

Here $A = -P\Delta$ is the Stokes operator with eigenvalues $0 < \lambda_1 \leq \lambda_2 \leq \dots$, $B(u, v) = P(\sum_{i=1}^2 u^i \partial_i v)$ is the nonlinear term, $f = Pf \in H$, and $P : (L_2(\Omega))^2 \rightarrow H$.

We restrict ourselves to the case $\gamma = 1$ and, in addition, assume that $\Omega = [0, 2\pi]^2$ (this simplifies the Fourier series below). The case of the square-shaped domain $\Omega = [0, L]^2$ reduces to this case by scaling. Furthermore, any domain with aspect ratio $\gamma < 1$ can be treated in the similar way, the absolute dimensionless constants c_1, c_2, \dots below will then depend on γ , however.

A vector field $u \in H$ has the Fourier series expansion

$$u = \sum_{j \in \mathbb{Z}^2} u_j e^{ij \cdot x}, \quad u_j \in \mathbb{C}^2, \quad u_{-j} = \bar{u}_j, \quad u_j \cdot j = 0, \quad u_0 = 0,$$

and

$$\|u\|^2 = \|u\|_{L_2}^2 = (2\pi)^2 \sum_{j \in \mathbb{Z}^2} |u_j|^2.$$

The eigenvalues of the Stokes operator A are the numbers $|j|^2$, and the domain of its powers is the set of vector functions u such that

$$(2\pi)^2 \sum_{j \in \mathbb{Z}^2} |j|^{4\alpha} |u_j|^2 = \|A^\alpha u\|^2 < \infty.$$

For $\tau, s > 0$ we define the Gevrey space $D(e^{\tau A^s})$ of functions u satisfying

$$(2\pi)^2 \sum_{j \in \mathbb{Z}^2} e^{2\tau |j|^{2s}} |u_j|^2 = \|e^{\tau A^s} u\|^2 < \infty. \quad (2.2)$$

We suppose that the forcing term f belongs to the Gevrey space of analytic functions

$$f \in D(e^{\sigma_1 A^{1/2}} A^{1/2}), \text{ so that } (2\pi)^2 \sum_{j \in \mathbb{Z}^2} |j|^2 e^{2\sigma_1 |j|} |u_j|^2 = \|e^{\sigma_1 A^{1/2}} A^{1/2} f\| < \infty \quad (2.3)$$

for some $\sigma_1 > 0$. We set

$$\varphi(t) = \min(\nu \lambda_1^{1/2} t, \sigma_1).$$

The norm and the scalar product in $D(e^{\varphi(t) A^{1/2}})$ are denoted by $\|\cdot\|_\varphi$ and $(\cdot, \cdot)_\varphi$, respectively.

We assume that $u_0 \in D(A^{1/2})$ and take the scalar product of (2.1) and Au in $D(e^{\varphi(t) A^{1/2}})$ for sufficiently small $t \leq \sigma_1 / (\nu \lambda_1^{1/2})$. Since

$$(e^{\varphi(t) A^{1/2}} \partial_t u(t), e^{\varphi(t) A^{1/2}} u(t)) = \frac{1}{2} \partial_t \|A^{1/2} u(t)\|_\varphi^2 - \nu \lambda_1^{1/2} (Au(t), A^{1/2} u(t))_\varphi,$$

we obtain

$$\frac{1}{2} \partial_t \|A^{1/2} u\|_\varphi^2 + \nu \|Au\|_\varphi^2 + \mu \|A^{1/2} u\|_\varphi^2 = (B(u, u) Au)_\varphi + \nu \lambda_1^{1/2} (Au, A^{1/2} u)_\varphi + (A^{1/2} f, A^{1/2} u)_\varphi. \quad (2.4)$$

Next we use the key estimate (see [15], [12], [40]) for the nonlinear term in Gevrey spaces

$$(B(u, u), Au)_\varphi \leq c_1 \|A^{1/2} u\|_\varphi^2 \|Au\|_\varphi \left(1 + \log \frac{\|Au\|_\varphi^2}{\lambda_1 \|A^{1/2} u\|_\varphi^2} \right)^{1/2}$$

and use Young's inequality for this estimate and for the last two terms in (2.4):

$$\begin{aligned} \partial_t \|A^{1/2}u\|_\varphi^2 + \nu \|Au\|_\varphi^2 &\leq \\ &\leq \frac{2c_1^2}{\nu} \|A^{1/2}u\|_\varphi^4 \left(1 + \log \frac{\|Au\|_\varphi^2}{\lambda_1 \|A^{1/2}u\|_\varphi^2}\right) + 2\nu\lambda_1 \|A^{1/2}u\|_\varphi^2 + \frac{\|A^{1/2}f\|_\varphi^2}{2\mu} \leq \\ &\leq \frac{c_2}{\nu} \|A^{1/2}u\|_\varphi^4 \left(1 + \log \frac{\|Au\|_\varphi^2}{\lambda_1 \|A^{1/2}u\|_\varphi^2}\right) + \nu^3\lambda_1^2 + \frac{\|A^{1/2}f\|_\varphi^2}{2\mu}, \end{aligned}$$

where $c_2 = 2c_1^2 + 1$. Next, using the inequality $-\alpha z + \beta(1 + \log z) \leq \beta \log \beta/\alpha$ (see [12], [13]), we find

$$-\nu \|Au\|_\varphi^2 + \frac{c_2}{\nu} \|A^{1/2}u\|_\varphi^4 \left(1 + \log \frac{\|Au\|_\varphi^2}{\lambda_1 \|A^{1/2}u\|_\varphi^2}\right) \leq \frac{c_2}{\nu} \|A^{1/2}u\|_\varphi^4 \log \frac{c_2 \|A^{1/2}u\|_\varphi^2}{\lambda_1 \nu^2},$$

and obtain the differential inequality

$$\partial_t \|A^{1/2}u\|_\varphi^2 \leq \frac{c_2}{\nu} \|A^{1/2}u\|_\varphi^4 \log \frac{c_2 \|A^{1/2}u\|_\varphi^2}{\lambda_1 \nu^2} + \nu^3\lambda_1^2 + \frac{\|A^{1/2}f\|_\varphi^2}{2\mu}.$$

Hence the function

$$y(t) = \frac{c_2 \|A^{1/2}u\|_\varphi^2}{\lambda_1 \nu^2} + \frac{\|A^{1/2}f\|_{\sigma_1}^2}{\lambda_1 \nu^{3/2} \mu^{1/2}} + e,$$

where $\ln e = 1$, satisfies

$$\partial_t y(t) \leq \nu \lambda_1 c_3 y^2 \log y, \quad c_3 = \max(1, c_2/2).$$

Therefore $y(t) \leq 2y(0)$ for as long as

$$t \leq (2\nu \lambda_1 c_3 y(0) \log 2y(0))^{-1}.$$

In other words,

$$\|A^{1/2}u\|_\varphi^2 \leq 2\|A^{1/2}u_0\|^2 + c_4(\nu/\mu)^{1/2} \|A^{1/2}f\|_{\sigma_1} + c_4\lambda_1\nu^2, \quad c_4 = e/c_2,$$

as long as $0 \leq t \leq T^*(\|A^{1/2}u_0\|)$, where

$$T^*(\|A^{1/2}u_0\|) = \frac{1}{2c_3\nu\lambda_1 \left(\frac{c_2\|A^{1/2}u_0\|^2}{\lambda_1\nu^2} + \frac{\|A^{1/2}f\|_{\sigma_1}}{\lambda_1\nu^{3/2}\mu^{1/2}} + e \right) \log \left(2 \left(\frac{c_2\|A^{1/2}u_0\|^2}{\lambda_1\nu^2} + \frac{\|A^{1/2}f\|_{\sigma_1}}{\lambda_1\nu^{3/2}\mu^{1/2}} + e \right) \right)}.$$

We now observe (see Lemma 3.1) that on the global attractor or in the absorbing ball we have, respectively,

$$\|A^{1/2}u(t)\| \leq \frac{\|A^{1/2}f\|}{\mu}, \quad t \in \mathbb{R}, \quad \|A^{1/2}u(t)\| \leq 2\frac{\|A^{1/2}f\|}{\mu}, \quad t \geq T_0(\|A^{1/2}u_0\|).$$

Therefore we have the following lower bound for T^* :

$$T^* \geq c_5 \left[\nu\lambda_1 \left(\frac{\|A^{1/2}f\|^2}{\lambda_1\nu^2\mu^2} + \frac{\|A^{1/2}f\|_{\sigma_1}}{\lambda_1\nu^{3/2}\mu^{1/2}} + 1 \right) \log \left(\frac{\|A^{1/2}f\|^2}{\lambda_1\nu^2\mu^2} + \frac{\|A^{1/2}f\|_{\sigma_1}}{\lambda_1\nu^{3/2}\mu^{1/2}} + 1 \right) \right]^{-1}$$

In the limit $\nu \rightarrow 0^+$ we have

$$\frac{\|A^{1/2}f\|^2}{\lambda_1\nu^2\mu^2} \gg \frac{\|A^{1/2}f\|_{\sigma_1}}{\lambda_1\nu^{3/2}\mu^{1/2}},$$

and we can write the lower bound for T^* as follows

$$T^* \geq c_6 [\nu\lambda_1 D^2 \log D]^{-1},$$

where

$$\frac{\|A^{1/2}f\|}{\lambda_1^{1/2}\nu\mu} = \frac{\|\operatorname{rot} f\| |\Omega|^{1/2}}{2\pi\nu\mu} \leq \frac{1}{2\pi}D, \quad \text{where} \quad D = \frac{\|\operatorname{rot} f\|_\infty |\Omega|}{\nu\mu}.$$

In terms of the analyticity radius l_a the lower bound for T^* takes the form

$$l_a \geq \frac{c_7 |\Omega|^{1/2}}{D^2 \log D}.$$

Thus, we have proved the following theorem.

Theorem 2.1. *Suppose that $f \in D(A^{1/2}e^{\sigma_1 A^{1/2}})$ for some $\sigma_1 > 0$. Then a solution u lying on the global attractor \mathcal{A} is analytic with analyticity radius*

$$l_a \geq \min \left(\frac{c_7 |\Omega|^{1/2}}{(D^2 + D_1 + 1) \log(D^2 + D_1 + 1)}, \sigma_1 \right),$$

where

$$D = \frac{\|\operatorname{rot} f\|_\infty |\Omega|}{\nu\mu}, \quad D_1 = \frac{\|A^{1/2}f\|_{\sigma_1}}{\lambda_1 \nu^{3/2} \mu^{1/2}}.$$

Moreover,

$$l_a \geq \frac{c_8 |\Omega|^{1/2}}{D^2 \log D} \quad \text{as} \quad \nu \rightarrow 0^+. \quad (2.5)$$

The constants c_7 and c_8 depend only on the aspect ratio of the periodic domain Ω .

Remark 2.1. We observe that the estimate (2.5) for the system (1.6) is of the order $\nu^{-2} \log(1/\nu)$ as far as the dependence on $\nu \rightarrow 0^+$ is concerned, while the estimate (1.1) for the classical Navier–Stokes system is, in this respect much larger; namely, is of the order $\nu^{-4} \log(1/\nu)$.

However, the estimate (2.5) is not sharp and will be improved in the next section. As has been demonstrated in [34] the Gevrey–Hilbert space technique does not always provide sharp estimates for the radius of analyticity. The mechanism explaining this has been reported in [34] by means of an explicitly solvable model equation.

3. SHARPER BOUNDS

In this section we obtain sharper lower bounds for the analyticity radius l_a . This is achieved by combining the ν -independent estimate for the vorticity contained in the following lemma and the L_p -technique developed in [21], [31] for the uniform analyticity radius of the solutions of the Navier–Stokes equations. We observe that similar technique has been earlier established in [2] for studying the analyticity of the Euler equations.

Applying the operator rot to (1.6) we obtain the well-known scalar vorticity equation

$$\partial_t \omega + u \cdot \nabla \omega = \nu \Delta \omega - \mu \omega + F, \quad (3.1)$$

where $\omega = \operatorname{rot} u$, $F = \operatorname{rot} f$, $u = \nabla^\perp \Delta^{-1} \omega$, so that $u \cdot \nabla \omega = \nabla^\perp \Delta^{-1} \omega \cdot \nabla \omega = J(\Delta^{-1} \omega, \omega)$, and $\nabla^\perp = (-\partial_2, \partial_1)$, $J(a, b) = \nabla^\perp a \cdot \nabla b$.

Lemma 3.1. (See [26].) *The solutions $u(t)$ lying on the global attractor \mathcal{A} satisfy the following bound:*

$$\|\omega(t)\|_{L_{2k}} \leq \frac{\|\operatorname{rot} f\|_{L_{2k}}}{\mu}, \quad t \in \mathbb{R}, \quad (3.2)$$

where $1 \leq k \leq \infty$.

Proof. We use the vorticity equation (3.1) and take the scalar product with ω^{2k-1} , where $k \geq 1$ is integer, and use the identity

$$(J(\psi, \varphi), \varphi^{2k-1}) = (2k)^{-1} \int J(\psi, \varphi^{2k}) dx = (2k)^{-1} \int \operatorname{div}(\varphi^{2k} \nabla^\perp \psi) dx = 0.$$

We obtain

$$\begin{aligned} \|\omega\|_{L_{2k}}^{2k-1} \partial_t \|\omega\|_{L_{2k}} + (2k-1)\nu \int |\nabla \omega|^2 \omega^{2k-2} dx + \mu \|\omega\|_{L_{2k}}^{2k} &= \\ &= (\operatorname{rot} f, \omega^{2k-1}) \leq \|\operatorname{rot} f\|_{L_{2k}} \|\omega\|_{L_{2k}}^{2k-1}. \end{aligned}$$

Hence, by Gronwall's inequality

$$\|\omega(t)\|_{L_{2k}} \leq \|\omega(0)\|_{L_{2k}} e^{-\mu t} + \mu^{-1} \|\operatorname{rot} f\|_{L_{2k}} (1 - e^{-\mu t}),$$

and passing to the limit as $k \rightarrow \infty$ we find

$$\|\omega(t)\|_\infty \leq \|\omega(0)\|_\infty e^{-\mu t} + \mu^{-1} \|\operatorname{rot} f\|_\infty (1 - e^{-\mu t}).$$

Now, we let $t \rightarrow \infty$ in the above inequalities and obtain

$$\limsup_{t \rightarrow \infty} \|\omega(t)\|_{L_{2k}} \leq \frac{\|\operatorname{rot} f\|_{L_{2k}}}{\mu}, \quad 1 \leq k \leq \infty,$$

which gives (3.2) since the solutions lying on the attractor are bounded for $t \in \mathbb{R}$. \square

As before we consider the square-shaped domain $\Omega = [0, L]^2$ and it is now convenient to write (1.6) in dimensionless form. We introduce dimensionless variables x' , t' , u' and p' by setting

$$x = Lx', \quad t = (L^2/\nu)t', \quad u = (\nu/L)u', \quad p = (\nu^2/L^2)p', \quad \mu = (\nu/L^2)\mu'.$$

We obtain

$$\begin{aligned} \partial_{t'} u' + \sum_{i=1}^2 u'^i \partial'_i u' &= -\mu' u' + \Delta' u' - \nabla' p' + f', \\ \operatorname{div}' u' &= 0, \end{aligned} \tag{3.3}$$

where $x' \in \Omega' = [0, 1]^2$, $f' = (L^3/\nu^2)f$. Accordingly, the dimensionless form of (3.1) is as follows (we omit the primes):

$$\partial_t \omega + u \cdot \nabla \omega = \Delta \omega - \mu \omega + F. \tag{3.4}$$

Remark 3.1. For dimensionless variables u' and ω' the estimate (3.2) with $k = \infty$ takes the form

$$\|\omega'\|_\infty = \|\operatorname{rot}' u'\|_\infty \leq \frac{\|\operatorname{rot}' f'\|_\infty}{\mu'} = \frac{\|\operatorname{rot} f\|_\infty L^2}{\nu \mu} = D. \tag{3.5}$$

The next lemma is similar to the main estimate for the space analyticity radius in [31].

Lemma 3.2. *Suppose that F is a restriction to Ω (that is, $y = 0$) of a bounded x -periodic analytic function $F(x + iy) + iG(x + iy)$ in the region $|y| \leq \delta_F$ and*

$$M_F^2 = \sup_{x \in \Omega, |y| \leq \delta_F} (F(x + iy)^2 + G(x + iy)^2). \tag{3.6}$$

Let $p \geq 3/2$ and let

$$t_0 = \frac{M_{2p}^2}{CM_F^2/\mu}.$$

Here (and throughout) C is a sufficiently large universal constant and $M_{2p} \geq \|\omega_0\|_{L_{2p}}$. Then the solution $\omega(t)$ is analytic for $t > 0$ and for $0 < t \leq t_0$ the space analyticity radius of $\omega(t)$ is greater than

$$\delta(t) = \min \left(\frac{t^{1/2}}{C}, \frac{1}{Cpt^{(2p-3)/4p}M_{2p}}, \frac{1}{Cpt^{(2p-3)/(4p+6)}M_{2p}^{2p/(2p+3)}}, \frac{1}{pt^{1/2}M_{2p}}, \delta_F \right).$$

Proof. We solve (3.4) by a sequence of approximating solutions (see [29], [31]). We set $u^{(0)} = 0$ and $\omega^{(0)} = 0$. Then for $\omega^{(n)}$, $u^{(n)}$ we have the equation

$$\begin{aligned} \partial_t \omega^{(n)} - \Delta \omega^{(n)} + u^{(n-1)} \cdot \nabla \omega^{(n)} + \mu \omega^{(n)} &= F \\ \omega^{(n)}(0) = \omega_0 = \text{rot } u_0, \quad u^{(n)} &= \nabla^\perp \Delta^{-1} \omega^{(n)}. \end{aligned} \quad (3.7)$$

The solutions $\omega^{(n)}$ and $u^{(n)}$ for $t > 0$ have analytic extensions $\omega^{(n)} + i\theta^{(n)}$ and $u^{(n)} + iv^{(n)}$ and since the system (3.7) is linear, their analyticity radius is at least δ_F . They satisfy the equation

$$\partial_t(\omega^{(n)} + i\theta^{(n)}) - \Delta(\omega^{(n)} + i\theta^{(n)}) + (u^{(n-1)} + iv^{(n-1)}) \cdot \nabla(\omega^{(n)} + i\theta^{(n)}) + \mu(\omega^{(n)} + i\theta^{(n)}) = F + iG,$$

or, equivalently, the system

$$\begin{aligned} \partial_t \omega^{(n)} - \Delta \omega^{(n)} + \mu \omega^{(n)} + u^{(n-1)} \cdot \nabla \omega^{(n)} - v^{(n-1)} \cdot \nabla \theta^{(n)} &= F, \\ \partial_t \theta^{(n)} - \Delta \theta^{(n)} + \mu \theta^{(n)} + u^{(n-1)} \cdot \nabla \theta^{(n)} + v^{(n-1)} \cdot \nabla \omega^{(n)} &= G, \end{aligned} \quad (3.8)$$

where, as before, $u^{(n)} = \nabla^\perp \Delta^{-1} \omega^{(n)}$, $v^{(n)} = \nabla^\perp \Delta^{-1} \theta^{(n)}$, and the differential operators are taken with respect to x . In view of the analyticity of the solutions we have the Cauchy–Riemann equations

$$\begin{aligned} \frac{\partial \omega^{(n)}}{\partial y_j} &= -\frac{\partial \theta^{(n)}}{\partial x_j}, \\ \frac{\partial \omega^{(n)}}{\partial x_j} &= \frac{\partial \theta^{(n)}}{\partial y_j}, \quad j = 1, 2, \end{aligned} \quad (3.9)$$

and the similar equations for $u^{(n)}$ and $v^{(n)}$.

Let $\varepsilon > 0$. We consider the functional

$$\psi_n(t) = \int_0^1 \int_\Omega (\omega^{(n)}(x, \alpha t s, t)^2 + \theta^{(n)}(x, \alpha t s, t)^2 + \varepsilon)^p dx ds. \quad (3.10)$$

We also set

$$Q_n(x, s, t) = \omega^{(n)}(x, \alpha t s, t)^2 + \theta^{(n)}(x, \alpha t s, t)^2 + \varepsilon.$$

Here $t \in \mathbb{R}^+$ and $\alpha \in \mathbb{R}^2$. The combination $\alpha t s$ will play the role of the variable y ; $p \geq 3/2$, and $\varepsilon > 0$ is arbitrary.

We differentiate $\psi_n(t)$ taking into account (3.8) and use the Cauchy–Riemann equations (3.9) to handle the derivatives with respect to y . We obtain

$$\frac{1}{2p} \partial_t \psi_n(t) + I_0 = I_1 + I_2 + I_3 + I_4, \quad (3.11)$$

where

$$\begin{aligned} I_0 &= \int_0^1 \int_\Omega Q_n^{p-1} (|\nabla \omega^{(n)}|^2 + |\nabla \theta^{(n)}|^2 + \mu(\omega^{(n)})^2 + \mu(\theta^{(n)})^2) dx ds + \\ &\quad + 2(p-1) \int_0^1 \int_\Omega Q_n^{p-2} (\omega^{(n)} \nabla \omega^{(n)} + \theta^{(n)} \nabla \theta^{(n)})^2 dx ds, \end{aligned}$$

and

$$\begin{aligned} I_1 &= \int_0^1 \int_{\Omega} Q_n^{p-1} (-\omega^{(n)} \nabla \theta^{(n)} + \theta^{(n)} \nabla \omega^{(n)}) \cdot \alpha s \, dx ds, \\ I_2 &= \int_0^1 \int_{\Omega} Q_n^{p-1} (\omega^{(n)} \nabla \omega^{(n)} + \theta^{(n)} \nabla \theta^{(n)}) \cdot u^{(n-1)} \, dx ds, \\ I_3 &= \int_0^1 \int_{\Omega} Q_n^{p-1} (-\omega^{(n)} \nabla \theta^{(n)} + \theta^{(n)} \nabla \omega^{(n)}) \cdot v^{(n-1)} \, dx ds, \\ I_4 &= \int_0^1 \int_{\Omega} Q_n^{p-1} (\omega^{(n)} F + \theta^{(n)} G) \, dx ds. \end{aligned}$$

The arguments of Q_n are x, s, t , and the arguments of $\omega^{(n)}, \theta^{(n)}, u^{(n)}$, and $v^{(n)}$ are $x, \alpha t s$, and t .

For an arbitrary $\eta > 0$ we have

$$\begin{aligned} I_1 &\leq \eta \int_0^1 \int_{\Omega} Q_n^{p-1} (|\nabla \omega^{(n)}|^2 + |\nabla \theta^{(n)}|^2) \, dx ds + \\ &C_{\eta} \int_0^1 \int_{\Omega} Q_n^{p-1} ((\omega^{(n)})^2 + (\theta^{(n)})^2) |\alpha|^2 s^2 \, dx ds \leq \eta I_0 + C_{\eta} |\alpha|^2 \psi_n(t). \end{aligned} \quad (3.12)$$

Next,

$$I_2 = \frac{1}{2p} \int_0^1 \int_{\Omega} \nabla Q_n^p \cdot u^{(n-1)} \, dx ds = 0. \quad (3.13)$$

For I_3 we have

$$I_3 \leq \eta I_0 + C_{\eta} \int_0^1 \int_{\Omega} Q_n^p |v^{(n-1)}|^2 \, dx ds \leq \eta I_0 + C_{\eta} I_3' I_3'', \quad (3.14)$$

where

$$I_3' = \left(\int_0^1 \int_{\Omega} Q_n(x, s, t)^{p^2/(p-1)} \, dx ds \right)^{(p-1)/p}, \quad I_3'' = \sum_{j=1}^2 \left(\int_0^1 \int_{\Omega} |v_j^{(n-1)}(x, \alpha t s, t)|^{2p} \, dx ds \right)^{1/p}.$$

We write I_3' as follows

$$I_3' = \|Q_n^{p/2}\|_{L_{\beta}(\Omega_0)}^2, \quad \Omega_0 = \Omega \times [0, 1] \subset \mathbb{R}^3, \quad \beta = 2p/(p-1), \quad 2 \leq \beta \leq 6,$$

and use in Ω_0 the Gagliardo–Nirenberg inequality

$$\|A\|_{L_{\beta}(\Omega_0)} \leq C \|A\|_{L_2(\Omega_0)}^{3/\beta-1/2} \|\nabla_{x,s} A\|_{L_2(\Omega_0)}^{3/2-3/\beta} + C \|A\|_{L_2(\Omega_0)}$$

for $A = A(x, s) = Q_n^{p/2}(x, s, t)$. We have

$$\begin{aligned} \|\nabla_{x,s} A\|_{L_2(\Omega_0)}^2 &= \|\nabla_{x,s} Q_n^{p/2}\|_{L_2(\Omega_0)}^2 = \\ p^2 \int_0^1 \int_{\Omega} Q_n^{p-2} ((\omega^{(n)} \nabla \omega^{(n)} + \theta^{(n)} \nabla \theta^{(n)})^2 + t^2 (\theta^{(n)} \alpha \cdot \nabla \omega^{(n)} - \omega^{(n)} \alpha \cdot \nabla \theta^{(n)})^2) \, dx ds &\leq \\ &\leq Cp^2(1 + |\alpha|^2 t^2) I_0. \end{aligned}$$

Hence,

$$\|\nabla_{x,s} A\|_{L_2(\Omega_0)}^{3/2-3/\beta} = \|\nabla_{x,s} Q_n^{p/2}\|_{L_2(\Omega_0)}^{3/2p} \leq C(1 + |\alpha|^2 t^2)^{3/4p} I_0^{3/4p}.$$

Next, $\|Q_n^{p/2}\|_{L_2(\Omega_0)}^2 = \psi_n(t)$,

$$\|A\|_{L_2(\Omega_0)}^{3/\beta-1/2} = \|Q_n^{p/2}\|_{L_2(\Omega_0)}^{(2p-3)/2p} = \psi_n(t)^{(2p-3)/4p}$$

and

$$I'_3 \leq C(1 + |\alpha|^2 t^2)^{3/2p} I_0^{3/2p} \psi_n(t)^{(2p-3)/2p} + C\psi_n(t). \quad (3.15)$$

We now consider I''_3 . Since $v_j^{(n-1)}(x, 0, t) = 0$ (the solution restricted to $y = 0$ is real-valued), we have (using the Cauchy–Riemann equations for v_j)

$$|v_j^{(n-1)}(x, \alpha t s, t)| = \left| \sum_{k=1}^2 \alpha_k t s \int_0^1 \partial_{y_k} v_j^{(n-1)}(x, \alpha t s \tau, t) d\tau \right| = \left| \sum_{k=1}^2 \alpha_k t s \int_0^1 \partial_k u_j^{(n-1)}(x, \alpha t s \tau, t) d\tau \right|.$$

Then

$$\begin{aligned} I''_3 &= \sum_{j=1}^2 \left(\int_0^1 \int_{\Omega} \left| \sum_{k=1}^2 \alpha_k t s \int_0^1 \partial_k u_j^{(n-1)}(x, \alpha t s \tau, t) d\tau \right|^{2p} dx ds \right)^{1/p} \leq \\ &C |\alpha|^2 t^2 \left(\int_0^1 \int_{\Omega} \int_0^1 |\nabla u^{(n-1)}(x, \alpha t s \tau, t)|^{2p} s^{2p} d\tau dx ds \right)^{1/p} = \\ &C |\alpha|^2 t^2 \left(\int_0^1 s^{2p} ds \int_0^1 d\tau \int_{\Omega} |\nabla u^{(n-1)}(x, \alpha t s \tau, t)|^{2p} dx \right)^{1/p}. \end{aligned}$$

Since $u = \nabla^\perp \Delta^{-1} \omega$, we have (see [20], [42])

$$\left(\int_{\Omega} |\nabla u(x)|^{2p} dx \right)^{1/2p} = \|\nabla \nabla^\perp \Delta^{-1} \omega\|_{L_{2p}} \leq \|\Delta^{-1} \omega\|_{W_{2p}^2} \leq Cp \|\omega\|_{L_{2p}}.$$

Therefore

$$\begin{aligned} I''_3 &\leq Cp^2 |\alpha|^2 t^2 \left(\int_0^1 \int_0^1 \int_{\Omega} |\omega^{(n-1)}(x, \alpha t s \tau, t)|^{2p} dx s^{2p} ds d\tau \right)^{1/p} \leq \\ &Cp^2 |\alpha|^2 t^2 \left(\int_0^1 \int_{\Omega} Q_{n-1}^p dx ds \right)^{1/p} \leq Cp^2 |\alpha|^2 t^2 \psi_{n-1}(t)^{1/p}, \end{aligned}$$

where we have used $\int_0^1 \int_0^1 h(s\tau) s^{2p} ds d\tau \leq (2p)^{-1} \int_0^1 h(s) ds$. Combining this with (3.14) and (3.15) we obtain

$$\begin{aligned} I_3 &\leq \\ &\eta' I_0 + C_\eta p^2 |\alpha|^2 t^2 (1 + |\alpha|^2 t^2)^{3/2p} I_0^{3/2p} \psi_n(t)^{(2p-3)/2p} \psi_{n-1}(t)^{1/p} + Cp^2 |\alpha|^2 t^2 \psi_{n-1}(t)^{1/p} \psi_n(t) \leq \\ &\eta I_0 + C_\eta p^2 (|\alpha| t)^{4p/(2p-3)} (1 + |\alpha|^2 t^2)^{3/(2p-3)} \psi_{n-1}(t)^{2/(2p-3)} \psi_n(t) + Cp^2 |\alpha|^2 t^2 \psi_{n-1}(t)^{1/p} \psi_n(t). \end{aligned} \quad (3.16)$$

Finally, we estimate I_4 :

$$\begin{aligned} I_4 &\leq \int_0^1 \int_{\Omega} Q_n^{p-1} ((\omega^{(n)})^2 + \theta^{(n)})^2 \eta \mu + (F^2 + G^2)/(4\eta \mu) dx ds \leq \\ &\eta I_0 + C_\eta (M_F^2/\mu) \int_0^1 \int_{\Omega} Q_n^{p-1} dx ds \leq \eta I_0 + C_\eta (M_F^2/\mu) \psi_n(t)^{(p-1)/p}, \end{aligned} \quad (3.17)$$

where M_F is defined in (3.6).

Taking $\eta > 0$ sufficiently small we infer from (3.11), (3.12), (3.13), (3.16), (3.17)

$$\begin{aligned} \partial_t \psi_n(t) &\leq Cp|\alpha|^2 \psi_n(t) + Cp^3 |\alpha|^{4p/(2p-3)} t^{4p/(2p-3)} \psi_{n-1}(t)^{2/(2p-3)} \psi_n(t) + \\ &\quad Cp^3 |\alpha|^{(4p+6)/(2p-3)} t^{(4p+6)/(2p-3)} \psi_{n-1}(t)^{2/(2p-3)} \psi_n(t) + \\ &\quad Cp^3 |\alpha|^2 t^2 \psi_{n-1}(t)^{1/p} \psi_n(t) + Cp \psi_n(t)^{(p-1)/p} M_F^2 / \mu, \end{aligned}$$

where

$$\psi_n(0) = \int_{\Omega} (\omega_0(x)^2 + \varepsilon)^p dx.$$

We set $\varphi_n(t) = \psi_n(t)^{1/2p}$ and obtain the differential inequality for φ_n :

$$\begin{aligned} \partial_t \varphi_n(t) &\leq C|\alpha|^2 \varphi_n(t) + Cp^2 |\alpha|^{4p/(2p-3)} t^{4p/(2p-3)} \varphi_{n-1}(t)^{4p/(2p-3)} \varphi_n(t) + \\ &\quad Cp^2 |\alpha|^{(4p+6)/(2p-3)} t^{(4p+6)/(2p-3)} \varphi_{n-1}(t)^{4p/(2p-3)} \varphi_n(t) + \\ &\quad Cp^2 |\alpha|^2 t^2 \varphi_{n-1}(t)^2 \varphi_n(t) + C \varphi_n(t)^{-1} M_F^2 / \mu. \end{aligned}$$

We now use the Gronwall-type Lemma 3.3 from [31] below and see that $\varphi(t) \leq 2\varphi(0)$ on the time interval specified in (3.18), (3.19), and letting $\varepsilon \rightarrow 0$ we obtain

$$\int_0^1 \int_{\Omega} (\omega^{(n)}(x, \alpha t s, t)^2 + \theta^{(n)}(x, \alpha t s, t)^2)^p dx ds \leq 2^{2p} M_{2p}^{2p},$$

for $t \geq 0$, $|\alpha|t \leq \delta_F$ and

$$t \leq \min(A_1, A_2, A_3, A_4, A_5), \quad (3.18)$$

where

$$\begin{aligned} A_1 &= \frac{1}{C|\alpha|^2}, \\ A_2 &= \frac{1}{Cp^{(4p-6)/(6p-3)} |\alpha|^{4p/(6p-3)} M_{2p}^{4p/(6p-3)}}, \\ A_3 &= \frac{1}{Cp^{(4p-6)/(6p+3)} |\alpha|^{(4p+6)/(6p+3)} M_{2p}^{4p/(6p+3)}}, \\ A_4 &= \frac{1}{Cp^{2/3} |\alpha|^{2/3} M_{2p}^{2/3}}, \\ A_5 &= \frac{M_{2p}^2}{CM_F^2 \mu^{-1}}. \end{aligned} \quad (3.19)$$

We now set

$$t_0 = \frac{M_{2p}^2}{CM_F^2 / \mu}. \quad (3.20)$$

Then the condition

$$t \leq \min(A_1, A_2, A_3, A_4)$$

can be written in terms of $y = \alpha t$ as follows

$$|y| \leq \min \left(\frac{t^{1/2}}{C}, \frac{1}{Cpt^{(2p-3)/4p} M_{2p}}, \frac{1}{Cpt^{(2p-3)/(4p+6)} M_{2p}^{2p/(2p+3)}}, \frac{1}{pt^{1/2} M_{2p}} \right). \quad (3.21)$$

Now for t_0 defined in (3.20) and

$$\delta(t) = \min \left(\frac{t^{1/2}}{C}, \frac{1}{Cpt^{(2p-3)/4p} M_{2p}}, \frac{1}{Cpt^{(2p-3)/(4p+6)} M_{2p}^{2p/(2p+3)}}, \frac{1}{pt^{1/2} M_{2p}}, \delta_F \right) \quad (3.22)$$

we have for $0 < t \leq t_0$ and $|y| \leq \delta(t)$

$$\int_0^1 \int_{\Omega} (\omega^{(n)}(x, sy, t)^2 + \theta^{(n)}(x, sy, t)^2) dx ds \leq 2^{2p} M_{2p}^{2p}$$

for all integer $n \geq 1$. Therefore for any $y \in \mathbb{R}^2$ with $|y| = 1$ this gives that

$$\int_0^{\delta(t)} \int_{\Omega} (\omega^{(n)}(x, sy, t)^2 + \theta^{(n)}(x, sy, t)^2) dx ds \leq 2^{2p} \delta(t) M_{2p}^{2p}$$

and since $\int_0^{\delta} f(sy) ds \leq B$, $|y| = 1$ implies $\int_{|y| \leq \delta} f(y) dy \leq 2\pi\delta B$, we obtain

$$\int_{|y| \leq \delta(t)} \int_{\Omega} (\omega^{(n)}(x, y, t)^2 + \theta^{(n)}(x, y, t)^2) dx ds \leq 2\pi 2^{2p} \delta(t)^2 M_{2p}^{2p}.$$

This estimate is uniform in n and as in [21], [31] we obtain the existence of an analytic solution of (3.4) with analyticity radius satisfying (3.22). The proof is complete. \square

Lemma 3.3. (See [31].) *Let $y_n(t) \in C^1[0, T]$ be a sequence of non-negative functions satisfying $y_0(t) \leq M$ for $0 \leq t \leq T$, and $y_n(0) \leq M$ for $n \geq 1$. Suppose that on the interval $0 \leq t \leq T$*

$$\partial_t y_n(t) \leq \sum_{j=1}^N K_j t^{\alpha_j} y_n(t)^{\beta_j} y_{n-1}(t)^{\gamma_j},$$

where $K_j > 0$, $\alpha_j > -1$, $\beta_j \in \mathbb{R}$, and $\gamma_j \geq 0$ are given constants. Then $y_n(t) \leq 2M$ for all $n = 0, 1, 2, \dots$ provided that

$$0 \leq t \leq \min \left(T, \min_{j=1, \dots, N} \left(\frac{\alpha_j + 1}{N K_j 2^{\beta_j^+ + \gamma_j} M^{\beta_j + \gamma_j - 1}} \right)^{1/(\alpha_j + 1)} \right),$$

where $\beta^+ = \max(\beta, 0)$.

We can now state the main result of this section.

Theorem 3.1. *The solutions on the 2D space-periodic damped-driven Navier–Stokes system (1.6) lying on the global attractor \mathcal{A} are analytic with space analyticity radius l_a satisfying the lower bound*

$$l_a \geq \frac{|\Omega|^{1/2}}{CD^{1/2}(1 + \log D)^{1/2}}, \quad \text{where} \quad D = \frac{\|\text{rot } f\|_{\infty} |\Omega|}{\mu\nu}. \quad (3.23)$$

Proof. We first observe that (3.23) is equivalent to the estimate

$$l_a \geq \frac{1}{CD^{1/2}(1 + \log D)^{1/2}} \quad (3.24)$$

for the equation written in dimensionless form.

Next, by Young's inequality

$$\begin{aligned} pt^{(2p-3)/4p} M &\leq CpM^{4p/(4p-3)} t^{1/2} + t^{-1/2}, \\ pt^{(2p-3)/(4p+6)} M^{2p/(2p+3)} &\leq CpMt^{1/2} + t^{-1/2}. \end{aligned}$$

Hence, the estimate (3.22) can be written as follows

$$\delta(t) \geq \min \left(\frac{t^{1/2}}{C}, \frac{1}{Cpt^{1/2}(M_{2p}^{4p/(4p-3)} + M_{2p})}, \delta_F \right). \quad (3.25)$$

The solutions lying on the attractor are bounded in L_{2p} :

$$\|\omega(t)\|_{L_{2p}} \leq M_{2p}, \quad M_{2p} \leq CM_\infty.$$

Setting

$$p = C(1 + \log M_\infty)$$

we see that

$$p(M_{2p}^{4p/(4p-3)} + M_{2p}) \Big|_{p=C(1+\log M_\infty)} \leq C(1 + \log M_\infty)M_\infty$$

and therefore

$$\delta(t) \geq \min \left(\frac{t^{1/2}}{C}, \frac{1}{C(1 + \log M_\infty)M_\infty t^{1/2}}, \delta_F \right).$$

At the moment of time

$$t^* = \frac{1}{C(1 + \log M_\infty)M_\infty},$$

which for sufficiently large M_∞ (the case of our interest) is smaller than t_0 defined in (3.20) (the details are given below) we have

$$\delta(t^*) \geq \frac{1}{CM_\infty^{1/2}(1 + \log M_\infty)^{1/2}}.$$

Since $M_\infty \leq D$ (see (3.5)), it follows that

$$\delta(t^*) \geq \frac{1}{CD^{1/2}(1 + \log D)^{1/2}}.$$

By the invariance property of the attractor we see that on the attractor the above estimate holds for all t^* , which proves (3.24).

To complete the proof it remains to show that

$$\frac{1}{C(1 + \log D)D} = t^* \leq t_0 = \frac{D^2}{CM_{F'}^2/\mu'}, \quad (3.26)$$

where in the expression for t_0 we reverted to the prime notation for the dimensionless damping coefficient μ' and the forcing F' . We relate the forcing term and its analytic extension by the equality

$$M_{F'} = K\|\text{rot}' f'\|_\infty, \quad K = K(F, \delta_F).$$

Recalling that $f' = (L^3/\nu^2)f$, $\mu' = (L^2/\nu)\mu$, and $x' = (1/L)x$ we see that

$$t_0 = \frac{\nu}{CK^2\mu L^2}.$$

Hence, (3.26) goes over to the condition

$$C(1 + \log D) \geq \frac{K^2\mu^2}{\|\text{rot} f\|_\infty},$$

which is obviously satisfied for all sufficiently small $\nu > 0$. The proof is complete. \square

4. CONCLUDING REMARKS

We have shown that the solutions lying on the attractor of the 2D space-periodic damped-driven Navier–Stokes system, the Stommel–Charney barotropic model of ocean circulation without rotation, with analytic forcing have space analyticity radius which up to a logarithmic term coincides with the small scale estimates both in terms of the sharp bounds for the fractal dimension of the global attractor, and in terms of the spatial lattice of determining nodes. The derivation of this lower bound for the analyticity radius essentially uses the techniques developed in [31].

ACKNOWLEDGMENTS

A.A.I. would like to thank the warm hospitality of the Mathematics Department at the University of California, Irvine, where this work was done.

This work was supported in part by the US Civilian Research and Development Foundation, grant no. RUM1-2654-MO-05 (A.A.I. and E.S.T.), by the Russian Foundation for Fundamental Research, grants nos. 06-01-00096 and 05-01-00429, and by the RAS Programme no.1 ‘Modern problems of theoretical mathematics’ (A.A.I.). The work of E.S.T. was supported in part by the National Science Foundation, grant no. DMS-0504619, the ISF grant no. 120/6, and the BSF grant no. 2004271.

REFERENCES

1. Babin A.V., Vishik M.I. *Attractors of Evolution Equations*, Nauka, Moscow (1988) English transl. North-Holland, Amsterdam (1992).
2. Bardos C., Benachour S. Domaine d’analyticit  des solutions de l’equation d’Euler dans un ouvert de \mathbb{R}^n . *Ann. Scuola Norm. Sup. Pisa* **4**, 647–687 (1977).
3. Chae D. On the ensemble average in the study of approximate inertial manifolds, II. *J. Math. Anal. Appl.* **164**, 337–349 (1992).
4. Charney J.G. The Gulf stream as an inertial boundary layer. *Proc. Nat. Acad. Sci. USA* **41**, 731–740 (1955).
5. Barcilon V., Constantin P., Titi E.S. Existence of solutions to the Charney model of the Gulf Stream. *SIAM J. Math. Anal.* **19**, 1355–1364 (1988).
6. Constantin P., Foias C. *Navier–Stokes Equations*, Univ. of Chicago Press, Chicago (1988).
7. Constantin P., Foias C., Temam R. *Attractors Representing Turbulent Flows*. Mem. Amer. Math. Soc. **53**, Providence, RI (1985).
8. Constantin P., Foias C., Temam R. On the dimension of the attractors in two–dimensional turbulence. *Physica D* **30**, 284–296 (1988).
9. Doering C.R., Gibbon J.D. *Applied Analysis of the Navier–Stokes Equations*, Cambridge, Cambridge University Press (1997).
10. Dymnikov V.P., Filatov A.N. *Mathematics of Climate Modelling*, Boston, Birkh user (1997).
11. Doering C.R., Titi E.S. Exponential decay rate of the power spectrum for solutions of the Navier–Stokes equations. *Phys. Fluids* **7**:6, 1384–1390 (1995).
12. Foias C., Manley O., Rosa R., and Temam R. *Navier–Stokes Equations and Turbulence*, Cambridge, UK, Cambridge Univ. Press (2001).
13. Foias C., Manley O., Temam R. Modelling of the interaction of small and large eddies in two dimensional turbulent flows. *Math. Mod. Num. Anal.* **22** (1988), 93–118.
14. Foias C. and Prodi G. Sur le comportement global des solutions non stationnaires des  quations de Navier–Stokes en dimension two. *Rend. Sem. Mat. Univ. Padova* **39**, 1–34 (1967).
15. Foias C., Temam R. Gevrey class regularity for the solutions of the Navier–Stokes equations. *J. Func. Anal.* **87** (1989), 359–369.
16. Foias C., Temam R. Determination of the solutions of the Navier–Stokes equations by a set of nodal values. *Mathematics of Computation* **43**, 117–133 (1984).
17. Friz P.K., Kukavica I., Robinson J.C. Nodal parameterizing of analytic attractors. *Disc. Contin. Dynam. Syst.* **7**, 643–657 (2001).

18. Friz P.K., Robinson J.C. Parameterizing the attractor of the two-dimensional Navier–Stokes system with a finite number of nodal values. *Physica D* **148**, 201–220 (2001).
19. Gibbon J.D., Titi E.S. Attractor dimension and small length scale estimates for the three-dimensional Navier–Stokes equations. *Nonlinearity* **10**, 109–119 (1997).
20. Gilbarg D., Trudinger N.S. *Elliptic Partial Differential Equations of Second Order. 2nd edition.* Springer-Verlag, Berlin, (1983).
21. Grujić Z., Kukavica I. Space analyticity for the Navier–Stokes and related equations with initial data in L^p . *J. Func. Anal.* **152**, 447–466 (1998).
22. Hauk S. *The Long-term Behavior of the Stommel–Charney Model of the Gulf Stream, an Analytical and Computational Study*, Ph.D. Thesis, Department of Mathematics, University of California, Irvine (1997).
23. Henshaw W.D., Kreiss H.-O., Reyna L.G. Smalles scale estimates for the Navier–Stokes equations for incompressible fluids. *Arch. Rat. Mech. Anal.* **112**, 21–44 (1990).
24. Ilyin A.A. The Euler equations with dissipation. *Mat. Sbornik* **182**, no. 12, 1729–1739 (1991) English transl. in *Math. USSR, Sbornik* **74**, no. 2, 475–486 (1993).
25. Ilyin A.A., Miranville A., Titi E.S. A small viscosity sharp estimate for the global attractor of the 2-D damped-driven Navier–Stokes equations. *Commun. Math. Sciences* **2**:3, 403–426 (2004).
26. Ilyin A.A., Titi E.S. Sharp estimates for the number of the degrees of freedom for the Damped-driven 2-D Navier–Stokes equations. *J. Nonlinear Sci.* **16**, 233–253 (2006).
27. Ilyin A.A., Titi E.S. The damped-driven 2D Navier–Stokes system on large elongated domains. *Journal of Mathematical Fluid Mechanics* (accepted).
28. Jones D., Titi E.S. Upper bounds on the number of determining modes, nodes, and volume elements for the Navier–Stokes equations. *Indiana Univ. Math. J.* **42**, 875–887 (1993).
29. Kato T. Strong L^p -solutions of the Navier–Stokes equation in \mathbb{R}^m , with applications to weak solutions. *Math. Z.* **187**, , 471-480 (1984).
30. Kraichnan R.H. Inertial ranges in two-dimensional turbulence. *Phys. Fluids* **10**, 1417–1423 (1967).
31. Kukavica I. On the dissipative scale for the Navier–Stokes equations. *Indiana Univ. Math. J.* **48**, 1057-1081 (1999) .
32. Ladyzhenskaya O.A. *The Mathematical Theory of Viscous Incompressible Flow*, Nauka, Moscow (1970); English transl. Gordon and Breach, New York (1969).
33. Liu V.X. A sharp lower bound for the Hausdorff dimension of the global attractors of the 2D Navier–Stokes equations. *Comm. Math. Phys.* **158** (1993), 327–339.
34. Oliver M., Titi E.S. On the domain of analyticity for solutions of second order analytic nonlinear differential equations. *J. Differential. Equations.* **174** (2001), 55–74.
35. Pedlosky J. *Geophysical Fluid Dynamics*, New York, Springer (1979).
36. Robinson J.C. *Infinite Dimensional Dynamical Systems*, Cambridge, Cambridge University Press, 2001.
37. Stommel H. The westward intensification of wind-driven ocean currents. *Trans. Amer. Geophys. Union* **29**, 202–206 (1948).
38. Temam R. *Navier–Stokes Equations. Theory and Numerical Analysis*, Amsterdam, North-Holland (1984).
39. Temam R. *Infinite Dimensional Dynamical Systems in Mechanics and Physics*, 2nd ed., New York, Springer-Verlag (1997).
40. Titi E.S. On a criterion for locating stable stationary solutions to the Navier–Stokes equations. *Nonlin. Anal.* **11**:9, 1085–1102 (1987).
41. Wolansky G. Existence, uniqueness, and stability of stationary barotropic flow with forcing and dissipation. *Comm. Pure Appl. Math* **41**, 19–46 (1988).
42. Yudovich V.I. On a bound for the solution of an elliptic equation. (Russian). *Uspehi Mat. Nauk* **20**:2, 213–219 (1965).