

Lawrence Berkeley National Laboratory

Recent Work

Title

IS THE MESON THE O- MESON?

Permalink

<https://escholarship.org/uc/item/0z62490f>

Authors

Rosenfeld, Arthur H.
Carmony, D. Duane.
Walle, Remy T. Van de

Publication Date

1962-02-20

University of California
Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-10055
Limited Distribution

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

IS THE η MESON THE 0^{-+} χ MESON?

Arthur H. Rosenfeld, D. Duane Carmony, and Remy T. Van de Walle

February 20, 1962

IS THE η MESON THE 0^{-+} χ MESON?*

Arthur H. Rosenfeld, D. Duane Carmony, and Remy T. Van de Walle†

Lawrence Radiation Laboratory
University of California
Berkeley, California

February 20, 1962

Pevsner et al.¹ discovered a meson of mass 550 Mev which decays into $\pi^+ \pi^- \pi^0$ and has isospin = 0.² They called it η , the name given by Sakurai to his proposed second vector meson (i. e., a meson whose spin, parity, and G-parity we write as 1^{--}).³ However, any $I=0$ configuration of three pions must be antisymmetric in all pairs of pions. The original data were insufficient to make this test, but Bastien et al. later reported additional η events whose Dalitz plot tended not to satisfy this $I=0$ symmetry.⁴ Their alternative postulate was that η had positive G-parity and that the G-forbidden decay $\eta \rightarrow 3\pi$ occurred only because virtual photons carry off one unit of isospin. They concluded that Dalitz plots did seem to rule out all assignments except 1^{--} and 0^{-+} , and that, of these two, 0^{-+} fitted their data slightly better. At about the same time, Feinberg (unaware of the 0^{-+} evidence) pointed out that the (1^{--}) η should be present in a so far unreported mode of ρ decay, $\rho \rightarrow \eta + \pi$.⁵ We find experimentally that this mode has a branching fraction $f_{\eta\pi}$ less than 0.6% and point out that this may be evidence that η has even G-parity, i. e., is 0^{-+} .

It is well-established that ρ is a 1^{-+} meson with mass 750 Mev, width $\Gamma \approx 100$ Mev, and a two-pion decay mode,

$$\rho^+ \rightarrow \pi^0 + \pi^+, \quad (1)$$

via a p-wave. Each π has a c. m. momentum $p_\pi = 350$ Mev/c. If η is 1^{--} , then the $\eta\pi$ decay mode of ρ is allowed,

$$\rho^+ \rightarrow \eta + \pi^+, \quad (2)$$

again via a p-wave with $p = 123 \text{ Mev}/c$. A crude estimate of the decay rate is $\Gamma(\rho \rightarrow \eta\pi) \approx \frac{p^2}{1+(pR)^2} \cdot \frac{p}{m_\rho} (2S_\eta + 1) R^3$, where the first factor accounts for p-wave barrier penetration, p/m_ρ is the Lorentz-invariant phase space, $(2S_\eta + 1) = 3$ is the multiplicity of the η spin, and R is some radius of interaction. We can write a similar expression for $\Gamma(\rho \rightarrow 2\pi)$. If we assume equal radii $R = \hbar/2m_\pi$ for both processes, then we find⁶

$$\Gamma(\rho \rightarrow \eta\pi)/\Gamma(\rho \rightarrow 2\pi) \approx 1/4. \quad (3)$$

To compare the two decay modes, we chose a sample of 2000 events



produced in the 72-in. hydrogen bubble chamber⁷ by 1.25-Bev/c π^+ .

Of these, 1684 fit the hypothesis $\pi^+ + p \rightarrow p + \pi^+ + \pi^0$, and 1100 of the 1684 have a two-pion mass $m_{\pi^+\pi^0}$ falling in the ρ peak. We attribute 500 of these to ρ production



and subsequent decay according to reaction (1).⁸ Corresponding to these 500 $\rho \rightarrow 2\pi$ events, our estimate (3) calls for $500/4 \approx 125 \rho \rightarrow \eta\pi$ events. Now the η decays with a branching fraction $f_{\text{neutral}} \approx 3/4$ entirely into neutrals, and with $f_{\text{charged}} \approx 1/4$ into $\pi^+\pi^-\pi^0$. Hence 125 η should produce $125 \times 3/4 \approx 100$ two-prong stars fulfilling the two conditions:

$$(a) \ m(\text{neutrals}) = 550 \pm 7 \text{ Mev}, \quad (b) \ m(+, \text{ neutrals}) = 750 \pm 50 \text{ Mev}.$$

Instead, if the events are plotted in a two-dimensional $m(\text{neutral}, m(+, \text{neutral}))$ space, we find a flat distribution containing in the area of interest only five events, all of which appear to be $p\pi^+2\pi^0$ background and of which at most a few ($\approx \sqrt{5}$) can be attributed to $\rho \rightarrow \eta + \pi^+$.⁹

In summary of the experimental situation: 500 ρ yield $\leq \sqrt{5} \eta$ decaying via f_{neutral} , and therefore $\leq \frac{4}{3} \sqrt{5} = 3 \eta$ altogether; i. e.,

$\Gamma(\rho \rightarrow \eta \pi) / \Gamma(\rho \rightarrow 2\pi) < 3/500 = 0.6\%$, with an uncertainty of 0.2%. Since our estimate (3) was 25% instead of 0.6%, we conclude either that our estimate must be too high by a factor ≈ 40 , or that $\rho \rightarrow \eta + \pi$ is forbidden by G parity and hence that η is 0^{++} .

In our experience, crude phase-space estimates such as expression (3) are seldom wrong by large factors: In other words, where strong interactions are involved, rates are usually controlled only by phase-space factors and selection rules. However, in this case we must point out that an independent estimate of the strength of the $\pi \rho \eta_{1--}$ vertex can be made based on the known π^0 decay rate $\Gamma(\pi^0 \rightarrow 2\gamma)$ via the dispersion theory diagram of Fig. 1. (If η is not a vector meson, the 1^{--} leg of the diagram can be only the ω , and Gell-Mann et al. already used this diagram to calculate the strength of of the $\pi \rho \omega$ vertex.¹⁰ If the η is a vector meson, then the $\pi \rho \eta_{1--}$ vertex will presumably be the dominant factor in π^0 decay). Glashow and Sakurai have used the $\pi \rho 1^{--}$ vertex as calculated by Gell-Mann et al. for the ω meson to calculate a width for the reaction we sought. They found $\Gamma(\rho \rightarrow \pi \eta_{1--} \approx 1 \text{ Mev})$. Since we know $\Gamma(\rho \rightarrow 2\pi) \approx 100 \text{ Mev}$, their version of our estimate (3) is about 1% instead of our 25%, and we would have expected to see only $500 \times 1\% \times 3/4 = 4$ events even for a $1^{--} \eta$ meson. Hence, as we warned, if all these considerations involving Fig. 1 are correct, then our experiment may not be sufficiently sensitive to rule out the vector η .

As an alternate explanation for the slow rate of π^0 decay, Chew points out that the coupling between a state of odd G-parity necessarily involves heavy intermediate particles, and that consequently the $\eta_{1--} \gamma$ matrix element could be substantially less than e.¹¹ Then the $\pi \rho \eta_{1--}$ coupling could be "normal," and our estimate (3) would still be justified.

We conclude that this experiment suggests that η has even G-parity, but theoretical uncertainties are such that, with our data alone, we can claim no proof.¹²

In the "8-fold way" Gell-Mann¹³ predicted the four strangeness-zero mesons listed in Table I,¹¹ and named them π - ρ , χ - ω . Accordingly, it begins to appear as though the η should be rechristened χ . Independent of the details of the eight-fold way, we note that these four mesons have spins, parities, and G-parity consistent with the model in which they can dissociate into $\bar{N}N$ pairs, bound in $1S_0$ or $3S_1$ states.

Table I. Strangeness of zero mesons

<u>Spin I</u>	<u>0^- (pseudoscalar)</u>	<u>1^- (Vector)</u>
0	$\chi(0^{-+})$	$\omega(1^{--})$
1	$\pi(0^{--})$	$\rho(1^{-+})$

We wish to thank Professors G. F. Chew, M. Gell-Mann, S. L. Glashow, and J. J. Sakurai for helpful discussions.

FIGURE LEGEND


Fig. 1. Decay of π^0 via vector mesons.

FOOTNOTES

- * Work sponsored by U. S. Atomic Energy Commission.
- † On leave of absence from the Inter-University Institute for Nuclear Sciences, Brussels, Belgium.
1. A. Pevsner, R. Kraemer, M. Nussbaum, C. Richardson, P. Schlein, R. Strand, T. Toohig, M. Block, A. Engler, R. Cessaroli, and C. Meltzer, *Phys. Rev. Letters* 8, 421 (1961).
 2. D. D. Carmony, A. H. Rosenfeld, and R. T. Van de Walle, *Phys. Rev. Letters* 8, 117 (1962).
 3. J. J. Sakurai, *Phys. Rev. Letters* 7, 355 (1961).
 4. P. L. Bastien, J. P. Berge, O. I. Dahl, M. Ferro-Luzzi, D. H. Miller, J. J. Murray, A. H. Rosenfeld, and M. B. Watson, *Phys. Rev. Letters*, 8, 114 (1962).
 5. G. Feinberg (submitted to *Phys. Rev. Letters*).
 6. G. Feinberg (reference 5) also assumes $R = \hbar/2m_{\pi}$, but assumes a specific matrix element and estimates $\Gamma_{\eta\pi}/\Gamma_{2\pi} \approx 3/4$.
 7. D. D. Carmony and R. T. Van de Walle, *Phys. Rev. Letters* 8, 73 (1962) and Lawrence Radiation Laboratory Report UCRL-9933, Feb. 1962 (submitted to *Phys. Rev.*); D. D. Carmony, Lawrence Radiation Laboratory Report UCRL-9886 (Thesis), Oct. 1961.
 8. That is, we apportion the events in the peak region into 600 background events and 500 due to ρ production. We then estimate that the number of ρ actually produced might be between 300 and 700. The large uncertainty arises because part of the background can interfere with the ρ events, changing the population of the peak and its central value.
 9. The weaker η decay mode into $\pi^+\pi^-\pi^0$ leads to four-prong stars, which

have not been measured; since they give a less sensitive test, it is doubtful that we would find any peak among them.

10. "Decay Rates of Neutral Mesons", M. Gell-Mann, D. Sharp, and W. G. Wagner, California Institute of Technology, Pasadena, Calif. (to be published).
11. B. R. Desai, Phys. Rev. 124, 1248 (1961).
12. We must also admit our surprise that the three-pion η decay modes are faster than the $\pi^+ \pi^- \gamma$ and $\gamma \gamma$ modes (see reference 4).
13. M. Gell-Mann, California Institute of Technology Scientific Laboratory Report CT-SI-20, Jan. 20, 1961 and Phys. Rev. (to be published).



MU-25910

Fig. 1

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.