UC San Diego

UC San Diego Previously Published Works

Title

Neutrino-induced fission of neutron-rich nuclei.

Permalink

<https://escholarship.org/uc/item/0z99315k>

Journal

Physical review letters, 92(11)

ISSN

0031-9007

Authors

Kolbe, E
Langanke, K
Fuller, GM

Publication Date

2004-03-01

DOI

10.1103/physrevlett.92.111101

Peer reviewed

Neutrino-Induced Fission of Neutron-Rich Nuclei

E. Kolbe,^{1,2} K. Langanke,^{3,2} and G. M. Fuller^{4,2}

¹*Departement für Physik, Universität Basel, Basel, Switzerland*

²*Institute for Nuclear Theory, University of Washington, Seattle, Washington 98195, USA*

³*Institut for Fysik og Astronomi, Århus Universitet, DK-8000 Århus C, Denmark*

⁴*Physics Department, University of California, San Diego, La Jolla, California 92093-0319, USA*

(Received 20 August 2003; published 17 March 2004)

We calculate neutrino-induced fission cross sections for selected nuclei with $Z = 84\text{--}92$. We show that these reactions populate the daughter nucleus at excitation energies where shell effects are significantly washed out, effectively reducing the fission barrier. If the r process occurs in the presence of a strong neutrino fluence, and electron neutrino average energies are sufficiently high, perhaps as a result of matter-enhanced neutrino flavor transformation, then neutrino-induced fission could lead to significant alteration in the r -process flow in slow outflow scenarios.

DOI: 10.1103/PhysRevLett.92.111101

PACS numbers: 95.30.Cq, 25.30.Pt, 25.85.-w, 26.30.+k

In this Letter we calculate neutrino capture-induced fission cross sections for heavy nuclei associated with rapid neutron capture (r -process) nucleosynthesis. In the r process, neutrons are captured on nuclei at rates in excess of the typical charge-changing weak interaction rates associated with these nuclei, leading to the production of heavy elements, e.g., iodine, gold, and uranium. Though roughly half of all nuclei with masses in excess of $A = 100$ are thought to have been made in the r process, the site of its origin, as well as much of the weak interaction physics history associated with it remain mysterious. A leading candidate site for r -process nucleosynthesis is the neutrino-heated ejecta following a core collapse supernova event. The intense neutrino fluxes of such an environment force a reassessment of the role of neutrino-nucleus neutral and charged-current interaction processes. Neutrino charged-current capture-induced fission, in particular, could be especially important in determining the nuclear reaction flow paths and nuclear abundances in the r process. Matter-enhanced neutrino flavor transformation could enhance this effect. Recently there has been an avalanche of new observational data on the abundance of elements made in the r process [1] which highlights the need for clarification of these weak interaction issues. For example, observations of the abundances of r -process-produced elements in low-metallicity, old galactic halo stars show patterns which agree with the solar r -process element abundance distribution for nuclides with mass numbers $A > 130$ but do not reproduce the solar r -process pattern for the lighter r -process elements.

It was recognized some time ago [2] that ν_e capture on heavy nuclei in the postcollapse supernova environment would leave the daughter nuclei at the high excitation energies characteristic of Gamow-Teller resonances. This leaves these nuclei vulnerable to fission. Recently, Qian has demonstrated [3] that ν_e capture-induced fission in neutrino-driven wind scenarios [4] can account for the observed abundance patterns. In this model it was pro-

posed that neutrino-induced fission occurs after the r process freezes out (i.e., all initial neutrons are exhausted) and the progenitor nuclei decay to stability.

Neutrino capture-induced fission cross sections have not been calculated before. Two aspects of nuclear physics conspire to make this process potentially important in dense environments with large neutrino fluxes: (i) the weak strength distribution in the charged-current (neutrino capture) channel is such that the postcapture daughter nucleus likely will be left in a highly excited state and (ii) fission barriers are lower at higher excitation energy.

It is expected that charged-current reactions on r -process nuclides will have larger partial fission cross sections than neutral-current reactions, despite the fact that the latter can be induced by $\nu_{\mu,\tau}$ neutrinos and their antiparticles, which, in a core bounce supernova explosion, *might* have larger average energies ($\langle E_\nu \rangle \sim 20\text{--}25$ MeV) than the ν_e neutrinos have ($\langle E_\nu \rangle \sim 10$ MeV). For the neutron-rich nuclei along the r -process path neutrino capture cross sections are quite large, as both allowed channels [Fermi and Gamow-Teller (GT)] are governed by sum rules which scale with the neutron excess $N - Z$ and the ν_e neutrino energy is large enough to excite the centroids of these allowed responses. Furthermore, the isobaric analog state and the GT centroid are located at energies in the daughter nucleus ($E \sim 20\text{--}30$ MeV) which are significantly above the fission barriers in these nuclei. Hence, fission can represent an important, even dominant decay mode following neutrino-induced reactions on neutron-rich nuclei. The principal competing decay mode is neutron emission, as neutron thresholds are also quite low in r -process nuclei.

Our calculations of neutrino-induced reactions proceed through two steps. First we calculate the neutrino cross sections as functions of excitation energy in the final nucleus and then determine the decay mode of the final nuclear state using a statistical approach. The neutrino cross sections are calculated with the random phase approximation (RPA), considering multipoles up to $J = 4$

and both parities. (See Ref. [5].) Our RPA scheme treats proton and neutron degrees of freedom separately and employs a partial occupancy formalism for nonclosed shell nuclei. We adopt a zero-range Migdal force as a residual interaction. We note that the RPA satisfies the Fermi and Ikeda sum rules, which fix the total strength for the allowed transitions.

In the second step we calculate for each final state with well-defined energy, angular momentum, and parity the branching ratios into the various decay channels using the statistical model code SMOKER [6], considering proton, neutron, α and γ emission, as well as fission. The fission barriers employed here were taken from the compilation of Howard and Möller [7] and the neutron separation energies from the mass table of Hilf *et al.* [8]. The final states in the residual nucleus were taken from the experimentally known levels supplemented at higher energies by an appropriate level density formula [6].

Assuming a typical supernova ν_e neutrino spectrum, i.e., a Fermi-Dirac spectrum with temperature $T_\nu = 4$ MeV and zero chemical potential, we have calculated the total (ν_e, e^-) cross section and the neutrino-induced fission cross section for selected even-even nuclei with charge numbers $Z = 84$ –92 (Fig. 1). As the total cross sections are dominated by allowed contributions, the cross sections increase linearly with neutron excess within the various isotope chains, simply reflecting the sum rules. The differences between the total and the fission cross sections are mainly accounted for by the partial (ν, e^-n) cross sections, although for the lighter Po and Rn isotopes the decay into the gamma channel

can compete with the fission decay branch. Because of the relatively high thresholds (Coulomb barriers), branchings into the proton and α channels are negligible. The competition between the two dominant decay modes, neutron emission and fission, are shown in Fig. 2 for selected Th and U isotopes. In our calculation, fission dominates the decay, except for the most neutron-rich nuclides shown. We note that this calculation considers only the decay branchings in the daughter nucleus and does not follow multiple decays; i.e., it represents the “first-chance” fission cross sections [9].

The competition between the dominant decay modes (fission and neutron emission) in a neutrino capture-excited daughter is governed by the relative values of the fission barrier B_f and the neutron separation energy S_n . The fission probability P_f is then approximately given by [10]

$$P_f = \frac{1}{1 + 4(m_n/\hbar^2)R^2 \text{Texp}\{(B_f - S_n)/T\}}, \quad (1)$$

where m_n is the nucleon mass and $R = 1.2A^{1/3}$ is the nuclear radius. This formula assumes that the decaying nucleus is at excitation energies E which are significantly larger than B_f and S_n , e.g., for $E \sim 25$ MeV. Such excitation energies correspond to nuclear temperatures of $T \approx 1$ MeV in nuclei with $A \sim 230$ –270. For simplicity we have assumed that $T = 1$ MeV in the following. Equation (1) yields $P_f \sim 1/6$ if $B_f = S_n$, and $P_f = 0.5$ if $B_f - S_n = -1.6$ MeV. The difference $U = B_f - S_n$ is strongly dependent on the excitation energy. In fact, Eq. (1) is derived from statistical considerations involving the level density at vanishing nuclear deformation (for the neutron emission probability) and at the saddle points of the double-humped fission barriers. The latter corresponds to a sizable nuclear deformation, where the level density increases faster than at vanishing deformation. This reduces U with increasing excitation energy,

FIG. 1 (color online). Neutrino-induced charged-current cross sections on selected Po (upper panel), Rn (second panel), Ra (third panel), Th (fourth panel), and U (lower panel) nuclides. The total cross sections are shown by circles and the partial fission cross section by squares. A Fermi-Dirac spectrum with temperature $T_\nu = 4$ MeV and zero chemical potential has been assumed for the ν_e neutrinos.

FIG. 2 (color online). Total (ν_e, e^-) (circles) and partial (ν_e, e^-n) (triangles) and neutrino-fission cross sections (squares) for selected Th (left panels) and U (right panels) nuclides. The calculations have been performed for Fermi-Dirac neutrino spectra with temperature $T_\nu = 4$ and 8 MeV and zero chemical potential.

enhancing the fission probability relative to neutron emission [9,11]. This is consistent with the fact that the fission barrier in heavy nuclei is strongly influenced by shell effects [10,12], and these are washed out with increasing excitation energy. Using Eq. (1) we have inverted our calculated fission probabilities to obtain $U(T) = B_f(T) - S_n(T)$, assuming $T = 1$ MeV. The desired quantity $\Delta U(T) = U(T=0) - U(T)$ is plotted in Fig. 3, where $U(0)$ has been derived from the compiled fission barriers [7] and neutron separation energies [8]. We note that the energy reduction is significant, amounting to about 4 MeV on average. This result is in good agreement with earlier estimates for heavy nuclei [9]. Although Fig. 3 shows some scatter among the studied nuclei caused by structure effects, we assume that $U(T)$ for supernova (ν_e, e^-) reactions on neutron-rich nuclei is lowered by 4 MeV compared to its ground state value. This allows for some interesting conclusions, which are rather independent of the chosen fission barriers and neutron separation energies.

For charged-current reactions with supernova ν_e neutrinos one then has $P_f = 0.5$ for $B_f - S_n \sim 2.4$ MeV and $P_f = 0.2$ for $B_f - S_n = 3.7$ MeV, where B_f and S_n are the tabulated values appropriate for low excitation energies. Note that the predicted fission barrier heights vary quite significantly where modern evaluations (i.e., [13]) give higher barriers for neutron-rich nuclei than did earlier work (i.e., [7]). The recent fission barriers of [13] predict a fission probability $P_f > 0.2$ for most nuclei with $Z > 92$ and $A > 230$, with the exception of nuclei with lower Z values around the potentially magic neutron number $N = 184$. The fission barriers of Möller and Howard [7] allow for significant spontaneous fission probabilities for nuclei with $Z > 87$ and $A > 230$, including those around $N = 184$. However, nuclei with $A < 230$

have a smaller fission probability in neutrino-induced reactions than tentatively assumed by Qian [3].

Nuclei on the r -process path have $S_n \sim 1.5$ – 2.5 MeV. Such nuclei fission after excitation by neutrinos with probability $P_f > 0.2$ if $B_f \sim 5.2$ – 6.2 MeV. This condition is satisfied for some nuclei on the r -process path with $Z \geq 96$ for the fission barriers of [13] and for $Z \geq 88$ for those of [7], which are, however, likely too small for neutron-rich nuclei [6]. This might suggest that ν -induced reactions in a sizable neutrino flux could initiate fission cycling. The fission barriers of [13] are clearly too high to allow for fission cycling during the r process by β -delayed or neutron-induced fission.

We note that the typical fission cross section for Th and U isotopes ($\sim 400 \times 10^{-42}$ cm²) corresponds to a half-life of ~ 0.08 s, assuming a neutrino reaction at a radius of 100 km above the neutron star and a typical supernova ν_e luminosity of 10^{52} ergs s⁻¹. Such a half-life is shorter than the expected half-lives for the r -process waiting point nuclei with $N = 126$ and $A > 195$ [14] (and also with $N = 184$ and $A > 280$ [15]). These typical half-lives may be shorter than the ~ 0.1 s expansion time scale in “slow” neutrino-driven wind models. Thus, if the r process occurs in a strong neutrino fluence neutrino-induced fission on the progenitor nuclei during the decay to stability might affect the relative Th/U r -process abundance. This abundance ratio is a necessary theoretical ingredient if one wants to deduce an age limit for the Universe from the observed Th/U abundance ratios in old galactic halo stars [16].

The leverage that neutrino capture-induced fission has in an r -process set in a neutrino-driven wind is dependent on the ν_e energy spectrum and on the neutrino fluxes at the position where the neutrons are captured. Models with an extremely fast outflow rate [17,18] generally have neutron capture occurring far from the neutron star where neutrino fluxes are low and, hence, neutrino capture-induced fission effects could be scant, though postprocessing fission could still be significant.

Models with a slow outflow rate suffer from a deficit of neutrons [2] associated with the “alpha effect.” However, these models can yield a viable r process close to the neutron star if neutrino flavor mixing effects are invoked [19,20]. A hierarchical neutrino energy spectrum, one where the μ and τ flavor neutrinos are more energetic than the electron neutrinos remains a possibility for at least some epochs following the bounce of the supernova core. In this case, matter-enhanced neutrino flavor transformation can play an important role in determining the efficacy of neutrino capture-induced fission, by making the average energies of the electron neutrinos larger and, hence, boosting fission probabilities in the region where neutrons are being captured in the r process. (This is demonstrated in Fig. 2 for a $T = 8$ MeV neutrino spectrum.) The relationship between radial distance r_6 from the neutron star’s center in units of 10 km and the

FIG. 3. The difference $\Delta U = U(0) - U(T)$ for ν capture-induced fission for ν_e with a Fermi-Dirac distribution with temperature $T_\nu = 4$ MeV and zero chemical potential. $U(T) = B_f(T) - S_n(T)$ is calculated as described in the text, while $U(0)$ is derived from the tabulated fission barriers and neutron separation energies.

temperature T_9 in 10^9 of Kelvins is $r_6 \approx 22.5/(S_{100}T_9)$, where S_{100} is the entropy per baryon in units of hundreds of Boltzmann's constant. Typically, neutron capture in the "slow outflow" schemes takes place in the region where $1 < T_9 < 3$. The location where a neutrino of energy E_r will transform its flavor is

$$T_9^{\text{MSW}} \approx 1.3(20 \text{ MeV}/E_r)^{1/3}[0.42/(Y_e + Y_\nu)]^{1/3}S_{100}^{1/3} \\ \times (\delta m^2 \cos 2\theta/3 \times 10^{-3} \text{ eV}^2)^{1/3}, \quad (2)$$

where δm^2 is the relevant difference of the squares of the vacuum neutrino mass eigenvalues and θ is the effective two-neutrino vacuum mixing angle, which for the $\nu_e \rightleftharpoons \nu_{\mu/\tau}$ transformation channel in a strictly three-neutrino mass/mixing scheme would be roughly $\theta_{13} < 0.15$. The experimental upper limit on this mixing angle precludes an adiabatic transformation at resonance in a straight Mikheyev-Smirnov-Wolfenstein (MSW) scheme for $\delta m^2 \sim 10^{-3} \text{ eV}^2$ and $E_\nu \sim 25 \text{ MeV}$, but we note that nevertheless large effective matter mixing could occur on account of the flavor basis off-diagonal neutrino-neutrino forward scattering contributions to the weak potential. Here Y_e is the electron fraction and Y_ν is the effective neutrino number fraction which enters into the neutrino forward scattering potential. The above expression for the neutrino transformation location is conservative: we may actually have a more energetic electron ν_e spectrum on account of "chaotic" maximal mixing below this position. Schemes with sterile neutrino mixing also can produce energetic ν_e 's. Crudely, the neutrino flux is $5 \times 10^{42} \text{ cm}^{-2} \text{ s}^{-1} r_6^{-2} (10 \text{ MeV}/\langle E_{\nu_e} \rangle) L_{\nu_e}^{51}$. Here $L_{\nu_e}^{51}$ is the effective electron neutrino luminosity in units of $10^{51} \text{ erg s}^{-1}$. If the entropy per baryon is $S_{100} = 2$, then the radius where a neutrino of energy $E_r = 20 \text{ MeV}$ transforms is $r_6 \approx 7$ (corresponding to $T_9 \approx 1.6$), and we expect the typical lifetime against fission per big nucleus to be $\lambda_f^{-1} \approx 0.05 \text{ s}/L_{\nu_\alpha}^{51}$, where $\alpha = e, \mu, \tau$ is the flavor of the progenitor of the electron neutrino when it leaves the neutrino sphere; whereas, if $Y_e + Y_\nu = 0.1$, a possibility if neutrino mixing has been augmented by the neutrino background potential(s), then $r_6 \approx 4.3$ and $T_9 \approx 2.6$ (for $S_{100} = 2$) so that $\lambda_f^{-1} \approx 0.02 \text{ s}/L_{\nu_\alpha}^{51}$. In either case, these lifetimes are shorter than typical waiting point r -process beta decay lifetimes and are shorter than at least a plausible range of expansion time scales, $\tau_{\text{dyn}} \sim 0.015 \text{ s}$. This implies that the neutron capture flow could proceed out to some threshold nuclide mass in the 195 peak or just beyond, whereupon fission sets in, producing two fission fragments in the 130 peak, as outlined by Qian.

To establish a steady state fission cycling scenario with neutrino capture-induced fission alone is problematic. If, in steady state flow, every seed nucleus is brought by neutron capture to a nuclear mass where the fission cross section is greater than some threshold value, σ_f^{th} , then fission of this nucleus will

result. Over a time $\Delta t \sim 2\tau_{\text{dyn}}$ there will be only some $\sim 72(\lambda_f/300 \text{ s}^{-1})(\Delta t/0.03 \text{ s})(N/8)$ neutrons liberated per threshold nuclear mass, where N is the assumed number of neutrons liberated per fission. Sustaining steady state fission cycle flow would require the liberation of some 70 to 100 neutrons per fission fragment and this is clearly untenable. Nevertheless, a more modest number of neutrons liberated per fission coupled with the large rate of mass 130 fission fragment production could represent a significant alteration in the r -process flow. At the very least it shows that the mass 130 and 195 peaks should have comparable abundances.

Discussions with Aksel S. Jensen are highly appreciated. We thank F.-K. Thielemann for making the SMOKER code available to us. This work is supported in part by the Danish Research Council, by NSF Grant No. PHY00-99499 at UCSD, and by a DoE SciDAC grant at UCSD.

-
- [1] C. Sneden *et al.*, *Astrophys. J.* **533**, L139 (2000).
 - [2] G. M. Fuller and B. S. Meyer, *Astrophys. J.* **453**, 792 (1995).
 - [3] Y.-Z. Qian, *Astrophys. J.* **569**, L103 (2002).
 - [4] S. E. Woosley, D. H. Hartmann, R. D. Hoffman, and W. C. Haxton, *Astrophys. J.* **356**, 272 (1990).
 - [5] E. Kolbe, K. Langanke, and P. Vogel, *Phys. Rev. C* **50**, 2576 (1994); *Nucl. Phys. A* **652**, 91 (1999).
 - [6] J. J. Cowan, F.-K. Thielemann, and J. W. Truran, *Phys. Rep.* **208**, 267 (1991).
 - [7] W. M. Howard and P. Möller, *At. Data Nucl. Data Tables* **25**, 219 (1980).
 - [8] E. R. Hilf, H. von Groote, and K. Takahashi, in *Proceedings of the 3rd Conference on Nuclei far from Stability, Cargese* (CERN, Geneva, 1976), p. 142.
 - [9] L. G. Moretto, *Nucl. Phys. A* **180**, 337 (1972).
 - [10] P. J. Siemens and A. S. Jensen, *Elements of Nuclei* (Addison-Wesley Publishers, Redwood City, CA, 1987).
 - [11] A. S. Jensen and J. Damgaard, *Nucl. Phys. A* **203**, 578 (1973); **A210**, 282 (1973).
 - [12] M. Brack, in *Proceedings on the Symposium of Physics and Chemistry of Fission* (IAEA, Jülich, Vienna, 1980), p. 227.
 - [13] A. Mamdouh, J. M. Pearson, M. Rayet, and F. Tondeur, *Nucl. Phys. A* **679**, 377 (2000).
 - [14] K. Langanke and G. Martinez-Pinedo, *Rev. Mod. Phys.* **75**, 819 (2003).
 - [15] P. Möller, J. R. Nix, and K.-L. Kratz, *At. Data Nucl. Data Tables* **66**, 131 (1997).
 - [16] R. Cayrel *et al.*, *Nature (London)* **409**, 691 (2001).
 - [17] C. Y. Cardall and G. M. Fuller, *Astrophys. J.* **486**, L111 (1997).
 - [18] T. Thompson, A. S. Burrows, and B. Meyer, *Astrophys. J.* **562**, 887 (2001).
 - [19] G. C. McLaughlin, J. Fetter, A. B. Balantekin, and G. M. Fuller, *Phys. Rev. C* **59**, 2873 (2001).
 - [20] J. Fetter, G. C. McLaughlin, A. B. Balantekin, and G. M. Fuller, *Astrophys. J.* **18**, 433 (2003).