UC Irvine

UC Irvine Previously Published Works

Title

Reply to comment by H. Vereecken et al. on "Field observations of soil moisture variability across scales"

Permalink

<https://escholarship.org/uc/item/0zk7v00m>

Journal

Water Resources Research, 44(12)

ISSN

0043-1397

Authors

Famiglietti, JS
Ryu, D
Berg, AA
[et al.](#)

Publication Date

2008-12-01

DOI

10.1029/2008WR007323

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Reply to comment by H. Vereecken et al. on “Field observations of soil moisture variability across scales”

James S. Famiglietti,¹ Dongryeol Ryu,² Aaron A. Berg,³
Matthew Rodell,⁴ and Thomas J. Jackson²

Received 30 July 2008; revised 30 July 2008; accepted 15 September 2008; published 4 December 2008.

Citation: Famiglietti, J. S., D. Ryu, A. A. Berg, M. Rodell, and T. J. Jackson (2008), Reply to comment by H. Vereecken et al. on “Field observations of soil moisture variability across scales,” *Water Resour. Res.*, 44, W12602, doi:10.1029/2008WR007323.

[1] We welcome and appreciate the comments of Vereecken et al. [2008] on our recent paper [Famiglietti et al., 2008]. Their comments provide the opportunity to reemphasize the complex nature of the processes driving the observed behavior of soil moisture variability across scales presented by Famiglietti et al. [2008].

[2] Vereecken et al. [2008] suggest that the spatial variability of soil moisture content versus its field mean, as presented in Famiglietti et al. [2008], can be largely explained by the shape of soil moisture retention curves for a homogeneous soil, and, for a heterogeneous soil, it is related to soil variability of the constitutive model parameters which control the shape of retention curves. As an example, Vereecken et al. [2008] show soil moisture content (θ) versus soil water capacity $C(\theta)$ and $\Delta\theta$ on the basis of the van Genuchten model, which forms concave curves with peaks exiting between the two end-members (i.e., residual water content and porosity) of soil moisture. Indeed, the behavioral similarity between the variability observed by Famiglietti et al. [2008] and the curves calculated using stochastic theory of unsaturated flow [Vereecken et al., 2007, 2008] has encouraged us to explore the fundamental causality that may exist between them. And we agree with Vereecken et al. [2008] in that, with given uniformly distributed suction head in a homogeneous soil, soil moisture variability peaks in the medium range of mean soil moisture content, and that progressively decreasing unsaturated hydraulic conductivity implied in the moisture retention curve plays an important role in reducing soil moisture variability toward the dry end of field mean soil moisture content.

[3] However, we would like to point out that some of the basic assumptions underlying the stochastic theory are rarely if ever met in the real field conditions, which makes it difficult to directly link our observations to the theory. For

instance, a heterogeneous soil in Vereecken et al. [2007] and Zhang et al. [1998] is defined as a uniform texture soil with hydraulic properties that are stationary random variables (e.g., $\ln K_s$, α , β). On the contrary, most regional-scale fields (e.g., 50-km-scale fields) in Famiglietti et al. [2008] contain more than one classes of soil texture, and the different drainage rates among the textural groups are an important source of high soil moisture variability in the midrange of mean soil moisture content. Another important source of peak variability in the midrange of soil moisture content is fractional rainfall coverage that typically occurs within regional-scale fields. In fact, Ryu and Famiglietti [2005] show that, at the regional scale, fractional rainfall has larger impact on soil moisture spatial variability than soil heterogeneity. Even at local (~ 800 -m) scales, lateral redistribution of soil moisture by topography can play a dominant role in creating soil moisture variability [Mohanty et al., 2000]. These examples, i.e., nonstationary heterogeneity in soil hydraulic parameters, highly heterogeneous atmospheric forcing, and lateral soil water flow, illustrate some of the frequently observed field conditions that are not considered by the stochastic theory.

[4] We agree with Vereecken et al. [2008] that existing theories in stochastic unsaturated flow can give an important insight into the behavioral features of soil moisture variability under many circumstances. However, the above-listed examples highlight a number of other important factors that contribute to the high variability observed in the midrange of mean soil moisture content. Additional examples can be found in section 5.1 of Famiglietti et al. [2008]. The complex interplay of processes driving soil moisture variations at larger scales was an important motivation for us to conduct a behavioral analysis rather than a process-oriented study. Owing to the existence of multiple controlling factors on observed soil moisture variability, we suggest that the similarity between the observed concavity of Famiglietti et al. [2008] and the simulated curves of Vereecken et al. [2007] (e.g., location of the peak) is only in part causal.

References

- Famiglietti, J. S., D. Ryu, A. A. Berg, M. Rodell, and T. J. Jackson (2008), Field observations of soil moisture variability across scales, *Water Resour. Res.*, 44, W01423, doi:10.1029/2006WR005804.
- Mohanty, B. P., T. H. Skaggs, and J. S. Famiglietti (2000), Analysis and mapping of field-scale soil moisture variability using high-resolution, ground-based data during the Southern Great Plains 1997 (SGP97)

¹Department of Earth System Science, University of California, Irvine, California, USA.

²Hydrology and Remote Sensing Laboratory, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, Maryland, USA.

³Department of Geography, University of Guelph, Guelph, Ontario, Canada.

⁴Hydrological Sciences Branch, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA.

- Hydrology Experiment, *Water Resour. Res.*, 36(4), 1023–1031, doi:10.1029/1999WR900360.
- Ryu, D., and J. S. Famiglietti (2005), Characterization of footprint-scale surface soil moisture variability using Gaussian and beta distribution functions during the Southern Great Plains 1997 (SGP97) hydrology experiment, *Water Resour. Res.*, 41, W12433, doi:10.1029/2004WR003835.
- Vereecken, H., T. Kamaï, T. Harter, R. Kasteel, J. Hopmans, and J. Vanderborght (2007), Explaining soil moisture variability as a function of mean soil moisture: A stochastic unsaturated flow perspective, *Geophys. Res. Lett.*, 34, L22402, doi:10.1029/2007GL031813.
- Vereecken, H., T. Kamaï, T. Harter, R. Kasteel, J. W. Hopmans, J. A. Huisman, and J. Vanderborght (2008), Comment on ‘Field observations of soil moisture variability across scales’ by James S. Famiglietti et al., *Water Resour. Res.*, 44, W12601, doi:10.1029/2008WR006911.
- Zhang, D. X., T. C. Wallstrom, and C. L. Winter (1998), Stochastic analysis of steady-state unsaturated flow in heterogeneous media: Comparison of the Brooks-Corey and Gardner-Russo models, *Water Resour. Res.*, 34(6), 1437–1449, doi:10.1029/98WR00317.
-
- A. A. Berg, Department of Geography, University of Guelph, Guelph, ON N1G 2W1, Canada.
- J. S. Famiglietti, Department of Earth System Science, University of California, Irvine, CA 92697-3100, USA.
- T. J. Jackson and D. Ryu, Hydrology and Remote Sensing Laboratory, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, MD 20705, USA.
- M. Rodell, Hydrological Sciences Branch, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA.