

To be published in the Proceedings
of the International School of
Physics, Enrico Fermi Course on
Nonlinear Optical Spectroscopy,
Varenna, Italy, June 30 - July 11, 1975

LBL-4178

c.1

NONLINEAR OPTICAL STUDY OF PRETRANSITIONAL
BEHAVIOR IN LIQUID CRYSTALLINE MATERIALS

Y. R. Shen

September 1975

RECEIVED
MAY 10 1976


DEC 10 1975

EX-100

Prepared for the U. S. Energy Research and
Development Administration under Contract W-7405-ENG-48

For Reference

Not to be taken from this room



LBL-4178

c.1

00004400943

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

To appear in The Proceedings of the
International School of Physics, Enrico Fermi
Course on Nonlinear Optical Spectroscopy,
June 30 thru July 11, 1975.

LBL-4178
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Berkeley Laboratory
Berkeley, California
AEC Contract No. W-7405-eng-48

NONLINEAR OPTICAL STUDY OF PRETRANSITIONAL
BEHAVIOR IN LIQUID CRYSTALLINE MATERIALS

Y. R. Shen

September, 1975

4460074000

NONLINEAR OPTICAL STUDY OF PRETRANSITIONAL
BEHAVIOR IN LIQUID CRYSTALLINE MATERIALS

by

Y. R. Shen

Department of Physics, University of California

and

Inorganic Materials Research Division

Lawrence Berkeley Laboratory, Berkeley, California 94720

00004400945

NONLINEAR OPTICAL STUDY OF PRETRANSITIONAL
BEHAVIOR IN LIQUID CRYSTALLINE MATERIALS

Nonlinear optics can sometimes be used as a probe to study phase transition in a material. For example, second-harmonic generation, which is very sensitive to symmetry change, has been used to monitor the phase transition in crystalline NH_4Cl ¹ and quartz². Here, we choose to discuss a case when the optical Kerr effect is used to probe the pretransitional behaviour of the isotropic \rightarrow nematic transition in a liquid crystal³. The same method has been applied to the study of pretransitional behaviour in plastic crystals⁴.

A nematic substance is composed of long molecules. They are randomly oriented in the isotropic phase, but become more or less aligned to a common axis in the nematic phase. We can consider the substance as an analog to a spin system with no spin polarity. Each molecule here plays the role of a spin. The liquid phase is then equivalent to the paramagnetic phase, and the nematic phase to the ferromagnetic phase. In a paramagnetic system, an applied magnetic field H can induce a spin alignment and hence a magnetization along the magnetic field. The induced magnetization M in the mean-field approximation obeys the well-known Curie's law

$$M = C_M H / (T - T_c) \quad (1)$$

which exhibits a critically divergent behaviour as the temperature approaches a critical temperature T_c , where C_M is a constant. A close

analogue exists in liquid crystalline materials. There, an applied optical field $E(\omega)$ can also induce a molecular alignment in the isotropic phase. However, since the field variation at optical frequencies is too fast for the molecules to respond, the induced molecular alignment is proportional to $|E(\omega)|^2$ rather than $E(\omega)$. We therefore have $|E|^2$ play the role of H , while the induced molecular alignment is manifested by the induced anisotropy δn in the optical refractive index. We then expect that δn should also obey the Curie's law

$$\delta n = C |E|^2 / (T - T_c) \quad (2)$$

where C is a constant and T_c is the critical temperature for a second-order isotropic \rightarrow nematic transition in the mean-field approximation. (We should note that δn can also be induced by a dc electric field. However, the results are often more difficult to interpret because of the presence of ionic conduction in the medium which also affects the molecular alignment.)

Let us now give a more rigorous derivation of Eq. (2). We use Landau's theory of second-order phase transition to describe the pre-transitional behaviour of the isotropic \rightarrow nematic transition^{3,5}. First, we define an order parameter Q by the relation

$$\delta \chi = \chi_1 - \chi_1^0 = Q \Delta \chi \quad (3)$$

where χ_{\perp} and χ_{\parallel} are the optical susceptibilities parallel and perpendicular to the applied linearly polarized field $\vec{E}(\omega)$ respectively, and $\Delta\chi$ is the anisotropy in χ when all molecules are perfectly aligned in one direction. Since the average susceptibility is $\chi_o = (\chi_{\perp} + 2\chi_{\parallel})/3$, we have $\chi_{\perp} = \chi_o + 2Q\Delta\chi/3$ and $\chi_{\parallel} = \chi_o - Q\Delta\chi/3$. Then, following Landau's theory, we can write the free energy per unit volume in the isotropic phase ($Q \ll 1$) as

$$F = F_o + \frac{1}{2} A Q^2 + B Q^3 + D Q^4 - \frac{1}{4} \chi_{\perp} |E(\omega)|^2 \quad (4)$$

where $A = a(T - T_o)$ and a , B , and D are constant coefficients. At temperatures sufficiently far above the transition, we expect Q to be so small that the BQ^3 and DQ^4 terms in Eq. (4) become negligible. We can then readily obtain the Curie's law of Eq. (2) from minimization of F with respect to Q . We find

$$\delta n = \frac{2\pi}{n} \delta\chi = \frac{\pi(\Delta\chi)^2}{3 a n} |E|^2 / (T - T_c). \quad (5)$$

It turns out that in nematic substances, the BQ^3 and DQ^4 terms in F , which are responsible for the observed first-order isotropic \rightarrow nematic transition in the medium, are negligibly small even when T is very close to the observed transition temperature T_k . In fact, in several substances, the observed T_k is only less than 1°C above the fictitious critical temperature T_c .

We can also discuss the dynamics of the optical field-induced anisotropy. It is described by the simple equation 3-5

$$\nu \delta Q / \delta t = -\delta F / \delta Q \quad (6)$$

where ν is a viscosity coefficient. The solution of the equation is

$$Q(t) = \int_{-\infty}^t \left(\frac{\Delta \chi |E(\omega)|^2}{6 \nu} \right) e^{-(t-t')/\tau} dt' \quad (7)$$

where

$$\tau = \nu / A = \nu / a(T - T_c) \quad (8)$$

is the relaxation time for the order parameter. Equation (7) shows that if $|E(\omega)|^2$ is a pulse shorter or comparable with τ , then at sufficiently large t , the order parameter Q , and hence the induced optical anisotropy $\delta n = 2\pi Q \Delta \chi / n$, will decrease exponentially with a time constant τ .

From Eq. (8), we note that as T approaches T_c , the relaxation time increases rapidly as $(T - T_c)^{-1}$. This is again analogous to the well-known critical slowing-down behaviour of the induced magnetization in a paramagnetic crystal near the paramagnetic \rightarrow ferromagnetic transition.

It is easy to see from Eq. (7) that by measuring the transient response of $\delta n(t)$ induced by $|E(\omega)|^2$ of a given pulse shape, we can find the constants $\chi_2^{(3)} \equiv (\Delta \chi)^2 / 6a(T - T_c)$ and τ from which we can deduce

$(\Delta \chi)^2/a, v/a,$ and T_c . Note that $\chi_Q^{(3)}$ is a third-order nonlinear optical susceptibility which is connected to the optical Kerr constant K by $K = 2\omega \chi_Q^{(3)}/nc$ and to δn by $\delta n = 2\pi \chi_Q^{(3)} |E|^2/n$. Because of the critically divergent behaviour of $\chi_Q^{(3)}$ and τ , the field-induced δn and its relaxation time for an isotropic liquid crystalline material are much larger than those for an ordinary liquid, especially when the material approaches the isotropic \rightarrow nematic transition.

The experimental arrangement for measuring $\delta n(t)$ is quite simple. An intense laser pulse is used to induce the optical anisotropy or Q , while a weak CW laser beam is used to probe the induced optical anisotropy. Since τ for liquid crystalline materials is of the order of a few nsec or longer, the laser pulse needed for the experiment can be either a Q-switched or a weakly mode-locked one. Such experiments have been carried out for a number of nematic substances. Figure 1 shows, as an example, the experimental results and theoretical comparison of $\chi^{(3)}$ and τ as functions of temperature for a nematic substance MBBA ($C_4H_9 - (C_6H_4) - CH=N - (C_6H_4) - CH_3$). The curves clearly demonstrate the critically divergent behaviour of $\chi_Q^{(3)}$ ($\beta \equiv 2\chi_Q^{(3)}$ in Fig. 1) and τ as the temperature approaches the isotropic \rightarrow nematic transition. We note that $\chi_Q^{(3)}$ and τ shown in Fig. are much larger than those of ordinary liquids. At $T - T_c = 5^\circ C$, $\chi_Q^{(3)}$ (MBBA) = 2.7×10^{-10} esu is almost 100 times larger than that of the well-known Kerr liquid CS_2 at room temperature and τ (MBBA) ≈ 100 nsec is 5×10^4 times longer than that of CS_2 . The large values of $\chi^{(3)}$ and τ seem to be the characteristic

properties of liquid crystalline materials as shown in table I for a number of nematic substances⁶. For MBBA, τ varies from 40 to 800 nsec as $T - T_c$ changes from 14.3 to 0.9°C.

Isotropic liquid crystalline materials with their large optical Kerr constants and the temperature-dependent long relaxation time turn out to be ideal systems for investigation of self-focusing of light. Because of the large δn , a Q-switched laser pulse will self-focus readily in such a medium. By varying the temperature of an appropriate liquid crystalline material, it is now possible to change the relaxation time τ from a value much larger than the laser pulsewidth to a value shorter than the laser pulse width. This means that with the same Q-switched pulse, we can now study not only the quasi-steady-state self-focusing phenomenon ($\tau \ll$ pulsewidth) but also the transient self-focusing phenomenon ($\tau \gg$ pulsewidth) and the intermediate cases between the two. In fact, study of self-focusing in MBBA has so far yielded the only quantitative experimental results on transient self-focusing in a liquid⁷. In ordinary liquids, τ is in the picosecond range, and therefore, picosecond laser pulses must be used to observe transient self-focusing. However, the present picosecond technology is not yet advanced enough for us to study transient self-focusing quantitatively.

For a third-order nonlinear optical effect governed by a third-order nonlinear polarization $P^{(3)} = \chi^{(3)} |E(\omega)|^2 E(\omega')$, there is always a counterpart in light scattering. Thus, corresponding to the optical Kerr effect due to induced ordering, we have light scattering by order -

000044000

parameter fluctuation in an isotropic liquid crystalline material^{3,8}
 Then, Eqs. (4) and (6) with $|E(\omega)|^2 = 0$ can again be used to describe
 the light scattering phenomenon. Since $|E(\omega)|^2 = 0$, there is no induced
 ordering. However, as a result of thermal fluctuations, the mean
 square value $\langle Q^2 \rangle$ is not zero, but from the equipartition theorem
 applied to Eq. (4), is equal to kT/A . Consequently, the scattering
 light intensity is

$$I \propto \left(\frac{d\epsilon}{dQ} \right)^2 \langle Q^2 \rangle = \left(\frac{d\epsilon}{dQ} \right)^2 \frac{kT}{a(T - T_c)} \quad (9)$$

which shows the critically divergent behaviour as $T \rightarrow T_c$. The dynamic
 equation for Q governs the spectrum of the scattered radiation. Equation
 6 with $|E(\omega)|^2 = 0$ is in the form of a relaxation equation, and
 therefore it leads to a Lorentzian spectral line centered at the incoming
 laser frequency. The half width of the Lorentzian line is given by

$$\Gamma = A/\nu' = a(T - T_c)/\nu. \quad (10)$$

We have, as expected, $\Gamma = 1/\tau$ which shows a critical narrowing behaviour
 as $T \rightarrow T_c$. The results in Eqs. (9) and (10) have been experimentally
 verified by Stinson and Litster⁷.

Light scattering by order-parameter fluctuation appears as an
 intense narrow spectral line (halfwidth $\lesssim 100$ MHz) centered at the laser
 frequency. However, as one looks more carefully at the scattering spectrum,

one would find another much broader and much weaker component also centered at the laser frequency but with its wings extended out to tens of a cm^{-1} .⁹ Such a broad component actually exists in all liquids and is believed to be due to light scattering by fluctuations of individual molecular orientations. This is known as Rayleigh -wing scattering.¹⁰ Now, for liquid crystals, there is apparently a dilemma, since the order parameter fluctuation is clearly also a fluctuation in the molecular orientation. How can the same fluctuation of molecular orientation give rise to both the narrow central component and the broad Rayleigh-wing component?

Qualitatively, the answer to the above question is simple. We remember that the Landau theory we have used is actually equivalent to the mean-field theory which takes into account only the average local-field correction. This can be seen as follows. Microscopically, the molecular orientation is induced by the local field

$$|E_{\text{loc}}|^2 = |E(\omega)|^2 + V \quad (11)$$

where V is due to molecular interaction. In the mean-field approximation, we have $V = \lambda Q$ with λ being a proportional constant. The average induced molecular orientation Q is proportional to $|E(\omega)|^2$, i.e., $Q = b(T) |E_{\text{loc}}|^2$. For $|T - T_c| \ll T_c$, by defining $b(T_c) = 1/\lambda$ and expressing $b(T) = b(T_c) + (T - T_c)(db/dT)_{T_c}$ we can write

00004400949

$$Q = \text{Constant } |E(\omega)|^2 / (T - T_c) \quad (12)$$

Equation (12) is of course the same as one would obtain from minimization of F in Eq. (4).

The above derivation essentially follows the derivation of the magnetic susceptibility for a paramagnetic crystal near phase transition taking into account the mean local-field correction. This shows that the results derived from the free energy F in Eq. (4) have not included the temporal and spatial dependence of the local field due to molecular interaction. Physically, the mean field approximation describes the molecular orientational fluctuations under the influence of an average local field and predicts only a single Lorentzian spectral line with critical behaviour for the scattered radiation. In a more rigorous treatment, we should take into account the local-field correction more rigorously. Then, the problem of N interacting molecules can be considered as a problem of N coupled oscillators (referring to the orientational motion) in the highly damped limit. It is well known that a system of N coupled harmonic oscillators has many resonant lines. In the highly damped limit, the resonant frequencies of these lines are negligible compared to the linewidths, but different lines have different linewidths. Consequently, we expect to find a series of Lorentzian lines with different linewidths centered at zero frequency. The narrowest line shows the critical behaviour arising from the average local-field correction, while the others are much broader and show no

critical behaviour but more or less characteristics of orientational fluctuations of individual molecules. The more formal derivation of the above discussion is given in Ref. 11.

ACKNOWLEDGMENT

This work supported by the U.S. Energy Research and Development Administration.

Table I.

Compound	R	T_c ($^{\circ}\text{C}$)	$\tau(T - T_c)$ (10^{-8} sec- $^{\circ}\text{K}$)	$\chi_Q^{(3)}(T - T_c)$ (10^{-10} esu- $^{\circ}\text{K}$)
$\text{C}_4\text{H}_9 - (\text{C}_6\text{H}_4) - \text{CH:N} - (\text{C}_6\text{H}_4) - \text{R}$	CH_3	41.7	67	13.2
	C_2H_5	67.6	21	7.8
	CH_3	130.7	8.5	15
	C_2H_5	161.0	7.5	10.2
$\text{RO} - (\text{C}_6\text{H}_4) - \text{N}_2\text{O} - (\text{C}_6\text{H}_4) - \text{OR}$	C_5H_{11}	115.0	25	5.6
	C_6H_{13}	126.0	15	4.7
	C_7H_{15}	121.3	15	5.4



Table I. Critical transition temperatures T_c , orientational relaxation times τ and nonlinear susceptibilities $\chi_Q^{(3)}$ for various nematic liquid crystalline compounds.

REFERENCES.

1. I. Freund, Phys. Rev. Lett. 19, 1288 (1967).
2. J. P. Bachheimer and G. Dolino, Phys. Rev. B11, 3195 (1975).
3. G. K. L. Wong and Y. R. Shen, Phys. Rev. Lett. 30, 895 (1973);
Phys. Rev. A10, 1277 (1974).
4. T. Bischofberger and E. Courtens 32, 163 (1974).
5. P. G. de Gennes, Phys. Lett. A30, 454 (1969); Mol. Cryst. Liq.
Cryst. 12, 193 (1971).
6. E. G. Hanson (to be published).
7. T. W. Stinson and J. D. Litster, Phys. Rev. Lett. 25, 503 (1970).
8. G. K. L. Wong and Y. R. Shen, Phys. Rev. Lett. 32, 527 (1974).
9. N. M. Amer, Y. S. Lin and Y. R. Shen, Solid State Comm. (1975).
10. See, for example, I. L. Fabellinski, "Molecular Scattering of
Light" (Plenum Press, New York, 1968) Chapters II and VIII.
11. C. Flytzanis and Y. R. Shen, Phys. Rev. Lett. 33, 14 (1974).

FIGURE CAPTION

Fig. 1 (a) Nonlinear refractive index ($\Delta n = \pi\beta|E|^2/n$) as a function of temperature for MBBA. Δ are experimental data from optical Kerr measurements and \circ are experimental data from ellipse-rotation measurements. The solid curve is given by $5.4 \times 10^{-9}/(T - T_c)$ with 314.7°K . (b) Relaxation time τ_L of the order parameter as a function of temperature for MBBA. The solid curve is the theoretical curve and the dots are the experimental data points (after G.K.L. Wong and Y.R. Shen, Phys. Rev. A10, 1277 (1974)).



XBL 759-8140

Fig. 1

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720