

UC Berkeley

Archaeological X-ray Fluorescence Reports

Title

Source Provenance of Obsidian Artifacts from a Selected Surface Sample at Piedras Marcadas (LA 290), Middle Rio Grande Valley, New Mexico

Permalink

<https://escholarship.org/uc/item/11q5k4gz>

Author

Shackley, M. Steven

Publication Date


2014-08-13

Supplemental Material

<https://escholarship.org/uc/item/11q5k4gz#supplemental>

License

[CC BY-NC 4.0](https://creativecommons.org/licenses/by-nc/4.0/)



GEOARCHAEOLOGICAL XRF LAB

GEOARCHAEOLOGICAL X-RAY FLUORESCENCE SPECTROMETRY LABORATORY

8100 WYOMING BLVD., SUITE M4-158

ALBUQUERQUE, NM 87113 USA

**SOURCE PROVENANCE OF OBSIDIAN ARTIFACTS FROM A SELECTED
SURFACE SAMPLE AT PIEDRAS MARCADAS (LA 290), MIDDLE RIO
GRANDE VALLEY, NEW MEXICO**

by

M. Steven Shackley, Ph.D.
Director
Geoarchaeological XRF Laboratory
Albuquerque, New Mexico

Report Prepared for

Dr. Matt Schmader
Albuquerque Open Space
City of Albuquerque, New Mexico

13 August 2014

INTRODUCTION

The analysis here of 50 visually selected obsidian and dacite artifacts from the surface of Piedras Marcadas (LA 290) in the middle Rio Grande River valley indicates a similar mix of sources as the previous analyses of surface and subsurface contexts, with the exception of an increase in the number of artifacts produced from one of the Mount Taylor sources (Shackley 2009, 2013a). In the test unit case, all the artifact quality sources of archaeological obsidian present in the Jemez Mountains, both pre-caldera and caldera event sources occurred in the assemblage. All these sources are present in the Rio Grande alluvium as far south as Albuquerque, although the Valles Rhyolite (Cerro del Medio) nodules are very small. No Mount Taylor obsidian was recovered sub-surface (Shackley 2013a). Mount Taylor is not available in Rio Grande Quaternary sediments this far north. The dacite artifacts are from northern New Mexico sources that have eroded into the Rio Grande Quaternary alluvium.

ANALYSIS AND INSTRUMENTATION

All archaeological samples are analyzed whole. The results presented here are quantitative in that they are derived from "filtered" intensity values ratioed to the appropriate x-ray continuum regions through a least squares fitting formula rather than plotting the proportions of the net intensities in a ternary system (McCarthy and Schamber 1981; Schamber 1977). Or more essentially, these data through the analysis of international rock standards, allow for inter-instrument comparison with a predictable degree of certainty (Hampel 1984).

The trace element analyses were performed in the Geoarchaeological XRF Laboratory, Albuquerque, New Mexico, using a Thermo Scientific *Quant'X* energy dispersive x-ray fluorescence spectrometer. The spectrometer is equipped with a ultra-high flux peltier air cooled Rh x-ray target with a 125 micron beryllium (Be) window, an x-ray generator that operates from 4-50 kV/0.02-1.0 mA at 0.02 increments, using an IBM PC based microprocessor and WinTraceTM 4.1 reduction software. The spectrometer is equipped with a 2001 min⁻¹ Edwards vacuum pump for the analysis of elements below titanium (Ti). Data is acquired through a pulse

processor and analog to digital converter. This is a significant improvement in analytical speed and efficiency beyond the former Spectrace 5000 and *QuanX* analog systems (see Davis et al. 2011; Shackley 2011a).

For Ti-Nb, Pb, Th elements the mid-Zb condition is used operating the x-ray tube at 30 kV, using a 0.05 mm (medium) Pd primary beam filter in an air path at 200 seconds livetime to generate x-ray intensity $K\alpha_1$ -line data for elements titanium (Ti), manganese (Mn), iron (as Fe^T), cobalt (Co), nickel (Ni), copper, (Cu), zinc, (Zn), gallium (Ga), rubidium (Rb), strontium (Sr), yttrium (Y), zirconium (Zr), niobium (Nb), lead (Pb), and thorium (Th). Not all these elements are reported since their values in many volcanic rocks is very low. Trace element intensities were converted to concentration estimates by employing a least-squares calibration line ratioed to the Compton scatter established for each element from the analysis of international rock standards certified by the National Institute of Standards and Technology (NIST), the US. Geological Survey (USGS), Canadian Centre for Mineral and Energy Technology, and the Centre de Recherches Pétrographiques et Géochimiques in France (Govindaraju 1994). Line fitting is linear (XML) for all elements but Fe where a derivative fitting is used to improve the fit for iron and thus for all the other elements. When barium (Ba) is acquired, the Rh tube is operated at 50 kV and 0.5 mA in an air path at 200 seconds livetime to generate x-ray intensity $K\alpha_1$ -line data, through a 0.630 mm Cu (thick) filter ratioed to the bremsstrahlung region (see Davis et al. 2011). Further details concerning the petrological choice of these elements in North American obsidians is available in Shackley (1988, 1990, 1995, 2005; also Mahood and Stimac 1991; and Hughes and Smith 1993). A suite of 17 specific standards used for the best fit regression calibration for elements Ti- Nb, Pb, and Th, include G-2 (basalt), AGV-2 (andesite), GSP-2 (granodiorite), SY-2 (syenite), BHVO-2 (hawaiite), STM-1 (syenite), QLO-1 (quartz latite), RGM-1 (obsidian), W-2 (diabase), BIR-1 (basalt), SDC-1 (mica schist), BCR-2 (basalt), TLM-1 (tonalite), SCO-1 (shale), all US Geological Survey standards, NBS-278 (obsidian) from the National Institute of Standards and Technology, BR-1 (basalt) from the Centre de Recherches Pétrographiques et

Géochimiques in France, and JR-1 and JR-2 (obsidian) from the Geological Survey of Japan (Govindaraju 1994).

The data from the WinTrace software were translated directly into Excel for Windows and into SPSS for statistical manipulation (Table 1). In order to evaluate these quantitative determinations, machine data were compared to measurements of known standards during each run (Table 1). RGM-1 is analyzed during each sample run for obsidian artifacts to check machine calibration (Table 1). Source assignments made by reference to source data at the lab, and Shackley (1995, 2005, 2011b).

DISCUSSION

Before a discussion of the source provenance of the samples, a short discussion of the Jemez Mountains sources is in order. Following this is a short discussion of the samples proper.

The Jemez Mountains and the Sierra de los Valles

A more complete discussion of the archaeological sources of obsidian in the Jemez Mountains is available in Shackley (2005:64-74; see Figure 1 here). Distributed in archaeological contexts over as great a distance as Government Mountain in the San Francisco Volcanic Field in northern Arizona, the Quaternary sources in the Jemez Mountains, most associated with the collapse of the Valles Caldera, are distributed at least as far south as Chihuahua through secondary deposition in the Rio Grande, and east to the Oklahoma and Texas Panhandles through exchange. And like the sources in northern Arizona, the nodule sizes are up to 10 to 30 cm in diameter; El Rechuelos, Cerro Toledo Rhyolite, and Valles Rhyolite (Valles Rhyolite derived from the Cerro del Medio dome complex) glass sources are as good a media for tool production as anywhere. Until the recent land exchange of the Baca Ranch properties, the Valles Rhyolite primary domes (i.e., Cerro del Medio) have been off-limits to most research. The discussion of this source group here is based on collections by Dan Wolfman and others, facilitated by Los Alamos National Laboratory, and the Museum of New Mexico, and recent

sampling of all the major sources courtesy of the Valles Caldera National Preserve (VCNP; Shackley 2005; Wolfman 1994).

There are at least four eruptive events in the last 8.7 million years that have produced the four chemical groups in the Jemez Mountains (Figure 1).




Figure 1. Generalized stratigraphic relations of the major volcanic and alluvial units in the Jemez Mountains (from Gardner et al. 1986). Note the near overlapping events at this scale for the Cerro Toledo and Valles Rhyolite members, and the position of Cerro Toledo Rhyolite at the upper termination of the Puye Formation.

The earliest is the Bear Springs Peak source, part of Canovas Canyon Rhyolite that is dated to about 8.7 mya, firmly in the Tertiary (Kempster et al. 2004; Figure 1 here). This source is a typical Tertiary marekanite source with remnant nodules embedded in a perlitic matrix. It is located in a dome complex including Bear Springs Peak on Santa Fe National Forest and radiating to the northeast through Jemez Nation land (Shackley 2009b). While the nodule sizes are small, the glass is an excellent media for tool production and has been found archaeologically

at Zuni and in secondary deposits as far south as Las Cruces (Church 2000; Shackley 2012). Four of the samples were produced from this source (Table 1 and 2 and Figure 2 here).

Part of the same Keres Member as Canovas Canyon Rhyolite is Paliza Canyon Rhyolite. They have similar elemental chemistry and are likely nearly contemporaneous (Shackley 2014). This source is rare in archaeological contexts, but occurs in Rio Grande alluvium, and is present as one sample here, as in the 2013 subsurface study.

The second relevant eruptive event that produced artifact quality obsidian is the El Rechuelos Rhyolite. This source, present as one sample here, is what I consider the best media for tool production of the group. It dates to about 2.4 million years ago, and nodules at least 10 cm in diameter are present in a number of domes north of dacite Polvadera Peak, the incorrect vernacular name for this source. El Rechuelos has eroded through the Rio Chama into the Rio Grande and has also been found in alluvium into southern New Mexico (Church 2000; Shackley 2013b).

About 1.4 mya, the first caldera collapse occurred in the Jemez Mountains, called Cerro Toledo Rhyolite. This very large event produced the Bandelier Tuffs and spread ash flows many kilometers into the area and horizontally southwest from what is now Rabbit Mountain and the Cerro Toledo domes to the east. These large ash flow sheets are responsible for the great quantity of Cerro Toledo obsidian that is present in the Quaternary Rio Grande alluvium all the way to Chihuahua (Church 2000; Shackley 2005, 2013b). Cerro Toledo Rhyolite secondary deposit nodules are present relatively near to Piedras Marcadas on Quaternary terraces above the east side of the Rio Grande, including Placitas and the Quaternary sands near Tijeras Wash south of the Albuquerque airport (Shackley 2013b).

The second caldera collapse that produced the Valles Rhyolite member of the Tewa Formation, called Valles Rhyolite here, occurred around one million years ago and created most of the geography of the current Valles Caldera. A number of rhyolite ring domes were produced on the east side of the caldera, but only Cerro del Medio produced artifact quality obsidian.

Indeed, the Cerro del Medio dome complex produced millions of tons of artifact quality glass, and is the volumetrically largest obsidian source in the North American Southwest challenged only by the Government Mountain dome complex in the San Francisco Volcanic Field. Cerro del Medio obsidian was apparently preferred by Folsom knappers, as well as those in all periods since. While Cerro Toledo probably appears in archaeological contexts in New Mexico sites with greater frequency, it is likely because it is distributed in secondary contexts. Valles Rhyolite (Cerro del Medio), present as three samples subsurface and one sample here importantly does not erode outside the caldera, in any quantity and size and likely had to be originally procured in the caldera proper (Shackley 2005).

Source Provenance Discussion

Most of the artifacts analyzed produced from all these sources are bipolar core or flake fragments and most appear to have waterworn cortex. This suggests that most of these raw materials were procured across the river somewhere. In the case of the Mount Taylor specimen the raw material had to be procured at Mount Taylor or in the Rio Puerco or the Rio Grande south of Socorro after the Rio Puerco joins the Rio Grande (Shackley 2005, 2013b). Mount Taylor sources (Grants Ridge, Horace and La Jara Mesas) are common in historic period contexts at Zuni and the source may have been “controlled” by the Zuni (Shackley 2005; Table 1 and Figure 3). The mix of sources in this selected assemblage mirrors the mix of sources recovered from the Rio Grande Quaternary Alluvium at Tijeras Wash almost identically (Shackley 2013a: Figure 2). This is the strongest argument for local procurement of obsidian toolstone at Piedras Marcadas as indicated by the test unit assemblage.

Surface versus Subsurface Results

While the samples are relatively small, the mix of sources recovered from surface contexts versus the subsurface test unit sample is somewhat different (see also Shackley 2013a). While both are dominated by Jemez Mountains secondary deposit sources, the presence of Mount Taylor sources on the surface indicates procurement through direct access to the Zuni

region or exchange with the Zuni. IF the subsurface material is earlier, and the surface material later, then one change seen is contact to the west rather than local procurement and/or contact north at an earlier period. It is possible that the Mount Taylor obsidian was procured by the Coronado Expedition knappers when they were at and around Zuni and transported the raw material to Piedras Marcadas during the siege as tool raw material. Again, the sample size is small.

DACITE ARTIFACTS

A number of dacite artifacts, mainly debitage or utilized flakes, were sampled from the surface. All but one of the artifacts were produced from either the San Antonio Mountain source in the Taos Plateau Volcanic Field, or the Cerros del Rio dacite in the Cerros del Rio Volcanic Field on the southern edge of Bandelier National Monument (see Shackley 2011b; Table 3 here). Both these sources erode into the Rio Grande and have been recovered in the alluvium at Tijeras Wash south of Piedras Marcadas. One sample is likely rhyolite or dacite, but doesn't match any known source.

REFERENCES CITED

- Church, T.
2000 Distribution and Sources of Obsidian in the Rio Grande Gravels of New Mexico. *Geoarchaeology* 15:649-678.
- Davis, K.D., T.L. Jackson, M.S. Shackley, T. Teague, and J.H. Hampel
2011 Factors Affecting the Energy-Dispersive X-Ray Fluorescence (EDXRF) Analysis of Archaeological Obsidian. In *X-Ray Fluorescence Spectrometry (XRF) in Geoarchaeology*, edited by M.S. Shackley, pp. 45-64. Springer, New York.
- Gardner, J. N., F. Goff, S. Garcia, R. Hagan
1986 Stratigraphic Relations and Lithologic Variations in the Jemez Volcanic Field, New Mexico. *Journal of Geophysical Research* 91B2:1763-1778.
- Govindaraju, K.
1994 1994 Compilation of Working Values and Sample Description for 383 Geostandards. *Geostandards Newsletter* 18 (special issue).
- Hampel, Joachim H.
1984 Technical Considerations in X-ray Fluorescence Analysis of Obsidian. In *Obsidian Studies in the Great Basin*, edited by R.E. Hughes, pp. 21-25. Contributions of the University of California Archaeological Research Facility 45. Berkeley.
- Hildreth, W.
1981 Gradients in Silicic Magma Chambers: Implications for Lithospheric Magmatism. *Journal of Geophysical Research* 86:10153-10192.
- Hughes, Richard E., and Robert L. Smith
1993 Archaeology, Geology, and Geochemistry in Obsidian Provenance Studies. In *Scale on Archaeological and Geoscientific Perspectives*, edited by J.K. Stein and A.R. Linse, pp. 79-91. Geological Society of America Special Paper 283.
- Kempton, K., G.R. Osburn, S. Kelley, M. Rampey, C. Ferguson, and J. Gardner
2004 Geology of the Bear Springs Peak 7.5' Quadrangle, Sandoval County, New Mexico. Open File Geologic Map OF-GM 74 (Draft), New Mexico Bureau of Mineral Resources, Socorro.
- Mahood, Gail A., and James A. Stimac
1990 Trace-Element Partitioning in Pantellerites and Trachytes. *Geochemica et Cosmochimica Acta* 54:2257-2276.
- McCarthy, J.J., and F.H. Schamber
1981 Least-Squares Fit with Digital Filter: A Status Report. In *Energy Dispersive X-ray Spectrometry*, edited by K.F.J. Heinrich, D.E. Newbury, R.L. Myklebust, and C.E. Fiori, pp. 273-296. National Bureau of Standards Special Publication 604, Washington, D.C.
- Schamber, F.H.
1977 A Modification of the Linear Least-Squares Fitting Method which Provides Continuum Suppression. In *X-ray Fluorescence Analysis of Environmental Samples*, edited by T.G. Dzubay, pp. 241-257. Ann Arbor Science Publishers.
- Shackley, M. Steven
1988 Sources of Archaeological Obsidian in the Southwest: An Archaeological, Petrological, and Geochemical Study. *American Antiquity* 53(4):752-772.

- 1990 *Early Hunter-Gatherer Procurement Ranges in the Southwest: Evidence from Obsidian Geochemistry and Lithic Technology*. Ph.D. dissertation, Arizona State University, Tempe.
- 1995 Sources of Archaeological Obsidian in the Greater American Southwest: An Update and Quantitative Analysis. *American Antiquity* 60(3):531-551.
- 2005 *Obsidian: Geology and Archaeology in the North American Southwest*. University of Arizona Press, Tucson.
- 2009 Source Provenance of Obsidian Artifacts from Piedras Marcadas (LA 290), Middle Rio Grande Valley, New Mexico. Report prepared for Matt Schmader, Albuquerque Open Space, City of Albuquerque, New Mexico.
- 2011a An Introduction to X-Ray Fluorescence (XRF) Analysis in Archaeology. In *X-Ray Fluorescence Spectrometry (XRF) in Geoarchaeology*, edited by M.S. Shackley, pp. 7-44. Springer, New York.
- 2011b Sources of Archaeological Dacite in Northern New Mexico. *Journal of Archaeological Science* 38:1001-1007
- 2013a Source Provenance of Obsidian Artifacts from Three Subsurface Test Units at Piedras Marcadas (LA 290), Middle Rio Grande Valley, New Mexico. Report prepared for Albuquerque Open Space, City of Albuquerque, New Mexico.
- 2013b The Secondary Distribution of Archaeological Obsidian in Rio Grande Quaternary Sediments, Jemez Mountains to San Antonito, New Mexico: Inferences for Paleoamerican Procurement and the Age of Sediments. Poster presented at the Paleoamerican Odyssey Conference, Santa Fe, New Mexico, October, 2013.
- 2014 The Paliza Canyon Source of Archaeological Obsidian in the Southern Jemez Mountains, Northern New Mexico. *IAOS Bulletin* 50: 6-9.
- Wolfman, Daniel
- 1994 Jemez Mountains Chronology Study. Report prepared by the Office of Archaeological Studies, Museum of New Mexico for the USDA Forest Service, Contract No. 53-8379-9-14.

Table 1. Elemental concentrations for the archaeological specimens and the USGS RGM-1 standard. All measurements in parts per million (ppm).

Sample	Ti	Mn	Fe	Rb	Sr	Y	Zr	Nb	Pb	Th	Source
100	689	46	1209	20	9	67	17	10	36	31	Cerro Toledo Rhy
		9	3	3			0	0			
101	354	57	1163	51	12	88	13	23	59	30	Horace Mesa (Mt Taylor)
		2	3	3			1	3			
103	514	46	1174	20	8	68	17	98	35	20	Cerro Toledo Rhy
		5	3	5			2				
104	417	58	1170	52	13	84	13	22	60	36	Horace Mesa (Mt Taylor)
		3	0	9			2	3			
105	537	48	1183	20	8	68	18	10	33	21	Cerro Toledo Rhy
		3	5	5			0	2			
106	548	46	1183	19	10	62	17	98	35	24	Cerro Toledo Rhy
		9	7	9			3				
107	634	48	1235	20	8	63	17	99	33	25	Cerro Toledo Rhy
		9	0	6			0				
108	646	56	1264	23	9	68	18	10	44	27	Cerro Toledo Rhy
		4	7	2			8	4			
110	601	54	1262	22	11	68	18	10	41	31	Cerro Toledo Rhy
		0	3	4			5	4			
111	551	50	1219	21	8	67	17	10	37	21	Cerro Toledo Rhy
		0	2	0			8	1			
112	678	47	1223	20	11	59	17	98	35	25	Cerro Toledo Rhy
		8	6	3			4				
113	581	45	1180	20	8	65	17	94	34	20	Cerro Toledo Rhy
		6	3	1			0				
114	605	48	1194	19	9	61	16	99	34	27	Cerro Toledo Rhy
		0	2	9			9				
115	708	40	1214	16	11	46	16	54	25	18	Valles Rhy (Cerro del Medio)
		4	5	6			7				
116	524	46	1164	19	11	65	17	10	31	28	Cerro Toledo Rhy
		6	3	7			2	0			
117	660	57	1283	22	9	68	18	10	42	26	Cerro Toledo Rhy
		7	5	7			3	4			
118	548	44	1154	19	8	66	17	97	32	30	Cerro Toledo Rhy
		1	4	9			1				
119	473	44	1161	20	9	65	17	96	35	24	Cerro Toledo Rhy
		9	4	4			4				
120	112	49	1099	10	88	23	12	33	20	16	Paliza Canyon
	1	6	9	5			6				
121-1	620	41	1147	18	9	59	15	96	30	25	Cerro Toledo Rhy
		9	0	5			7				
121-2	526	51	1211	21	9	64	17	96	36	23	Cerro Toledo Rhy
		2	9	5			7				
122-1	599	40	1023	15	11	23	71	46	24	16	El Rechuelos
		1	9	4							
122-2	524	47	1173	19	10	61	17	94	33	24	Cerro Toledo Rhy
		5	0	9			3				
123-1	501	49	1186	20	10	66	16	98	33	21	Cerro Toledo Rhy
		8	5	7			8				
123-2	535	53	1224	21	8	67	17	96	38	22	Cerro Toledo Rhy
		8	2	5			7				
124	571	45	1166	20	16	61	16	97	33	26	Cerro Toledo Rhy
		7	1	7			3				
125	461	36	1096	17	8	59	16	92	26	18	Cerro Toledo Rhy
		6	1	9			1				
128	668	52	1221	19	10	61	17	97	33	26	Cerro Toledo Rhy
		9	9	8			1				

129	553	47	1190	20	10	65	17	99	33	24	Cerro Toledo Rhy
		2	1	2			4				
130	497	48	1182	20	8	67	16	99	35	22	Cerro Toledo Rhy
		1	8	1			9				
131	579	49	1188	20	8	65	17	10	32	23	Cerro Toledo Rhy
		0	5	7			7	1			
132	494	45	1164	20	10	61	16	10	32	18	Cerro Toledo Rhy
		6	8	0			7	0			
135	310	72	1084	54	12	78	11	19	59	23	Grants Ridge (Mt Taylor)
		0	5	4			8	4			
137-1	594	47	1172	20	10	65	17	98	35	26	Cerro Toledo Rhy
		9	7	6			2				
137-2	483	45	1163	20	10	66	16	98	33	23	Cerro Toledo Rhy
		9	8	6			9				
138-1	493	46	1181	20	11	63	17	99	35	30	Cerro Toledo Rhy
		5	8	8			9				
138-2	522	49	1200	21	8	63	17	10	38	29	Cerro Toledo Rhy
		5	5	1			9	3			
139-1	502	47	1172	20	8	61	17	97	32	23	Cerro Toledo Rhy
		5	8	2			3				
139-2	535	51	1231	22	11	70	17	10	39	20	Cerro Toledo Rhy
		5	3	0			7	0			
140	456	48	1185	20	9	63	17	98	35	21	Cerro Toledo Rhy
		2	1	6			2				
141	574	45	1194	21	8	65	17	96	38	26	Cerro Toledo Rhy
		7	5	0			6				
142	559	47	1180	20	9	62	17	10	33	23	Cerro Toledo Rhy
		2	0	7			4	0			
143	508	46	1173	20	8	64	17	92	34	25	Cerro Toledo Rhy
		3	6	2			3				
144	545	52	1206	20	10	62	17	93	36	25	Cerro Toledo Rhy
		4	7	9			4				
145	569	46	1182	20	9	61	17	91	33	31	Cerro Toledo Rhy
		5	7	6			2				
RGM1-S4	155	28	1374	14	10	26	21	8	20	13	standard
	4	5	6	9	5		2				
RGM1-S4	159	28	1377	14	10	22	21	8	20	12	standard
	0	3	2	8	6		8				
RGM1-S5	153	28	1368	15	10	22	21	10	17	11	standard
	4	5	6	0	8		4				

Table 2. Frequency distribution of sources in the selected surface assemblage.

	Frequency	Percent
Cerro Toledo Rhy	39	86.7
El Rechuelos	1	2.2
Mt Taylor	3	6.7
Paliza Canyon	1	2.2
Valles Rhy (Cerro del Medio)	1	2.2
Total	45	100.0

Table 3. Elemental concentrations for the dacite/rhyolite samples.

Sample	Ti	Mn	Fe	Rb	Sr	Y	Zr	Nb	Pb	Th	Source
102	3851.467	432.498	19294.69	48.634	596.563	18.476	174.996	23.081	9.519	7.867	San Antonio Mtn dacite
109	6314.205	916.665	31600.83	93.064	666.718	33.45	242.255	15.493	8.621	6.723	San Antonio Mtn dacite
126	4646.919	626.298	26011.9	48.329	859.167	17.505	208.616	21.244	12.64	3.096	Cerros del Rio dacite
133	1099.237	706.971	13527.73	147.663	15.872	66.155	397.993	39.266	29.116	21.144	rhyolite or dacite
134	4252.501	570.218	25260.16	42.539	840.703	17.807	200.52	22.429	12.526	9.303	Cerros del Rio dacite




Figure 2. Zr versus Rb bivariate plot of the samples. Cerro Toledo Rhyolite and Valles Rhyolite samples discriminated by Y and Nb, not plotted here.




Figure 3. Frequency histograms of source provenance in the test units (top left), general surface (top right), and selected surface (bottom center)