

UC Riverside

UC Riverside Previously Published Works

Title

Intercellular and systemic trafficking of RNAs in plants

Permalink

<https://escholarship.org/uc/item/11w7854f>

Journal

Nature Plants, 4(11)

ISSN

2055-026X

Authors

Liu, Lin
Chen, Xuemei

Publication Date

2018-11-01

DOI

10.1038/s41477-018-0288-5

Peer reviewed



Published in final edited form as:

Nat Plants. 2018 November ; 4(11): 869–878. doi:10.1038/s41477-018-0288-5.

Intercellular and systemic trafficking of RNAs in plants

Lin Liu^{1,2}, Xuemei Chen^{3,*}

¹Guangdong Provincial Key Laboratory for Plant Epigenetics, College of Life Sciences and Oceanography, Shenzhen University, Shenzhen, China.

²Longhua Bioindustry and Innovation Research Institute, Shenzhen University, Shenzhen, China.

³Department of Botany and Plant Sciences, Institute of Integrative Genome Biology, University of California, Riverside, CA, USA.

Abstract

Plants have evolved dynamic and complex networks of cell-to-cell communication to coordinate and adapt their growth and development to a variety of environmental changes. In addition to small molecules, such as metabolites and phytohormones, macromolecules such as proteins and RNAs also act as signalling agents in plants. As information molecules, RNAs can move locally between cells through plasmodesmata, and over long distances through phloem. Non-cell-autonomous RNAs may act as mobile signals to regulate plant development, nutrient allocation, gene silencing, antiviral defence, stress responses and many other physiological processes in plants. Recent work has shed light on mobile RNAs and, in some cases, uncovered their roles in intercellular and systemic signalling networks. This review summarizes the current knowledge of local and systemic RNA movement, and discusses the potential regulatory mechanisms and biological significance of RNA trafficking in plants.

Cell-to-cell communication plays a critical role in plant development, disease resistance and responses to various stresses from the external environment. As a strategy for efficient intercellular communications, plants have evolved a plant-specific symplasmic pathway mediated by plasmodesmata (PD) and phloem to transport signalling molecules between cells¹. Various types of plant RNA species, including messenger RNAs (mRNAs), small interfering RNAs (siRNAs), microRNAs (miRNAs), ribosomal RNAs (rRNAs) and transfer RNAs (tRNAs), can move from cell to cell (short-range) or systemically (long-range) to potentially regulate whole-plant physiological processes^{2–5}. The non-cell-autonomous nature of RNA molecules suggests that RNAs may function beyond the cells in which they are synthesized. Regulatory roles of mobile RNAs in cell differentiation, organ formation and patterning, nutrient homeostasis, stress adaptations, and plant-microorganism and plant-

Reprints and permissions information is available at www.nature.com/reprints.

*Correspondence should be addressed to X.C., xuemei.chen@ucr.edu.

Author contributions

L. L. and X. C. wrote this article.

Competing interests

The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

plant interactions have been discovered, prompting further studies to understand the scope and impact of RNA trafficking in signalling networks. Recently, thanks to advances in genomics technologies, an abundance of mobile RNAs has been discovered^{6–10}, further underscoring the question of the functional significance of mobile RNAs. Here, we review recent advances in intercellular and systemic RNA trafficking in plants and discuss the possible regulatory mechanisms and biological functions of RNA trafficking.

Routes for RNA trafficking between plant cells

Together, PD and phloem form a symplasmic pathway that links nearly all plant cells. RNAs can move cell-to-cell through PD and long-distance through phloem. A vesicle-mediated pathway is also a potential route for RNA trafficking between plant cells.

PD as intercellular micro-channels of mobile RNAs.

PD are membrane-lined micro-channels that cross the cell wall and connect neighbouring cells¹¹. They are bordered by plasma membrane, and contain an appressed form of endoplasmic reticulum (ER), called desmotubule, in the central region (Fig. 1a). The space between these membranes forms a cytosolic sleeve that allows cellular molecules to migrate between cells. Mobile molecules may also move through the lumen or membrane of the desmotubule^{12–14}. The key feature of PD is that they establish cytosolic and endomembrane continuity between adjacent cells, thus forming a symplasmic pathway to mediate the transportation of molecules between adjacent cells^{1,15}. Various molecules are able to move through PD, including small molecules such as water, ions and phytohormones, as well as large molecules such as proteins and RNAs¹⁶. The transport of molecules through PD depends on their size, shape and tissue types, along with the development stages in which they are present. As highly dynamic channels, PD are tightly regulated and undergo various structural and functional changes during plant development. The numbers and size exclusion limit of PD vary among different tissues at different development stages^{17–19}. The structure of PD ranges from simple channels to twinned and branched channels²⁰. The size exclusion limit of PD is regulated by reversible callose (β -1,3-glucan) deposition and by certain PD-associated proteins and mobile proteins^{11,16,21,22}. A number of mobile RNAs, such as miR390 (ref.³), miR165/166 (ref.⁵), sucrose transporter *SUC1* mRNA (ref.²³), and transcription factor *KNOTTED1* mRNA (ref.²⁴), have been reported to move through PD, indicating the importance of these channels for RNA trafficking in plants.

Long-distance movement of RNAs through phloem.

While PD mediate the cell-to-cell movement of cellular molecules, the vascular system—consisting of xylem and phloem—facilitates long-distance trafficking in plants. Water and mineral nutrients are transported from roots to the aerial parts of plants through the xylem. In contrast, phloem supports the movement of photosynthates and macromolecules from source to sink tissues²⁵. The phloem is composed of living enucleated sieve elements assisted by companion cells (Fig. 1b). Mature sieve-elements are connected to adjacent companion cells by highly modified, funnel-like PD called plasmodesmata pore units (PPUs). These specialized PPU are structurally distinct from regular PD. They consist of multiple channels on the companion-cell side, which merge into a single pore on the sieve-

element side²⁶. The sieve element cells stack together to form the sieve tube, which allows for rapid flow of molecules over long distances in plants²⁷. Macromolecules, including proteins, RNAs and ribonucleoprotein complexes, have been found in the phloem stream^{4,28,29}. Plant phloem appears to be an ideal route for RNA trafficking because no RNase activity is detectable in phloem sap^{30,31}. As discussed in detail in the section ‘Mobile RNAs in plants’, many lines of evidence support that various RNA species are transported through the phloem to distant tissues^{6,8,32}.

Potential RNA trafficking routes mediated by vesicles.

In eukaryotes, proteins are generally secreted into the extracellular space through the classic ER-Golgi route. However, cytosolic proteins lacking signal peptides are also found outside of cells, indicating the existence of unconventional protein secretion pathways³³. In animals, one such secretion pathway involves the release of exosomes to the extracellular compartment. Exosomes are membrane-bound vesicles 30–100 nm in diameter and derived from multivesicular bodies (MVBs) that fuse with the plasma membrane, resulting in the release of their intraluminal vesicles as exosomes³⁴. Interestingly, in addition to proteins, various RNAs including mRNAs, miRNAs and other non-coding RNAs, have also been identified in exosomes and have been shown to be critical for intercellular communication between animal cells³⁴. The secretion of exosomes appears to be an evolutionarily conserved process, and accumulating evidence indicates that exosome-like vesicles also exist in plants. Structures similar to exosomes have been isolated from the apoplastic fluids of sunflower (*Helianthus annuus*) seeds, olive (*Olea europaea*) pollen grains and *Arabidopsis* leaves, and these exosome-like vesicles are enriched with leaderless secretory proteins^{35–37}.

Ultrastructural data has shown that barley leaves release vesicles resembling exosomes under pathogenic fungal attack³⁸. When invaded by filamentous oomycetes pathogens, a series of structural and biochemical responses takes place in the host plant cells, including the re-organization of subcellular structures. Numerous organelles accumulate in the vicinity of haustorium infected sites, particularly MVBs, Golgi stacks, ER and secretory vesicles, indicating that the secretion process of the host plant cell may be activated at the penetration site to defend against pathogenic infection^{39,40}. Animal exosomes can mediate RNA trafficking for cell-to-cell communication and affect the phenotypes of recipient cells³⁴. It was recently revealed that plants also employ exosome-like vesicles to transport small RNAs to a fungal pathogen⁴¹. It is unknown whether exosomes or other vesicles are used in RNA trafficking between plant cells, as plant cells are separated by cell walls, unlike the highly specialized haustoria at fungal infection sites. However, given that exosomes can be detected from plant tissues or organs, it is a formal possibility that RNA trafficking between neighbouring cells occurs through exosomes. Figure 1c depicts imaginary scenarios whereby exosomes or vesicles may transport RNAs intercellularly, either through exocytosis/endocytosis or through the PD channels between cells. Vesicle-mediated RNA trafficking deserves attention in future research.

Mobile RNAs in plants

Various types of plant RNA species, including mRNAs, small RNAs, ribosomal RNAs and transfer RNAs, have been found to travel beyond the cells in which they are synthesized.

Mobile mRNAs.

Intercellular trafficking of plant mRNAs through PD was first shown in microinjection experiments with the maize *KNOTTED1* (*KN1*) transcription factor²⁴. Fluorescently labelled sense *kn1* RNA and KN1 protein were coinjected into tobacco mesophyll cells, and the fluorescent probes were observed to move rapidly from the injected cell into neighbouring cells, indicating that the transcript of *KN1* can be transported through PD with the assistance of KN1 protein²⁴. Another strong evidence for mRNA being mobile is from localization studies of potato sucrose transporter *suc1* mRNA (ref.²³). *Suc1* mRNA was transcribed in companion cells, however in situ hybridization revealed the presence of the transcript in both companion cells and sieve elements. Because enucleated sieve elements lack the transcription machinery, the presence of *suc1* mRNA was attributed to cell-to-cell movement of the mRNA from companion cells to adjacent sieve elements, probably through PD (ref.²³). Subsequent studies investigated the distribution of RNAs in the phloem sap of various plant species, including *Arabidopsis*⁴², rice³¹, barley^{30,43}, pumpkin⁴⁴, melon⁴⁵, *Ricinus communis*⁴⁶, *Lupinus albus*⁴⁷, watermelon and cucumber³². These studies identified numerous phloem transcripts encoding various types of proteins, such as transcription factors, phytohormone regulators, stress response factors and proteins involved in a wide range of plant developmental processes. Although the presence in phloem sap alone was insufficient evidence of RNA systemic movement or functionality, the mobility of some phloem mRNAs was subsequently confirmed by grafting studies^{44,48–52} and host-parasite interaction analysis^{8,9,53,54}.

Grafting connects two or more living tissues from different plants into one single plant⁵⁵. Grafting has been widely used to test the translocation of molecules across the graft junctions and has proven to be a useful tool for characterizing long-distance mobile RNAs. Examples of mRNAs whose mobility was demonstrated by grafting experiments (Table 1) include *Arabidopsis FT* (ref.⁵⁶) and *Aux/IAA* (ref.⁴⁸), pumpkin *PP16* (ref.⁴⁹) and *NACP* (ref.⁴⁴), potato *BEL5* (ref.⁵⁰) and *POTH1* (ref.⁵¹), apple *SLR/IAA14* (ref.⁵²), and tomato *PPF-T6* (ref.²) and *PS* (ref.⁵⁷). Some recent studies have identified large numbers of graft-transmissible mRNAs using high-throughput sequencing. From the heterograft of *Arabidopsis* and tobacco, 138 transcripts from the stock of *Arabidopsis* were found to move into the tobacco scion⁵⁸. A separate study reported a total number of 2,006 mobile RNAs by grafting shoots and roots of different *Arabidopsis* ecotypes⁸. The difference in the numbers of mobile RNAs identified in these two studies may be attributable to differences in sequencing coverage and plant growth conditions. Indeed, even 2,006 mobile RNAs may be an underestimate, given that mobile RNA detection is dependent on the availability of single nucleotide polymorphisms (SNPs), mRNA stability, sequencing depth, materials sampled and other biological factors. In addition to the model plant *Arabidopsis*, high-throughput sequencing has identified mobile RNAs across graft junctions in agricultural crops, including grape¹⁰ and cucumber⁷ from which the numbers of identified mobile transcripts were 3,333 and 3,546, respectively.

Besides the grafting approach, host-parasite interactions analyses have also provided strong evidence for RNA movement. In host-parasite interactions, mRNAs can move from host plants to parasitic plants across the parasite's haustorium. Examples of mRNAs capable of

moving from host plants to their parasitic plants (Table 1) include tomato *GAI* and *PFP* (refs^{53,54}). Genome-wide analysis of mRNAs exchanged between host and parasitic plants revealed thousands of mobile transcripts^{8,9,59}. In an *Arabidopsis-Cuscuta* host-parasite system, nearly half of the expressed transcriptome of the *Arabidopsis* host was found to move into the *Cuscuta* parasite⁹.

These mobile RNAomics data reveal a large number of mobile transcripts and raise the possibility of RNA-based systemic signalling in plants. Future studies are needed to identify the patterns of RNA movement in plants, the underlying regulatory mechanisms and the functional impacts of mobile RNAs.

Mobile small RNAs: siRNAs and miRNAs.

Small RNAs are a population of 21 to 24 nucleotide (nt) RNAs that guide RNA silencing. They can be classified into two major categories in plants, siRNAs and miRNAs, based on their biogenesis and molecular features⁶⁰. miRNA genes are transcribed by RNA polymerase II (Pol II), generating imperfectly paired hairpin precursors that are cleaved by DICER-LIKE1 (DCL1) to produce mature miRNAs. These mature miRNAs are incorporated into ARGONAUTE 1 (AGO1) to form the RNA-induced silencing complex (RISC), which catalyses the cleavage or translational repression of target RNAs⁶¹. SiRNAs are further classified as trans-acting siRNAs (ta-siRNAs) and heterochromatic siRNAs (hc-siRNAs). These siRNAs are generated from long double-stranded RNAs (dsRNAs) produced from single-stranded RNAs by RNA-dependent RNA polymerases (RDRs) or formed from the transcription of inverted-repeat (IR) sequences. In *Arabidopsis*, the dsRNAs are mainly processed by DCL2, DCL3 or DCL4 and generate 21-, 22- or 24-nt siRNAs, which are sorted into AGO1, AGO2, AGO3, AGO4, AGO6 or AGO9 RISCs to silence target genes⁶². Silencing mediated by small RNAs can be classified as transcriptional gene silencing (TGS) or post-transcriptional gene silencing (PTGS). TGS regulates transposable elements and repetitive DNA sequences through DNA methylation or histone modifications in the nucleus, whereas PTGS functions primarily to eliminate exogenous invading RNAs or to regulate endogenous genes through RNA cleavage or translation inhibition in the cytoplasm⁶³.

Over the past few decades, extensive studies have uncovered non-cell-autonomous features of RNA silencing. Systemic spreading of silencing was initially shown in tobacco plants using grafting experiments, in which a scion containing a non-silenced transgene was grafted onto a stock containing a silenced transgene⁶⁴. In a parallel study, green fluorescent protein (GFP) silencing was observed in the upper leaves of tobacco plants stably expressing a *GFP* transgene after the induction of GFP silencing by transient *Agrobacterium* infiltration of the lower leaves⁶⁵. Subsequent studies further confirmed the systemic nature of RNA silencing and indicated the existence of mobile sequence-specific signals^{66–68}. It was hypothesized that small RNAs or their precursors act as mobile signalling molecules that mediate the spreading of RNA silencing. To further investigate the non-cell-autonomous nature of RNA silencing, two artificial siRNA reporter systems, SUC-SUL and SUC-PDS, were developed. In both systems, long inverted-repeat dsRNAs are expressed in phloem companion cells but target the chlorophyll biosynthesis gene *SULPHUR* (*SUL*) or the

carotenoid biosynthesis gene *PHYTOENE DESATURASE (PDS)* in leaf mesophyll cells^{69,70}. Processing of the inverted repeat dsRNAs generates siRNAs and results in non-cell-autonomous RNA silencing spreading from the vasculature to the surrounding 10 to 15 neighbouring cells, reflecting local cell-to-cell transmission of RNA silencing. Genetic screens based on these two reporter systems were performed, leading to identification of a number of genes that affect RNA silencing, including *AGO1*, *HEN1*, *CLSY1*, *DCL1*, *DCLA*, *NRPD1A*, *RDR2* and *HPR1*; these findings have been well reviewed in ref.⁶⁸.

Both siRNAs and miRNAs have been observed in phloem exudates from various plant species, including pumpkin⁷¹, oilseed rape (*Brassica napus*)^{72,73}, apple⁷⁴ and white lupin (*Lupinus albus*)⁴⁷, further supporting the hypothesis that small RNAs act as long-distance transmitters of RNA silencing in plants.

The mobility of many siRNAs has been documented (Table 1). The first case of endogenous mobile siRNAs was that of the low-abundance, conserved group of *TAS3*-derived ta-siRNAs, termed tasiR-ARFs, which target auxin response factors. The biogenesis of tasiR-ARFs is restricted to the upper adaxial side of leaves by the localized expression of *AGO7* and *TAS3A*. However, in situ hybridization shows that tasiR-ARFs accumulate outside this defined domain of biogenesis and form an adaxial-to-abaxial gradient across leaves. This gradient shapes the expression pattern of the abaxial determinant *AUXIN RESPONSE FACTOR3 (ARF3)*. Thus, tasiR-ARFs function as a mobile signal in the establishment of adaxial-abaxial leaf polarity³. Transposable-element-derived siRNAs could be transported from the pollen vegetative cell to sperm cells to inhibit transposable element activity and stabilize the genome during reproduction^{75,76}. Recently, elegant grafting experiments have demonstrated that both transgene-derived siRNAs and endogenous siRNAs can move long distances^{6,77-79}. By grafting roots from the siRNA biogenesis-defective Dicer triple mutant *dc12 dc13 dc14* to wild-type *Arabidopsis* shoots, a substantial population of mobile siRNAs was identified using high-throughput sequencing⁷⁷. Moreover, these mobile siRNAs were found to target thousands of genomic loci, predominately transposons located in euchromatic regions, for DNA methylation⁶.

In contrast to siRNAs, miRNAs are thought to be relatively less mobile as the sites of their transcription and function are well correlated^{80,81}. However, a few miRNAs have been reported to act non-cell-autonomously (Table 1). For example, the *miR390* precursor localizes to the vasculature and pith region below the shoot apical meristem but not in the meristem or youngest leaf primordia, based on in situ hybridization. However, the accumulation of mature miR390 throughout the vegetative apex, including the meristem and the youngest leaf primordia, strongly suggests that miR390 is capable of cell-to-cell short-distance movement³. miR165/166 species are probably mobile as well; the gene promoters are active in the single-cell-layer root endodermis, while the mature miRNAs accumulate across all radial layers of the root in *Arabidopsis*⁵. Similarly, miR394 was found to spread from the site of biogenesis in the epidermal layer to internal cells in the shoot apical meristem⁸². In addition to short-range movement, long-distance movement of miRNAs has been demonstrated by grafting experiments. Using *Arabidopsis* overexpressing miR399 as the scion and wild type as the rootstock, miR399 was found to exhibit long-range movement from shoot to root to degrade its target transcript *PHOSPHATE 2 (PHO2)* and to regulate

plant phosphate homeostasis^{83,84}. The mobility of miR399 was confirmed in an independent grafting experiment using wild-type *Arabidopsis* as the scion and the miRNA-deficient mutant *hen1* as the rootstock. The same study also revealed that miR395 could traverse graft junctions to target the *APS* gene in the root⁷³. Using the same approach of wild-type-*hen1* micrografting, another study further showed that, in addition to miR399, the corresponding near-complementary miR399* species was graft-transmissible between shoots and roots. The analysis also revealed that miR827 and miR2111 were capable of long-distance movement, while their respective miRNA* species were not, indicating that the mobility of miRNA species is selective⁸⁵. miR2111 in the legume *Lotus japonicus* has also been found to undergo shoot-to-root long-distance translocation to regulate rhizobial infection. While *miR2111:GUS*-expressing lines showed that miR2111 was synthesized in shoots, mature miR2111 was detected in both shoots and roots. In addition, the identification of miR2111 from phloem sap further supports the long-distance transport of this miRNA (ref.⁸⁶).

Small RNAs can traffic between parasites and host plants. The parasitic plant *Cuscuta* was found to induce many miRNAs at the haustorium when it parasitizes *Arabidopsis* and tobacco. The majority of these miRNAs are 22-nt long and they can hijack the silencing machinery of the host cells to produce secondary siRNAs, resulting in the degradation of host mRNAs⁸⁷. In addition, small RNAs were observed to move in a trans-kingdom manner from host plants to parasitic pathogens. miR166 and miR159 were found to be exported from host cotton plants to a fungal pathogen and were shown to down-regulate their fungal target genes *Clp-1* and *HiC-15*, which are essential for fungal virulence⁸⁸. Using the *Arabidopsis-Botrytis cinerea* pathosystem, another study⁴¹ profiled small RNAs from pathogen *B. cinerea* protoplasts and extracellular vesicles of infected *Arabidopsis* leaves. A number of siRNAs and miRNAs have been identified to transfer from *Arabidopsis* to *B. cinerea* through extracellular vesicles to silence fungal virulence-related genes and contribute to host immunity.

The findings described here have helped establish that siRNAs and miRNAs can serve as mobile agents that function in recipient cells (Fig. 2). Thus, non-cell-autonomous RNA silencing may be a signalling mechanism that coordinates developmental and physiological processes in plants.

Other mobile RNAs: rRNAs and tRNAs.

Transcriptome analyses have also revealed the presence of rRNAs and tRNAs in phloem sap (Table 1). All plant rRNA species, including *5S*, *5.8S*, *18S*, and *25S* rRNAs, were found in the phloem exudate of *B. napus* and pumpkin^{4,28,72}. Large quantities of tRNAs have been identified from an RNA pool derived from pumpkin phloem, and high-throughput sequencing revealed that tRNA species distributed non-equally in the phloem sap^{4,89}. For example, while *tRNA-Asp* and *tRNA-Lys* were highly abundant, *tRNA-Ile* and *tRNA-Thr* were barely detected. Moreover, a considerable fraction of the phloem-sap-derived tRNAs was in truncated forms, or tRNA halves, derived from cleavage of tRNAs in the anticodon loop. In vitro assays showed that these phloem-specific tRNA halves effectively inhibited translation in a non-specific manner⁴. In addition to rRNAs and tRNAs, several ribosomal proteins and ribonucleoprotein complexes were also identified in the phloem sap, however,

some important ribosomal proteins or translation factors essential for ribosomal function are missing²⁸. According to these observations, it was suggested that mRNA translation cannot take place in phloem. Further strong evidence supporting this notion is that in a hypocotyl-grafted *GUS:tRNA_{Met}*/wild-type (Col-0) *Arabidopsis* plant, the mobile *GUS:tRNA_{Met}* fusion can move to the root tip of wild-type plants and show GUS activity there, but GUS activity could not be detected in the phloem of the root close to the graft junction, indicating that mobile transcripts are translated after transport⁹⁰. Other non-coding RNA species identified in phloem sap include small nuclear RNAs (snRNAs), mitochondrial and chloroplastic rRNAs and tRNAs, signal recognition particle RNAs and RNAs with unknown function, but it is unclear whether they may unload from the phloem to serve functions elsewhere⁴.

Regulatory mechanisms of RNA movement

Although a large number of RNAs have been found to be mobile in the past few decades, the mechanisms that regulate their movement have only recently begun to emerge. Based on current knowledge, here we discuss the possible mechanisms of RNA trafficking in plants.

RNA sequence, length, abundance and stability are possible factors impacting mobility.

The identification of numerous mobile RNAs raises questions about the mechanisms that determine their mobility. A recent study investigated the potential link between mRNA abundance and mobility using a computational diffusion-based model and concluded that mRNA abundance is a key determinant of mobility. The statistical analyses also indicated that mRNA stability and transcript length might contribute to mobility—mRNAs with predicted longer half-lives and smaller size seem to be more mobile⁹¹. These findings might lead to the assumption that mRNAs traffic in a passive, non-selective manner. However, evidence exists for an active, selective mechanism. For example, *GUS-YFP* transcripts from the strong *35S* promoter do not move to distant plant tissues, suggesting that high levels of expression do not necessarily induce mobility of RNAs^{8,90}. Some highly expressed GFP protein fusions have been shown to be graft-mobile, but their mRNAs are not⁹². Besides, it was observed that mobile transcripts can move from shoot to root (source to sink), or from root to shoot against the source-sink gradient, and certain mobile mRNAs may be transported preferentially to specific tissues, such as flowers or leaves^{8,93}. Specific motifs have also been found to affect RNA mobility. In potato, the 5' and 3' untranslated regions (UTRs) of the BEL1-like transcription factor *BEL5* are required for its long-distance transport into roots, where *BEL5* regulates tuber formation⁹⁴. Some low-abundance mobile transcripts were found to be enriched for three sequence motifs, indicating a sequence-specific selective mechanism for the movement of this set of transcripts⁹¹. In addition, fusion of the phloem enriched tRNA-like structures (TLSs) with immobile RNAs can make the fused transcripts mobile across graft junctions, while deletion of the TLS motif from a dicistronic mRNA-tRNA transcript made it immobile, indicating that TLS motifs can trigger the mobility of mRNAs⁹⁰. By analysing the graft-mobile transcriptome data from *Arabidopsis*⁸ and grape¹⁰, it has been revealed that a large number of mobile transcripts harbour a TLS motif in the coding sequences or UTRs, or are transcribed from genes that are in close proximity to annotated tRNA genes, which further supports that TLSs are necessary

for RNA mobility and indicates the existence of a selective mRNA delivery mechanism⁹⁰. This evidence indicates two possible transport pathways for mRNAs in plants: a passive, nonselective pathway and an active, selective pathway.

RNA trafficking mediated by non-cell-autonomous proteins with RNA-binding activity.

The existence of specific motifs associated with mobile RNAs implies that RNA-binding proteins may interact with RNAs and modulate their trafficking. Phloem exudate analyses have shown the presence of many RNA-binding proteins^{29,95,96}. Some of these proteins were found to associate with RNAs to form ribonucleoprotein complexes and might be involved in phloem-mediated RNA movement. Microinjection assays in tobacco revealed that fluorescein-labelled maize KN1 (ref.²⁴) and pumpkin PP16 (ref.⁴⁹) interact with their own RNAs and specifically facilitate their cell-to-cell or long-distance translocation by regulating the size-exclusion limit of PD. Pumpkin RBP50, a polypyrimidine tract binding (PTB) protein, acts as the core of a ribonucleoprotein (RNP) complex that moves in the phloem translocation stream, as shown by heterografting assays. Co-immunoprecipitation showed that RBP50 associates with six mRNAs containing PTB motifs and incorporates them into the RNP complex to mediate their long-range transport in the phloem⁹⁷. Potato PTB proteins PTB1 and PTB6 have also been found to bind the 3' UTR of the transcript of *BEL5* (ref.⁹⁸). Overexpression of *PTB1/6* resulted in both enhanced stability and long-distance movement of the *BEL5* mRNA, whereas suppression of *PTB1/6* led to decreased stability and movement⁹⁸. Small RNAs may also require RNA-binding proteins for their mobility. The pumpkin phloem protein PSRP1 was found to preferentially bind 25-nt single-stranded RNA (ssRNA) species by gel mobility-shift assays. Co-injection of PSRP1 and fluorescein-labelled 25-nt ssRNAs in tobacco showed that PSRP1 could mediate the trafficking of ssRNAs to neighbouring cells⁷¹. PSRP1 interacts with a specific set of proteins and forms a complex⁹⁹. In vivo co-immunoprecipitation showed that the PSRP1-based complex contains endogenous 24-nt small RNAs. Dephosphorylation of PSRP1 results in disassembly of this small RNA-protein complex, which presumably needs to happen to release small RNAs in target cells⁹⁹. However, orthologues of PSRP1 have yet to be characterized in other plant species.

Control of RNA trafficking by regulating PD permeability.

PD permeability can be regulated to facilitate or block the trafficking of RNA and other molecules during plant development and stress responses, and PD apertures may be temporarily dilated by mobile proteins, such as the aforementioned KN1 and PP16 (refs^{24,49,100}). Many phloem sap proteins were reported to interact with PD to increase their size exclusion limit, whereas isoforms of the proteins absent in phloem were incapable of such interactions^{101,102}. Reversible callose deposition is another mechanism employed by plants to regulate PD permeability in response to various environmental stresses, such as pathogen invasion, wounding and nutrient deficiency^{17,100}. A gain-of-function mutation in *Arabidopsis CALLOSE SYNTHASE 3 (CALS3)* abolishes the intercellular movement of miR165/166, indicating that the regulation of PD permeability is an effective way to control small RNA trafficking¹⁰³. However, research on PD is challenging as mutants lacking PD are lethal and biochemical isolation of PD is difficult^{11,104}.

Specific mechanisms for small RNA mobility.

Given that the biogenesis machinery is different for the different classes of small RNAs, it is possible that the biogenesis pathway affects the mobility of small RNAs. To investigate the movement capability of different types of small RNAs, de Felippes et al.¹⁰⁵ generated artificial miRNA and ta-siRNA reporter lines, in which a small RNA was expressed in phloem companion cells but exerted its silencing effect in leaf mesophyll cells. When the miRNAs and ta-siRNAs were designed to have identical sequences, ta-siRNA-based silencing spread into a much broader area than miRNA-based silencing¹⁰⁵, indicating an influence of small RNA biogenesis on the non-cell-autonomous effects of RNA silencing.

A more recent study revealed that miRNA mobility is regulated through a gating mechanism different from that regulating mobile proteins¹⁰⁶. By using a miR-GFP sensor system, in which the artificial miRNA miR-GFP driven by tissue specific promoters silences a constitutively expressed GFP reporter, miRNA movement was shown to be directional across specific cell-cell interfaces¹⁰⁶. This unidirectional movement generates selectivity in long-distance shoot-to-root trafficking and leads to domain-autonomous behaviours within plant stem cell niches. It should be noted that the two studies^{105,106} both employed artificial miRNAs. Further research is necessary to determine if endogenous miRNAs behave similarly.

Biological functions of RNA trafficking

Against the backdrop of a large set of mobile RNAs, only a relatively small number of them have been demonstrated to function non-cell-autonomously. In these cases, the mobile RNAs play roles in a broad range of physiological processes in plants. For example, long-distance trafficking of tomato *PPF-T6* (ref.²) and pumpkin *GAI* (ref.¹⁰⁷) transcripts was found to affect leaf morphology. miR166 and tasiR-ARFs act non-cell-autonomously in a concentration-dependent manner to generate sharp developmental boundaries to pattern leaf polarity¹⁰⁸. Through grafting experiments, phloem-mobile *FT* was shown to function in systemic floral signalling to regulate flowering time in *Arabidopsis*⁵⁶. RNA detection methods and heterografting experiments demonstrated that potato transcription factor *BEL5* transcripts could be transported to stolon tips and induce tuber formation^{50,94}. The transcripts of two other BEL1-like genes, *BEL11* and *BEL29*, are also phloem-mobile and function antagonistically to *BEL5* to fine-tune the development processes of potato tuberization¹⁰⁹. In *Arabidopsis*, mobile *AUX/IAA* transcripts (ref.⁴⁸) and miR165/166 (ref.⁵) regulate root development. Under stress conditions, miR395 (ref.⁷³) and miR399 (refs^{73,83,84}) act as long-distance signals to regulate sulfate and phosphate homeostasis, respectively. In addition, mobile siRNAs travel systemically to direct DNA methylation of transposable elements in target tissues, including meristematic and meiotically active cells^{6,76,77,79}, which may contribute to epigenetic memory. The non-cell-autonomous nature of RNA silencing also facilitates plant defence against pathogens. On viral infection, siRNAs may act as mobile signals and move in advance of the spread of infection to prime antiviral silencing in uninfected cells to restrict viral spreading^{110,111}. Certain types of small RNAs can even be transferred from plants to pathogens to induce cross-kingdom gene silencing to inhibit virulence gene expression^{41,88,112}. These accumulating pieces of

evidence indicate that RNA trafficking has a role in intercellular and systemic information communication that regulates fundamental biological processes in plants (Fig. 3).

Grafting is routinely used in the cultivation of horticultural crops¹¹³. The large genomic-scale RNA exchange between scions and stocks raises the possibility that RNA trafficking benefits graft performance. The combination of two genetically different graft partners would increase the diversity of the RNA pool accessible to both scion and stock, and this merged RNA pool may provide plants with more genetic resources to achieve better traits, such as enhanced quality or yield. However, it is also possible that mobile RNAs underlie genetic incompatibility leading to graft failure. It has been demonstrated that siRNAs can move into germ-line tissues to methylate target loci^{75,76,79}, so DNA methylation patterns may be altered by grafting and changes could be transmitted to future progeny.

Transcriptome analysis of parasitic plants and their hosts revealed that a large population of mRNAs moves in a bidirectional manner across species^{8,9,59}. The host-parasite interaction has been compared to perfect heterografts, since both of them connect separate plants to form a chimaera. The movement of RNAs between hosts and parasites may function as a means of communication to coordinate their mutual development. The ability for translocation of RNAs from hosts to parasites raises the possibility of using RNA silencing as a strategy to control parasitic weeds¹¹⁴.

Concluding remarks

There have been many advances in documenting and understanding the non-cell autonomy of RNAs in recent years. Thousands of mobile RNAs have been identified through genomics approaches. The recently released database PlaMOM (Plant Mobile Macromolecules) collects and organizes published data on mobile RNAs¹¹⁵. A fluorescence-based RNA labelling system based on the bacteriophage coat protein MS2 has been recently optimized in plants and may potentially serve as a powerful approach for visualizing the real-time trafficking of specific mobile RNAs in living cells¹¹⁶. The discovery of RNAs functioning as mobile molecules is an exciting revelation in plant biology, since RNAs were traditionally believed to function in the same cell in which they are synthesized. With the spatial uncoupling of RNA synthesis and action, at least for some RNAs, the site of action of a given RNA cannot be definitively determined from promoter assays alone.

Despite these advances, many challenges remain. A considerable number of RNAs, but still a tiny portion of the mobile RNAome, have been shown to act as active, mobile signals with roles in coordinating plant growth, development and stress responses^{6,8-10}. Although it has been confirmed that some mobile mRNAs can be translated into functional proteins after transport⁹⁰, it remains unclear to what extent translation of mobile RNAs occurs in recipient tissues. Proteomic data on grafted plants revealed the presence of heterologous protein products that are perhaps encoded by mobile mRNAs⁸, however, the identified proteins are few in number due to their low abundance in recipient tissues. It is also difficult to distinguish between the proteins translated by mobile RNAs and proteins themselves that are mobile. Moving forward, many questions need to be addressed, such as: What are the underlying mechanisms of RNA trafficking? How are RNAs selected for export and import, and do mobile RNAs have other unknown common features? What are the factors that

mediate RNA transport and how do they function? How do plants control the trafficking of RNAs into specific cells or tissues, and what is the fate of mobile RNAs in those recipient tissues? Are they translated and degraded in the same manner as local RNAs in recipient tissues? Are mobile small RNAs transported in single- or double-stranded forms, and are their precursors mobile? Are mobile RNAs conserved in different plant species? Is RNA movement a widespread and functional phenomenon in plant-parasite relationships? Answers to these questions will have broad impacts on our understanding of RNA trafficking and its role in intercellular signalling in plants.

Acknowledgements

We apologize for not being able to discuss or cite many studies due to limited space. Research in the Liu and Chen labs is supported by National Key Research and Development Program of China (91740202), National Institutes of Health (GM061146), Guangdong Innovation Research Team Fund (2014ZT05S078), National Science Foundation (IOS-1340001) and National Natural Science Foundation of China (31600982).

Reference

- Otero S, Helariutta Y & Benitez-Alfonso Y Symplastic communication in organ formation and tissue patterning. *Curr. Opin. Plant Biol* 29, 21–28 (2016). [PubMed: 26658335]
- Kim M, Canio W, Kessler S & Sinha N Developmental changes due to long-distance movement of a homeobox fusion transcript in tomato. *Science* 293, 287–289 (2001). [PubMed: 11452121]
- Chitwood DH et al. Pattern formation via small RNA mobility. *Genes Dev.* 23, 549–554 (2009). [PubMed: 19270155]
- Zhang S, Sun L & Kragler F The phloem-delivered RNA pool contains small noncoding RNAs and interferes with translation. *Plant Physiol.* 150, 378–387 (2009). [PubMed: 19261735]
- Carlsbecker A et al. Cell signalling by microRNA165/6 directs gene dose-dependent root cell fate. *Nature* 465, 316–321 (2010). [PubMed: 20410882]
- Lewsey MG et al. Mobile small RNAs regulate genome-wide DNA methylation. *Proc. Natl Acad. Sci. USA* 113, E801–810 (2016).
- Zhang Z et al. Vascular-mediated signalling involved in early phosphate stress response in plants. *Nat. Plants* 2, 16033 (2016). [PubMed: 27249565]
- Thieme CJ et al. Endogenous *Arabidopsis* messenger RNAs transported to distant tissues. *Nat. Plants* 1, 15025 (2015). [PubMed: 27247031]
- Kim G, LeBlanc ML, Wafula EK, dePamphilis CW & Westwood JH Genomic-scale exchange of mRNA between a parasitic plant and its hosts. *Science* 345, 808–811 (2014). [PubMed: 25124438]
- Yang Y et al. Messenger RNA exchange between scions and rootstocks in grafted grapevines. *BMC Plant Biol.* 15, 251 (2015). [PubMed: 26480945]
- Brunkard JO, Runkel AM & Zambryski PC The cytosol must flow: intercellular transport through plasmodesmata. *Curr. Opin. Cell Biol* 35, 13–20 (2015). [PubMed: 25847870]
- Cantrill LC, Overall RL & Goodwin PB Cell-to-cell communication via plant endomembranes. *Cell Biol. Int* 23, 653–661 (1999). [PubMed: 10736188]
- Barton DA et al. Cell-to-cell transport via the lumen of the endoplasmic reticulum. *Plant J.* 66, 806–817 (2011). [PubMed: 21332847]
- Grabski S, De Feijter AW & Schindler M Endoplasmic reticulum forms a dynamic continuum for lipid diffusion between contiguous soybean root cells. *Plant Cell* 5, 25–38 [PubMed: 12271014]
- Jackson D Plasmodesmata spread their influence. *F1000 Prime Rep.* 7, 25 (2015).
- Lee JY Plasmodesmata: a signaling hub at the cellular boundary. *Curr. Opin. Plant Biol* 27, 133–140 (2015). [PubMed: 26247123]
- Sager R & Lee JY Plasmodesmata in integrated cell signalling: insights from development and environmental signals and stresses. *J. Exp. Bot* 65, 6337–6358 (2014). [PubMed: 25262225]

18. Oparka KJ et al. Simple, but not branched, plasmodesmata allow the nonspecific trafficking of proteins in developing tobacco leaves. *Cell* 97, 743–754 (1999). [PubMed: 10380926]
19. Crawford KM & Zambryski PC Non-targeted and targeted protein movement through plasmodesmata in leaves in different developmental and physiological states. *Plant Physiol.* 125, 1802–1812 (2001). [PubMed: 11299360]
20. Burch-Smith TM, Stonebloom S, Xu M & Zambryski PC Plasmodesmata during development: re-examination of the importance of primary, secondary, and branched plasmodesmata structure versus function. *Protoplasma* 248, 61–74 (2011). [PubMed: 21174132]
21. Haywood V, Kragler F & Lucas WJ Plasmodesmata: pathways for protein and ribonucleoprotein signaling. *Plant Cell* 14, S303–325 (2002). [PubMed: 12045285]
22. Levy A, Erlanger M, Rosenthal M & Epel BL A plasmodesmata-associated beta-1,3-glucanase in *Arabidopsis*. *Plant J* 49, 669–682 (2007). [PubMed: 17270015]
23. Kuhn C, Franceschi VR, Schulz A, Lemoine R & Frommer WB Macromolecular trafficking indicated by localization and turnover of sucrose transporters in enucleate sieve elements. *Science* 275, 1298–1300 (1997). [PubMed: 9036853]
24. Lucas WJ et al. Selective trafficking of KNOTTED1 homeodomain protein and its mRNA through plasmodesmata. *Science* 270, 1980–1983 (1995). [PubMed: 8533088]
25. Notaguchi M & Okamoto S Dynamics of long-distance signaling via plant vascular tissues. *Front. Plant Sci* 6, 161 (2015). [PubMed: 25852714]
26. van Bel AJ Interaction between sieve element and companion cell and the consequences for photoassimilate distribution. Two structural hardware frames with associated physiological software packages in dicotyledons? *J. Exp. Botany* 47, 1129–1140 (1996). [PubMed: 21245242]
27. Heo JO, Roszak P, Furuta KM & Helariutta Y Phloem development: current knowledge and future perspectives. *Am. J. Bot* 101, 1393–1402 (2014). [PubMed: 25253700]
28. Ostendorp A et al. Functional analysis of *Brassica napus* phloem protein and ribonucleoprotein complexes. *New Phytol.* 214, 1188–1197 (2017). [PubMed: 28052459]
29. Giavalisco P, Kapitza K, Kolasa A, Buhtz A & Kehr J Towards the proteome of *Brassica napus* phloem sap. *Proteomics* 6, 896–909 (2006). [PubMed: 16400686]
30. Doering-Saad C, Newbury HJ, Bale JS & Pritchard J Use of aphid stylectomy and RT-PCR for the detection of transporter mRNAs in sieve elements. *J. Exp. Bot* 53, 631–637 (2002). [PubMed: 11886882]
31. Sasaki T, Chino M, Hayashi H & Fujiwara T Detection of several mRNA species in rice phloem sap. *Plant Cell Physiol.* 39, 895–897 (1998). [PubMed: 9787465]
32. Guo S et al. The draft genome of watermelon (*Citrullus lanatus*) and resequencing of 20 diverse accessions. *Nat. Genet* 45, 51–58 (2013). [PubMed: 23179023]
33. Rabouille C Pathways of unconventional protein secretion. *Trends Cell Biol.* 27, 230–240 (2017). [PubMed: 27989656]
34. Colombo M, Raposo G & Thery C Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles. *Annu. Rev. Cell. Dev. Biol* 30, 255–289 (2014). [PubMed: 25288114]
35. Regente M et al. Vesicular fractions of sunflower apoplastic fluids are associated with potential exosome marker proteins. *FEBS Lett.* 583, 3363–3366 (2009). [PubMed: 19796642]
36. Prado N et al. Nanovesicles are secreted during pollen germination and pollen tube growth: a possible role in fertilization. *Mol. Plant* 7, 573–577 (2014). [PubMed: 24177685]
37. Rutter BD & Innes RW Extracellular vesicles isolated from the leaf apoplast carry stress-response proteins. *Plant Physiol.* 173, 728–741 (2017). [PubMed: 27837092]
38. An Q, Huckelhoven R, Kogel KH & van Bel AJ Multivesicular bodies participate in a cell wall-associated defence response in barley leaves attacked by the pathogenic powdery mildew fungus. *Cell. Microbiol* 8, 1009–1019 (2006). [PubMed: 16681841]
39. Takemoto D, Jones DA & Hardham AR GFP-tagging of cell components reveals the dynamics of subcellular re-organization in response to infection of *Arabidopsis* by oomycete pathogens. *Plant J.* 33, 775–792 (2003). [PubMed: 12609049]

40. Lu YJ et al. Patterns of plant subcellular responses to successful oomycete infections reveal differences in host cell reprogramming and endocytic trafficking. *Cell. Microbiol* 14, 682–697 (2012). [PubMed: 22233428]
41. Cai Q et al. Plants send small RNAs in extracellular vesicles to fungal pathogen to silence virulence genes. *Science* 8, 1126–1129 (2018).
42. Deeken R et al. Identification of *Arabidopsis thaliana* phloem RNAs provides a search criterion for phloem-based transcripts hidden in complex datasets of microarray experiments. *Plant J.* 55, 746–759 (2008). [PubMed: 18485061]
43. Gaupels F et al. Adaptation of aphid stylectomy for analyses of proteins and mRNAs in barley phloem sap. *J. Exp. Bot* 59, 3297–3306 (2008). [PubMed: 18632729]
44. Ruiz-Medrano R, Xoconostle-Cazares B & Lucas WJ Phloem long-distance transport of *CmNACP* mRNA: implications for supracellular regulation in plants. *Development* 126, 4405–4419 (1999). [PubMed: 10498677]
45. Omid A, Keilin T, Glass A, Leshkowitz D & Wolf S Characterization of phloem-sap transcription profile in melon plants. *J. Exp. Bot* 58, 3645–3656 (2007). [PubMed: 17928373]
46. Doering-Saad C, Newbury HJ, Couldridge CE, Bale JS & Pritchard J A phloem-enriched cDNA library from *Ricinus*: insights into phloem function. *J. Exp. Bot* 57, 3183–3193 (2006). [PubMed: 16936221]
47. Rodriguez-Medina C, Atkins CA, Mann AJ, Jordan ME & Smith PM Macromolecular composition of phloem exudate from white lupin (*Lupinus albus* L.). *BMC Plant Biol.* 11, 36 (2011). [PubMed: 21342527]
48. Notaguchi M, Wolf S & Lucas WJ Phloem-mobile *Aux/IAA* transcripts target to the root tip and modify root architecture. *J. Integr. Plant Biol* 54, 760–772 (2012). [PubMed: 22925478]
49. Xoconostle-Cazares B et al. Plant paralog to viral movement protein that potentiates transport of mRNA into the phloem. *Science* 283, 94–98 (1999). [PubMed: 9872750]
50. Banerjee AK et al. Dynamics of a mobile RNA of potato involved in a long-distance signaling pathway. *Plant Cell* 18, 3443–3457 (2006). [PubMed: 17189340]
51. Mahajan A, Bhogale S, Kang IH, Hannapel DJ & Banerjee AK The mRNA of a Knotted1-like transcription factor of potato is phloem mobile. *Plant Mol. Biol* 79, 595–608 (2012). [PubMed: 22638904]
52. Kanehira A et al. Apple phloem cells contain some mRNAs transported over long distances. *Tree Genet. Genom* 6, 635–642 (2010).
53. Roney JK, Khatibi PA & Westwood JH Cross-species translocation of mRNA from host plants into the parasitic plant dodder. *Plant Physiol.* 143, 1037–1043 (2007). [PubMed: 17189329]
54. David-Schwartz R, Runo S, Townsley B, Machuka J & Sinha N Long-distance transport of mRNA via parenchyma cells and phloem across the host-parasite junction in *Cuscuta*. *New Phytol* 179, 1133–1141 (2008). [PubMed: 18631294]
55. Goldschmidt EE Plant grafting: new mechanisms, evolutionary implications. *Front. Plant Sci* 5, 727 (2014). [PubMed: 25566298]
56. Lu KJ, Huang NC, Liu YS, Lu CA & Yu TS Long-distance movement of *Arabidopsis FLOWERING LOCUS TRNA* participates in systemic floral regulation. *RNA Biol.* 9, 653–662 (2012). [PubMed: 22614833]
57. Zhang H et al. Expression of tomato prosystemin gene in *Arabidopsis* reveals systemic translocation of its mRNA and confers necrotrophic fungal resistance. *New Phytol.* 217, 799–812 (2018). [PubMed: 29105094]
58. Notaguchi M, Higashiyama T & Suzuki T Identification of mRNAs that move over long distances using an RNA-Seq analysis of *Arabidopsis/Nicotiana benthamiana* heterografts. *Plant Cell Physiol.* 56, 311–321 (2015). [PubMed: 25527829]
59. LeBlanc M, Kim G, Patel B, Stromberg V & Westwood J Quantification of tomato and *Arabidopsis* mobile RNAs trafficking into the parasitic plant *Cuscuta pentagona*. *New Phytol.* 200, 1225–1233 (2013). [PubMed: 23914903]
60. Bologna NG & Voinnet O The diversity, biogenesis, and activities of endogenous silencing small RNAs in *Arabidopsis*. *Annu. Rev. Plant Biol* 65, 473–503 (2014). [PubMed: 24579988]

61. Rogers K & Chen X Biogenesis, turnover, and mode of action of plant microRNAs. *Plant Cell* 25, 2383–2399 (2013). [PubMed: 23881412]
62. Molnar A, Melnyk C & Baulcombe DC Silencing signals in plants: a long journey for small RNAs. *Genome Biol.* 12, 215 (2011). [PubMed: 21235831]
63. Kim YJ, Maizel A & Chen X Traffic into silence: endomembranes and post-transcriptional RNA silencing. *EMBO J.* 33, 968–980 (2014). [PubMed: 24668229]
64. Palauqui JC, Elmayan T, Pollien JM & Vaucheret H Systemic acquired silencing: transgene-specific post-transcriptional silencing is transmitted by grafting from silenced stocks to non-silenced scions. *EMBO J.* 16, 4738–4745 (1997). [PubMed: 9303318]
65. Voinnet O & Baulcombe DC Systemic signalling in gene silencing. *Nature* 389, 553 (1997). [PubMed: 9335491]
66. Voinnet O, Vain P, Angell S & Baulcombe DC Systemic spread of sequence-specific transgene RNA degradation in plants is initiated by localized introduction of ectopic promoterless DNA. *Cell* 95, 177–187 (1998). [PubMed: 9790525]
67. Palauqui JC & Balzergue S Activation of systemic acquired silencing by localised introduction of DNA. *Curr. Biol* 9, 59–66 (1999). [PubMed: 10021363]
68. Melnyk CW, Molnar A & Baulcombe DC Intercellular and systemic movement of RNA silencing signals. *EMBO J.* 30, 3553–3563 (2011). [PubMed: 21878996]
69. Dunoyer P, Himber C, Ruiz-Ferrer V, Alioua A & Voinnet O Intra- and intercellular RNA interference in *Arabidopsis thaliana* requires components of the microRNA and heterochromatic silencing pathways. *Nat. Genet* 39, 848–856 (2007). [PubMed: 17558406]
70. Smith LM et al. An SNF2 protein associated with nuclear RNA silencing and the spread of a silencing signal between cells in *Arabidopsis*. *Plant Cell* 19, 1507–1521 (2007). [PubMed: 17526749]
71. Yoo BC et al. A systemic small RNA signaling system in plants. *Plant Cell* 16, 1979–2000 (2004). [PubMed: 15258266]
72. Buhtz A, Springer F, Chappell L, Baulcombe DC & Kehr J Identification and characterization of small RNAs from the phloem of *Brassica napus*. *Plant J.* 53, 739–749 (2008). [PubMed: 18005229]
73. Buhtz A, Pieritz J, Springer F & Kehr J Phloem small RNAs, nutrient stress responses, and systemic mobility. *BMC Plant Biol.* 10, 64 (2010). [PubMed: 20388194]
74. Varkonyi-Gasic E, Gould N, Sandanayaka M, Sutherland P & MacDiarmid RM Characterisation of microRNAs from apple (*Malus domestica* ‘Royal Gala’) vascular tissue and phloem sap. *BMC Plant Biol.* 10, 159 (2010). [PubMed: 20682080]
75. Slotkin RK et al. Epigenetic reprogramming and small RNA silencing of transposable elements in pollen. *Cell* 136, 461–472 (2009). [PubMed: 19203581]
76. Martinez G, Panda K, Kohler C & Slotkin RK Silencing in sperm cells is directed by RNA movement from the surrounding nurse cell. *Nat. Plants* 2, 16030 (2016). [PubMed: 27249563]
77. Molnar A et al. Small silencing RNAs in plants are mobile and direct epigenetic modification in recipient cells. *Science* 328, 872–875 (2010). [PubMed: 20413459]
78. Melnyk CW, Molnar A, Bassett A & Baulcombe DC Mobile 24 nt small RNAs direct transcriptional gene silencing in the root meristems of *Arabidopsis thaliana*. *Curr. Biol* 21, 1678–1683 (2011). [PubMed: 21962713]
79. Zhang W et al. Graft-transmissible movement of inverted-repeat-induced siRNA signals into flowers. *Plant J.* 80, 106–121 (2014). [PubMed: 25039964]
80. Parizotto EA, Dunoyer P, Rahm N, Himber C & Voinnet O *In vivo* investigation of the transcription, processing, endonucleolytic activity, and functional relevance of the spatial distribution of a plant miRNA. *Genes Dev.* 18, 2237–2242 (2004). [PubMed: 15371337]
81. Alvarez JP et al. Endogenous and synthetic microRNAs stimulate simultaneous, efficient, and localized regulation of multiple targets in diverse species. *Plant Cell* 18, 1134–1151 (2006). [PubMed: 16603651]
82. Knauer S et al. A protodermal miR394 signal defines a region of stem cell competence in the *Arabidopsis* shoot meristem. *Dev. Cell* 24, 125–132 (2013). [PubMed: 23333352]

83. Pant BD, Buhtz A, Kehr J & Scheible WR MicroRNA399 is a long-distance signal for the regulation of plant phosphate homeostasis. *Plant J.* 53, 731–738 (2008). [PubMed: 17988220]
84. Lin SI et al. Regulatory network of microRNA399 and PHO2 by systemic signalling. *Plant Physiol.* 147, 732–746 (2008). [PubMed: 18390805]
85. Huen AK, Rodriguez-Medina C, Ho AY, Atkins CA & Smith PM Long-distance movement of phosphate starvation-responsive microRNAs in *Arabidopsis*. *Plant Biol* 19, 643–649 (2017). [PubMed: 28322489]
86. Tsikou D et al. Systemic control of legume susceptibility to rhizobial infection by a mobile microRNA. *Science* <http://doi.org/ct72>. (2018)
87. Shahid S et al. MicroRNAs from the parasitic plant *Cuscuta campestris* target host messenger RNAs. *Nature* 553, 82–85 (2018). [PubMed: 29300014]
88. Zhang T et al. Cotton plants export microRNAs to inhibit virulence gene expression in a fungal pathogen. *Nat. Plants* 2, 16153 (2016). [PubMed: 27668926]
89. Kragler F RNA in the phloem: A crisis or a return on investment? *Plant Sci.* 178, 99–104 (2010).
90. Zhang W et al. tRNA-related sequences trigger systemic mRNA transport in plants. *Plant Cell* 28, 1237–1249 (2016). [PubMed: 27268430]
91. Calderwood A, Kopriva S & Morris RJ Transcript abundance explains mRNA mobility data in *Arabidopsis thaliana*. *Plant Cell* 28, 610–615 (2016). [PubMed: 26952566]
92. Paultre DS, Gustin MP, Molnar A & Oparka KJ Lost in transit: long-distance trafficking and phloem unloading of protein signals in *Arabidopsis* homografts. *Plant Cell* 28, 2016–2025 (2016). [PubMed: 27600534]
93. Westwood JH RNA transport: Delivering the message. *Nat. Plants* 1, 15038 (2015). [PubMed: 27247038]
94. Banerjee AK, Lin T & Hannapel DJ Untranslated regions of a mobile transcript mediate RNA metabolism. *Plant Physiol.* 151, 1831–1843 (2009). [PubMed: 19783647]
95. Carella P, Wilson DC, Kempthorne CJ & Cameron RK Vascular sap proteomics: Providing insight into long-distance signaling during stress. *Front. Plant Sci* 7, 651 (2016). [PubMed: 27242852]
96. Gomez G, Torres H & Pallas V Identification of translocatable RNA-binding phloem proteins from melon, potential components of the long-distance RNA transport system. *Plant J.* 41, 107–116 (2005). [PubMed: 15610353]
97. Ham BK et al. A polypyrimidine tract binding protein, pumpkin RBP50, forms the basis of a phloem-mobile ribonucleoprotein complex. *Plant Cell* 21, 197–215 (2009). [PubMed: 19122103]
98. Cho SK et al. Polypyrimidine tract-binding proteins of potato mediate tuberization through an interaction with *StBEL5* RNA. *J. Exp. Bot* 66, 6835–6847 (2015) [PubMed: 26283046]
99. Ham BK, Li G, Jia W, Leary JA & Lucas WJ Systemic delivery of siRNA in pumpkin by a plant PHLOEM SMALL RNA-BINDING PROTEIN 1-ribonucleoprotein complex. *Plant J.* 80, 683–694 (2014). [PubMed: 25227635]
100. Seville I, Yadav SR & Helariutta Y Plasmodesmata: channels for intercellular signaling during plant growth and development. *Methods Mol. Biol* 1217, 3–24 (2015). [PubMed: 25287193]
101. Balachandran S, Xiang Y, Schobert C, Thompson GA & Lucas WJ Phloem sap proteins from *Cucurbita maxima* and *Ricinus communis* have the capacity to traffic cell to cell through plasmodesmata. *Proc. Natl Acad. Sci. USA* 94, 14150–14155 (1997). [PubMed: 9391168]
102. Aoki K, Kragler F, Xoconostle-Cazares B & Lucas WJ A subclass of plant heat shock cognate 70 chaperones carries a motif that facilitates trafficking through plasmodesmata. *Proc. Natl Acad. Sci. USA* 99, 16342–16347 (2002). [PubMed: 12456884]
103. Vaten A et al. Callose biosynthesis regulates symplastic trafficking during root development. *Dev. Cell* 21, 1144–1155 (2011). [PubMed: 22172675]
104. Grison MS, Fernandez-Calvino L, Mongrand S & Bayer EM Isolation of plasmodesmata from *Arabidopsis* suspension culture cells. *Methods Mol. Biol* 1217, 83–93 (2015). [PubMed: 25287197]
105. de Felippes FF, Ott F & Weigel D Comparative analysis of non-autonomous effects of tasiRNAs and miRNAs in *Arabidopsis thaliana*. *Nucleic Acids Res.* 39, 2880–2889 (2011). [PubMed: 21134910]

106. Skopelitis DS et al. Gating of miRNA movement at defined cell-cell interfaces governs their impact as positional signals. *Nat. Commun* 9, 3107 (2018). [PubMed: 30082703]
107. Haywood V, Yu TS, Huang NC & Lucas WJ Phloem long-distance trafficking of *GIBBERELLIC ACID-INSENSITIVE* RNA regulates leaf development. *Plant J.* 42, 49–68 (2005). [PubMed: 15773853]
108. Skopelitis DS, Benkovics AH, Husbands AY & Timmermans MCP Boundary formation through a direct threshold-based readout of mobile small RNA gradients. *Dev. Cell* 43, 265–273 (2017). [PubMed: 29107557]
109. Ghatge TH, Sharma P, Kondhare KR, Hannapel DJ & Banerjee AK The mobile RNAs, *StBEL11* and *StBEL29*, suppress growth of tubers in potato. *Plant Mol. Biol* 93, 563–578 (2017). [PubMed: 28084609]
110. Dunoyer P, Melnyk C, Molnar A & Slotkin RK Plant mobile small RNAs. *Cold Spring Harb. Perspect. Biol* 5, a017897 (2013). [PubMed: 23818501]
111. Sarkies P & Miska EA Small RNAs break out: the molecular cell biology of mobile small RNAs. *Nat. Rev. Mol. Cell Biol* 15, 525–535 (2014). [PubMed: 25053358]
112. Weiberg A & Jin H Small RNAs—the secret agents in the plant-pathogen interactions. *Curr. Opin. Plant Biol* 26, 87–94 (2015). [PubMed: 26123395]
113. Melnyk CW & Meyerowitz EM Plant grafting. *Curr. Biol* 25, 183–188 (2015).
114. Yoder JI, Gunathilake P, Wu B, Tomilova N & Tomilov AA Engineering host resistance against parasitic weeds with RNA interference. *Pest Manag. Sci* 65, 460–466 (2009). [PubMed: 19235710]
115. Guan D et al. PlaMoM: a comprehensive database compiles plant mobile macromolecules. *Nucleic Acids Res.* 45, 1021–1028 (2017).
116. Luo KR, Huang NC & Yu TS Selective targeting of mobile mRNAs to plasmodesmata for cell-to-cell movement. *Plant Physiol.* <http://doi.org/gdrsfz>. (2018).




Fig. 1 | Routes for RNA trafficking between plant cells.

a, PD as micro-channels for cell-to-cell movement of mobile molecules. PD are membrane-lined channels that traverse the cell wall and connect neighbouring cells. Cell wall, plasma membrane, endoplasmic reticulum (ER), desmotubule and callose are indicated. Red circles represent soluble molecules capable of moving through the desmotubule or cytoplasmic sleeve of PD. **b**, Phloem as a conduit for long-distance movement of molecules. Phloem is composed of stacked enucleated sieve elements assisted by companion cells. Mature sieve elements are connected to adjacent companion cells by highly modified and funnel-like plasmodesmata pore units (PPUs). Mobile molecules (red circles) are primarily transported from source to sink tissues over long distances through phloem. Arrows indicate the direction of movement. Gaps in the line indicate multiple, stacked sieve elements (one cell in the diagram) mediating long-distance transport. **c**, Putative vesicle-mediated RNA trafficking in plants. Vesicles containing RNAs are taken into MVBs, which subsequently fuse with the plasma membrane and release their intraluminal vesicles into the extracellular space as exosomes. These exosomes fuse with the plasma membrane of the recipient cell

through endocytosis and unload the cargo RNAs. Alternatively, vesicles may transport RNAs to adjacent cells, either through exocytosis/endocytosis or through the PD channels between cells. Note that these are purely hypothetical events, as indicated by dashed lines in the diagram.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 2 |. Schematic drawing of non-cell-autonomous RNA silencing in plants.

SiRNAs (red) and miRNAs (blue) can act as mobile signals and move from cell to cell or over long distances to mediate non-cell-autonomous RNA silencing in plants. In the destination cells, siRNAs mediate RNA-directed DNA methylation or guide the cleavage of target RNAs, whereas miRNAs guide the cleavage or translational repression of target mRNAs. Filled orange circles indicate methyl groups on DNA or histones.




Fig. 3 |. Dynamic network of intercellular communication.

A hypothetical model of how RNA trafficking influences plant development and physiological processes. Internal or external stimuli may trigger the movement of plant RNAs, including mRNAs, siRNAs, miRNAs, rRNAs and tRNAs, from their synthesis sites (incipient cells) to distant tissues (recipient cells). These mobile RNAs may act non-cell-autonomously. In the recipient cells, mRNAs, rRNAs and tRNAs may participate in translation, while miRNAs and siRNAs may mediate the silencing and regulation of genes. These mobile RNAs may therefore act as signalling molecules for intercellular information exchange to regulate plant development, nutrient allocation, stress responses and many other physiological processes in plants.

Table | 1

Representative examples of mobile RNAs in plants

RNA	Function	Plant Species	References
mRNA			
<i>SUC1</i>	Sucrose transport	Potato	23
<i>FT1</i>	Flowering induction	<i>Arabidopsis</i>	56
<i>Aux/IAA</i>	Root development	<i>Arabidopsis</i>	48
<i>PP16</i>	RNA transport	Pumpkin	49
<i>NACP</i>	Meristem maintenance	Pumpkin	44
<i>BEL5</i>	Tuber formation	Potato	50
<i>POTH1</i>	Leaf development	Potato	51
<i>SLR/IAA14</i>	Lateral root formation	Apple	52
<i>PPF-T6</i>	Leaf development	Tomato	2
<i>PS</i>	Pathogen resistance	Tomato	57
<i>GAI</i>	Leaf development	Tomato	53,54
<i>PPF</i>	Leaf development	Tomato	54
miRNA			
miR165/166	Root development	<i>Arabidopsis</i>	5
miR390	Leaf polarity	<i>Arabidopsis</i>	3
miR394	Shoot meristem formation	<i>Arabidopsis</i>	82
miR395	Sulfate homeostasis	<i>Arabidopsis</i>	73
miR399	Phosphate homeostasis	<i>Arabidopsis</i>	73, 83,84
siRNA			
ta-siRNA	Leaf polarity	<i>Arabidopsis</i>	3
hc-siRNA	DNA methylation	<i>Arabidopsis</i> , <i>N. tabacum</i> , <i>N. benthamiana</i>	6,75–79
rRNA			
5S rRNA	Translation	<i>B. napus</i> , pumpkin	4,72
5.8S rRNA	Translation	<i>B. napus</i> , pumpkin	4,72
18S rRNA	Translation	<i>B. napus</i> , pumpkin	4,72
25S rRNA	Translation	<i>B. napus</i> , pumpkin	4,72
tRNA ^a	Translation	Pumpkin	4

* Not all tRNA species are detected in the phloem sap, and some tRNA are present in truncated forms or as tRNA halves.